

An Introduction to Chaos in Topological Dynamical Systems

Fraser Robert Love

School of Mathematics and Statistics
University of St Andrews

March 27, 2023

Table of Contents

Topological Dynamics

Introduction to Chaos

Topological Characteristics of Chaos

Devaney Chaos

Concluding Remarks

Bibliography

Topological Dynamics

Definition (Topological Dynamical System).

- ▶ Let X be a non-empty compact metric space. A *topological dynamical system*, denoted (X, f) , is given by a continuous map $f : X \rightarrow X$.
- ▶ The system starts at an initial point $x \in X$ and evolves through successive iterations of the map f .
- ▶ After $k \in \mathbb{N}$ iterations of f , the system can be described by $f^k := f \circ f \circ \cdots \circ f$, where x is mapped to the point $f^k(x)$.

Topological Dynamics

Definition (Orbit).

Let (X, f) be a topological dynamical system. The *orbit* of $x \in X$ under f is the set $\mathcal{O}_f(x) = \{f^n(x) : n \geq 0\} = \{x, f(x), f^2(x), \dots\}$ of iterates of x under the map f .

Definition (Periodic Point, Cycle).

- ▶ Let (X, f) be a topological dynamical system. A point $x \in X$ is *periodic* if $f^n(x) = x$ for some $n \in \mathbb{N}$.
- ▶ The *period* of a point x is the least positive integer k such that $f^k(x) = x$. If x has a period of k we say that x is a *period- k* point.
- ▶ The orbit $\mathcal{O}_f(x) = \{x, f(x), \dots, f^{k-1}(x)\}$ of a periodic point is a finite set of unique points, called a *periodic orbit* of period k or simply a *k -cycle*.

Topological Dynamics

Example (Logistic Map).

Define $F_\mu : [0, 1] \rightarrow [0, 1]$ to be the *logistic map*, where $F_\mu(x) = \mu x(1 - x)$ and $\mu > 0$.




Figure: Logistic map F_μ with $\mu = 3$.

Topological Dynamics

Example (Doubling Map).

Define $\mathcal{D} : S^1 \rightarrow S^1$ to be the *doubling map* on $S^1 = \{z \in \mathbb{C} : |z| = 1\}$, where $\mathcal{D}(z) = z^2$, or equivalently $\mathcal{D}(e^{i\theta}) = e^{2i\theta}$ for some $\theta \in \mathbb{R}$.




Figure: First ten iterations of the doubling map \mathcal{D} .

Topological Dynamics

Definition (Sequence Space).

- ▶ Let $\Sigma_2 = \{(s_1, s_2, \dots) : s_i \in \{0, 1\}\}$ be the set of sequences of ones and zeros.
- ▶ Define (Σ_2, d) to be the *sequence space* where $d(\underline{s}, \underline{t}) = \sum_{i=1}^{\infty} |s_i - t_i| 2^{-i}$ is a metric for $\underline{s} = (s)_{i=1}^{\infty}$, $\underline{t} = (t)_{i=1}^{\infty} \in \Sigma_2$.
- ▶ The sequence space (Σ_2, d) is compact.

Definition (Shift Map).

- ▶ The *shift map* $\sigma : \Sigma_2 \rightarrow \Sigma_2$ given by $\sigma((s)_{i=1}^{\infty}) = (s)_{i=2}^{\infty}$.
- ▶ This map is continuous, hence (Σ_2, σ) is a topological dynamical system.

Topological Dynamics

Proposition.

The *shift map* $\sigma : \Sigma_2 \rightarrow \Sigma_2$ given by $\sigma((s)_{i=1}^\infty) = (s)_{i=2}^\infty$ is continuous.

Proof.

Let $\varepsilon > 0$ and choose $\underline{s} = (s_i)_{i=1}^\infty$, $\underline{t} = (t_i)_{i=1}^\infty \in \Sigma_2$ such that $d(\underline{s}, \underline{t}) = \sum_{i=1}^\infty |s_i - t_i| 2^{-i} < \delta$. Choose n such that $2^{-n} \leq \varepsilon$ and let $\delta = 2^{-(n+1)}$. Hence \underline{s} and \underline{t} agree on the first $n+1$ symbols and $\sigma(\underline{s})$ and $\sigma(\underline{t})$ agree on the first n symbols. Then $d(\sigma(\underline{s}), \sigma(\underline{t})) = d((s)_{i=n+1}^\infty, (t)_{i=n+1}^\infty) = \sum_{i=n+1}^\infty |s_i - t_i| 2^{-i} = 2^{-n} \leq \varepsilon$. □

Hence (Σ_2, σ) defines a topological dynamical system.

Introduction to Chaos

- ▶ Many different definitions of chaos exist for topological dynamical systems.
- ▶ Devaney chaos, Li-Yorke chaos, Topological chaos, etc.
- ▶ These definitions rely on many different topological characteristics to define chaos in a natural way.
- ▶ We will focus on Devaney chaos and the characteristics of topological transitivity / existence of a dense orbit, sensitive dependence on initial conditions and dense periodic points.

Topological Characteristics of Chaos

Definition (Topological Transitivity).

Let (X, f) be a topological dynamical system. The map f is *topologically transitive* if for every pair of non-empty open sets $U, V \subseteq X$ there exists $k > 0$ such that $f^k(U) \cap V \neq \emptyset$.

Proposition.

Let (X, f) be a topological dynamical system and suppose X has no isolated points. The map f is topologically transitive if and only if there exists some $x \in X$ such that $\mathcal{O}(x)$ is dense in X . [2]

Topological Characteristics of Chaos

Proposition.

The shift map (Σ_2, σ) is topologically transitive. [1]

Proof.

Let $\underline{t} = (t_i)_{i=0}^{\infty} \in \Sigma_2$ be arbitrary and $\varepsilon > 0$. Consider the sequence $\underline{s} = (0, 1, \{0, 0\}, \{0, 1\}, \{1, 0\}, \{1, 1\}, \{0, 0, 0\}, \{0, 0, 1\}, \dots)$ of 0s and 1s sorted in len-lex order. By construction we can perform some $k \in \mathbb{N}$ iterations of σ such that the first n symbols of $\sigma^k(\underline{s})$ and \underline{t} agree. Choose $N \geq \log_2 \frac{1}{\varepsilon}$. Therefore for $n \geq N - 1$ we have that, $d(\sigma^k(\underline{s}), \underline{t}) = \sum_{i=n}^{\infty} |\sigma^k(\underline{s})_i - t_i| 2^{-i} \leq 2^{-(n+1)} < 2^{-N} \leq \varepsilon$. Hence $\mathcal{O}(\underline{s})$ is dense in Σ_2 and so (Σ_2, σ) is topologically transitive. \square

Topological Characteristics of Chaos

Definition (Sensitive Dependence on Initial Conditions).

Let (X, f) be a topological dynamical system and $\varepsilon > 0$. A point $x \in X$ is ε -unstable if, for every neighbourhood U of x , there exists a point $y \in U$ and $k \geq 0$ such that $d(f^k(x), f^k(y)) \geq \varepsilon$. The map f has *sensitive dependence on initial conditions* if for all points $x \in X$, x is ε -unstable.

Topological Characteristics of Chaos

Example.

The shift map (Σ_2, σ) has sensitive dependence on initial conditions.

Proof.

Take $\delta = 1$. Let $\underline{s} = (s)_{i=1}^{\infty} \in \Sigma_2$ and let $\varepsilon > 0$. Choose n such that $2^{-n} \leq \varepsilon$. Pick $\underline{t} = (t)_{i=1}^{\infty} \in \Sigma_2$ such that $d(\underline{s}, \underline{t}) \leq 2^{-n} \leq \varepsilon$. Hence \underline{s} and \underline{t} agree on the first $n + 1$ symbols. Now there exists a $k > n + 1$ such that $s_k \neq t_k$. The first term of $\sigma^k(\underline{s})$ is s_k and the first term of $\sigma^k(\underline{t})$ is t_k . Therefore

$$d(\sigma^k(\underline{s}), \sigma^k(\underline{t})) = \sum_{i=0}^{\infty} |s_{i+k} - t_{i+k}| 2^{-i} \geq |s_k - t_k| 2^0 = 1 = \delta. \quad \square$$

Topological Characteristics of Chaos

Proposition.

The periodic points of the shift map (Σ_2, σ) are dense in Σ_2 . [1]

Proof.

Let $\underline{s} = (s_i)_{i=1}^{\infty}$ be an arbitrary point in Σ_2 and let $\varepsilon > 0$. Pick an n such that $2^{-n} \leq \varepsilon$ and define $\underline{t} = (s_0, \dots, s_n, s_0, \dots, s_n, \dots)$ to be an infinite repeating sequence where $t_i = s_i$ for $1 \leq i \leq n$. Hence \underline{s} and \underline{t} agree on the first n symbols. Therefore $d(\underline{s}, \underline{t}) = \sum_{i=n+1}^{\infty} |s_i - t_i| 2^{-i} \leq \sum_{i=n+1}^{\infty} 2^{-i} = 2^{-n} \leq \varepsilon$. Hence as $n \rightarrow \infty$ we have $\underline{t}_n \rightarrow \underline{s}$. Therefore the periodic points of σ are dense. □

Devaney Chaos

Definition (Devaney Chaos).

A topological dynamical system (X, f) is *chaotic in the sense of Devaney* if it is topologically transitive, has sensitive dependence on initial conditions, and if the periodic points of f are dense in X .

Devaney's definition considers:

- ▶ Unpredictability (sensitive dependence on initial conditions).
- ▶ Repetitiveness (dense periodic points).
- ▶ Indecomposability (topological transitivity).

Devaney Chaos

Proposition.

The shift map (Σ_2, σ) is chaotic in the sense of Devaney.

Proof.

We previously showed that (Σ_2, σ) is topologically transitive (has a dense orbit), has sensitive dependence on initial conditions and has dense periodic points in Σ_2 . Hence (Σ_2, σ) is Devaney chaotic. \square

Concluding Remarks

- ▶ We have defined chaos to be a mixture of unpredictability, repetitiveness and indecomposability.
- ▶ This was achieved using the properties of topological transitivity, sensitive dependence on initial conditions and dense periodic points.
- ▶ We have shown that the shift map (Σ_2, σ) is a Devaney chaotic topological dynamical system.
- ▶ Using topological conjugacy we can use the fact that (Σ_2, σ) is Devaney chaotic to prove that $([0, 1], F_\mu)$ and various other systems exhibit Devaney chaos.

Bibliography

R. L. Devaney.

An introduction to Chaotic Dynamical Systems.

Addison-Wesley Publishing Company Advanced Book Program,
Redwood City, California, second edition, 1989.

S. Silverman.

On maps with dense orbits and the definition of chaos.

Rocky Mountain Journal of Mathematics, 22(1):353–375, 1992.