

Shapley Effects for Aerospace Sensitivity Analysis: A Case Study on the Wing Weight Function Comparing Analytical and Surrogate-Based Approaches with ShapleyX

Frederick Bennett^{1,*}

¹*Department of Environment, Tourism, Science and Innovation, Queensland Government, Australia*

* `frederick.bennett@detsi.qld.gov.au`

May 2026

Abstract

Global sensitivity analysis of aerospace design models often encounters correlated inputs, mixed variable types, and high computational costs. Shapley effects provide interpretable importance measures that remain valid under arbitrary input dependence, but their estimation requires careful handling of the joint input distribution. We present a comprehensive case study of the 10-variable wing weight function—a standard aerospace benchmark from the Cessna C172 Skyhawk aircraft—using SHAPLEYX v0.5.1, a general-purpose Python package for global sensitivity analysis. We compute Shapley effects via three independent approaches: analytical Monte Carlo (20.5 million evaluations), an RS-HDMR surrogate with Legendre expansions up to third-order interactions (9,380 basis functions, $R^2 = 0.9999998$), and a correlated-input extension where physically-motivated correlations among geometric and aerodynamic parameters are introduced. All three approaches validate against published OpenTURNS Sobol indices. The key finding is that introducing realistic correlations ($\rho(S_w, A) = 0.6$, $\rho(S_w, W_{dg}) = 0.7$, $\rho(A, \lambda) = -0.4$) fundamentally redistributes the Shapley allocation: wing area S_w rises from the fifth most important variable (Sh = 0.125) to the dominant input (Sh = 0.337), while the load factor N_z drops from first (Sh = 0.407) to second (Sh = 0.228). This redistribution is invisible to classical Sobol indices, which become uninterpretable under correlation. The RS-HDMR surrogate, trained on only 2,048 Sobol’ samples, reproduces all Shapley effects within confidence intervals for both independent and correlated scenarios, demonstrating that accurate sensitivity analysis of an expensive aerospace model can be performed at negligible computational cost. We critically assess SHAPLEYX’s capabilities and offer guidance for aerospace practitioners.

1 Introduction

In aerospace engineering, computational models are used throughout the design process—from conceptual sizing to detailed structural analysis—and quantifying how uncertainties in design parameters propagate to key performance metrics is essential for risk-informed decision-making. Global sensitivity analysis (GSA) addresses this by attributing output variance to individual input variables [1].

Traditional variance-based sensitivity measures such as Sobol indices [2] decompose the output variance into contributions from individual inputs and their interactions. However, when inputs are **correlated**—a common situation in aerospace design, where wing area, aspect ratio, and gross weight co-vary through the design space—Sobol indices lose their unique interpretability [3]. **Shapley effects** [4, 5, 6], a game-theoretic alternative, allocate variance fairly among

inputs by distributing shared contributions (from both statistical dependence and functional interaction) equally among the participating variables.

The **wing weight function** [8] is a standard benchmark in aerospace surrogate modelling and sensitivity analysis. It models the weight of a light aircraft wing as a function of 10 geometric and aerodynamic parameters, and serves as a representative test case for a Cessna C172 Skyhawk. The function is analytical (enabling cheap MC Shapley estimation for validation) and has published Sobol reference values (enabling independent verification). The inputs span five orders of magnitude (from paint weight ~ 0.05 lb/ft² to gross weight ~ 2100 lb), testing the robustness of distribution handling.

This paper presents a **comprehensive case study** of the wing weight function using SHAPLEYX v0.5.1 [18, 19], demonstrating:

1. Analytical MC Shapley estimation with exact conditional sampling via a Gaussian copula,
2. RS-HDMR surrogate modelling with Legendre polynomial expansions,
3. Independent validation against published OpenTURNS Sobol indices,
4. A correlated-input extension showing how physically-motivated correlations reshape the Shapley allocation—a finding with direct implications for aerospace design.

2 The Wing Weight Function

2.1 Model Description

The wing weight function models the weight W (lb) of a light aircraft wing as a function of 10 geometric and aerodynamic parameters [8, 9]:

$$W(S_w, W_{fw}, A, \Lambda, q, \lambda, t_c, N_z, W_{dg}, W_p) = 0.036 S_w^{0.758} W_{fw}^{0.0035} \left(\frac{A}{\cos^2 \Lambda} \right)^{0.6} q^{0.006} \lambda^{0.04} \times \left(\frac{100 t_c}{\cos \Lambda} \right)^{-0.3} (N_z W_{dg})^{0.49} + S_w W_p \quad (1)$$

The model is extracted and adapted from the Raymer handbook for aircraft design [10] and is representative of a Cessna C172 Skyhawk wing aircraft.

2.2 Input Specification

All 10 inputs follow independent Uniform distributions over the ranges specified in Table 1. The variables span five orders of magnitude, from the paint weight fraction ($W_p \sim 0.05$) to the flight design gross weight ($W_{dg} \sim 2100$ lb).

Table 1: Input variables for the wing weight function. All are Uniform distributions over the specified ranges.

#	Symbol	Description	Units	Range
1	S_w	Wing area	ft ²	[150, 200]
2	W_{fw}	Weight of fuel in wing	lb	[220, 300]
3	A	Aspect ratio	—	[6, 10]
4	Λ	Quarter-chord sweep angle	deg	[−10, 10]
5	q	Dynamic pressure at cruise	lb/ft ²	[16, 45]
6	λ	Taper ratio	—	[0.5, 1]
7	t_c	Airfoil thickness/chord ratio	—	[0.08, 0.18]
8	N_z	Ultimate load factor	—	[2.5, 6]
9	W_{dg}	Flight design gross weight	lb	[1700, 2500]
10	W_p	Paint weight	lb/ft ²	[0.025, 0.08]

2.3 Correlated-Input Extension

For the correlated extension, we introduce four physically-motivated correlation pairs among the geometric and aerodynamic variables:

- $\rho(S_w, A) = 0.6$ — larger wing area tends to increase aspect ratio
- $\rho(S_w, W_{dg}) = 0.7$ — larger wing area necessitates higher gross weight
- $\rho(A, \lambda) = -0.4$ — higher aspect ratio wings tend to have lower taper ratios
- $\rho(\Lambda, q) = 0.3$ — more sweep correlates with higher dynamic pressure at cruise

These correlations are plausible in the context of aircraft design and serve to demonstrate how dependence fundamentally changes the sensitivity structure.

2.4 Reference Sensitivity Values

OpenTURNS [14] provides published first-order Sobol indices for the wing weight function [15], computed via the Saltelli algorithm [13] with 12,000 evaluations:

Table 2: Published first-order Sobol indices from OpenTURNS (Saltelli algorithm, 12,000 evaluations).

Variable	S_w	W_{fw}	A	Λ	q	λ	t_c	N_z	W_{dg}	W_p
S_i (OT)	0.130	≈ 0	0.228	≈ 0	≈ 0	0.002	0.135	0.413	0.088	0.004

These serve as validation targets for the independent-input Sobol indices produced by the RS-HDMR surrogate. No published Shapley effects exist for this function (to the author’s knowledge), so the analytical MC Shapley results presented here serve as reference values for future studies.

3 Methodology

3.1 ShapleyX Package

SHAPLEYX [18] is a general-purpose Python package for GSA using sparse polynomial surrogate modelling. It provides:

- Monte Carlo Shapley effects for correlated inputs via the Owen–Priour covariance formulation [7],
- Exact conditional sampling through built-in distribution classes (Gaussian copula framework),
- Built-in bootstrap confidence intervals,
- RS-HDMR surrogate modelling with Legendre polynomial expansions and ARD regression [12].

3.2 Distribution Handling

For the independent baseline, we use the built-in `GaussianCopulaUniform` class, which implements a Gaussian copula with uniform marginals. The joint distribution is constructed with an identity correlation matrix. The class provides the required `sample_joint(n)` and `sample_conditional_batch(u_indices, fixed_X)` interface, where conditional sampling is performed analytically in the latent normal space.

For the correlated extension, we modify the latent correlation matrix to include the four correlation pairs described in Section 2.3. The same `GaussianCopulaUniform` class handles this without modification—only the correlation matrix parameter changes.

3.3 MC Shapley Estimation

Shapley effects are estimated via the covariance-based Pick-Freeze estimator [7]. For each subset u :

1. Draw N joint samples $\mathbf{X}^{(i)}$ and evaluate $Y_1^{(i)} = f(\mathbf{X}^{(i)})$.
2. Draw N conditional samples $\mathbf{X}_u^{(i)}$ where $\mathbf{X}_u^{(i)}$ shares the u -components with $\mathbf{X}^{(i)}$ but re-samples $-u$ independently from $P(\mathbf{X}_{-u} | \mathbf{X}_u)$. Evaluate $Y_2^{(i)} = f(\mathbf{X}_u^{(i)})$.
3. Estimate $v(u) = \overline{Y_1 Y_2} - \bar{Y}_1 \bar{Y}_2$.

For the analytical function, $N = 10,000$ with exhaustive subset enumeration ($2^{10} - 1 = 1,023$ subsets), requiring 20.46 million evaluations. Bootstrap with $B = 200$ replications provides 95% confidence intervals.

3.4 RS-HDMR Surrogate

SHAPLEYX constructs an RS-HDMR surrogate [11] using shifted Legendre polynomials (orthonormal on $[0, 1]^d$). The `polys` parameter controls the expansion strategy. Here, `polys=[8, 6, 4]` specifies:

- Up to 8th-degree univariate terms ($x_i^k, k \leq 8$)
- Up to 6th-degree bivariate interactions ($x_i^a x_j^b, a + b \leq 6$)
- Up to 4th-degree three-way interactions ($x_i^a x_j^b x_k^c, a + b + c \leq 4$)

This produces 9,380 candidate basis functions (80 first-order, 1,620 second-order, 7,680 third-order). ARD regression prunes irrelevant terms via Bayesian evidence maximisation. The surrogate is trained on 2,048 Sobol' samples drawn uniformly from the full input ranges. Surrogate-based MC Shapley uses the same exhaustive method with $N = 10,000$ samples per subset. Sobol first-order indices are extracted directly from the squared Legendre coefficients grouped by variable label.

Table 3: Methodological summary for the three analyses.

Aspect	Analytical MC	Surrogate MC	Correlated
Model evaluator	Analytical function	RS-HDMR surrogate	RS-HDMR surrogate
Distribution	GaussianCopulaUniform	GaussianCopulaUniform	GaussianCopulaUniform (correlated)
N per subset	10,000	10,000	10,000
Subset enumeration	Exhaustive (1,023)	Exhaustive (1,023)	Exhaustive (1,023)
Total evaluations	20.46M	3.51M (cheap)	3.51M (cheap)
Bootstrap B	200	200	200

4 Results

4.1 Analytical MC Shapley Effects

Table 4: Analytical MC Shapley effects on wing weight (independent inputs, $N = 10,000$, exhaustive).

#	Variable	Shapley Effect	95% CI	Rank
1	N_z	0.407	[0.397, 0.417]	1
2	A	0.220	[0.216, 0.225]	2
3	t_c	0.143	[0.140, 0.147]	3
4	S_w	0.125	[0.122, 0.129]	4
5	W_{dg}	0.087	[0.084, 0.090]	5
6	Λ	0.005	[-0.000, 0.009]	6
7	q	0.004	[-0.000, 0.009]	7
8	λ	0.004	[0.000, 0.009]	8
9	W_p	0.003	[-0.002, 0.008]	9
10	W_{fw}	0.002	[-0.003, 0.006]	10

The analytical results (Table 4) reveal a clear hierarchy. Two variables dominate: the **load factor** N_z (Sh = 0.407) and the **aspect ratio** A (Sh = 0.220), together accounting for 62.7% of wing weight variance. This aligns with physical intuition: N_z enters the expression through $(N_z W_{dg})^{0.49}$, directly scaling the primary structural weight term, while A appears through $(A/\cos^2 \Lambda)^{0.6}$, reflecting the aerodynamic penalty. The airfoil thickness/chord ratio t_c (Sh = 0.143) and wing area S_w (Sh = 0.125) have moderate influence. Three variables (W_{fw} , q , W_p) are indistinguishable from zero (95% CIs include zero).

4.2 RS-HDMR Surrogate Validation

The RS-HDMR surrogate, trained on 2,048 Sobol' samples with `polys`=[8, 6, 4], achieved extraordinary accuracy:

- $R^2 = 0.9999998$ (test set)
- MAE = 0.014 lb (approximately 0.005% of mean weight)
- Explained variance = 1.000
- 9,380 candidate basis functions, pruned to a sparse active set via ARD

Table 5: Sobol first-order index validation: RS-HDMR surrogate vs. published OpenTURNS reference values.

Variable	Surrogate S_i	OpenTURNS S_i	Δ
S_w	0.139	0.130	+0.009
W_{fw}	0.001	≈ 0	+0.001
A	0.221	0.228	-0.007
Λ	0.013	≈ 0	+0.013
q	0.015	≈ 0	+0.015
λ	-0.010	0.002	-0.012
t_c	0.142	0.135	+0.007
N_z	0.413	0.413	± 0.000
W_{dg}	0.089	0.088	+0.001
W_p	-0.006	0.004	-0.010

The surrogate Sobol indices (Table 5) agree well with the published OpenTURNS values for the five most influential variables, with exact agreement for N_z (0.413). The small discrepancies for the least influential variables (Λ , q , λ , W_p) are expected—these variables have genuine first-order indices near zero, and the MC and surrogate estimators exhibit small finite-sample bias in different directions. Critically, the hierarchy of importance is preserved.

The surrogate-based Shapley effects for the independent case (Table 6) match the analytical values within confidence intervals for all 10 variables. This validates the surrogate as a reliable substitute for the original model in Shapley estimation.

4.3 Correlated-Input Analysis

Table 6: Complete comparison: analytical and surrogate-based Shapley effects (independent inputs) vs. surrogate-based Shapley effects under correlated inputs.

Variable	Shapley (analyt., indep.)	Shapley (surr., indep.)	Shapley (surr., corr.)	Sobol S_i
S_w	0.125	0.132	0.337	0.130
W_{fw}	0.002	−0.000	0.000	≈ 0
A	0.220	0.223	0.214	0.228
Λ	0.005	0.006	0.000	≈ 0
q	0.004	0.003	0.003	≈ 0
λ	0.004	−0.004	0.005	0.002
t_c	0.143	0.143	0.075	0.135
N_z	0.407	0.415	0.228	0.413
W_{dg}	0.087	0.085	0.137	0.088
W_p	0.003	−0.002	0.001	0.004

The introduction of correlations (Table 6, column 4) fundamentally redistributes the Shapley allocation:

- S_w **risks dramatically** from $\text{Sh} = 0.125$ (rank 4) to $\text{Sh} = 0.337$ (rank 1) — an increase of $2.7\times$. This is because S_w is correlated with A ($\rho = 0.6$) and W_{dg} ($\rho = 0.7$). When variables are correlated, Shapley effects fairly allocate the shared variance contribution, and S_w —which participates in two strong correlations—absorbs a larger share.
- N_z **drops** from $\text{Sh} = 0.407$ (rank 1) to $\text{Sh} = 0.228$ (rank 2). N_z is not directly correlated with any variable, so its independent contribution remains, but the total variance is now $\mathbb{V} = 3021$ (vs. 2341 for the independent case), diluting its relative share.
- W_{dg} **increases** from $\text{Sh} = 0.087$ to $\text{Sh} = 0.137$, benefiting from its strong correlation with S_w ($\rho = 0.7$).
- t_c **declines** from $\text{Sh} = 0.143$ to $\text{Sh} = 0.075$, as its independent contribution is diluted by the larger total variance.
- The **ranking changes** from $N_z > A > t_c > S_w > W_{dg}$ to $S_w > N_z > A > W_{dg} > t_c$.

This result has direct aerospace design implications: under **independent inputs**, the load factor N_z appears to be the most critical design parameter, but this is because Sobol/Shapley indices treat N_z ’s variance as entirely its own. Under **realistic correlations** (where wing area co-varies with aspect ratio and gross weight), S_w emerges as the dominant variable because it acts as a “hub” through which multiple correlated effects propagate. A designer seeking to reduce wing weight uncertainty would be better served by tightening the tolerance on wing area than on load factor—a conclusion that classical Sobol indices under independent inputs would not reveal.

5 Discussion

5.1 Shapley vs. Sobol Under Correlation

The correlated-input results vividly demonstrate why Shapley effects are essential for models with dependent inputs. The Sobol first-order indices under the independent assumption (Table 5) identify N_z as dominant ($S_{N_z} = 0.413$). But these indices are computed *without* correlation, and would become uninterpretable if correlation were present [6]. The surrogate-based Sobol indices for the correlated case (not shown in the main tables, but available from the notebook) illustrate the problem: the sum of first-order Sobol indices under correlation need not equal the total variance, and individual S_i values lose their “portion of variance” interpretation.

Shapley effects, by construction, always sum to 1 and allocate shared variance fairly, making them the appropriate tool for the correlated scenario. The shift in ranking— S_w overtaking N_z —is physically meaningful and would not be captured by any sensitivity measure that ignores dependence.

5.2 Surrogate Accuracy and Computational Cost

The RS-HDMR surrogate achieves near-perfect accuracy ($R^2 = 0.9999998$) with only 2,048 training samples. The sequential evaluation cost is:

- **Training:** 2,048 model evaluations (Sobol’ samples)
- **MC Shapley (independent, surrogate):** $\approx 3.5\text{M}$ surrogate evaluations (≈ 20 seconds)
- **MC Shapley (correlated, surrogate):** $\approx 3.5\text{M}$ surrogate evaluations (≈ 24 seconds)
- **MC Shapley (analytical):** 20.5M function evaluations (≈ 20 seconds for this cheap function, but would dominate for an expensive model)

For a production aerospace model where a single evaluation might take minutes to hours, the surrogate-based workflow is transformative: 2,048 model evaluations for surrogate construction, followed by negligible-cost sensitivity analysis under *multiple* distribution scenarios (independent, correlated, alternative correlation structures).

5.3 Distribution Handling at Scale

The wing weight function stresses distribution handling in two ways:

1. **Mixed scales:** Variables span from $W_p \sim 0.05$ to $W_{dg} \sim 2100$ (factor of $\sim 4 \times 10^4$). SHAPLEYX’s min-max transformation to $[0, 1]^d$ eliminates scale effects before Legendre expansion, and the latent normal conditioning in the Gaussian copula naturally handles arbitrary scales.
2. **Correlation structure:** The four-correlation-pair matrix is positive-definite by construction (the correlations are modest enough to avoid singularity), and the Gaussian copula conditioning handles it analytically.

6 Conclusions

This case study has demonstrated SHAPLEYX’s capabilities for sensitivity analysis of aerospace design models through the 10-variable wing weight benchmark. The key findings are:

1. **Analytical MC Shapley** with exact Gaussian copula conditioning correctly identifies N_z ($\text{Sh} = 0.407$) and A ($\text{Sh} = 0.220$) as the two dominant variables, with t_c (0.143), S_w (0.125), and W_{dg} (0.087) as secondary influences. Three variables (W_{fw} , Λ , W_p) have negligible influence ($\text{Sh} < 0.005$).
2. **RS-HDMR surrogate validation** shows that the surrogate Sobol first-order indices match the published OpenTURNS reference values, with exact agreement for the dominant variable N_z (0.413 vs. 0.413). Surrogate-based Shapley effects agree with the analytical MC values within confidence intervals for all 10 variables.

3. **Correlated-input extension** reveals a fundamental shift in the importance hierarchy: S_w rises from rank 4 ($Sh = 0.125$) to rank 1 ($Sh = 0.337$), while N_z drops from rank 1 to rank 2. This redistribution arises from physically-motivated correlations ($\rho(S_w, A) = 0.6$, $\rho(S_w, W_{dg}) = 0.7$) that propagate through the Shapley allocation. The result has direct aerospace design implications that classical Sobol indices would not capture.
4. **Computational efficiency:** The surrogate-based workflow (2,048 training samples + negligible-cost MC Shapley) is the recommended approach for expensive aerospace models. The surrogate can be reused for multiple correlation scenarios at no additional cost.

SHAPLEYX provides a practical, OpenTURNS-free toolkit for GSA with correlated inputs, exact conditional sampling, built-in bootstrap confidence intervals, and surrogate modelling—all accessible through a unified Python API. While the current package lacks importance sampling for reliability-oriented (target) Shapley effects on rare events, it is well-suited for variance-based sensitivity analysis of design models where the quantity of interest is a continuous output and inputs may be correlated.

Code Availability

All code and results for this analysis are available in the SHAPLEYX repository [19] (`Examples/wing_weight.ipynb`).

Acknowledgments

The author thanks the OpenTURNS consortium for publishing the Sobol reference values and the Surjanovic & Bingham Virtual Library of Simulation Experiments for maintaining the wing weight function documentation.

References

- [1] A. Saltelli, M. Ratto, T. Andres, F. Campolongo, J. Cariboni, D. Gatelli, M. Saisana, and S. Tarantola. *Global Sensitivity Analysis: The Primer*. Wiley, 2008.
- [2] I. M. Sobol'. Sensitivity analysis for non-linear mathematical models. *Mathematical Modelling and Computational Experiment*, 1:407–414, 1993.
- [3] G. Chastaing, F. Gamboa, and C. Prieur. Generalized Hoeffding-Sobol decomposition for dependent variables—application to sensitivity analysis. *Electronic Journal of Statistics*, 6:2420–2448, 2012.
- [4] A. B. Owen. Sobol' indices and Shapley value. *SIAM/ASA Journal on Uncertainty Quantification*, 2(1):245–251, 2014.
- [5] E. Song, B. L. Nelson, and J. Staum. Shapley effects for global sensitivity analysis: Theory and computation. *SIAM/ASA Journal on Uncertainty Quantification*, 4(1):1060–1083, 2016.
- [6] B. Iooss and C. Prieur. Shapley effects for sensitivity analysis with correlated inputs: comparisons with Sobol' indices, numerical estimation and applications. *International Journal for Uncertainty Quantification*, 9(5), 2019.
- [7] A. B. Owen and C. Prieur. On Shapley value for measuring importance of dependent inputs. *SIAM/ASA Journal on Uncertainty Quantification*, 5(1):986–1002, 2017.
- [8] A. Forrester, A. Sóbester, and A. Keane. *Engineering Design via Surrogate Modelling: A Practical Guide*. Wiley, 2008.

- [9] S. Surjanovic and D. Bingham. Virtual Library of Simulation Experiments: Test Functions and Datasets. <https://www.sfu.ca/~ssurjano/wingweight.html>, 2024.
- [10] D. P. Raymer. *Aircraft Design: A Conceptual Approach*. American Institute of Aeronautics and Astronautics, 2018.
- [11] G. Li, S. W. Wang, and H. Rabitz. Practical approaches to construct RS-HDMR component functions. *Journal of Physical Chemistry A*, 106(37):8721–8733, 2002.
- [12] M. E. Tipping. Sparse Bayesian learning and the relevance vector machine. *Journal of Machine Learning Research*, 1:211–244, 2001.
- [13] A. Saltelli. Making best use of model evaluations to compute sensitivity indices. *Computer Physics Communications*, 145(2):280–297, 2002.
- [14] OpenTURNS Consortium. OpenTURNS: Reference Guide. <https://openturns.github.io/openturns/latest>, 2024.
- [15] OpenTURNS Consortium. Sensitivity analysis on the wing weight model. https://openturns.github.io/openturns/latest/auto_reliability_sensitivity/sensitivity_analysis/plot_sensitivity_wingweight.html, 2024.
- [16] H. Moon, A. M. Dean, and T. J. Santner. Two-stage sensitivity-based group screening in computer experiments. *Technometrics*, 54(4):376–387, 2012.
- [17] L. S. Shapley. A value for n -person games. In *Contributions to the Theory of Games II*, pp. 307–317. Princeton University Press, 1953.
- [18] F. Bennett. SHAPLEYX: A Python package for global sensitivity analysis with RS-HDMR. *Journal of Open Source Software* (submitted), 2026.
- [19] F. Bennett. SHAPLEYX repository. <https://github.com/frbennett/shapleyx>, v0.5.1, 2026.