# Cell



Volume 171, Issue 2, 5 October 2017, Pages 321–330.e14

#### **Article**


# Single-Cell Analysis of Human Pancreas Reveals Transcriptional Signatures of Aging and Somatic Mutation Patterns

Martin Enge<sup>1, 6</sup>, H. Efsun Arda<sup>2</sup>, Marco Mignardi<sup>1, 5</sup>, John Beausang<sup>1</sup>, Rita Bottino<sup>4</sup>, Seung K. Kim<sup>2</sup>, Stephen R. Quake<sup>1, 3, 4, 7,</sup> ▲ □

- <sup>1</sup> Department of Bioengineering and Applied Physics, Stanford University, Stanford, CA 94305, USA
- <sup>2</sup> Department of Developmental Biology, Stanford University School of Medicine, CA 94305, USA
- <sup>3</sup> Chan Zuckerberg Biohub, San Francisco, CA 94158, USA
- <sup>4</sup> Institute of Cellular Therapeutics, Allegheny Health Network, 320 East North Avenue, Pittsburgh, PA 15212, USA
- <sup>5</sup> Department of Information Technology, Uppsala University, Sweden and SciLifeLab, Uppsala, Sweden SE-751 05

Received 23 March 2017, Revised 2 July 2017, Accepted 30 August 2017, Available online 28 September 2017






RNA-seq of single cells from donors allows detection of stochastic age-related errors:

- Cells from older donors have increased transcriptional noise and signs of fate drift
- •Endocrine pancreas cells display an oxidative stress-related mutational signature
- •Cellular stress and metabolic genes are high in cells with accumulation of errors

## Some definitions

- <u>Transcriptional Noise</u>: *transcriptome instability* or *Increased disregulation of gene expression*.
- <u>ERCC</u>: RNA spike-in is an RNA transcript of known sequence and quantity used to calibrate measurements in RNA hybridization assays, such as RNA-Seq.
- <u>Linear regression</u>: In statistics, **linear regression** is a linear approach for **modeling the relationship** between **a variable y** and **one or more explanatory variables** (or independent variables) denoted X.
- CPM: Counts per million. Unit to count gene expression level

## A Comprehensive Survey of Single Pancreatic Cells from Human Donors across Different Ages





#### Pancreas function:

Endocrine gland: alpha- (GluCaGon), beta- (INSulin) et delta- cells and pancreatic PP.


- Glucagon: UP glucose levels in blood.
- Insulin: DOWN glucose levels in blood.

Type2 Diabetes (age-related disease)






#### Transcriptional Instability and Fate Drift in Cells from Older Donors




pancreatic islet containing cells with atypical hormone expression




Aging is accompanied by the accumulation of somatic mutations


#### Somatic Mutation Profiles Derived from Single Primary Human Cells



## Endocrine Cells Display a Specific Mutational Signature Related to Oxidative Stress



## Mutational Load of Signature S1 Is Higher in Endocrine Cells from Older Donors and Correlate with Induction of Protein Synthesis-Related Genes



### Conclusion

- Aging is accompanied of increase in transcriptome noise and accumulation of genetic errors.
- Absence of causal link between transcriptional instability and mutational load.
- The cellular heterogeneity suggests that agingdependent changes are due to events in a subset of cells.
- Age specific mutational signature observed in endocrine is due to ROS-dependent lesions of DNA.
- They defined a method to determine transcriptome instability and mutations signatures from scRNAseq on arbitrary cells on primary tissue.