

Machine Learning

Week 3

Section 2


Regularization

The Problem of Overfitting

When algorithms are overfit, they perform poorly on new instances.

Overfitting is also known as 'high variance'

Example: Linear regression (housing prices)




Overfitting: If we have too many features, the learned hypothesis may fit the training set very well ($J(\theta) = \frac{1}{2m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})^2 \approx 0$), but fail to generalize to new examples (predict prices on new examples).

Addressing overfitting:

Options:

- Reduce number of features
 - Manually select which features to keep
 - Model selection algorithm (later in this course)
 - Combine features?
- Regularization
 - Keep all the features, but reduce the magnitude/values of parameters theta
 - Works well when we have lots of feature, each of which contributes a bit to predicting y

Consider the problem of predicting y from $x \in \mathbb{R}$. The leftmost figure below shows the result of fitting a $y = \theta_0 + \theta_1 x$ to a dataset. We see that the data doesn't really lie on straight line, and so the fit is not very good.



Instead, if we had added an extra feature x^2 , and fit $y = \theta_0 + \theta_1x + \theta_2x^2$, then we obtain a slightly better fit to the data (See middle figure). Naively, it might seem that the more features we add, the better. However, there is also a danger in adding too many features: The rightmost figure is the result of fitting a 5th order polynomial $y = \sum_{j=0}^5 \theta_j x^j$. We see that even though the fitted curve passes through the data perfectly, we would not expect this to be a very good predictor of, say, housing prices (y) for different living areas (x). Without formally defining what these terms mean, we'll say the figure on the left shows an instance of **underfitting**—in which the data clearly shows structure not captured by the model—and the figure on the right is an example of **overfitting**.

Underfitting, or high bias, is when the form of our hypothesis function h maps poorly to the trend of the data. It is usually caused by a function that is too simple or uses too few features. At the other extreme, overfitting, or high variance, is caused by a hypothesis function that fits the available data but does not generalize well to predict new data. It is usually caused by a complicated function that creates a lot of unnecessary curves and angles unrelated to the data.

This terminology is applied to both linear and logistic regression. There are two main options to address the issue of overfitting:

1) Reduce the number of features:


- Manually select which features to keep.
- Use a model selection algorithm (studied later in the course).

2) Regularization

- Keep all the features, but reduce the magnitude of parameters θ_j .
- Regularization works well when we have a lot of slightly useful features.

Cost Function

Intuition



Suppose we penalize and make θ_3, θ_4 really small.

$$\rightarrow \min_{\theta} \frac{1}{2m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})^2 + 1000 \underline{\theta_3^2} + 1000 \underline{\theta_4^2}$$

$\underline{\theta_3 \approx 0} \quad \underline{\theta_4 \approx 0}$

More generally,

- Small values for parameters of theta 1 ... n
 - "Simpler" hypothesis
- Less prone to overfitting

Since we don't know which values need to be regularized, we add a regularization term to the end of our cost function to penalize all values of theta > 0, lambda is the regularization parameter.

Housing:

$$\lambda \approx 0$$

- Features: x_1, x_2, \dots, x_{100}
- Parameters: $\theta_0, \theta_1, \theta_2, \dots, \theta_{100}$

$$J(\theta) = \frac{1}{2m} \left[\sum_{i=1}^m (h_\theta(x^{(i)}) - y^{(i)})^2 + \lambda \sum_{j=1}^n \theta_j^2 \right]$$

** exclude theta 0, as noted by the summation starting at 1

If lambda is too large, ex 10^{10} , then the hypothesis won't fit the problem at all and be useless. (Under-fitting)
Also known as 'too high a bias'

Later we'll cover how to automatically find a value for lambda

Note: [5:18 - There is a typo. It should be $\sum_{j=1}^n \theta_j^2$ instead of $\sum_{i=1}^n \theta_i^2$]

If we have overfitting from our hypothesis function, we can reduce the weight that some of the terms in our function carry by increasing their cost.

Say we wanted to make the following function more quadratic:

$$\theta_0 + \theta_1 x + \theta_2 x^2 + \theta_3 x^3 + \theta_4 x^4$$

We'll want to eliminate the influence of $\theta_3 x^3$ and $\theta_4 x^4$. Without actually getting rid of these features or changing the form of our hypothesis, we can instead modify our **cost function**:

$$\min_{\theta} \frac{1}{2m} \sum_{i=1}^m (h_\theta(x^{(i)}) - y^{(i)})^2 + 1000 \cdot \theta_3^2 + 1000 \cdot \theta_4^2$$

We've added two extra terms at the end to inflate the cost of θ_3 and θ_4 . Now, in order for the cost function to get close to zero, we will have to reduce the values of θ_3 and θ_4 to near zero. This will in turn greatly reduce the values of $\theta_3 x^3$ and $\theta_4 x^4$ in our hypothesis function. As a result, we see that the new hypothesis (depicted by the pink curve) looks like a quadratic function but fits the data better due to the extra small terms $\theta_3 x^3$ and $\theta_4 x^4$.

We could also regularize all of our theta parameters in a single summation as:

$$\min_{\theta} \frac{1}{2m} \sum_{i=1}^m (h_\theta(x^{(i)}) - y^{(i)})^2 + \lambda \sum_{j=1}^n \theta_j^2$$

The λ , or lambda, is the **regularization parameter**. It determines how much the costs of our theta parameters are inflated.

Using the above cost function with the extra summation, we can smooth the output of our hypothesis function to reduce overfitting. If lambda is chosen to be too large, it may smooth out the function too much and cause underfitting. Hence, what would happen if $\lambda = 0$ or is too small ?

Regularized Linear Regression

Recall we worked out two learning algorithms for linear regression:

- Gradient Descent
- Normal Equation

Now we'll generalize them for Regularized Linear Regression

Gradient Descent:

Gradient descent

$$\frac{\partial J(\theta)}{\partial \theta_0} = \frac{\partial J(\theta)}{\partial \theta_1, \theta_2, \dots, \theta_n}$$

Repeat {

$$\theta_0 := \theta_0 - \alpha \frac{1}{m} \sum_{i=1}^m (h_\theta(x^{(i)}) - y^{(i)}) x_0^{(i)}$$

$$\theta_j := \theta_j - \alpha \left[\frac{1}{m} \sum_{i=1}^m (h_\theta(x^{(i)}) - y^{(i)}) x_j^{(i)} - \frac{\lambda}{m} \theta_j \right] \quad (j = 1, 2, 3, \dots, n)$$

$$\theta_j := \theta_j (1 - \alpha \frac{\lambda}{m}) - \alpha \frac{1}{m} \sum_{i=1}^m (h_\theta(x^{(i)}) - y^{(i)}) x_j^{(i)}$$

$$|1 - \alpha \frac{\lambda}{m}| < 1 \quad 0.99 \quad \theta_j \times 0.99$$

$\rightarrow J(\theta)$

θ_j

Normal Equation:

Normal equation

$$X = \begin{bmatrix} (x^{(1)})^T \\ \vdots \\ (x^{(m)})^T \end{bmatrix}_{m \times (n+1)} \quad y = \begin{bmatrix} y^{(1)} \\ \vdots \\ y^{(m)} \end{bmatrix} \in \mathbb{R}^m$$

$$\min_{\theta} J(\theta) \quad \frac{\partial J(\theta)}{\partial \theta_j} \stackrel{\text{set } 0}{=} 0 \quad \Rightarrow$$

$$\theta = (X^T X + \lambda \begin{bmatrix} 0 & 1 & 0 & \dots & 0 \\ 1 & 0 & 1 & \dots & 0 \\ 0 & 1 & 0 & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \dots & 1 \end{bmatrix})^{-1} X^T y$$

E.g. $n=2$ $\begin{bmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \quad (n+1) \times (n+1)$

Non-invertibility issue:

Even if the non-invertibility issue arises, for the regularized normal equation there won't be a problem and the inner matrix will be invertible.

Non-invertibility (optional/advanced).

Suppose $m \leq n$, \leftarrow

(#examples) (#features)

$$\theta = \frac{(X^T X)^{-1} X^T y}{\text{non-invertible / singular}} \quad \underline{\text{pinv}} \quad \underline{\frac{\text{inv}}{\kappa}}$$

If $\lambda > 0$,

$$\theta = \left(X^T X + \lambda \begin{bmatrix} 0 & & & \\ & 1 & & \\ & & 1 & \\ & & & \ddots \\ & & & & 1 \end{bmatrix} \right)^{-1} X^T y$$

invertible.

Using this method you will be able to avoid overfitting even if you have a high number of features.

Note: [8:43 - It is said that X is non-invertible if $m \leq n$. The correct statement should be that X is non-invertible if $m < n$, and may be non-invertible if $m = n$.

We can apply regularization to both linear regression and logistic regression. We will approach linear regression first.

Gradient Descent

We will modify our gradient descent function to separate out θ_0 from the rest of the parameters because we do not want to penalize θ_0 .

Repeat {

$$\begin{aligned} \theta_0 &:= \theta_0 - \alpha \frac{1}{m} \sum_{i=1}^m (h_\theta(x^{(i)}) - y^{(i)}) x_0^{(i)} \\ \theta_j &:= \theta_j - \alpha \left[\left(\frac{1}{m} \sum_{i=1}^m (h_\theta(x^{(i)}) - y^{(i)}) x_j^{(i)} \right) + \frac{\lambda}{m} \theta_j \right] \quad j \in \{1, 2, \dots, n\} \\ \} \end{aligned}$$

The term $\frac{\lambda}{m} \theta_j$ performs our regularization. With some manipulation our update rule can also be represented as:

$$\theta_j := \theta_j \left(1 - \alpha \frac{\lambda}{m} \right) - \alpha \frac{1}{m} \sum_{i=1}^m (h_\theta(x^{(i)}) - y^{(i)}) x_j^{(i)}$$

The first term in the above equation, $1 - \alpha \frac{\lambda}{m}$ will always be less than 1. Intuitively you can see it as reducing the value of θ_j by some amount on every update. Notice that the second term is now exactly the same as it was before.

Normal Equation

Now let's approach regularization using the alternate method of the non-iterative normal equation.

To add in regularization, the equation is the same as our original, except that we add another term inside the parentheses:

$$\theta = (X^T X + \lambda \cdot L)^{-1} X^T y$$

where $L = \begin{bmatrix} 0 & & & \\ & 1 & & \\ & & 1 & \\ & & & \ddots & \\ & & & & 1 \end{bmatrix}$

L is a matrix with 0 at the top left and 1's down the diagonal, with 0's everywhere else. It should have dimension $(n+1) \times (n+1)$. Intuitively, this is the identity matrix (though we are not including x_0), multiplied with a single real number λ .

Recall that if $m < n$, then $X^T X$ is non-invertible. However, when we add the term $\lambda \cdot L$, then $X^T X + \lambda \cdot L$ becomes invertible.


Regularized Logistic Regression

We'll go over how to adapt both gradient descent and more advanced optimization techniques to work for Regularized Logistic Regression:

Recall: Using many high order polynomial or many features will result in overfitting:

We can adapt our cost function as follows:

Regularized logistic regression.



Cost function:

$$\rightarrow J(\theta) = - \left[\frac{1}{m} \sum_{i=1}^m y^{(i)} \log h_{\theta}(x^{(i)}) + (1 - y^{(i)}) \log (1 - h_{\theta}(x^{(i)})) \right] + \frac{\lambda}{2m} \sum_{j=1}^n \theta_j^2$$

$\theta_1, \theta_2, \dots, \theta_n$

We will change our gradient descent algorithm as follows to use the Regularized Cost Function:

Gradient descent

Repeat {

$$\rightarrow \theta_0 := \theta_0 - \alpha \frac{1}{m} \sum_{i=1}^m (h_\theta(x^{(i)}) - y^{(i)}) x_0^{(i)}$$

$$\rightarrow \theta_j := \theta_j - \alpha \left[\frac{1}{m} \sum_{i=1}^m (h_\theta(x^{(i)}) - y^{(i)}) x_j^{(i)} + \frac{\lambda}{m} \theta_j \right] \leftarrow$$

$(j = \cancel{0}, 1, 2, 3, \dots, n)$

$\theta_0, \dots, \theta_n$

$$\frac{\partial}{\partial \theta_j} J(\theta)$$

$$h_\theta(x) = \frac{1}{1 + e^{-\theta^T x}}$$

How to get Regularized Logistic Regression to work with Advanced Optimization:

Advanced optimization

```


function [jVal, gradient] = costFunction(theta)
    jVal = [ code to compute  $J(\theta)$  ];
     $\rightarrow J(\theta) = \left[ -\frac{1}{m} \sum_{i=1}^m y^{(i)} \log(h_\theta(x^{(i)})) + (1 - y^{(i)}) \log(1 - h_\theta(x^{(i)})) \right] + \left[ \frac{\lambda}{2m} \sum_{j=1}^n \theta_j^2 \right]$ 
     $\rightarrow \text{gradient}(1) = [ \text{code to compute } \frac{\partial}{\partial \theta_0} J(\theta) ];$ 
     $\frac{1}{m} \sum_{i=1}^m (h_\theta(x^{(i)}) - y^{(i)}) x_0^{(i)} \leftarrow$ 
     $\rightarrow \text{gradient}(2) = [ \text{code to compute } \frac{\partial}{\partial \theta_1} J(\theta) ];$ 
     $\left( \frac{1}{m} \sum_{i=1}^m (h_\theta(x^{(i)}) - y^{(i)}) x_1^{(i)} \right) + \frac{\lambda}{m} \theta_1 \leftarrow$ 
     $\rightarrow \text{gradient}(3) = [ \text{code to compute } \frac{\partial}{\partial \theta_2} J(\theta) ];$ 
     $\vdots \quad \left( \frac{1}{m} \sum_{i=1}^m (h_\theta(x^{(i)}) - y^{(i)}) x_2^{(i)} \right) + \frac{\lambda}{m} \theta_2 \leftarrow$ 
     $\text{gradient}(n+1) = [ \text{code to compute } \frac{\partial}{\partial \theta_n} J(\theta) ];$ 

```

Regularized Logistic Regression

We can regularize logistic regression in a similar way that we regularize linear regression. As a result, we can avoid overfitting. The following image shows how the regularized function, displayed by the pink line, is less likely to overfit than the non-regularized function represented by the blue line:

Regularized logistic regression.



Cost function:

$$\rightarrow J(\theta) = - \left[\frac{1}{m} \sum_{i=1}^m y^{(i)} \log h_{\theta}(x^{(i)}) + (1 - y^{(i)}) \log(1 - h_{\theta}(x^{(i)})) \right] + \frac{\lambda}{2m} \sum_{j=1}^n \theta_j^2 \quad | \boxed{\theta_1, \theta_2, \dots, \theta_n}$$

Regularization term

Cost Function

Recall that our cost function for logistic regression was:

$$J(\theta) = -\frac{1}{m} \sum_{i=1}^m [y^{(i)} \log(h_{\theta}(x^{(i)})) + (1 - y^{(i)}) \log(1 - h_{\theta}(x^{(i)}))]$$

We can regularize this equation by adding a term to the end:

$$J(\theta) = -\frac{1}{m} \sum_{i=1}^m [y^{(i)} \log(h_{\theta}(x^{(i)})) + (1 - y^{(i)}) \log(1 - h_{\theta}(x^{(i)}))] + \frac{\lambda}{2m} \sum_{j=1}^n \theta_j^2$$

The second sum, $\sum_{j=1}^n \theta_j^2$ means to explicitly exclude the bias term, θ_0 . I.e. the θ vector is indexed from 0 to n (holding n+1 values, θ_0 through θ_n), and this sum explicitly skips θ_0 , by running from 1 to n, skipping 0. Thus, when computing the equation, we should continuously update the two following equations:

Gradient descent

Repeat {

$$\rightarrow \theta_0 := \theta_0 - \alpha \frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)}) x_0^{(i)}$$

$$\rightarrow \theta_j := \theta_j - \alpha \left[\frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)}) x_j^{(i)} + \frac{\lambda}{m} \theta_j \right] \leftarrow$$

$(j = \boxed{1, 2, 3, \dots, n})$

$\theta_1, \dots, \theta_n$

$$\frac{\partial}{\partial \theta_j} J(\theta) \quad h_{\theta}(x) = \frac{1}{1 + e^{-\theta^T x}}$$

