




Hohlaubgletscher Length Change Over Time




Hohlaubgletscher Cumulative Length Change Over Time




Hohlaubgletscher Annual Mass Balance Over Time



Hohlaubgletscher Cumulative Annual Mass Balance Over Time



Annual Mass Balance for each Elevation Bin over Time - Hohlaubgletscher




Sion Summer Mean Temperature



Sion Winter Total Precipitation



Hohlaubgletscher Summer Mass Balance with relation to Temperature





Regression: Monthly 1961-1990

=====
MONTHLY DEVIATIONS ANALYSIS USING 1961-1990 CLIMATE NORMS
=====

=====
MONTHLY DEVIATIONS for Hohlaubgletscher (1961-1990 norms)
=====

Correlation Analysis with Significance Testing:
Table with 5 columns: Variable, Correlation Coefficient, P-value, Significant (p < 0.05). Rows include months (october\_pd to april\_pd) and time periods (may\_td to july\_td), plus a constant.

Number of observations: 70

Regression Summary:

OLS Regression Results
Table with 2 columns: Dep. Variable: annual mass balance (mm w.e.), and various statistics (R-squared, Adj. R-squared, F-statistic, etc.).

Table with 7 columns: coef, std err, t, P>|t|, [0.025, 0.975]. Rows include coefficients for various months (const, may\_td, june\_td, etc.) and time periods.

Table with 4 columns: Omnibus, Prob(Omnibus), Skew, Kurtosis; Durbin-Watson, Jarque-Bera (JB), Prob(JB), Cond. No.

Regression: Optimal 1961-1990

=====
OPTIMAL SEASONAL DEVIATIONS ANALYSIS USING 1961-1990 CLIMATE NORMS
=====

=====
OPTIMAL SEASONAL DEVIATIONS for Hohlaubgletscher (1961-1990 norms)
=====

Correlation Analysis with Significance Testing:

	Variable	Correlation Coefficient	P-value	Significant (p < 0.05)
2	opt_season_pd	0.272364	2.254596e-02	True
1	opt_season_td	-0.604964	2.908402e-08	True
0	const	NaN	NaN	False

Number of observations: 70

Regression Summary:

OLS Regression Results						
=====						
Dep. Variable:	annual mass balance (mm w.e.)			R-squared:	0.410	
Model:	OLS			Adj. R-squared:	0.393	
Method:	Least Squares			F-statistic:	23.30	
Date:	Mon, 08 Dec 2025			Prob (F-statistic):	2.08e-08	
Time:	12:08:42			Log-Likelihood:	-556.88	
No. Observations:	70			AIC:	1120.	
Df Residuals:	67			BIC:	1126.	
Df Model:	2					
Covariance Type:	nonrobust					
=====						
	coef	std err	t	P> t	[0.025	0.975]
-----						
const	1.41e+04	2336.689	6.033	0.000	9434.146	1.88e+04
opt_season_td	-415.0115	67.175	-6.178	0.000	-549.094	-280.929
opt_season_pd	2.3416	1.045	2.241	0.028	0.256	4.427
=====						
Omnibus:	8.191		Durbin-Watson:		1.533	
Prob(Omnibus):	0.017		Jarque-Bera (JB):		7.858	
Skew:	-0.655		Prob(JB):		0.0197	
Kurtosis:	3.989		Cond. No.		2.29e+03	
=====						

Notes:
[1] Standard Errors assume that the covariance matrix of the errors is correctly specified.
[2] The condition number is large, 2.29e+03. This might indicate that there are strong multicollinearity or other numerical problems.

Coefficient Interpretation:
Intercept (normal mass balance): 14098.20 (p=0.0000)
opt\_season\_td: -415.01 (p=0.0000)
opt\_season\_pd: 2.34 (p=0.0284)

Variance Inflation Factors (VIF):

	Variable	VIF
0	const	768.775289
1	opt_season_td	1.011060
2	opt_season_pd	1.011060

R-squared: 0.4102
Adjusted R-squared: 0.3926

Regression: Seasonal 1961-1990

=====
SUMMER/WINTER SEASONAL DEVIATIONS ANALYSIS USING 1961-1990 CLIMATE NORMS
=====

=====
SUMMER/WINTER SEASONAL DEVIATIONS for Hohlaubgletscher (1961-1990 norms)
=====

Correlation Analysis with Significance Testing:
Table with 5 columns: Variable, Correlation Coefficient, P-value, Significant (p < 0.05)
Rows: winter\_pd, summer\_td, const

Number of observations: 70

Regression Summary:

OLS Regression Results
Table with 7 columns: Dep. Variable, annual mass balance (mm w.e.), R-squared, Adj. R-squared, Method, Least Squares, F-statistic, Date, Mon, 08 Dec 2025, Prob (F-statistic), Time, 12:08:42, Log-Likelihood, No. Observations, 70, AIC, Df Residuals, 67, BIC, Df Model, 2, Covariance Type, nonrobust
Table with 7 columns: coef, std err, t, P>|t|, [0.025, 0.975]
Rows: const, summer\_td, winter\_pd
Omnibus, Prob(Omnibus), Skew, Kurtosis, Durbin-Watson, Jarque-Bera (JB), Prob(JB), Cond. No.

Notes:
[1] Standard Errors assume that the covariance matrix of the errors is correctly specified.
[2] The condition number is large, 2.76e+03. This might indicate that there are strong multicollinearity or other numerical problems.

Coefficient Interpretation:
Intercept (normal mass balance): 15711.42 (p=0.0000)
summer\_td: -474.42 (p=0.0000)
winter\_pd: 1.82 (p=0.0472)

Variance Inflation Factors (VIF):
Table with 2 columns: Variable, VIF
Rows: const, summer\_td, winter\_pd

R-squared: 0.4326
Adjusted R-squared: 0.4156

# Regression: Monthly 1991-2020

## MONTHLY DEVIATIONS ANALYSIS USING 1991-2020 CLIMATE NORMS

## MONTHLY DEVIATIONS for Hohlaubgletscher (1991-2020 norms)

### Correlation Analysis with Significance Testing:

	Variable	Correlation Coefficient	P-value	Significant (p < 0.05)
6	october_pd	0.192638	1.101076e-01	False
10	february_pd	0.154829	2.006150e-01	False
7	november_pd	0.101388	4.036308e-01	False
8	december_pd	0.082763	4.957804e-01	False
11	march_pd	0.064021	5.985187e-01	False
9	january_pd	0.014279	9.066076e-01	False
12	april_pd	-0.141633	2.421798e-01	False
1	may_td	-0.388732	8.821633e-04	True
5	september_td	-0.414801	3.566138e-04	True
2	june_td	-0.444657	1.150260e-04	True
4	august_td	-0.467402	4.515425e-05	True
3	july_td	-0.551877	7.340182e-07	True
0	const	NaN	NaN	False

Number of observations: 70

### Regression Summary:

#### OLS Regression Results

Dep. Variable:	annual mass balance (mm w.e.)	R-squared:	0.518
Model:	OLS	Adj. R-squared:	0.417
Method:	Least Squares	F-statistic:	5.105
Date:	Mon, 08 Dec 2025	Prob (F-statistic):	1.05e-05
Time:	12:08:42	Log-Likelihood:	-549.81
No. Observations:	70	AIC:	1126.
Df Residuals:	57	BIC:	1155.
Df Model:	12		
Covariance Type:	nonrobust		

	coef	std err	t	P> t	[0.025	0.975]
const	-761.0987	100.998	-7.536	0.000	-963.343	-558.854
may_td	-51.7296	66.025	-0.783	0.437	-183.942	80.483
june_td	-16.4373	62.241	-0.264	0.793	-141.073	108.198
july_td	-161.9487	66.779	-2.425	0.018	-295.671	-28.226
august_td	-132.5787	78.796	-1.683	0.098	-290.364	25.207
september_td	-121.8690	62.697	-1.944	0.057	-247.418	3.680
october_pd	7.0823	3.071	2.306	0.025	0.932	13.232
november_pd	2.8305	2.282	1.240	0.220	-1.739	7.400
december_pd	4.4549	1.907	2.336	0.023	0.636	8.274
january_pd	0.6375	2.321	0.275	0.785	-4.010	5.285
february_pd	1.2869	1.802	0.714	0.478	-2.322	4.896
march_pd	-0.7323	2.653	-0.276	0.784	-6.045	4.581
april_pd	-4.2385	4.053	-1.046	0.300	-12.354	3.877

Omnibus:	3.799	Durbin-Watson:	1.573
Prob(Omnibus):	0.150	Jarque-Bera (JB):	3.408
Skew:	-0.540	Prob(JB):	0.182
Kurtosis:	3.006	Cond. No.	65.8

Regression: Optimal 1991-2020

=====
OPTIMAL SEASONAL DEVIATIONS ANALYSIS USING 1991-2020 CLIMATE NORMS
=====

=====
OPTIMAL SEASONAL DEVIATIONS for Hohlaubgletscher (1991-2020 norms)
=====

Correlation Analysis with Significance Testing:

	Variable	Correlation Coefficient	P-value	Significant (p < 0.05)
2	opt_season_pd	0.272364	2.254596e-02	True
1	opt_season_td	-0.605611	2.786038e-08	True
0	const	NaN	NaN	False

Number of observations: 70

Regression Summary:

OLS Regression Results						
=====						
Dep. Variable:	annual mass balance (mm w.e.)			R-squared:	0.409	
Model:	OLS			Adj. R-squared:	0.392	
Method:	Least Squares			F-statistic:	23.21	
Date:	Mon, 08 Dec 2025			Prob (F-statistic):	2.19e-08	
Time:	12:08:42			Log-Likelihood:	-556.93	
No. Observations:	70			AIC:	1120.	
Df Residuals:	67			BIC:	1127.	
Df Model:	2					
Covariance Type:	nonrobust					
=====						
	coef	std err	t	P> t	[0.025	0.975]
-----						
const	-749.6822	101.557	-7.382	0.000	-952.390	-546.974
opt_season_td	-414.5566	67.243	-6.165	0.000	-548.774	-280.339
opt_season_pd	2.2984	1.047	2.196	0.032	0.209	4.387
=====						
Omnibus:	7.664		Durbin-Watson:		1.534	
Prob(Omnibus):	0.022		Jarque-Bera (JB):		7.203	
Skew:	-0.628		Prob(JB):		0.0273	
Kurtosis:	3.944		Cond. No.		107.	
=====						

Notes:
[1] Standard Errors assume that the covariance matrix of the errors is correctly specified.

Coefficient Interpretation:
Intercept (normal mass balance): -749.68 (p=0.0000)
opt\_season\_td: -414.56 (p=0.0000)
opt\_season\_pd: 2.30 (p=0.0316)

Variance Inflation Factors (VIF):
Variable VIF
0 const 1.449949
1 opt\_season\_td 1.012557
2 opt\_season\_pd 1.012557

R-squared: 0.4093
Adjusted R-squared: 0.3917

# Regression: Seasonal 1991-2020

=====

SUMMER/WINTER SEASONAL DEVIATIONS ANALYSIS USING 1991-2020 CLIMATE NORMS

=====

=====

SUMMER/WINTER SEASONAL DEVIATIONS for Hohlaubgletscher (1991-2020 norms)

=====

Correlation Analysis with Significance Testing:

	Variable	Correlation Coefficient	P-value	Significant (p < 0.05)
2	winter_pd	0.227639	5.806108e-02	False
1	summer_td	-0.628791	5.584849e-09	True
0	const	NaN	NaN	False

Number of observations: 70

Regression Summary:

OLS Regression Results						
=====						
Dep. Variable:	annual mass balance (mm w.e.)			R-squared:	0.431	
Model:	OLS			Adj. R-squared:	0.414	
Method:	Least Squares			F-statistic:	25.33	
Date:	Mon, 08 Dec 2025			Prob (F-statistic):	6.41e-09	
Time:	12:08:42			Log-Likelihood:	-555.64	
No. Observations:	70			AIC:	1117.	
Df Residuals:	67			BIC:	1124.	
Df Model:	2					
Covariance Type:	nonrobust					
=====						
	coef	std err	t	P> t	[0.025	0.975]
-----						
const	-763.8050	99.610	-7.668	0.000	-962.627	-564.983
summer_td	-472.5222	70.779	-6.676	0.000	-613.797	-331.247
winter_pd	1.8323	0.900	2.036	0.046	0.036	3.629
=====						
Omnibus:	6.431		Durbin-Watson:		1.533	
Prob(Omnibus):	0.040		Jarque-Bera (JB):		5.631	
Skew:	-0.599		Prob(JB):		0.0599	
Kurtosis:	3.704		Cond. No.		124.	
=====						

Notes:

[1] Standard Errors assume that the covariance matrix of the errors is correctly specified.

Coefficient Interpretation:

Intercept (normal mass balance): -763.81 (p=0.0000)

summer\_td: -472.52 (p=0.0000)

winter\_pd: 1.83 (p=0.0457)

Variance Inflation Factors (VIF):

	Variable	VIF
0	const	1.447098
1	summer_td	1.004137
2	winter_pd	1.004137

R-squared: 0.4306

Adjusted R-squared: 0.4136