




Claridenfirn Length Change Over Time




Claridenfirn Cumulative Length Change Over Time




Claridenfirn Annual Mass Balance Over Time




Claridenfirn Cumulative Annual Mass Balance Over Time




Annual Mass Balance for each Elevation Bin over Time - Claridenfirn




Altdorf Summer Mean Temperature




Aldorf Winter Total Precipitation



Claridenfirn Summer Mass Balance with relation to Temperature



Claridenfirn Winter Mass Balance with relation to Precipitation



Regression: Monthly 1961-1990

=====
MONTHLY DEVIATIONS for Claridenfirn using 1961-1990 climate norms
=====

Correlation Analysis with Significance Testing:
Skipping constant column: const
Table with 5 columns: Variable, Correlation Coefficient, P-value, Significant (p < 0.05), and an unlabeled column. Rows include months from july_td to november_pd.

Number of observations: 109

Regression Summary:

OLS Regression Results
Table with 2 columns: Label and Value. Rows include Dep. Variable, Model, Method, Date, Time, No. Observations, Df Residuals, Df Model, Covariance Type, R-squared, Adj. R-squared, F-statistic, Prob (F-statistic), Log-Likelihood, AIC, and BIC.

Table with 7 columns: coef, std err, t, P>|t|, [0.025, 0.975]. Rows include const, may_td, june_td, july_td, august_td, september_td, october_pd, november_pd, december_pd, january_pd, february_pd, march_pd, and april_pd.

Table with 4 columns: Label, Value, Label, Value. Rows include Omnibus, Prob(Omnibus), Skew, Kurtosis, Durbin-Watson, Jarque-Bera (JB), Prob(JB), and Cond. No.

Notes:
[1] Standard Errors assume that the covariance matrix of the errors is correctly specified.

Regression: Optimal 1961-1990

=====

OPTIMAL SEASONAL DEVIATIONS for Claridenfirn using 1961-1990 climate norms

=====

Correlation Analysis with Significance Testing:
Skipping constant column: const

	Variable	Correlation Coefficient	P-value	Significant (p < 0.05)
0	opt_season_td	-0.729232	2.449264e-19	True
1	opt_season_pd	0.333928	3.873699e-04	True

Number of observations: 109

Regression Summary:

OLS Regression Results						
=====						
Dep. Variable:	annual mass balance (mm w.e.)			R-squared:	0.640	
Model:	OLS			Adj. R-squared:	0.633	
Method:	Least Squares			F-statistic:	94.27	
Date:	Wed, 17 Dec 2025			Prob (F-statistic):	3.00e-24	
Time:	20:25:16			Log-Likelihood:	-842.09	
No. Observations:	109			AIC:	1690.	
Df Residuals:	106			BIC:	1698.	
Df Model:	2					
Covariance Type:	nonrobust					
=====						
	coef	std err	t	P> t	[0.025	0.975]

const	83.3856	60.955	1.368	0.174	-37.464	204.236
opt_season_td	-604.5066	48.447	-12.478	0.000	-700.557	-508.456
opt_season_pd	2.8935	0.512	5.649	0.000	1.878	3.909
=====						
Omnibus:	1.729		Durbin-Watson:		1.776	
Prob(Omnibus):	0.421		Jarque-Bera (JB):		1.287	
Skew:	-0.250		Prob(JB):		0.526	
Kurtosis:	3.180		Cond. No.		133.	
=====						

Notes:
[1] Standard Errors assume that the covariance matrix of the errors is correctly specified.

Regression: Seasonal 1961-1990

=====
SUMMER/WINTER SEASONAL DEVIATIONS for Claridenfirn using 1961-1990 climate norms
=====

Correlation Analysis with Significance Testing:
Skipping constant column: const
Variable Correlation Coefficient P-value Significant (p < 0.05)
0 summer_td -0.772507 7.814267e-23 True
1 winter_pd 0.344108 2.485777e-04 True

Number of observations: 109

Regression Summary:

OLS Regression Results						
=====						
Dep. Variable:	annual mass balance (mm w.e.)			R-squared:	0.717	
Model:	OLS			Adj. R-squared:	0.712	
Method:	Least Squares			F-statistic:	134.6	
Date:	Wed, 17 Dec 2025			Prob (F-statistic):	8.03e-30	
Time:	20:25:16			Log-Likelihood:	-828.90	
No. Observations:	109			AIC:	1664.	
Df Residuals:	106			BIC:	1672.	
Df Model:	2					
Covariance Type:	nonrobust					
=====						
	coef	std err	t	P> t	[0.025	0.975]

const	113.6658	53.563	2.122	0.036	7.471	219.860
summer_td	-708.5002	47.257	-14.993	0.000	-802.191	-614.809
winter_pd	2.6683	0.396	6.730	0.000	1.882	3.454
=====						
Omnibus:	3.151		Durbin-Watson:		1.635	
Prob(Omnibus):	0.207		Jarque-Bera (JB):		2.632	
Skew:	-0.367		Prob(JB):		0.268	
Kurtosis:	3.201		Cond. No.		155.	
=====						

Notes:
[1] Standard Errors assume that the covariance matrix of the errors is correctly specified.

Regression: Monthly 1991-2020

=====
MONTHLY DEVIATIONS for Claridenfirn using 1991-2020 climate norms
=====

Correlation Analysis with Significance Testing:
Skipping constant column: const
Table with 5 columns: Variable, Correlation Coefficient, P-value, Significant (p < 0.05), and an unlabeled column. Rows include months from July to November.

Number of observations: 109

Regression Summary:

OLS Regression Results
Table with 7 columns: Variable, coef, std err, t, P>|t|, [0.025, 0.975]. Rows include const, months, and diagnostic statistics (Omnibus, Prob(Omnibus), Skew, Kurtosis).

Notes:
[1] Standard Errors assume that the covariance matrix of the errors is correctly specified.

Regression: Optimal 1991-2020

=====

OPTIMAL SEASONAL DEVIATIONS for Claridenfirn using 1991-2020 climate norms

=====

Correlation Analysis with Significance Testing:
Skipping constant column: const

	Variable	Correlation Coefficient	P-value	Significant (p < 0.05)
0	opt_season_td	-0.731612	1.638598e-19	True
1	opt_season_pd	0.333928	3.873699e-04	True

Number of observations: 109

Regression Summary:

OLS Regression Results

=====

Dep. Variable:	annual mass balance (mm w.e.)	R-squared:	0.642
Model:	OLS	Adj. R-squared:	0.635
Method:	Least Squares	F-statistic:	94.89
Date:	Wed, 17 Dec 2025	Prob (F-statistic):	2.40e-24
Time:	20:25:16	Log-Likelihood:	-841.86
No. Observations:	109	AIC:	1690.
Df Residuals:	106	BIC:	1698.
Df Model:	2		
Covariance Type:	nonrobust		

=====

	coef	std err	t	P> t	[0.025	0.975]
const	-844.3329	70.849	-11.917	0.000	-984.798	-703.868
opt_season_td	-610.0771	48.721	-12.522	0.000	-706.672	-513.482
opt_season_pd	2.8672	0.511	5.609	0.000	1.854	3.881

=====

Omnibus:	2.136	Durbin-Watson:	1.779
Prob(Omnibus):	0.344	Jarque-Bera (JB):	1.580
Skew:	-0.252	Prob(JB):	0.454
Kurtosis:	3.307	Cond. No.	156.

=====

Notes:

[1] Standard Errors assume that the covariance matrix of the errors is correctly specified.

Regression: Seasonal 1991-2020

=====
SUMMER/WINTER SEASONAL DEVIATIONS for Claridenfirn using 1991-2020 climate norms
=====

Correlation Analysis with Significance Testing:
Skipping constant column: const
Variable Correlation Coefficient P-value Significant (p < 0.05)
0 summer_td -0.772507 7.814267e-23 True
1 winter_pd 0.344108 2.485777e-04 True

Number of observations: 109

Regression Summary:

OLS Regression Results						
Dep. Variable:	annual mass balance (mm w.e.)			R-squared:	0.717	
Model:	OLS			Adj. R-squared:	0.712	
Method:	Least Squares			F-statistic:	134.6	
Date:	Wed, 17 Dec 2025			Prob (F-statistic):	8.03e-30	
Time:	20:25:16			Log-Likelihood:	-828.90	
No. Observations:	109			AIC:	1664.	
Df Residuals:	106			BIC:	1672.	
Df Model:	2					
Covariance Type:	nonrobust					
	coef	std err	t	P> t	[0.025	0.975]
const	-864.8931	62.383	-13.864	0.000	-988.574	-741.212
summer_td	-708.5002	47.257	-14.993	0.000	-802.191	-614.809
winter_pd	2.6683	0.396	6.730	0.000	1.882	3.454
Omnibus:	3.151	Durbin-Watson:		1.635		
Prob(Omnibus):	0.207	Jarque-Bera (JB):		2.632		
Skew:	-0.367	Prob(JB):		0.268		
Kurtosis:	3.201	Cond. No.		181.		

Notes:
[1] Standard Errors assume that the covariance matrix of the errors is correctly specified.