

CHAPTER 5 REVIEW

When the vessel breaks

Pathophysiology of non-traumatic
intracerebral haemorrhage

Oxford Textbook of Stroke & Cerebrovascular Disease, OUP 2014


Chapter authors: Constanza Rossi & Charlotte Cordonnier



10 December 1996

7 a.m. · Jill Bolte Taylor, neuroanatomist, aged 37

“I woke up to a pounding pain behind my left eye.”



Source: Jill Bolte Taylor, "My stroke of insight", TED 2008.

10 December 1996

7 a.m. · Jill Bolte Taylor, neuroanatomist, aged 37

Within four hours she could not **walk, talk, read or write.**



Source: Jill Bolte Taylor, "My stroke of insight", TED 2008.


40%

**die within a month of an
intracerebral haemorrhage**

No improvement from 1980 to 2008.

van Asch et al., Lancet Neurol 2010 (36 population studies, 8145 patients): median 1-month case fatality 40.4%.

ICH is a process, not a moment




Each stage is a window to act.


01

The vessel

Why does a small artery give way?




Hypertension and amyloid cause over half



55%
two small-vessel
diseases


SMASH-U: Meretoja et al., Stroke 2012 (n = 1013, Helsinki). Full breakdown on the handout.

Two diseases, two sets of vessels




Healthy

Thin wall, wide channel



Hypertension


Deep arteries: thick, brittle wall



Amyloid angiopathy

Surface arteries: amyloid replaces muscle

Where it bleeds tells you why




Deep

→ hypertension

putamen • thalamus • pons • cerebellum

Schematic axial slice. The pons and cerebellum sit below this level.

Where it bleeds tells you why



Deep

→ hypertension

putamen • thalamus • pons • cerebellum

Lobar

→ amyloid (older adults)

Young or atypical? Look for a lesion.

Schematic axial slice. The pons and cerebellum sit below this level.

Microbleeds are the footprints



Surface
→ amyloid



Deep
→ hypertension

60%

**of people with ICH have
microbleeds on MRI**

Healthy adults: 5%

Cordonnier, Al-Shahi Salman & Wardlaw, Brain 2007 (systematic review). Boston criteria on the handout.

Jill: primary or secondary?



37 years old



No hypertension



Lobar bleed



Secondary: an arteriovenous malformation



37 years old



No hypertension



Lobar bleed

65%

of patients like Jill had a lesion on angiography

0%

if over 45, hypertensive, with a deep bleed

Zhu et al., Stroke 1997 (prospective; 206 patients, all had catheter angiography). Other secondary causes on the handout.


02

The first hours

Rupture, growth and pressure




One rupture can trigger more



Fisher CM, J Neuropathol Exp Neurol 1971: secondary bleeding points at the edge of the haematoma.

How many bleeds keep growing?




Each dot = one patient, scanned within 3 hours of onset

A 5 in 100

B 15 in 100

C 40 in 100

38 in 100 keep growing




Each dot = one patient, scanned within 3 hours of onset

- 26 grew within the first hour
- 12 more grew within 20 hours
- 62 did not grow substantially

Jill: four hours
from headache to losing her speech

Brott et al., Stroke 1997 (n = 103; growth > 33%). Every 10% of growth: +5% hazard of death (Davis et al., Neurology 2006).

Jill's clot was the size of a golf ball




A golf ball $\approx$ 40 mL

60 mL + GCS $\leq$ 8: **91% predicted 30-day mortality**

A clot that grows by a third looks only 10% wider on a scan.

Volume: Broderick et al., Stroke 1993 (Cincinnati, n = 188). A regulation golf ball is 42.7 mm across (about 40.7 mL).

The skull is a closed box



≈45%

break into the ventricles

→ hydrocephalus, worse outcome

Hanley, Stroke 2009.


03

The days after

When the blood becomes a toxin



Oedema builds in waves



+75% oedema in the first 24 hours

Jill: surgery 17 days later

Xi, Keep & Hoff, Lancet Neurol 2006; Gebel et al., Stroke 2002.

Three toxins, one result



Thrombin



Iron



Inflammation



Leaky barrier → more oedema → neuronal death

Xi, Keep & Hoff, Lancet Neurol 2006. Mostly from animal models.

True or false?


“The rim around the clot is starving tissue.”

False: the rim is resting, not starving

“The rim around the clot is starving tissue.”

Lowering BP below 150 did not reduce flow around the clot (ICH ADAPT).

Rim vs the same area on the other side (PET)



Zazulia et al., J Cereb Blood Flow Metab 2001 (n = 19, 5–22 h after onset); Butcher et al., Stroke 2013.

04

At the bedside

Every stage is a window



Every stage is a window to act



The vessel

Before and after

Control blood pressure. Review antithrombotics.



The first hours

First 24 hours

Image fast. Reverse anticoagulants. Watch the GCS closely.



The days after

Days to weeks

Expect fluctuation. Escalate new drowsiness. Pace therapy.

If you remember three things

1

The vessel decides where it bleeds.

2

The first hours decide how big it gets.

3


The blood keeps injuring the brain for days.

EIGHT YEARS LATER

Jill walked onto the TED stage holding a human brain.

**ICH is a process,
not a moment.**

What would you like me to clarify?



SOURCES

References

1. Rossi C, Cordonnier C. Pathophysiology of non-traumatic intracerebral haemorrhage. In: Norrving B, ed. Oxford Textbook of Stroke and Cerebrovascular Disease. OUP; 2014: ch. 5.
2. Taylor JB. My stroke of insight. TED 2008; and My Stroke of Insight (book), 2006.
3. van Asch CJ, et al. Lancet Neurol 2010;9:167–176.
4. Meretoja A, et al. SMASH-U. Stroke 2012;43:2592–2597.
5. Cordonnier C, Al-Shahi Salman R, Wardlaw J. Brain 2007;130:1988–2003.
6. Zhu XL, Chan MS, Poon WS. Stroke 1997;28:1406–1409.
7. Fisher CM. J Neuropathol Exp Neurol 1971;30:536–550.
8. Brott T, et al. Stroke 1997;28:1–5.
9. Davis SM, et al. Neurology 2006;66:1175–1181.
10. Broderick JP, et al. Stroke 1993;24:987–993.
11. Hanley DF. Stroke 2009;40:1533–1538.
12. Gebel JM, et al. Stroke 2002;33:2631–2635.
13. Xi G, Keep RF, Hoff JT. Lancet Neurol 2006;5:53–63.
14. Zazulia AR, et al. J Cereb Blood Flow Metab 2001;21:804–810.
15. Butcher KS, et al. (ICH ADAPT). Stroke 2013;44:620–626.