

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/329231444>

# Solar Activities and Its Impact on Space Weather

Article in *Proceedings of the International Astronomical Union* · February 2018

DOI: 10.1017/S1743921318001606

CITATION

1

READS

305

4 authors:



**Drabindra Pandit**  
St. Xavier's College

23 PUBLICATIONS 8 CITATIONS

[SEE PROFILE](#)



**N. P. Chapagain**  
Tribhuvan University

129 PUBLICATIONS 593 CITATIONS

[SEE PROFILE](#)



**Binod Adhikari**  
St. Xavier's College

103 PUBLICATIONS 316 CITATIONS

[SEE PROFILE](#)



**Roshan Kumar Mishra**  
University of Maryland, Baltimore County

33 PUBLICATIONS 72 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Factors influencing aviation Visibility [View project](#)



Analysis of solar, interplanetary, and geomagnetic parameters during solar cycles 22, 23, and 24 [View project](#)

# Solar Activities and Its Impact on Space Weather

Drabindra Pandit<sup>1,2,3</sup>, Narayan P. Chapagain<sup>2</sup>, Binod Adhikari<sup>1,3</sup> and Roshan K. Mishra<sup>1</sup>

<sup>1</sup>Dept. of Physics, St. Xavier's College, Tribhuvan University  
Maitighar, Kathmandu, Nepal

<sup>2</sup>Dept. of Physics, Patan Multiple Campus, Tribhuvan University  
Patandhoka, Lalitpur, Nepal

<sup>3</sup>Central Dept. of Physics, Tribhuvan University  
Maitighar, Kathmandu, Nepal

email: drabindrapandit087@gmail.com, npchapagain@gmail.com, binod.adhi@gmail.com, roshanmishra.phy@gmail.com

**Abstract.** The Sun is an active star and its magnetic field fluctuates from a fraction of a second to a long period of time. The solar wind, CME, solar prominence, solar flares, solar particle and solar filament are the direct result of solar magnetic activity effects on the interplanetary space, Earth's magnetosphere and ionosphere. The intensity of irruption of these phenomena from the Sun's surface depends upon its phases. The extreme events affect technology both in space and on the ground. The data obtained from series of observations can help to predict solar activities and safekeeping to the space technology. In this study the cross correlations between IMF Bz, solar wind velocity ( $V_{sw}$ ) and interplanetary electric field ( $E_y$ ) with AE and SYM-H are studied. The results reveal that strong geomagnetic disturbances have high impact on the components of space weather than weak disturbances have.


**Keywords.** Interplanetary space, Earth's magnetosphere, Geomagnetic disturbances etc.

## 1. Introduction


The Sun has nearly 11 year of solar cycle which has maximum, minimum, ascending and descending phases. It is more active during maximum phases of solar cycle where it has maximum sunspots number and during minimum phases it is less active and has minimum number of sunspots. During active phases of the Sun tremendous amount of plasma and magnetic field are released into interplanetary space (Parker 1958), which interacts with the Earth's magnetosphere and a part of plasma particles and their kinetic energy enters into magnetosphere resulting into geomagnetic storm (Gonzalez *et al.*, 1987), substorm, auroral and Australian borealis in the northern and southern sky respectively that ultimately affect the technology borne in space or on the ground (Eastwood *et al.*, 2017). The parameters used to study their strength are solar wind parameters and geomagnetic indices (Adhikari *et al.*, 2017).

## 2. Result and Discussion

During intense magnetic storm the southward component of interplanetary magnetic field (IMF Bz) is strong and there occurs magnetic reconnection between this field and geomagnetic field, which produces open field lines, allowing mass, energy and momentum pervade into the Earth's magnetosphere. Such events affect more on intense storm day than during quiet day.



**Figure 1.** The panel from top to bottom show the variation of interplanetary magnetic field ( $B_z$  in nT), SYM-H (nT), solar wind speed ( $V_{sw}$  in Km/s), interplanetary electric field ( $E_y$  in mV/m), Aurora index (AE in nT) on 17 March 2015 intense storm and 16 January 2017 quiet day (left) and 22 June 2015 intense storm and 16 January 2017 quiet day (right) respectively.



**Figure 2.** Showing variation in cross-correlation coefficients versus time(min) for  $B_z$ (nT)-AE(nT),  $B_z$ (nT)-SYM-H(nT),  $V_{sw}$ (Km/s)-AE(nT),  $V_{sw}$ (Km/s)-SYM-H(nT),  $E_y$ (mV/m)-AE(nT),  $E_y$ (mV/m)-SYM-H(nT) on 17 March 2015 (left) and 22 June 2015 (right) respectively.

### 3. CONCLUSION

During geomagnetic storms, the interaction occurs between the Sun's and the Earth's magnetic field. Energy and charged particles are injected into the Earth's magnetosphere, which decrease the Earth's magnetic field and affect the ground and space-based technologies. It is also found that cross correlations between  $V_{sw}$ -AE and  $E_y$ -AE are positive and between  $E_y$ -SYM-H is negative whereas the cross correlations between  $B_z$ -AE,  $B_z$ -SYM-H and  $V_{sw}$ -SYM-H vary with time, taking positive and negative values.

### References

- Parker, E. N. 1958, *Astrophysical Journal*, 128, 664  
 Gonzalez, W. D. & Tsurutani, B. T. 1987, *Planet Space Sci.*, 1987, 35,1101  
 Eastwood, J. P., Biffis, E., Hapgood, M. A., Green, L., Bisi, M. M., Bentley, R. D., Wicks, R., Mckinnell, L. A., & Gibbs, M. and Burnett, C. 2017, *Risk Analysis*, 2  
 Adhikari, B., Mishra, R. K., Pandit, D., & Bhatteirai, B., Chapagain N. P. 2017, *JIST*, 22 (1), 45