

| Index Value Range                | Clinical Meaning & Accuracy                                                                  | Recommended Next Step                                                                                                                |
|----------------------------------|----------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------|
| <b>Under 0.90</b> Negative       | No antibodies detected. Highly reliable if tested <b>12+ weeks</b> post-exposure.            | No action needed.                                                                                                                    |
| <b>0.90 to 1.09</b> Equivocal    | Borderline result. The test cannot definitively determine a positive or negative status.     | Repeat the standard IgG test in <b>4 to 6 weeks</b> , or proceed to a confirmatory test.                                             |
| <b>1.10 to 3.50</b> Low Positive | Antibodies detected, but has a <b>significant false-positive rate</b> (up to 50% for HSV-2). | Seek a second confirmation via the <b>University of Washington Western Blot</b> or an inhibition assay before accepting a diagnosis. |
| <b>Above 3.50</b> High Positive  | Highly reliable positive result. False positives are rare at this level.                     | Confirmed infection. Speak to your provider about management or daily suppression if you experience symptoms.                        |

### How can I tell the difference between detox flare-ups and a medical emergency?

Detox reactions are generally mild, familiar, and transient; emergencies feature severe, sudden pain, breathing failure, or rapidly deteriorating vital signs. In the event of a medical emergency, stop the protocol and **immediately** contact your local emergency services or go to the nearest emergency room.

### Can I participate this protocol while pregnant or taking other medications?

Do not participate in this protocol if you are pregnant, breastfeeding, or taking blood thinners, diabetes medications, lithium or diuretics. Consult your doctor or other medical professional before participating.

### Are there other supplements that I can add?

Yes! You can aid the process by adding super strength olive leaf extract, lemon balm, elderberry, and echinacea to your routine. These herbs pair exceptionally well with the oil of oregano.