

The Whole Food Guide

To prepare, you'll be focusing on opening your detox pathways to aid the the next phases in advance. Your diet will consist of foods that improve bile flow as follows:

Food Categories & Examples

- **Cruciferous Vegetables (for Liver Support):** Broccoli, sprouts, kale, cauliflower, and cabbage contain glucosinolates (produces sulforaphane) that support detoxification enzymes.
- **Alliums (as Sulfur Boosters):** Raw garlic, onions, leeks, and chives provide sulfur, which is essential for liver detoxification.
- **Bitter Greens & Herbs (for Gallbladder/Liver):** Dandelion greens, beet greens, artichokes, arugula, parsley, and cilantro improve bile production, aiding in the digestion and elimination of fat-soluble toxins.
- **Fiber-Rich Foods (for Intestinal Binding):** Chia seeds, flaxseeds, legumes, raisins, oats, quinoa, brown rice, and root vegetables bind to toxins in the gut, ensuring they are excreted.
- **Amino acids:** Eggs, red meat, low mercury fish (e.g. wild cod, wild salmon), raw dairy, raw dairy kefir to evacuate Phase One toxic load. Wild caught salmon, sardines, anchovies, herring, and Atlantic mackerel are all great fatty fish that are low in mercury. For those who are strictly plant-based, you can have high amino acid protein substitutes like lentils, edamame, green peas, asparagus, brussel sprouts, spinach, quinoa, amaranth, and chickpeas.
- **Hydrating Foods (for Kidney Function):** Cucumbers, watermelons, and citrus fruits support renal filtration.
- **Nuts & Seeds (for Liver Support):** Hemp seeds, pumpkin seeds, walnuts, almonds (2–3 single almonds per day), pecans, and brazil nuts to supply healthy Omega 3 and Omega 6 fat ratio. Do not exceed 1/4 cup of nuts per day. Aim for a 4:1 ratio of omega 3 to omega 6. Too much omega 6 can be inflammatory.
- **Healthy Fats (for Cellular Function):** Avocados, unrefined coconut oil, extra virgin olive oil, and grass-fed ghee, butter, or tallow. The liver loves ghee and coconut oil. Avoid seed oils.
- **Glutathione-Producing Fruits:** Berries (strawberries, raspberries, blackberries, blueberries, black grapes or black raisins), citrus fruits (lemons, limes, oranges, grapefruits), avocados, apples, pomegranates. Berries are to be introduced only when Phase One begins for anthocyanins to support cytochrome P450 enzymes. Seeded fruits only.