

FIN3380 Final Review

Dec 9, 2019

dongliyaun@cuhk.edu.cn


zhaidaojing@cuhk.edu.cn

Lecture 3 Risk Management

- VaR
 - Gaussian VaR, historical VaR: steps to calculate VaR
 - Property for Gaussian VaR: subadditivity
- Expected Shortfall
 - Gaussian ES, historical ES: steps to calculate ES
- Drawdown
 - How much an investment or trading account is down from the peak before it recovers back to the peak
 - Calculate drawdown: `cummax()`
 - Find out the longest dd period and the maximum dd

Lecture 3 Risk Management

- Co-drawdown
 - The probability that two assets simultaneously experience drawdowns taking into account of the size of drawdown
 - Steps to calculate co-drawdown



Lecture 4 Factor Model

- Simple excess return model
- CAPM model
 - Formula: $r_i = r_f + \beta_i(r_m - r_f)$,
- Multi-Factor model – FF 3 Factor model
 - Formula: $r_i = r_f + \beta_i(r_m - r_f) + h_i HML + s_i SMB$

Lecture 4 Factor Model

- Regression
 - Simple linear regression
 - what is alpha and beta
 - t statistic and p value
 - Make prediction
 - Rolling window regression
 - What is rolling window
 - Implement: define a rolling function - `tibbletime::rollify()`
 - Summarize the regression results - `broom::tidy()`; `purrr::map()` and `tidyr::unnest()`
 - Visualize regression results

Lecture 5 Times Series

- Stationary
 - Definition of weak stationary
 - How to test whether a time series is stationary
 - How to deal with the non-stationary series
- ACF
 - Calculation: $\hat{\rho}_k = \frac{\sum_{t=k+1}^T (x_t - \bar{x})(x_{t-k} - \bar{x})}{\sum_{t=1}^T (x_t - \bar{x})^2}, \quad 0 \leq k < T - 1.$
 - Test
 - Test individual ACF
 - Joint test: `Box.test()`

Lecture 5 Times Series

- PACF

$$x_t = \phi_{0,1} + \phi_{1,1}x_{t-1} + e_{1t},$$

$$x_t = \phi_{0,2} + \phi_{1,2}x_{t-1} + \phi_{2,2}x_{t-2} + e_{2t},$$

$$x_t = \phi_{0,3} + \phi_{1,3}x_{t-1} + \phi_{2,3}x_{t-2} + \phi_{3,3}x_{t-3} + e_{3t},$$

$$x_t = \phi_{0,4} + \phi_{1,4}x_{t-1} + \phi_{2,4}x_{t-2} + \phi_{3,4}x_{t-3} + \phi_{4,4}x_{t-4} + e_{4t},$$

\vdots

- Estimate the parameters in the above equations to get the PACF
- For an AR(p) model, the lag-p sample PACF should not be 0, but the PACF of lag-j with $j > p$ should be 0.

Lecture 5 Times Series

- AR(p) model
 - AR(p): $x_t = \phi_0 + \phi_1 x_{t-1} + \cdots + \phi_p x_{t-p} + a_t$,
 - Detect lag order p
 - Use the PACF
 - Use information criteria: AIC, BIC
 - The long term average of a stationary AR(p) model: $E(x_t) = \frac{\phi_0}{1 - \phi_1 - \cdots - \phi_p}$
 - Forecast
 - Make one step/ multiple steps prediction

Lecture 5 Times Series

- MA(q) model
 - MA(q): $x_t = c_0 + a_t - \theta_1 a_{t-1} - \dots - \theta_q a_{t-q}$
 - Detect lag order q
 - Use the ACF
 - Use information criteria: AIC, BIC
 - The long term average of a stationary MA(q) model: $E(x_t) = c_0$
 - Forecast
 - Make one step/ multiple steps prediction

Lecture 5 Times Series

- ARIMA(p,d,q) model

- ARIMA(p,d,q): $y'_t = c + \phi_1 y'_{t-1} + \cdots + \phi_p y'_{t-p} + \theta_1 \varepsilon_{t-1} + \cdots + \theta_q \varepsilon_{t-q} + \varepsilon_t,$

- Detect lag orders

- Use `auto.arima()`

- Use information criteria: AIC, BIC

- Forecast

- Make one step/ multiple steps prediction

Lecture 5 Times Series

– Co-integration

- Definition: two time series is cointegrated if their linear difference is stationary.

$$u_t = Y_t - (\alpha + \beta X_t)$$

$$u_t : \sim I(0), \text{stationary process}$$

$$X_t, Y_t : \sim I(1)$$

– Construct Pair trading

- S1: test whether the log price of each pair is cointegrated – Johansen test (urca::ca.jo)
- S2: Calculate the spread of log prices of paired stocks and test whether the spread is stationary
- S3: Examine whether the spread is too high or too low and generate 'buy' signals.
- S4: calculate return on each day and the cumulative returns