



# A guide to deciphering microbial interactions and metabolic fluxes in microbiome communities

Maciek R Antoniewicz

Microbiomes occupy nearly all environments on Earth. These communities of interacting microorganisms are highly complex, dynamic biological systems that impact and reshape the molecular composition of their habitats by performing complex biochemical transformations. The structure and function of microbiomes are influenced by local environmental stimuli and spatiotemporal changes. In order to control the dynamics and ultimately the function of microbiomes, we need to develop a mechanistic and quantitative understanding of the ecological, molecular, and evolutionary driving forces that govern these systems. Here, we describe recent advances in developing computational and experimental approaches that can promote a more fundamental understanding of microbial communities through comprehensive model-based analysis of heterogeneous data types across multiple scales, from intracellular metabolism, to metabolite cross-feeding interactions, to the emergent macroscopic behaviors. Ultimately, harnessing the full potential of microbiomes for practical applications will require developing new predictive modeling approaches and better tools to manipulate microbiome interactions.

## Address

Department of Chemical Engineering, Metabolic Engineering and Systems Biology Laboratory, University of Michigan, Ann Arbor, MI 48109, USA

Corresponding author: Antoniewicz, Maciek R ([mranton@umich.edu](mailto:mranton@umich.edu))

**Current Opinion in Biotechnology** 2020, **64**:230–237

This review comes from a themed issue on **Analytical biotechnology**

Edited by **Yinjie Tang** and **Ludmilla Aristilde**

<https://doi.org/10.1016/j.copbio.2020.07.001>

0958-1669/© 2020 Elsevier Ltd. All rights reserved.


## Introduction

Microbial communities are ubiquitous in nature and their impact on Earth's ecosystems is pervasive, from biogeochemical cycles to agriculture and human health [1–3]. Studying microbial communities and their impact on natural environments is therefore essential to understand living systems [4,5]. Moreover, microbial communities have a wide range of practical applications, including in

biodegradation of organic chemicals [6], conversion of toxic compounds [7,8], and production of biofuels and other products from single or multiple substrates [9]. Microbial consortia carry out efficient bio-transformations that result from multiple complementary metabolic systems working together [10,11]. The capabilities of multi-microorganism systems often cannot be predicted by the sum of their parts [12]. Rather, complex interactions at multiple levels result in improved overall performance of these systems [7]. The emerging field of microbiome synthetic and systems biology promises the assembly of complementary metabolic pathways into functional systems [13–17,18], where the diversity of metabolic capabilities and the ability to exchange metabolites dramatically increases the possible metabolic space [17,19]. This alleviates limitations encountered in single-organism systems [14,20]. For example, integration of complementary metabolic systems can result in more efficient substrate conversions [7], make communities more robust to fluctuations [16], and accomplish transformations that would be thermodynamically infeasible with single-organism systems [21,22].

Given the prevalence and importance of microbiomes in nature, much effort has been invested in the past decade to unravel the members, structures, functions, interactions, and governing principles of microbial communities [23] (Figure 1). With the advent of next-generation sequencing, metagenomics first emerged as an important tool in the study of microbial communities. It allowed quantitative analysis of the diversity, composition, and dynamics of these systems [24]. Statistical modeling approaches, such as multiple linear regression and multi-dimensional cluster analysis, were then used to interpret the wealth of metagenomics data to identify microbial community trends and correlations between metagenomics data and other observable system variables [25–29]. These statistical modeling approaches provided new hypotheses regarding potential interactions and functions of communities. However, metagenomics data alone could not predict causal relationships in microbial communities. For this, new computational models were needed that allowed a more systematic and rigorous interpretation and interrogation of the huge amount of heterogeneous data that was generated [30,31]. Once a model was constructed, systems biological properties could be analyzed by comparing model simulations with experimentally observed data. Mathematical models of microbial communities thus provided critical tools for generating and testing biological hypotheses to better

Figure 1



The two primary approaches used to model microbial communities are ecological models and metabolic models. Ecological models focus on describing population dynamics and evaluating the stability of communities under various conditions or perturbations. Metabolic models focus on cellular metabolism and explicit cross-feeding interactions to predict population properties of microbiomes. Recent advances in  $^{13}\text{C}$ -metabolic flux analysis also allow intracellular metabolic fluxes and metabolite cross-feeding rates to be measured experimentally in co-cultures.


understand and predict the dynamics and interactions among community members [32] (Figure 2). Looking forward, mathematical models will play an increasingly important role in identifying new targets for the design and optimization of synthetic microbial communities for practical applications in biotechnology, agriculture and medicine [33]. In this review, we provide an overview of the wide range of computational and experimental techniques that have been developed to analyze microbial communities and highlight recent advances that can further enhance our understanding of these complex biological systems.

### Ecological models describe population dynamics in communities

The two primary approaches that have been applied to model microbial communities are ecological models and

metabolic (e.g. genome-scale) models. Ecological models focus on species abundances and how they change over time given certain relationships between the species. In these models, species abundances are the only system variables, and thus all interactions between community members must be described by suitable mathematical functions to capture the wide range of possible interactions, including mutualistic, neutral, and competitive interactions. In practice, ecological models are often set up as a set of differential equations that can be studied as a dynamic system. The advantage of ecological models is that they can directly use available time-series species abundance data to quantify various properties of communities, such as the stability of biological systems under various conditions and perturbations. The best known ecological model is the generalized Lotka-Volterra model (gLTV, also known as predator-prey model), first proposed

Figure 2



Mathematical models of microbial communities provide critical tools for generating and testing biological hypotheses. New computational approaches and software tools are needed that can promote fundamental understanding of microbial communities through comprehensive model-based analysis of omics data sets across multiple scales, from intracellular metabolism, to metabolite cross-feeding interactions between cells, to the emergent behaviors, structures and functions of microbial communities.

in 1920s. In this model, the dynamics of populations of any number of species  $N$  ( $X_1, X_2, \dots, X_N$ ) is described by the following set of ordinary differential equations:

$$\frac{dX_i}{dt} = X_i \cdot \left( \mu_i + \sum A_{i,j} \cdot X_j \right)$$

Here,  $\mu_i$  is the growth rate of species  $i$ , and  $\mathbf{A} = [A_{i,j}]$  is the 'community interactions matrix', where the fields represent interactions between species, such that positive  $A_{i,j}$  values capture positive interactions (e.g. mutualism), while negative values represent negative interactions (e.g. competition). By integrating the gLV equations, temporal dynamics of populations can be simulated under various interaction assumptions, or perturbations. Inversely, by fitting time-series data of species abundances (e.g. time-resolved metagenomics data) to the gLV model, interaction parameters can be estimated [34]. In the past decade, important systems level properties of microbial communities have been investigated using gLV models such as microbiome stability, resistance to perturbations, and the speed at which systems can reach steady states [30,32,35]. While gLV models have proven their value in explaining global systems properties of communities, they also have important limitations [36]. A notable disadvantage of ecological models is that only certain simplified aggregate interactions between organisms are possible. These interactions

are often static and do not relate directly to cellular characteristics such as the metabolic state of cells. Since population abundances are the only system variables in these models, all interactions must be described as a function of these variables. As a result, ecological models typically cannot capture cross-feeding interactions of nutrients between species and provide no information regarding metabolism within each species. Moreover, biochemical production of products and community responses to various substrates are not easily captured by ecological models.

### Ecological models of syntrophic co-cultures

Cross-feeding of nutrients is a key characteristic of microbial communities, whereby certain members of the community feed off the products of others, and vice versa [10,11,37]. To better understand the interactions in microbial communities, and in the future to allow rational engineering of microbiomes, it is critical to elucidate syntrophic phenotypes [12,38\*,39,40\*\*,41]. For this purpose, auxotrophic *Escherichia coli* strains provide useful model systems to study syntrophic interactions. In a pioneering study, Wintermute *et al.* [12] evaluated 1035 co-cultures of 46 conditionally lethal auxotrophic *E. coli* strains. While none of the 46 strains could grow in minimal medium with glucose as the sole substrate, a surprisingly large fraction (17%) of the paired co-cultures were able to grow on glucose. It was hypothesized that the *E. coli* mutants complemented one another's growth by

cross-feeding essential metabolites, possibly amino acids, although this was not experimentally verified. To describe the observed dynamics a simple empirical ecological model was used to capture the essential features of these co-cultures:

$$\frac{dX_1}{dt} = C_{1,2} \cdot \left( \frac{X_2}{X_1 + X_2} \right) \cdot \left( 1 - \frac{X_1 + X_2}{k} \right)$$

$$\frac{dX_2}{dt} = C_{2,1} \cdot \left( \frac{X_1}{X_1 + X_2} \right) \cdot \left( 1 - \frac{X_1 + X_2}{k} \right)$$

Here,  $X_1$  and  $X_2$  represent the respective strain abundances,  $C_{1,2}$  and  $C_{2,1}$  quantify the synergistic cooperation on the part of the two strains, and  $k$  is a logistic carrying capacity of the batch culture related to the initial glucose concentration and overall biomass yield. With this model, the population dynamics and final co-culture compositions could be predicted. Later, Mee *et al.* [42] investigated a smaller set of 14 amino acid auxotrophic *E. coli* strains in dual and triple co-cultures. To quantitatively model the co-cultures, slightly revised model equations were used that better captured the observed dynamics according to the authors:

$$\frac{dX_1}{dt} = C_{1,2} \cdot X_2 \cdot \left( \frac{X_1}{X_1 + b} \right) \cdot \left( 1 - \frac{X_1 + X_2}{k} \right)$$

$$\frac{dX_2}{dt} = C_{2,1} \cdot X_1 \cdot \left( \frac{X_2}{X_2 + b} \right) \cdot \left( 1 - \frac{X_1 + X_2}{k} \right)$$

These equations were then generalized to describe triple co-cultures, and eventually to model a complex 14-member consortium. Through these investigations, the authors identified that biosynthetically costly amino acids tended to promote stronger cooperative interactions, and more generally, that amino acid auxotrophies could be an evolutionarily optimizing strategy that reduces biosynthetic burden while promoting cooperative interactions between bacteria in microbiomes [42,43].

### Genome-scale metabolic models of microbial communities

In contrast to ecological models, genome-scale metabolic models focus on cellular metabolism and explicit cross-feeding interactions to predict population properties of microbiomes [44,45]. These models take advantage of the extensive knowledge accumulated over the past two decades regarding metabolic reactions from whole-genome sequencing. In these models, interactions between species are specified at the molecular level by allowing cross-feeding of specific metabolites between community


members [46\*\*]. The central mathematical framework for analyzing microbial communities are constraint-based modeling and optimization techniques such as flux balance analysis (FBA) [47], in which a given cellular or community objective is optimized to estimate intracellular metabolic fluxes and cross-feeding rates. Various objective functions have been proposed to model microbial communities [48]. For example, microbial communities have been modeled as a single superorganism with one overall objective function [49,50], or as a set of interacting species each with their own objective functions [51–53].

Based on this framework, various extended modeling approaches have been developed for modeling microbial communities, including the dynamic multispecies metabolic modeling framework (DMMM) [54], the community and systems-level interactive optimization (CASINO) approach [55], the computation of microbial ecosystems in time and space (COMETS) framework [56], OptCom [57], and d-OptCom [58]. While these models can predict certain aspects of microbial communities, important challenges remain that must be addressed in the future. Perhaps the most significant concern with these approaches is their inherent reliance on a specific optimality assumption to predict global system behavior and specific metabolic fluxes and cross-feeding rates. Since different modeling frameworks use different optimality criteria, model predictions become highly dependent on the specific set of assumptions used. Optimality assumptions are notoriously difficult to verify experimentally, even for single-organism systems [59,60], let alone for complex microbial communities. Moreover, metabolic flux predictions based on constraint based approaches can be inaccurate, as they often disagree with experimentally measured fluxes using techniques such as  $^{13}\text{C}$  metabolic flux analysis ( $^{13}\text{C}$ -MFA), as was shown recently for *E. coli* [61]. To improve quantitative accuracy of these models it would be valuable to incorporate kinetic information, which can significantly improve metabolic flux predictions [62].

### $^{13}\text{C}$ -Metabolic flux analysis for analysis of microbial communities

$^{13}\text{C}$ -MFA is the gold standard technique for quantifying metabolic fluxes in living cells. In the past decade,  $^{13}\text{C}$ -MFA has become an indispensable tool in metabolic engineering and synthetic biology [63,64]. In contrast to FBA,  $^{13}\text{C}$ -MFA does not rely on optimality assumptions to elucidate metabolism. Instead, precise metabolic fluxes are determined through model-based analysis of stable-isotope labeling measurements using well established protocols and modeling approaches [65\*,66,67]. Thus far, only a few studies have focused on measuring metabolism in microbial communities using  $^{13}\text{C}$ -MFA. In these studies, physical separation of either proteins or cells was required to measure species-specific isotopic labeling, from which then species-specific fluxes were

Figure 3



Microorganisms can engage in a wide range of possible interactions, including parasitism, commensalism, neutralism, amensalism, competition and mutualism. Even for a simple two-member system there is a large number of positive, neutral and negative interactions possible based on metabolite exchanges and substrate competition.

calculated [68,69]. Separation was accomplished either by direct separation of cells via centrifugation or fluorescence-assisted sorting, or indirect separation through purification of an overexpressed 'reporter protein', where the labeling of the overexpressed reporter protein was used as a proxy for whole-cell protein labeling. However, these approaches for  $^{13}\text{C}$ -MFA have a few important limitations. Notably, incomplete separation of cells or proteins was shown to produce inaccurate results [68]. Additionally, fluorescence-assisted cell sorting is a rather slow separation technique, thus making it impractical for routine use in  $^{13}\text{C}$ -MFA, while methods based on an overexpressed 'reporter protein' are limited to organisms with well-developed genetic tools. Moreover, these approaches require large sample sizes. Lastly, none of these approaches was able to quantify metabolite cross-feeding fluxes that are of significant interest.

Recently, a new  $^{13}\text{C}$ -MFA framework for microbial communities was developed that overcomes previous limitations [70]. This approach doesn't require any physical separation of cells or proteins. Instead, isotopic labeling of the entire sample can be used to resolve species-specific

fluxes with high precision and determine inter-species metabolite exchange. This approach for conducting  $^{13}\text{C}$ -MFA is based on the idea that different populations in a community will have distinct isotopic labeling patterns and that the measured labeling can be represented as an average population-weighted labeling state. The advantage of this approach is that isotopic labeling of different populations can be computationally deconvoluted from total labeling and species-specific metabolic fluxes and cross-feeding rates between the populations can be quantified with high precision [70]. This approach was recently applied to quantify intracellular fluxes and metabolite cross-feeding interactions between distinct microbial subpopulations in a biofilm [71•]. In the future, by building upon this framework and developing additional tools, for example, for dynamic  $^{13}\text{C}$ -MFA and incorporating additional multi-omics data, we envision that understanding of the general principles that govern the structure and function of microbial communities can be further enhanced.

## Conclusions and future outlook

One of the key challenges in microbiome research is identifying and manipulating cross-feeding interactions



between community members that drive system dynamics and functions [72,73,74\*]. Even for a simple two-member system there are a large number of possible interactions [75\*\*] (Figure 3). To better understand microbiomes and make quantitative predictions, more advanced methods are needed for identifying and experimentally validating these cross-feeding interactions [39,76]. Moreover, to further advance our understanding of the role of metabolism in microbial communities, new mathematical approaches are needed that can span multiple length scales and time scales to enable integrated multiscale analysis of metabolite cross-feeding interactions and community metabolic fluxes [77,78,79\*,80\*]. With respect to population models, the scope of these models needs to be expanded to allow multiple heterogeneous data sets to be integrated for systems biology analysis of microbial communities. Cross-feeding interactions and other substrate-level effects should be directly incorporated into these new models. The basis for these models could kinetic growth models that have been used extensively in the past to simulate and optimize large-scale industrial fermentations. These predictive models are based on mass balances for all relevant system components (i.e. biomass, substrate, product, oxygen), combined with kinetic expressions for cell growth, product formation and gas-liquid mass transfer. For modeling microbial communities, the kinetic growth rate expressions of individual species must be carefully chosen, since in microbial communities cell growth is often directly linked to cross-feeding rates, rather than to substrate concentrations as in traditional Monod kinetic growth models. To further advance the use of quantitative modeling and analysis techniques for interrogating microbial communities [81,82], new software tools are also needed that can seamlessly integrate the various modeling strategies and analysis functions, so that these approaches are available to a broader cross-section of the scientific community. Ultimately, we envision that integrated models of microbiomes will enhance our fundamental understanding of microbial communities and establish new theories and mechanisms that govern the structure and function of these complex biological systems.

### Conflict of interest statement

Nothing declared.

### Acknowledgement

This work was supported by grant NSFMCB-1616332.

### References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Martiny JB, Bohannan BJ, Brown JH, Colwell RK, Fuhrman JA, Green JL, Horner-Devine MC, Kane M, Krumins JA, Kuske CR et al.: **Microbial biogeography: putting microorganisms on the map.** *Nat Rev Microbiol* 2006, **4**:102-112.
2. Falkowski PG, Fenchel T, Delong EF: **The microbial engines that drive Earth's biogeochemical cycles.** *Science* 2008, **320**:1034-1039.
3. Cho I, Blaser MJ: **The human microbiome: at the interface of health and disease.** *Nat Rev Genet* 2012, **13**:260-270.
4. Fuhrman JA: **Microbial community structure and its functional implications.** *Nature* 2009, **459**:193-199.
5. Fuhrman JA, Cram JA, Needham DM: **Marine microbial community dynamics and their ecological interpretation.** *Nat Rev Microbiol* 2015, **13**:133-146.
6. Hays SG, Patrick WG, Ziesack M, Oxman N, Silver PA: **Better together: engineering and application of microbial symbioses.** *Curr Opin Biotechnol* 2015, **36**:40-49.
7. Sabra W, Dietz D, Tjahjajari D, Zheng AP: **Biosystems analysis and engineering of microbial consortia for industrial biotechnology.** *Eng Life Sci* 2010, **10**:407-421.
8. Harishankar MK, Sasikala C, Ramya M: **Efficiency of the intestinal bacteria in the degradation of the toxic pesticide, chlorpyrifos.** *3 Biotech* (3):2013:137-142.
9. Sgobba E, Wendisch VF: **Synthetic microbial consortia for small molecule production.** *Curr Opin Biotechnol* 2020, **62**:72-79
- This paper reviews recent progress in applying synthetic biology approaches for engineering microbial consortia for industrial applications.
10. Bader J, Mast-Gerlach E, Popovic MK, Bajpai R, Stahl U: **Relevance of microbial coculture fermentations in biotechnology.** *J Appl Microbiol* 2010, **109**:371-387.
11. Shong J, Jimenez Diaz MR, Collins CH: **Towards synthetic microbial consortia for bioprocessing.** *Curr Opin Biotechnol* 2012, **23**:798-802.
12. Wintermute EH, Silver PA: **Emergent cooperation in microbial metabolism.** *Mol Syst Biol* 2010, **6**:407.
13. Zhang H, Pereira B, Li Z, Stephanopoulos G: **Engineering *Escherichia coli* coculture systems for the production of biochemical products.** *Proc Natl Acad Sci U S A* 2015, **112**:8266-8271.
14. Zhou K, Qiao K, Edgar S, Stephanopoulos G: **Distributing a metabolic pathway among a microbial consortium enhances production of natural products.** *Nat Biotechnol* 2015, **33**:377-383.
15. Sheth RU, Cabral V, Chen SP, Wang HH: **Manipulating Bacterial communities by in situ microbiome engineering.** *Trends Genet* 2016, **32**:189-200.
16. Brenner K, You L, Arnold FH: **Engineering microbial consortia: a new frontier in synthetic biology.** *Trends Biotechnol* 2008, **26**:483-489.
17. McCarty NS, Ledesma-Amaro R: **Synthetic biology tools to engineer microbial communities for biotechnology.** *Trends Biotechnol* 2019, **37**:181-197.
18. Roell GW, Zha J, Carr RR, Koffas MA, Fong SS, Tang YJ: **Engineering microbial consortia by division of labor.** *Microb Cell Fact* 2019, **18**:35
- Another good review article that focuses on new engineering strategies for microbial consortia through the concept of division of labor as a means to address metabolic burden.
19. Kato S, Haruta S, Cui ZJ, Ishii M, Igarashi Y: **Stable coexistence of five bacterial strains as a cellulose-degrading community.** *Appl Environ Microbiol* 2005, **71**:7099-7106.
20. Tsoi R, Wu F, Zhang C, Bewick S, Karig D, You L: **Metabolic division of labor in microbial systems.** *Proc Natl Acad Sci U S A* 2018, **115**:2526-2531
- This paper explores the trade-off between division of labor in microbial consortia and the introduction of additional transport limitations for metabolites and enzymes. The authors propose general criteria when microbial consortia are expected to outperform a single population.

21. Zhang H, Wang X: **Modular co-culture engineering, a new approach for metabolic engineering.** *Metab Eng* 2016, **37**:114-121.
  22. Jones JA, Vernacchio VR, Collins SM, Shirke AN, Xiu Y, Englaender JA, Cress BF, McCutcheon CC, Linhardt RJ, Gross RA et al.: **Complete biosynthesis of anthocyanins using *E. coli* polycultures.** *mBio* 2017, **8**.
  23. Friedman J, Higgins LM, Gore J: **Community structure follows simple assembly rules in microbial microcosms.** *Nat Ecol Evol* 2017, **1**:109.
  24. Wooley JC, Godzik A, Friedberg I: **A primer on metagenomics.** *PLoS Comput Biol* 2010, **6**:e1000667.
  25. Hamady M, Knight R: **Microbial community profiling for human microbiome projects: Tools, techniques, and challenges.** *Genome Res* 2009, **19**:1141-1152.
  26. White JR, Nagarajan N, Pop M: **Statistical methods for detecting differentially abundant features in clinical metagenomic samples.** *PLoS Comput Biol* 2009, **5**:e1000352.
  27. Flint HJ, Scott KP, Louis P, Duncan SH: **The role of the gut microbiota in nutrition and health.** *Nat Rev Gastroenterol Hepatol* 2012, **9**:577-589.
  28. Arumugam M, Raes J, Pelletier E, Le Paslier D, Yamada T, Mende DR, Fernandes GR, Tap J, Bruls T, Batto JM et al.: **Enterotypes of the human gut microbiome.** *Nature* 2011, **473**:174-180.
  29. Faith JJ, McNulty NP, Rey FE, Gordon JL: **Predicting a human gut microbiota's response to diet in gnotobiotic mice.** *Science* 2011, **333**:101-104.
  30. Faust K, Raes J: **Microbial interactions: from networks to models.** *Nat Rev Microbiol* 2012, **10**:538-550.
  31. Zuniga C, Zaramela L, Zengler K: **Elucidation of complexity and prediction of interactions in microbial communities.** *Microb Biotechnol* 2017, **10**:1500-1522.
  32. Coyte KZ, Schluter J, Foster KR: **The ecology of the microbiome: networks, competition, and stability.** *Science* 2015, **350**:663-666.
  33. Lindemann SR, Bernstein HC, Song HS, Fredrickson JK, Fields MW, Shou W, Johnson DR, Beliaev AS: **Engineering microbial consortia for controllable outputs.** *ISME J* 2016, **10**:2077-2084.
  34. Fisher CK, Mehta P: **Identifying keystone species in the human gut microbiome from metagenomic timeseries using sparse linear regression.** *PLoS One* 2014, **9**:e102451.
  35. Stein RR, Bucci V, Toussaint NC, Buffie CG, Ratsch G, Pamer EG, Sander C, Xavier JB: **Ecological modeling from time-series inference: insight into dynamics and stability of intestinal microbiota.** *PLoS Comput Biol* 2013, **9**:e1003388.
  36. Momeni B, Xie L, Shou W: **Lotka-Volterra pairwise modeling fails to capture diverse pairwise microbial interactions.** *eLife* 2017, **6**.
  37. D'Souza G, Shitut S, Preussger D, Yousif G, Waschina S, Kost C: **Ecology and evolution of metabolic cross-feeding interactions in bacteria.** *Nat Prod Rep* 2018, **35**:455-488.
  38. Long CP, Antoniewicz MR: **How adaptive evolution reshapes metabolism to improve fitness: recent advances and future outlook.** *Curr Opin Chem Eng* 2018, **22**:209-215.
- This paper discusses the interplay between metabolism and evolution. Understanding the molecular and evolutionary driving in microbial communities will be important in the future to steer the function of microbial communities in desired directions.
39. LaSarre B, McCully AL, Lennon JT, McKinlay JB: **Microbial mutualism dynamics governed by dose-dependent toxicity of cross-fed nutrients.** *ISME J* 2017, **11**:337-348.
  40. Lv Y, Qian S, Du G, Chen J, Zhou J, Xu P: **Coupling feedback genetic circuits with growth phenotype for dynamic population control and intelligent bioproduction.** *Metab Eng* 2019, **54**:109-116

This paper discusses emerging engineering strategies based on feed-back genetic circuits to control population dynamics, eliminate cheaters, and enhance microbial cooperation. Applying these approaches could improve the cost-competitiveness of industrial fermentations.

41. Thommes M, Wang T, Zhao Q, Paschalidis IC, Segre D: **Designing metabolic division of labor in microbial communities.** *mSystems* 2019, **4**.
  42. Mee MT, Collins JJ, Church GM, Wang HH: **Syntrophic exchange in synthetic microbial communities.** *Proc Natl Acad Sci U S A* 2014, **111**:E2149-2156.
  43. Germerodt S, Bohl K, Luck A, Pande S, Schroter A, Kaleta C, Schuster S, Kost C: **Pervasive selection for cooperative cross-feeding in bacterial communities.** *PLoS Comput Biol* 2016, **12**:e1004986.
  44. Biggs MB, Medlock GL, Kolling GL, Papin JA: **Metabolic network modeling of microbial communities.** *Wiley Interdiscip Rev Syst Biol Med* 2015, **7**:317-334.
  45. Stolyar S, Van Dien S, Hillesland KL, Pintel N, Lie TJ, Leigh JA, Stahl DA: **Metabolic modeling of a mutualistic microbial community.** *Mol Syst Biol* 2007, **3**:92.
  46. Machado D, Andrejev S, Tramontano M, Patil KR: **Fast automated reconstruction of genome-scale metabolic models for microbial species and communities.** *Nucleic Acids Res* 2018, **46**:7542-7553.
- In this work CarveMe is introduced, an automated open-source tool for reconstruction of species and community level metabolic models. A database of thousands of bacterial models allows fast generation of microbial community models.
47. Orth JD, Thiele I, Palsson BO: **What is flux balance analysis?** *Nat Biotechnol* 2010, **28**:245-248.
  48. Bauer E, Zimmermann J, Baldini F, Thiele I, Kaleta C: **BacArena: individual-based metabolic modeling of heterogeneous microbes in complex communities.** *PLoS Comput Biol* 2017, **13**:e1005544.
  49. Greenblum S, Turnbaugh PJ, Borenstein E: **Metagenomic systems biology of the human gut microbiome reveals topological shifts associated with obesity and inflammatory bowel disease.** *Proc Natl Acad Sci U S A* 2012, **109**:594-599.
  50. Tobalina L, Bargiela R, Pey J, Herbst FA, Lores I, Rojo D, Barbas C, Pelaez AI, Sanchez J, von Bergen M et al.: **Context-specific metabolic network reconstruction of a naphthalene-degrading bacterial community guided by metaproteomic data.** *Bioinformatics* 2015, **31**:1771-1779.
  51. Heinken A, Thiele I: **Systematic prediction of health-relevant human-microbial co-metabolism through a computational framework.** *Gut Microbes* 2015, **6**:120-130.
  52. Heinken A, Thiele I: **Anoxic conditions promote species-specific mutualism between gut microbes in silico.** *Appl Environ Microbiol* 2015, **81**:4049-4061.
  53. Shoaie S, Karlsson F, Mardinoglu A, Nookaew I, Bordel S, Nielsen J: **Understanding the interactions between bacteria in the human gut through metabolic modeling.** *Sci Rep* 2013, **3**:2532.
  54. Zhuang K, Izallalen M, Mouser P, Richter H, Risso C, Mahadevan R, Lovley DR: **Genome-scale dynamic modeling of the competition between *Rhodospirillum rubrum* and *Geobacter* in anoxic subsurface environments.** *ISME J* 2011, **5**:305-316.
  55. Shoaie S, Ghaffari P, Kovatcheva-Datchary P, Mardinoglu A, Sen P, Pujos-Guillot E, de Wouters T, Juste C, Rizkalla S, Chilloux J et al.: **Quantifying diet-induced metabolic changes of the human gut microbiome.** *Cell Metab* 2015, **22**:320-331.
  56. Harcombe WR, Riehl WJ, Dukovski I, Granger BR, Betts A, Lang AH, Bonilla G, Kar A, Leiby N, Mehta P et al.: **Metabolic resource allocation in individual microbes determines ecosystem interactions and spatial dynamics.** *Cell Rep* 2014, **7**:1104-1115.
  57. Zomorodi AR, Maranas CD: **OptCom: a multi-level optimization framework for the metabolic modeling and analysis of microbial communities.** *PLoS Comput Biol* 2012, **8**:e1002363.

58. Zomorodi AR, Islam MM, Maranas CD: **d-OptCom: dynamic multi-level and multi-objective metabolic modeling of microbial communities**. *ACS Synth Biol* 2014, **3**:247-257.
59. Schuetz R, Kuepfer L, Sauer U: **Systematic evaluation of objective functions for predicting intracellular fluxes in *Escherichia coli***. *Mol Syst Biol* 2007, **3**:119.
60. Feist AM, Palsson BO: **The biomass objective function**. *Curr Opin Microbiol* 2010, **13**:344-349.
61. Long CP, Antoniewicz MR: **Metabolic flux responses to deletion of 20 core enzymes reveal flexibility and limits of *E. coli* metabolism**. *Metab Eng* 2019, **55**:249-257.
62. Foster CJ, Gopalakrishnan S, Antoniewicz MR, Maranas CD: **From *E. coli* mutant <sup>13</sup>C labeling data to a core kinetic model: a kinetic model parameterization pipeline**. *PLoS Comput Biol* 2019, **15**:e1007319.
63. Gonzalez JE, Antoniewicz MR: **Tracing metabolism from lignocellulosic biomass and gaseous substrates to products with stable-isotopes**. *Curr Opin Biotechnol* 2017, **43**:86-95.
64. Long CP, Antoniewicz MR: **Metabolic flux analysis of *Escherichia coli* knockouts: lessons from the Keio collection and future outlook**. *Curr Opin Biotechnol* 2014, **28**:127-133.
65. Long CP, Antoniewicz MR: **High-resolution (<sup>13</sup>C) metabolic flux analysis**. *Nat Protoc* 2019, **14**:2856-2877
- <sup>13</sup>C-metabolic flux analysis is an indispensable tool for quantifying metabolic fluxes in biological systems. This paper describes current state-of-the-art methods for <sup>13</sup>C-MFA. These methods were recently extended to measure fluxes and cross-feeding rates in microbial co-cultures.
66. Antoniewicz MR: **Parallel labeling experiments for pathway elucidation and <sup>13</sup>C metabolic flux analysis**. *Curr Opin Biotechnol* 2015, **36**:91-97.
67. Au J, Choi J, Jones SW, Venkataramanan KP, Antoniewicz MR: **Parallel labeling experiments validate *Clostridium acetobutylicum* metabolic network model for (<sup>13</sup>C) metabolic flux analysis**. *Metab Eng* 2014, **26**:23-33.
68. Ruhl M, Hardt WD, Sauer U: **Subpopulation-specific metabolic pathway usage in mixed cultures as revealed by reporter protein-based <sup>13</sup>C analysis**. *Appl Environ Microbiol* 2011, **77**:1816-1821.
69. Shaikh AS, Tang YJ, Mukhopadhyay A, Keasling JD: **Isotopomer distributions in amino acids from a highly expressed protein as a proxy for those from total protein**. *Anal Chem* 2008, **80**:886-890.
70. Gebreselassie NA, Antoniewicz MR: **(<sup>13</sup>C)-metabolic flux analysis of co-cultures: a novel approach**. *Metab Eng* 2015, **31**:132-139.
71. Wolfsberg E, Long CP, Antoniewicz MR: **Metabolism in dense microbial colonies: (<sup>13</sup>C) metabolic flux analysis of *E. coli* grown on agar identifies two distinct cell populations with acetate cross-feeding**. *Metab Eng* 2018, **49**:242-247
- *E. coli* grown in biofilm separates into two distinct cell populations that engage in metabolite cross-feeding. In this work, a new <sup>13</sup>C-MFA approach was applied to elucidate the metabolism of the two distinct subpopulations and quantify the cross-feeding of acetate between the populations. This <sup>13</sup>C-MFA approaches can also be applied to elucidate cross-feeding rates in microbial communities.
72. Phelan VV, Liu WT, Pogliano K, Dorrestein PC: **Microbial metabolic exchange—the chemotype-to-phenotype link**. *Nat Chem Biol* 2011, **8**:26-35.
73. Braga RM, Dourado MN, Araujo WL: **Microbial interactions: ecology in a molecular perspective**. *Braz J Microbiol* 2016, **47** (Suppl. 1):86-98.
74. Hsu RH, Clark RL, Tan JW, Ahn JC, Gupta S, Romero PA, Venturelli OS: **Microbial interaction network inference in microfluidic droplets**. *Cell Syst* 2019, **9**:229-242 e224
- Identification of microbial interactions is an inherently challenging problem. In this work, the authors have applied fluorescence microscopy to quantify strain abundances in microfluidic droplets and infer pairwise and higher-order interactions in synthetic consortia.
75. Kong W, Meldgin DR, Collins JJ, Lu T: **Designing microbial consortia with defined social interactions**. *Nat Chem Biol* 2018, **14**:821-829
- In this study, six two-strain consortia were constructed, each possessing a different mode of interaction: commensalism, amensalism, neutralism, cooperation, competition and predation. The population dynamics of these consortia displayed distinct characteristics determined by their underlying interaction modes.
76. Lawson CE, Harcombe WR, Hatzepichler R, Lindemann SR, Löffler FE, O'Malley MA, Garcia Martin H, Pfleger BF, Raskin L, Venturelli OS et al.: **Common principles and best practices for engineering microbiomes**. *Nat Rev Microbiol* 2019, **17**:725-741.
77. Gonze D, Coyte KZ, Lahti L, Faust K: **Microbial communities as dynamical systems**. *Curr Opin Microbiol* 2018, **44**:41-49.
78. Harcombe WR, Chacon JM, Adamowicz EM, Chubiz LM, Marx CJ: **Evolution of bidirectional costly mutualism from byproduct consumption**. *Proc Natl Acad Sci U S A* 2018, **115**:12000-12004.
79. Kumar M, Ji B, Zengler K, Nielsen J: **Modelling approaches for studying the microbiome**. *Nat Microbiol* 2019, **4**:1253-1267
- This paper provides an excellent overview of the wide range of mathematical modeling approaches that have been developed and applied to study gut microbiomes.
80. Saleski TE, Kerner AR, Chung MT, Jackman CM, Khasbaatar A, Kurabayashi K, Lin XN: **Syntrophic co-culture amplification of production phenotype for high-throughput screening of microbial strain libraries**. *Metab Eng* 2019, **54**:232-243
- The authors developed a high-throughput co-culture screening platform based on cross-feeding auxotrophs to select for high-production phenotypes of a wide range of target molecules. As a proof-of-concept, the approach was used to select *Escherichia coli* strains for the overproduction of two target molecules.
81. Ghosh A, Nilmeier J, Weaver D, Adams PD, Keasling JD, Mukhopadhyay A, Petzold CJ, Martin HG: **A peptide-based method for <sup>13</sup>C metabolic flux analysis in microbial communities**. *PLoS Comput Biol* 2014, **10**:e1003827.
82. Choi J, Antoniewicz MR: **Tandem mass spectrometry for (<sup>13</sup>C) metabolic flux analysis: methods and algorithms based on emu framework**. *Front Microbiol* 2019, **10**:31.