

From Cellular Automata to the Lattice Boltzmann Method

BGCE Honours Project

Theresa Pollinger, Marcial Gaißert
Ferienakademie Sarntal 2017
September 5, 2017

Outline

The Lattice Boltzmann Method

The Lattice Boltzmann Method

Stream Step


Discrete
particle distribution function

$$f_i = c_j \cdot w_j$$

Stream Step

Discrete
particle distribution function

$$f_i = \mathbf{c}_i \cdot \mathbf{w}_i$$




One streaming step for a D2Q9-stencil

Stream Step

Discrete
particle distribution function

$$f_i = \mathbf{c}_i \cdot \mathbf{w}_i$$



One streaming step for a D2Q9-stencil

The Lattice Boltzmann Equation

Applied as the collide step:

$$f_i(\mathbf{x} + \mathbf{c}_i \Delta t, t + \Delta t) = f_i(\mathbf{x}, t) + \Omega_i(\mathbf{x}, t)$$

Moments - the Interesting Outputs in LBM

Density and velocity are given locally

$$\rho(\mathbf{x}, t) = \sum_i f_i(\mathbf{x}, t)$$

$$\rho \mathbf{u}(\mathbf{x}, t) = \sum_i \mathbf{c}_i f_i(\mathbf{x}, t)$$

Source: **kruger.2016**

Earth Mantle Convection II

- Creeping flow, according to **weismueller.2015**

$$\left. \begin{array}{l} \mu \approx 10^{21} \text{ Pa} \cdot \text{s} \\ u \approx 3 \text{ cm/a} \\ h = 2867 \text{ km} \\ \rho \approx 4000 \text{ kg/m}^3 \end{array} \right\} Re = \frac{\rho \cdot u \cdot h}{\mu} \approx 10^{-20} \ll 1$$

- Dominant viscous forces, negligible inertial forces
- *Navier-Stokes* equations simplify to *Stokes* equations

$$\begin{aligned} \mu \Delta \mathbf{u} - \nabla p + \mathbf{f} &= 0 \\ \nabla \cdot \mathbf{u} &= 0 \end{aligned}$$

Earth Mantle Convection II

- Creeping flow, according to **weismueller.2015**

$$\left. \begin{array}{l} \mu \approx 10^{21} \text{ Pa} \cdot \text{s} \\ u \approx 3 \text{ cm/a} \\ h = 2867 \text{ km} \\ \rho \approx 4000 \text{ kg/m}^3 \end{array} \right\} Re = \frac{\rho \cdot u \cdot h}{\mu} \approx 10^{-20} \ll 1$$

- Dominant viscous forces, negligible inertial forces
- *Navier-Stokes* equations simplify to *Stokes* equations

$$\begin{aligned} \mu \Delta \mathbf{u} - \nabla p + \mathbf{f} &= 0 \\ \nabla \cdot \mathbf{u} &= 0 \end{aligned}$$

Derivation of a Benchmark II

- Construct a divergence-free velocity field

$$\begin{bmatrix} u \\ v \end{bmatrix} = \nabla^\perp \Phi = \begin{bmatrix} \sin(2\pi x) \cos(\pi y) \\ -2 \cos(2\pi x) \sin(\pi y) \end{bmatrix}$$

- Construct the pressure field according to the *Stokes* equations with $\mu = 1$

$$\mathbf{F} = \begin{bmatrix} f_x \\ f_y \end{bmatrix} = \begin{bmatrix} -\Delta u \\ -\Delta v \end{bmatrix} + \nabla p = \begin{bmatrix} 0 \\ ? \end{bmatrix}$$

$$\Rightarrow p = \frac{5}{2} \pi \cos(2\pi x) \cos(\pi y)$$

- Calculate the force field

$$\mathbf{F} = \begin{bmatrix} f_x \\ f_y \end{bmatrix} = \begin{bmatrix} 0 \\ -12.5\pi^2 \cos(2\pi x) \sin(\pi y) \end{bmatrix}$$

- Extend the domain to $\Omega = (0, 1)^3$ to obtain a pseudo-3D benchmark

Derivation of a Benchmark II

- Construct a divergence-free velocity field

$$\begin{bmatrix} u \\ v \end{bmatrix} = \nabla^\perp \Phi = \begin{bmatrix} \sin(2\pi x) \cos(\pi y) \\ -2 \cos(2\pi x) \sin(\pi y) \end{bmatrix}$$

- Construct the pressure field according to the *Stokes* equations with $\mu = 1$

$$\mathbf{F} = \begin{bmatrix} f_x \\ f_y \end{bmatrix} = \begin{bmatrix} -\Delta u \\ -\Delta v \end{bmatrix} + \nabla p = \begin{bmatrix} 0 \\ ? \end{bmatrix}$$

$$\Rightarrow p = \frac{5}{2} \pi \cos(2\pi x) \cos(\pi y)$$

- Calculate the force field

$$\mathbf{F} = \begin{bmatrix} f_x \\ f_y \end{bmatrix} = \begin{bmatrix} 0 \\ -12.5\pi^2 \cos(2\pi x) \sin(\pi y) \end{bmatrix}$$

- Extend the domain to $\Omega = (0, 1)^3$ to obtain a pseudo-3D benchmark

Derivation of a Benchmark II

- Construct a divergence-free velocity field

$$\begin{bmatrix} u \\ v \end{bmatrix} = \nabla^\perp \Phi = \begin{bmatrix} \sin(2\pi x) \cos(\pi y) \\ -2 \cos(2\pi x) \sin(\pi y) \end{bmatrix}$$

- Construct the pressure field according to the *Stokes* equations with $\mu = 1$

$$\mathbf{F} = \begin{bmatrix} f_x \\ f_y \end{bmatrix} = \begin{bmatrix} -\Delta u \\ -\Delta v \end{bmatrix} + \nabla p = \begin{bmatrix} 0 \\ ? \end{bmatrix}$$

$$\Rightarrow p = \frac{5}{2} \pi \cos(2\pi x) \cos(\pi y)$$

- Calculate the force field

$$\mathbf{F} = \begin{bmatrix} f_x \\ f_y \end{bmatrix} = \begin{bmatrix} 0 \\ -12.5\pi^2 \cos(2\pi x) \sin(\pi y) \end{bmatrix}$$

- Extend the domain to $\Omega = (0, 1)^3$ to obtain a pseudo-3D benchmark

Derivation of a Benchmark II

- Construct a divergence-free velocity field

$$\begin{bmatrix} u \\ v \end{bmatrix} = \nabla^\perp \Phi = \begin{bmatrix} \sin(2\pi x) \cos(\pi y) \\ -2 \cos(2\pi x) \sin(\pi y) \end{bmatrix}$$

- Construct the pressure field according to the *Stokes* equations with $\mu = 1$

$$\mathbf{F} = \begin{bmatrix} f_x \\ f_y \end{bmatrix} = \begin{bmatrix} -\Delta u \\ -\Delta v \end{bmatrix} + \nabla p = \begin{bmatrix} 0 \\ ? \end{bmatrix}$$

$$\Rightarrow p = \frac{5}{2} \pi \cos(2\pi x) \cos(\pi y)$$

- Calculate the force field

$$\mathbf{F} = \begin{bmatrix} f_x \\ f_y \end{bmatrix} = \begin{bmatrix} 0 \\ -12.5\pi^2 \cos(2\pi x) \sin(\pi y) \end{bmatrix}$$

- Extend the domain to $\Omega = (0, 1)^3$ to obtain a pseudo-3D benchmark

Derivation of a Benchmark II

- Construct a divergence-free velocity field

$$\begin{bmatrix} u \\ v \end{bmatrix} = \nabla^\perp \Phi = \begin{bmatrix} \sin(2\pi x) \cos(\pi y) \\ -2 \cos(2\pi x) \sin(\pi y) \end{bmatrix}$$

- Construct the pressure field according to the *Stokes* equations with $\mu = 1$

$$\mathbf{F} = \begin{bmatrix} f_x \\ f_y \end{bmatrix} = \begin{bmatrix} -\Delta u \\ -\Delta v \end{bmatrix} + \nabla p = \begin{bmatrix} 0 \\ ? \end{bmatrix}$$

$$\Rightarrow p = \frac{5}{2} \pi \cos(2\pi x) \cos(\pi y)$$

- Calculate the force field

$$\mathbf{F} = \begin{bmatrix} f_x \\ f_y \end{bmatrix} = \begin{bmatrix} 0 \\ -12.5\pi^2 \cos(2\pi x) \sin(\pi y) \end{bmatrix}$$

- Extend the domain to $\Omega = (0, 1)^3$ to obtain a pseudo-3D benchmark

Preparations

- Use CLOCK_MONOTONIC from the system environment for runtime measurement
- Implement “random-noise” initialization
- Implement an approximated L2 norm to ensure comparability with LBM
⇒ Verification by taking the norm of the analytical velocity field
- Calculate the accuracy based on the approximated L2 norm

$$\frac{\| \mathbf{u} - \mathbf{u}_{\text{exact}} \|}{\| \mathbf{u}_{\text{exact}} \|}$$

- Use the tolerance 10^{-12} between two iterations to identify the steady state

LBM - very short overview

Collide step: application of the lattice Boltzmann equation

$$f_i(\mathbf{x} + \mathbf{c}_i \Delta t, t + \Delta t) = f_i(\mathbf{x}, t) + \Omega_i(\mathbf{x}, t)$$

Density and velocity are given locally

$$\rho(\mathbf{x}, t) = \sum_i f_i(\mathbf{x}, t)$$

$$\rho \mathbf{u}(\mathbf{x}, t) = \sum_i \mathbf{c}_i f_i(\mathbf{x}, t)$$

Source: **kruger.2016**

LBM - very short overview

Collide step: application of the lattice Boltzmann equation

$$f_i(\mathbf{x} + \mathbf{c}_i \Delta t, t + \Delta t) = f_i(\mathbf{x}, t) + \Omega_i(\mathbf{x}, t)$$

Density and velocity are given locally

$$\rho(\mathbf{x}, t) = \sum_i f_i(\mathbf{x}, t)$$

$$\rho \mathbf{u}(\mathbf{x}, t) = \sum_i \mathbf{c}_i f_i(\mathbf{x}, t)$$

Source: **kruger.2016**

Boundary Conditions : Expectations

- Free-slip boundary conditions (Neumann Boundary Conditions)
- No-slip / UBB boundary conditions (Dirichlet boundary conditions)


Both are expected to converge with order 2 in Δx

Boundary Conditions : Expectations

- Free-slip boundary conditions (Neumann Boundary Conditions)
- No-slip / UBB boundary conditions (Dirichlet boundary conditions)

Both are expected to converge with order 2 in Δx

Thermal LBM : Principle



Thanks for listening.
Any questions?

References

References