

From Cellular Automata to the Lattice Boltzmann Method

Theresa Pollinger, Marcial Gaißert
Ferienakademie Sarntal 2017
September 6, 2017

Outline

The Lattice Boltzmann Method


The Lattice Boltzmann Method

Collide Step

Particle distribution function f_i :
discrete in space and time,
continuous in probability values

Collide Step


Particle distribution function f_i :
discrete in space and time,
continuous in probability values



One collide step for a D2Q9-stencil

Collide Step


Particle distribution function f_i :
discrete in space and time,
continuous in probability values



One collide step for a D2Q9-stencil

Stream Step


Particle distribution function f_i :
discrete in space and time,
continuous in probability values



One stream step for a D2Q9-stencil

Stream Step

Particle distribution function f_i :
discrete in space and time,
continuous in probability values



One stream step for a D2Q9-stencil

The Lattice Boltzmann Equation

...is what we get when we apply both steps:

$$f_i(\mathbf{x} + \mathbf{c}_i \Delta t, t + \Delta t) = f_i(\mathbf{x}, t) + \Omega_i(\mathbf{x}, t)$$

With collision operator $\Omega_i(\mathbf{x}, t)$, e.g.:

- BGK: $\Omega(f) = \frac{1}{\tau}(f_i^{\text{eq}} - f_i) = \omega(f_i^{\text{eq}} - f_i)$
- TRT:

and the equilibrium particle distribution function $f_i^{\text{eq}} =$

Cf. Boltzmann Equation

$$\frac{\partial f}{\partial t} + \frac{\mathbf{p}}{m} \cdot \nabla f = \left(\frac{\partial f}{\partial t} \right)_{\text{coll}}$$

The Lattice Boltzmann Equation

...is what we get when we apply both steps:

$$f_i(\mathbf{x} + \mathbf{c}_i \Delta t, t + \Delta t) = f_i(\mathbf{x}, t) + \Omega_i(\mathbf{x}, t)$$

With collision operator $\Omega_i(\mathbf{x}, t)$, e.g.:

- BGK: $\Omega(f) = \frac{1}{\tau}(f_i^{\text{eq}} - f_i) = \omega(f_i^{\text{eq}} - f_i)$
- TRT:

and the equilibrium particle distribution function $f_i^{\text{eq}} =$

Cf. Boltzmann Equation

$$\frac{\partial f}{\partial t} + \frac{\mathbf{p}}{m} \cdot \nabla f = \left(\frac{\partial f}{\partial t} \right)_{\text{coll}}$$

The Lattice Boltzmann Equation

...is what we get when we apply both steps:

$$f_i(\mathbf{x} + \mathbf{c}_i \Delta t, t + \Delta t) = f_i(\mathbf{x}, t) + \Omega_i(\mathbf{x}, t)$$

With collision operator $\Omega_i(\mathbf{x}, t)$, e.g.:

- BGK: $\Omega(f) = \frac{1}{\tau}(f_i^{\text{eq}} - f_i) = \omega(f_i^{\text{eq}} - f_i)$
- TRT:

and the equilibrium particle distribution function $f_i^{\text{eq}} =$

Cf. Boltzmann Equation

$$\frac{\partial f}{\partial t} + \frac{\mathbf{p}}{m} \cdot \nabla f = \left(\frac{\partial f}{\partial t} \right)_{\text{coll}}$$

The Lattice Boltzmann Equation

...is what we get when we apply both steps:

$$f_i(\mathbf{x} + \mathbf{c}_i \Delta t, t + \Delta t) = f_i(\mathbf{x}, t) + \Omega_i(\mathbf{x}, t)$$

With collision operator $\Omega_i(\mathbf{x}, t)$, e.g.:

- BGK: $\Omega(f) = \frac{1}{\tau}(f_i^{\text{eq}} - f_i) = \omega(f_i^{\text{eq}} - f_i)$
- TRT:

and the equilibrium particle distribution function $f_i^{\text{eq}} =$

Cf. Boltzmann Equation

$$\frac{\partial f}{\partial t} + \frac{\mathbf{p}}{m} \cdot \nabla f = \left(\frac{\partial f}{\partial t} \right)_{\text{coll}}$$

Moments - the Interesting Outputs in LBM

Density and velocity are given locally

$$\rho(\mathbf{x}, t) = \sum_i f_i(\mathbf{x}, t)$$

$$\rho \mathbf{u}(\mathbf{x}, t) = \sum_i \mathbf{c}_i f_i(\mathbf{x}, t)$$

Nondimensionalization – setting the step size

= Normalization in the three physical dimensions time, space, mass

Usual choice:

$$\Delta x^* = 1 = C_x \cdot \Delta x$$

$$\Delta t^* = 1 = C_t \cdot \Delta t$$

$$\Delta \rho^* = 1 = C_\rho \cdot \Delta \rho$$

The kinematic velocity in lattice units is directly linked to ω/τ :

Nondimensionalization – setting the step size

= Normalization in the three physical dimensions time, space, mass

Usual choice:

$$\Delta x^* = 1 = C_x \cdot \Delta x$$

$$\Delta t^* = 1 = C_t \cdot \Delta t$$

$$\Delta \rho^* = 1 = C_\rho \cdot \Delta \rho$$

The kinematic velocity in lattice units is directly linked to ω/τ :

Nondimensionalization – setting the step size

= Normalization in the three physical dimensions time, space, mass

Usual choice:

$$\Delta x^* = 1 = C_x \cdot \Delta x$$

$$\Delta t^* = 1 = C_t \cdot \Delta t$$

$$\Delta \rho^* = 1 = C_\rho \cdot \Delta \rho$$

The kinematic velocity in lattice units is directly linked to ω/τ :

Thanks for listening.
Any questions?

References

References