

## DO LY $\alpha$ ABSORBERS CO-ROTATE WITH GALAXY DISKS?

DAVID M. FRENCH, BART P. WAKKER

Department of Astronomy, University of Wisconsin, Madison, WI 53706, USA

### ABSTRACT

We present results of a study comparing the relative velocity of Ly $\alpha$  absorbers to the rotation direction and velocity of nearby galaxy disks. We find...

*Keywords:* galaxies:intergalactic medium, galaxies:evolution, galaxies:halos, quasars: absorption lines

### 1. INTRODUCTION

Our current  $\Lambda$ CDM cosmology picture describes galaxies forming hierarchically out of overdensities in the underlying dark matter distribution. As matter is funneled toward a growing galaxy, conservation of angular momentum redistributes the angular momentum in this gas to match that of the halo and underlying dark matter as the gas is shock-heated and slowly cools. As this infalling gas is responsible for birthing and continuing to feed the galaxies, it is expected that the extended gaseous halos should rotate in the same sense as both the galactic disks and dark matter halos. Galaxy rotation curves have been observed to extend at constant velocity out to... (cite...). It becomes increasingly difficult to measure gas rotation much farther from this however as the density rapidly decreases. Within this region the galaxy disks transition into circumgalactic medium (CGM), and eventually the CGM merges with the intergalactic medium (IGM). At what point, however, does the surrounding medium cease to circulate with the galaxy?

HYDRO? simulations such as those by ?? suggests that the bulk CGM kinematics out to (WHAT DISTANCE) may circulate, and that absorption in intervening QSO sightlines should be able to accurately capture this rotation signature. Observational confirmation, however, has been inconclusive. Côté et al. 2005 probed the halos of nine galaxies using *HST* observed background QSOs, finding large warps would be needed to explain the velocity of H I absorbers by an extended rotating disk. ? compiled a sample of 4 galaxy-QSO systems from the literature, finding only 1/4 of Ly $\alpha$  absorbers appeared to co-rotate with the associated galaxy disk. Approaching the question from a different angle, ? probed the halo of a single galaxy, NGC1097, with 4 nearby QSO sightlines, and suggests that an extended, slowly rotating disk with additional inflowing IGM material best matches observations.

Numerous studies have shown a correlation between equivalent width and decreasing velocity difference between galaxies and IGM absorbers (e.g., ?, MORE).

To make progress here, we have obtained rotation

curves for 12 nearby spiral galaxies which are located within 500 kpc of a background QSO observed by the Cosmic Origins Spectrograph (COS) on *HST*. A literature search yielded an additional 17 galaxies with published rotation curves and known orientations. Each of these is probed by at least one QSO within 500 kpc.

We have augmented this new sample with additional galaxies with known rotation velocity and orientation from the literature. In Section 2 we describe the selection and reduction of both SALT and COS spectra. We then discuss each galaxy-QSO system in detail in Section 3, and introduce our halo-velocity model for interpreting these systems in Section 4. In Section 5 we discuss the overall results of this exercise and present a physically-motivated interpretation of these results. See Section 6 for a summary of our results and conclusions.

### 2. DATA AND ANALYSIS

#### 2.1. SALT Data

Our sample contains 12 galaxies observed with the Southern African Large Telescope (SALT) Robert Stobie Spectrograph (RSS) in longslit mode. These 12 were selected from a larger pool of 48 submitted targets by the SALT observing queue. These 48 possible targets were chosen for their proximity to background QSOs whose spectra contained promising Ly $\alpha$  lines. Finally, we only included galaxies with  $z \leq 0.33$  ( $cz \leq 10,000 \text{ km s}^{-1}$ ), angular sizes less than 6' to enable easy sky subtraction without taking additional exposures, and surface brightnesses sufficient to keep exposure times below  $\sim 1300s$ . Table 2 summarizes these observations. Data was taken for 2 additional galaxies, NGC3640 and NGC2962, but proved unusable due to issues with spectral identification and low signal-to-noise (respectively).

All SALT galaxy spectra were reduced and extracted using the standard PySALT reduction package (CITATION), which includes procedures to prepare the data, correct for gain, cross-talk, bias, and overscan, and finally mosaic the images from the 3 CCDs. Next, we rectify the images with wavelength solutions found via Ne and Ar arc lamp spectra line identification. Finally, we perform a basic sky subtraction using an off-

Target	Galaxy	R.A.	Dec.	$z$	Program	Cyl. Vel. Range [km s <sup>-1</sup> ]	NFW Vel. Range [km s <sup>-1</sup> ]	$T_{exp}$ * [ks]
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)
1H0419-577	NGC1566	04 26 00.7	-57 12 02.0	0.10400	11686	[49, 77]	[-2, 31]	20429
HE0429-5343	NGC1566	04 30 40.0	-53 36 56.0	0.04001	12275	[-54, -2]	[-22, 17]	2067
HE0435-5304	NGC1566	04 36 50.9	-52 58 47.0	0.42616	11520	[-45, -43]	[-16, -15]	8372
RBS567	NGC1566	04 39 38.7	-53 11 31.0	0.24300	11520	[N/A]	[N/A]	8176
HE0439-5254	NGC1566	04 40 12.0	-52 48 18.0	1.05300	11520	[N/A]	[N/A]	8402
H1101-232	NGC3513	11 03 37.7	-23 29 31.0	0.18600	12025	[-36, 29]	[-35, 33]	13341
SDSSJ112005.00+041323.0	NGC3633	11 20 05.0	+04 13 23.0	0.54689	12603	[-155, -12]	[-77, 13]	4708
RX_J1121.2+0326	CGCG039-137	11 21 14.0	+03 25 47.0	0.15200	12248	[-36, 138]	[-36, 166]	2695
RX_J1121.2+0326	NGC3633	11 21 14.0	+03 25 47.0	0.15200	12248	[-155, -12]	[-77, 13]	2695
SDSSJ112224.10+031802.0	NGC3633	11 22 24.1	+03 18 02.0	0.47528	12603	N/A	N/A	7588
SDSSJ112224.10+031802.0	CGCG039-137	11 22 24.1	+03 18 02.0	0.47528	12603	[-130, -114]	[-116, -98]	7588
3C273.0	NGC4536	12 29 06.7	+02 03 09.0	0.15834	12038	[67, 134]	[-6, 43]	4002
HE1228+0131	NGC4536	12 30 50.0	+01 15 23.0	0.11700	11686	[23, 131]	[7, 33]	11036
PG1302-102	NGC4939	13 05 33.0	-10 33 19.0	0.27840	12038	[-219, 14]	[-119, 36]	5979
SDSSJ135726.27+043541.4	NGC5364	13 57 26.3	+04 35 41.0	1.23453	12264	[-38, 123]	[-41, 80]	14148
QSO1500-4140	NGC5786	15 03 34.0	-41 52 23.0	0.33500	11659	[120, 175]	[26, 69]	9258
SDSSJ151237.15+012846.0	UGC09760	15 12 37.2	+01 28 46.0	0.26625	12603	[-30, 30]	[-30, 86]	7590
RBS1768	ESO343-G014	21 38 49.9	-38 28 40.0	0.18299	12936	[-203, 10]	[-122, 30]	6962
MRC2251-178	MCG-03-58-009	22 54 05.9	-17 34 55.0	0.06609	12029	[-52, 166]	[-61, 110]	5515
RBS2000	IC5325	23 24 44.7	-40 40 49.0	0.17359	13448	[-46, 5]	[-24, 17]	5046
RX_J1017.5+4702	NGC3198	10 17 31.0	+47 02 25.0	0.33544	13314	[-154, -21]	[-91, 6]	8655
SDSSJ101622.60+470643.0	NGC3198	10 16 22.6	+47 06 43.0	0.82222	11598	[-155, -17]	[-89, 9]	4906
RX_J1236.0+2641	NGC4565	12 36 04.0	+26 41 36.0	0.20920	12248	[-1, 250]	[-29, 146]	4235
PG0804+761	UGC04238	08 10 58.7	+76 02 43.0	0.10200	11686	[-3, 86]	[-6, 94]	5510
SDSSJ104335.90+115129.0	NGC3351	10 43 35.9	+11 05 29.0	0.79400	14071	[-99, 12]	[-68, 20]	4736
MRK771	NGC4529	12 32 03.6	+20 09 30.0	0.06301	12569	[-103, -40]	[-105, -35]	1868
MRK876	NGC6140	16 13 57.2	+65 43 11.0	0.12900	11524	[40, 101]	[35, 102]	12579
SBS1503+570	NGC5907	15 04 55.6	+56 49 20.0	0.35894	12276	[31, 228]	[-24, 101]	5163
SDSSJ152053.59+571122.1	NGC5907	15 20 53.7	+57 11 23.0	0.02952	13654	[-18, 228]	[-42, 114]	3753
RBS1503	NGC5907	15 29 7.5	+56 16 07.0	0.09900	12276	[-228, -9]	[-96, 33]	1964
SDSSJ112448.30+531818.0	UGC06446	11 24 48.3	+53 18 19.0	0.53151	14240	[-9, 65]	[-15, 61]	7920
RX_J1117.6+5301	UGC06446	11 17 40.5	+53 01 51.0	0.15871	14240	[35, 47]	[25, 35]	4943
SBS1116+523	UGC06399	11 19 47.9	+52 05 53.0	0.35568	14240	[-91, -54]†	[-80, -42]†	4949
CSO1208	NGC3726	11 40 47.9	+46 22 05.0	0.11500	14729	[-27, 29]	[-28, 21]	3052
RX_J1142.7+4625	NGC3726	11 42 41.2	+46 24 36.0	0.11500	14772	[-34, -14]	[-30, -7]	2368
3C232	NGC3067	09 58 20.9	+32 24 02.0	0.5306 0	8596	[-121, 25]	[-139, 26]	44662
SDSSJ095914.80+320357.0	NGC3067	09 59 14.8	+32 03 57.0	0.56462	12603	[11, 138]	[-12, 81]	2273
RX_J1002.9+3240	NGC3067	10 02 54.5	+32 40 39.0	0.83000	12603	[115, 147]	[42, 65]	7713
FBQSJ0908+3246	NGC2770	09 08 38.8	+32 46 20.0	0.25989	14240	[-146, -4]	[-132, 6]	7430
SDSSJ091052.80+333008.0	NGC2770	09 10 52.8	+33 30 08.0	0.11631	14240	[6, 145]	[-4, 127]	7442
SDSSJ091127.30+325337.0	NGC2770	09 11 27.3	+32 53 37.0	0.29038	14240	[-150, -43]	[-132, -25]	10028
TON1009	NGC2770	09 09 06.2	+32 36 30.0	0.81028	12603	[-146, -39]	[-125, -20]	4740
TON1015	NGC2770	09 10 37.0	+33 29 24.0	0.35400	14240	[3, 146]	[-7, 130]	4774
CSO295	NGC3432	10 52 05.6	+36 40 40.0	0.60900	14772	[-37, 48]	[-37, 134]	1088
MS1047.3+3518	NGC3432	10 50 10.9	+35 02 02.0	0.04125	8316	[35, 124]	[19, 109]	8301
RX_J1054.2+3511	NGC3432	10 54 16.2	+35 11 24.0	0.20300	14772	[0, 123]	[-9, 111]	533
SDSSJ112439.50+113117.0	NGC3666	11 24 39.4	+11 31 17.0	0.14300	14071	[-87, 20]	[-136, 20]	10427
SDSSJ112632.90+120437.0	NGC3666	11 26 32.9	+12 04 37.0	0.97700	13314	[-59, -29]	[-45, -18]	8289
SDSSJ112756.70+115427.0	NGC3666	11 27 56.8	+11 54 27.0	0.50900	14145	[-127, -80]	[-100, -57]	5146
2E1530+1511	NGC5951	15 33 14.3	+15 01 03.0	0.09000	14071	[-31, 114]	[-32, 136]	9348
MRK335	NGC7817	00 06 19.5	+20 12 11.0	0.02578	11524	[-26, -24]	[-26, -24]	5122
PG1259+593	UGC08146	13 01 12.9	+59 02 07.0	0.4778 0	11541	[-13, 82]	[-15, 91]	9200

**Table 1.** Summary of COS targets in this sample. \*Total exposure time and S/N ratio is given for multi-orbit exposures. † Velocity range based on a  $2 \times 4 R_{vir}$  halo.

sky portion of the spectrum, and extract 5-10 pixel wide 1-D strips from the reduced 2-D spectrum.

For each 1-D spectrum, we identify the H $\alpha$  emission lines and perform a non-linear least-squares Voigt profile fit using the Python package LMFIT<sup>1</sup>. The line centroid and  $1\sigma$  standard errors are returned, and these fits are then shifted to rest-velocity based on the galaxy systemic redshift and heliocentric velocity corrections are calculated with the IRAF rvcorrect procedure. The final rotation velocity is calculated by then applying the inclination correction,  $v_{rot} = v/\sin(i)$ . Final errors are calculated as a quadrature sum of  $1\sigma$  fit errors, systemic redshift error, and inclination uncertainty as follows:

$$\sigma^2 = \left(\frac{\partial v_{rot}}{\partial \lambda_{obs}}\right)^2 (\Delta \lambda_{obs})^2 + \left(\frac{\partial v_{rot}}{\partial v_{sys}}\right)^2 (\Delta v_{sys})^2 + \left(\frac{\partial v_{rot}}{\partial i}\right)^2 (\Delta i)^2, \quad (1)$$

where  $\Delta \lambda_{obs}$ ,  $\Delta v_{sys}$ , and  $\Delta i$  are the errors in observed line center, galaxy redshift, and inclination, respectively.

We determine the inclination error by calculating the standard deviation of the set of all axis ratio values available in NED for each galaxy. The final physical scale is calculated using the SALT image scale of 0.1267 arcsec/pixel, multiplied by the 4-pixel spatial binning, and converted to physical units using a redshift-independent distance if available, and a Hubble flow estimate if not. We adopt a Hubble constant of  $H_0 = 71 \text{ km s}^{-1} \text{ Mpc}^{-1}$  throughout.

Finally, we calculate our approaching and receding velocities via a weighted mean of the outer 1/2 of each rotation curve, with errors calculated as weighted standard errors in the mean. Our final redshifts are calculated by forcing symmetric rotation, such that the outer 1/2 average velocity for each side matches in magnitude. See Figure ?? for an example.

## 2.2. COS Spectra

The Barbara A. Mikulski Archive for Space Telescopes (MAST) archives yield 19 QSO targets observed by COS which lie within 500 kpc of our SALT galaxies. These targets vary widely in signal-to-noise from approximately 5 to 100 due to our choosing them based only on their proximity to galaxies with known rotation. The reduction procedure for these spectra follow those described by ? and ?. In short, spectra are processed with CALCOS vXXXX? and combined via the method of ?, which helps corrects the COS wavelength scale misalignments produced by CALCOS. Multiple exposures are combined via alignment with Galactic 21cm absorp-

tion spectra and summing total counts per pixel before converting to flux. The COS instrument is described in detail by ?.

## 3. HALO ROTATION MODEL

In order to better understand how QSO sightlines probe intervening velocity structure we have developed a simple halo gas rotation model. This model is seeded by an observed rotation curve (or whatever rotation curve-like data suits ones fancy). This input curve is then interpolated and extended out to  $2R_{vir}$  based on the average velocity of the outer 1/2 radius. Next, we project this interpolated rotation curve onto a plane oriented to a faux QSO sightline identically to the input galaxy-QSO pair orientation. By stacking multiple rotation-planes in the galaxy z-axis direction, we then create a simple cylindrical rotating halo model. Finally, each rotation-plane in the stack is projected onto the faux sightline. The result is a function representing the rotation velocity encountered by the sightline as a function of velocity (or distance) along it.

For each galaxy-QSO pair we created 2 rotation models: 1) a purely cylindrical halo extending  $2R_{vir}$  in height and  $3R_{vir}$  in radius, and 2) a cylindrical model extending  $2R_{vir}$  in height and  $3R_{vir}$  in radius with rotation velocities which smoothly decline based on a NFW profile fit (??). Each model produces the velocity a co-rotating absorber would project onto the spectrum as a function of velocity along the sightline. We then collapse this into a simple range of possible observed velocities by summing the x- and y-components along the allowed range.

## 4. SALT GALAXIES

### 4.1. CGCG039-137

CGCG039-137 is an isolated Scd type galaxy with a measured systemic velocity of  $6918 \pm 24 \text{ km s}^{-1}$  and inclination of  $63^\circ$ . There are two associated sightlines: RX\_J1121.2+0326 at an impact parameter of 99 kpc and azimuth angle of  $71^\circ$  on the receding side, and SDSSJ112224.10+031802.0 at 491 kpc and  $24^\circ$  on the approaching side. Ly $\alpha$  absorption is detected in both sightlines within  $400 \text{ km s}^{-1}$  of CGCG039-137.

Towards RX\_J1121.2+0326 we detect Ly $\alpha$  at  $6975 \text{ km s}^{-1}$ , which, at  $\Delta v = 57 \text{ km s}^{-1}$ , lies well within the range of projected velocities consistent with co-rotation. The absorber detected toward SDSSJ112224.10+031802.0 occurs at a more distant  $6606 \text{ km s}^{-1}$  ( $\Delta v = -312 \text{ km s}^{-1}$ ). Although this absorber has the correct sign for co-rotation (blue-ward on the approaching side of the disk), the large velocity difference and impact parameter make it unlikely that this absorption can be linked to coherent halo rotation.

### 4.2. ESO343-G014

ESO343-G014 is an edge on spiral galaxy with a measured systemic velocity of  $9138.9 \pm 31.7 \text{ km s}^{-1}$ . It has a smaller neighboring galaxy, 2MASXJ21372816-3824412,

<sup>1</sup> <http://cars9.uchicago.edu/software/python/lmfit/contents.html>

Galaxy	R.A.	Dec.	$cz$ ( $\text{km s}^{-1}$ )	Type	Grating	$V_{\text{rot}}$ [ $\text{km s}^{-1}$ ]	$V_{\text{rot}}/\sin(i)$ [ $\text{km s}^{-1}$ ]	Obs Date	$T_{\text{exp}}$ [ks]
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)
CGCG039-137	11 21 26.95	+03 26 41.68	$6918 \pm 24$	Scd	PG2300	$132 \pm 16$	$139 \pm 26$	05 11 2016	700
ESO343-G014	21 37 45.18	-38 29 33.22	$9139 \pm 32$	S	PG2300	$203 \pm 32$	$203 \pm 32$	05 16 2016	1000
IC5325	23 28 43.43	-41 20 04.9	$1512 \pm 8$	SAB(rs)bc	PG2300	$53 \pm 5$	$125 \pm 39$	05 17 2016	600
MCG-03-58-009	22 53 40.85	-17 28 44.00	$9015 \pm 19$	Sc	PG2300	$150 \pm 12$	$171 \pm 23$	05 16 2016	1200
NGC1566	04 20 04.2	-54 56 16.12	$1502 \pm 15$	SAB(rs)bc	PG2300	$64 \pm 8$	$195 \pm 47$	10 18 2016	400
NGC3513	11 03 46.08	-23 14 43.8	$1204 \pm 12$	SB(s)c	PG2300	$11 \pm 10$	$22 \pm 24$	05 26 2016	600
NGC3633	11 20 26.22	+03 35 8.20	$2587 \pm 7$	SAa	PG2300	$149 \pm 6$	$157 \pm 9$	05 11 2016	1200
NGC4536	12 34 27.05	+02 11 17.30	$1867 \pm 33$	SAB(rc)bc	PG2300	$129 \pm 9$	$148 \pm 41$	05 11 2016	1300
NGC4939	13 04 14.39	-10 20 22.60	$3093 \pm 33$	SA(s)bc	PG2300	$204 \pm 25$	$275 \pm 66$	05 14 2016	500
NGC5364	13 56 12.00	+05 00 52.09	$1238 \pm 17$	SA(rs)bc pec	PG2300	$130 \pm 13$	$155 \pm 22$	05 11 2016	700
NGC5786	14 58 56.26	-42 00 48.10	$2975 \pm 22$	SAB(s)bc	PG2300	$156 \pm 10$	$172 \pm 25$	05 11 2016	250
UGC09760	15 12 02.44	+01 41 55.46	$2094 \pm 16$	Sd	PG2300	$46 \pm 10$	$46 \pm 16$	05 11 2016	500

**Table 2.** SALT targeted galaxies. Columns are as follows: 1) the galaxy name, 2), 3) R.A., Dec. in J2000, 4) galaxy systemic velocity, 5) morphological type (RC3), 6) RSS grating used, 7) approaching side velocity, 8) receding side velocity, 9) observation date, 10) exposure time, and 11) S/N of the H $\alpha$  or Ca H&K lines.

located north of it's major axis at a projected distance of 216 kpc and velocity of 9129. The nearest sightline is towards RBS1768 at an impact parameter of 466kpc and  $74^\circ$  azimuth angle on the approaching side. We detect 3 Ly $\alpha$  absorption lines within  $300 \text{ km s}^{-1}$  of ESO343-G014 (at 9308, 9360, and 9434  $\text{km s}^{-1}$ ). All of these are anti-aligned with the rotation of ESO343-G014, but unfortunately the presence of 2MASXJ21372816-3824412 makes it difficult to attribute this gas solely to ESO343-G014. Additionally, this gas could be attributed to either the approaching or receding side of the disk due to the large impact parameter and high azimuth angle of the sightline.

#### 4.3. IC5325

IC5325 is a face-on SAB(rs)bc type galaxy with a measured velocity of  $1511.9 \pm 8.4 \text{ km s}^{-1}$ . It's inclination is just high enough ( $25^\circ$ ) to obtain a reasonable rotation curve. The closest neighboring galaxy is ESO347-G020 to the Southeast at 306 kpc and a heliocentric velocity of  $1745 \text{ km s}^{-1}$ . Three other much smaller galaxies are also located  $\sim 450$  kpc to the Southwest. We detect Ly $\alpha$  absorption at  $1598 \text{ km s}^{-1}$ ,  $\Delta v = 86 \text{ km s}^{-1}$  in the spectrum towards RBS2000 at an impact parameter of 314 kpc and azimuth angle of  $64^\circ$  on the approaching side. While this velocity is anti-aligned with the rotation the disk gas, the low inclination angle of IC5325 leads to a highly uncertain position angle. Without additional observations, we cannot say for certain if the location of RBS2000 actually lies on the approaching or receding side. This position angle uncertainty also means our SALT rotation curve is a lower limit on the true rotation velocity of IC5325.

#### 4.4. MCG-03-58-009

MCG-03-58-009 is a massive and very isolated Sc type galaxy at a measured velocity of  $9015 \pm 19 \text{ km s}^{-1}$  and

inclination angle of  $49^\circ$ . A weak Ly $\alpha$  absorber is detected at  $9029 \text{ km s}^{-1}$  towards MRC2251-178, which lies 355 kpc away at an azimuth angle of  $71^\circ$  on the receding side. Although this absorber matches the velocity direction expected for co-rotation, the velocity difference ( $\Delta v = 14 \text{ km s}^{-1}$ ) is within the systemic velocity uncertainty. The relative weakness of this absorber ( $\text{EW} = 62 \pm 4 \text{ m\AA}$ ) is somewhat surprising given it's proximity (just outside of  $1 R_{\text{vir}}$ ) to a massive galaxy. If this is representative of an isolated system such as MCG-03-58-009, then we should expect the halo rotational velocity to approach systemic by  $1 R_{\text{vir}}$ .

#### 4.5. NGC1566

NGC1566 is well sampled (5 nearby QSO sightlines), but unfortunately also part of a complex environment of neighboring galaxies. We detect Ly $\alpha$  in all 5 of these sightlines. The farthest three, HE0439-5254, RBS567, and HE0435-5304, are clustered close together toward the northeast of NGC1566 at 459, 423, and 396 kpc and  $\sim 60^\circ$  azimuth angle. HE0439-5254 and RBS567 are both located beyond our  $2 \times 3 R_{\text{vir}}$  model limits and therefore are only included here for the sake of completeness. The slightly closer HE0435-5304 sneaks under the limit.

HE0429-5343 is in the same direction and azimuth angle but closer at  $\rho = 256$  kpc, and shows Ly $\alpha$  absorption at 1167 and  $1358 \text{ km s}^{-1}$ . These absorbers both have the correct velocity *sign*, but we would expect a smaller velocity for co-rotation (approximately  $\Delta v \sim \pm 40 \text{ km s}^{-1}$  projected). This difference could be explained by invoking either a warped extended disk, or perhaps inflowing gas.

1H0419-577 is located to the south at 303 kpc and just east of the receding side of the major axis at an azimuth angle of  $10^\circ$ . We detect Ly $\alpha$  at 1071, 1123, 1188, and  $1264 \text{ km s}^{-1}$ , all of which are the wrong sign for co-rotation or distant in velocity. This sightline is actually

closer to a small group of galaxies including NGC1549, NGC1546 and NGC1536, all with systemic velocities near  $1200 \text{ km s}^{-1}$ . We expect all of these lines to be associated with this group rather than with NGC1566.

#### 4.6. NGC3513

NGC3513 is a mostly face-on SB(rs)c galaxy with heliocentric velocity  $V_{hel} = 1204 \pm 12 \text{ km s}^{-1}$ . It has a companion galaxy in NGC3511 at an impact parameter of 44 kpc at  $v_{hel} = 1109 \text{ km s}^{-1}$ . We detect Ly $\alpha$  at  $1182 \text{ km s}^{-1}$  toward background QSO H1101-232, which is located directly south at 60 kpc and azimuth angle of  $67^\circ$  on the receding side. NGC3513 appears to be rotating slowly, with a maximal inclination-corrected rotation velocity of  $22 \pm 24 \text{ km s}^{-1}$ . The  $\Delta v = -22 \text{ km s}^{-1}$  for this absorber matches well with the magnitude of this rotation, but is opposite in sign for co-rotation. Given that NGC3511 is so close, this absorber's velocity is probably subject to a complex velocity field influenced by both NGC3511 and NGC3513.

#### 4.7. NGC3633

NGC3633 is an isolated, edge-on SAa type galaxy at a velocity of  $2587 \pm 7 \text{ km s}^{-1}$ . Several locations along the disk of NGC3633 show two velocities for emission. We have combined these into a single velocity measurement via a weighted average.

There are three nearby sightlines: SDSSJ112005.00+041323.0 is straight north at 468 kpc and  $78^\circ$  azimuth, RXJ1121.2+0326 is to the southeast at 184 kpc and  $58^\circ$  azimuth, and SDSSJ112224.10+031802.0 at 413 kpc and  $50^\circ$  azimuth. Toward RXJ1121.2+0326 we detect a Ly $\alpha$  absorber at  $2605 \text{ km s}^{-1}$  on the approaching side, which is essentially systemic velocity for NGC3633. The spectrum of SDSSJ112224.10+031802.0 shows absorbers at  $2285$  and  $2578 \text{ km s}^{-1}$ , both of which are of the correct sign for co-rotation. We do not detect any Ly $\alpha$  towards the third sightline, SDSSJ112224.10+031802.0.

SDSSJ112005.00+041323.0 and SDSSJ112224.10+031802.0 are both outside of our  $2 \times 3 R_{vir}$  model limits.

#### 4.8. NGC4536

NGC4536 is a SAB(rs)bc type galaxy located in a complex environment with many other nearby galaxies. The data on the receding side of NGC4536 is quite messy, and may include contamination from background sources. Hence, our measured systemic velocity of  $1867 \pm 33 \text{ km s}^{-1}$ , and thus rotation velocity of  $139 \pm 37 \text{ km s}^{-1}$ , have relatively high uncertainty. Other published redshift values available from NED and rotation velocities from the HyperLEDA database are broadly consistent with our values, albeit biased slightly lower and higher in velocity, respectively.

There are 2 sightlines to the southwest of NGC4536, both on the receding side of the galaxy. HE1228+0131 at 338 kpc and  $86^\circ$  azimuth has 5 Ly $\alpha$  lines: 1495,

1571, 1686, 1721, and  $1854 \text{ km s}^{-1}$ . None of these are of the correct orientation for co-rotation, and all are more likely to be associated with other nearby galaxies, such as NGC4517A, which is slightly closer to these absorbers in impact parameter and velocity than is NGC4536. The second nearby sightline is toward 3C273 at 344 kpc and  $46^\circ$  azimuth angle, and shows 3 Ly $\alpha$  lines at velocities of 1580, 2156,  $2267 \text{ km s}^{-1}$ . Two of these are correctly oriented for co-rotation, but are too high in velocity to make this scenario probable. Overall, given the number of nearby galaxies and their locations, we would expect these absorbers to trace the overall velocity field instead of the halo rotation of any particular galaxy.

#### 4.9. NGC4939

NGC4939 is a large, fast rotating ( $V_{rot} = 275 \pm 49 \text{ km s}^{-1}$ ) SA(s)bc type galaxy at systemic velocity  $V_{hel} = 3093 \pm 33 \text{ km s}^{-1}$ . We detect a single Ly $\alpha$  absorber at  $3448 \text{ km s}^{-1}$  towards PG1302-102 at 254 kpc and  $61^\circ$  azimuth angle towards the southeast. This absorber is located on the approaching side of this galaxy, so we can easily rule out co-rotation in this case. NGC4939 does not have any close neighbors, so represents strong case against co-rotation for gas near or past  $1 R_{vir}$ .

#### 4.10. NGC5364

NGC5364 is a SA(rs)bc pec type galaxy at a measured systemic velocity of  $1238 \pm 17 \text{ km s}^{-1}$ . It is located in a group environment with 5 other large, nearby galaxies. The sightline toward SDSSJ135726.27+043541.4 at 165 kpc and  $84^\circ$  azimuth angle contains Ly $\alpha$  absorbers at  $1124$  and  $1296 \text{ km s}^{-1}$  on the receding side. However, because of the orientation of NGC5364 on the sky with respect to this sightline, these absorbers lie extremely close to the inflection point where projected rotation velocities flip to approaching instead of receding. For example, shifting the location of SDSSJ135726.27+043541.4 east by a tenth of a degree ( $\sim 20 \text{ kpc}$ ) is sufficient to put these absorbers on the approaching side of NGC5364. Hence, both of these absorbers could be co-rotating with NGC5364 given very reasonable assumptions on the shape of an extended disk. Nonetheless, the fact that this system lives in galaxy group environment likely dominates the surrounding velocity field.

#### 4.11. NGC5786

NGC5786 is a large, strongly-barred spiral galaxy at  $V_{sys} = 2975 \pm 22$ . The sole nearby sightline lies along the major axis on the receding side of NGC5786 at 453 kpc toward QSO1500-4140. Unfortunately for the purpose of this exercise, there are two neighboring galaxies, both of which are closer than this. ESO327-G038 and ESO327-G039 are both located directly south at 62 kpc and 296 kpc, respectively. Toward QSO1500-4140 we detect Ly $\alpha$  absorption at  $3138 \text{ km s}^{-1}$ , corresponding to a velocity difference  $\Delta v = 163 \text{ km s}^{-1}$ . Although this velocity aligns nicely with expected co-rotation velocities, the nearby neighboring galaxies and large distance

to the absorption ( $\sim 2.5R_{vir}$ ) make it difficult to believe this as evidence of an extended disk.

#### 4.12. UGC09760

UGC09760 is an edge-on, slow-rotating Sd galaxy with a single sightline toward SDSSJ151237.15+012846.0 located 123 kpc away along the minor axis. Our measured systemic velocity,  $V_{sys} = 2094 \pm 16$ , deviates slightly from other published redshifts, such as the The Updated Zwicky Catalog value of  $V_{sys} = 2023 \pm 2$  (?). This is likely due to our method of imposing rotation symmetry and averaging the approaching and receding velocities to derive  $V_{sys}$ . If we do not sample the rotation curve far enough out, a systematic offset is not unreasonable. Indeed, we do not detect the rotation curve turnover or flattening point.

We detect Ly $\alpha$  absorption at  $2051 \text{ km s}^{-1}$  ( $\Delta v = -43 \text{ km s}^{-1}$ ). This velocity matches well with the rotation velocity of UGC09760 ( $V_{rot} = 46 \pm 16 \text{ km s}^{-1}$ ), but unfortunately this sightline lies almost exactly at an azimuth of  $90^\circ$ . Hence, the motion of this gas could easily be either co-rotating or counter-rotating depending on a minute change in the position angle assigned to UGC09760. This is especially true if we assume our measured  $V_{sys}$  is erroneously high, and indeed closer to the values obtained by other observations.

It is worth noting that there are several small satellite galaxies nearby, including SDSSJ151208.16+013508.5, SDSSJ151121.63+013637.6, SDSSJ151241.38+013723.7 and UGC09746 (impact parameters of 53, 88, 82, 230 kpc respectively). All of these galaxies lie slightly blueward of UGC09760, and thus *farther* away in velocity from the Ly $\alpha$  absorber at  $2051 \text{ km s}^{-1}$ .

### 5. ANCILLARY DATA

To increase our sample size we have also searched the literature for galaxies with published rotation curves and orientations. Unfortunately, while the rotation velocity is available for thousands of galaxies, only a handful also include the *orientation* of the rotation on the sky. Of these, we were able to find 18 additional galaxies which have a systemic velocity greater than  $\sim 500 \text{ km s}^{-1}$ , and are near to a COS or STIS sightline with available data.

#### 5.1. NGC3198

NGC3198 is a well studied SB(rs)c type galaxy which is included in the detailed THINGS rotation curve study of ?. We extracted the raw rotation curve derived by ? using the plot digitization software WebPlotDigitizer<sup>2</sup>. NGC3198 has a systemic velocity of  $660 \text{ km s}^{-1}$ , and is located 370 kpc away from a sightline toward RX.1017.5+4702 at an azimuth angle of  $55^\circ$  on the approaching side of NGC3198 (northeast). NGC3198 has an even and flat rotation curve, with an average velocity of  $v_{rot} = 152 \text{ km s}^{-1}$  and an expected velocity range of

$[-82, -23]$  based on our projected NFW profile fit. We detect Ly $\alpha$  absorption in the spectrum of RX.1017.5+4702 at  $629 \text{ km s}^{-1}$  ( $\Delta v = -31 \text{ km s}^{-1}$ ). Hence, the velocity of this absorber can nicely be described by a co-rotating disk.

SDSSJ101622.60+470643.0 is also located in the same direction but at a more distant 401 kpc away (**NOT IDENTIFIED**).

The small dwarf galaxy SDSSJ101848.77+452137.0 is located 65 kpc away toward the southwest.

#### 5.2. NGC3351

At  $V_{sys} = 778 \text{ km s}^{-1}$  NGC3351 is a mostly face-on ( $i = 29^\circ$ ) SB(r)b type galaxy located  $\sim 200$  kpc southwesterly of the core of the Leo I group. We take the rotation curve and orientation produced by ?. While we expect any extended disk rotation to be quickly disrupted due to the complex Leo I environment, this galaxy also has the closest sightline of our sample with SDSSJ104335.90+115129.0 at 31 kpc and  $13^\circ$  azimuth on the northwest, approaching side. We detect Ly $\alpha$  at 717, 882, and  $1030 \text{ km s}^{-1}$  ( $\Delta v = -61, 104, 252 \text{ km s}^{-1}$ ) toward this sightline. The lowest velocity absorber agrees nicely with both models for co-rotation, while the other two are too high in velocity.

SDSSJ104335.90+115129.0 at 31kpc, 13 az (Ly $\alpha$  at 717, 882, 1030)

SDSSJ104816.30+120735.0 at 198kpc, 85 az (G130M but not in reduce)

SDSSJ104709.80+130454.0 at 277kpc, 47 az (G130M but not in reduce)

SDSSJ104843.50+130605.0 at 317kpc, 57 az (G130M but not in reduce)

SDSSJ105220.60+101751.0 at 435kpc, 39 az (not ID'd, possible line)

SDSSJ104341.53+085558.2 at 484kpc, 18 az (G130M but not in reduce)

#### 5.3. NGC5907

NGC5907 is a large, edge-on SA(s)c type galaxy at  $V_{sys} = 667 \text{ km s}^{-1}$ . We have taken the rotation curve and orientation information from ?. The closest sightline resides toward SDSSJ152053.59+571122.1 on the northwest (receding) side, at 286 kpc and  $63^\circ$  azimuth.

SDSSJ152053.59+571122.1 at 286 kpc, 63 az (possible line around  $880 \text{ km s}^{-1}$ , messy spectrum, not ID'd)  
SBS1503+570 at 413 kpc, 47 az (Ly $\alpha$  at 708kms)  
RBS1503 at 478 kpc, 63 az (no lines)

#### 5.4. NGC4565

NGC4565 is an edge-on ( $i = 75^\circ$ ) SA(s)b type galaxy at a heliocentric velocity of  $1230 \text{ km s}^{-1}$ , for which we have taken the rotation curve and orientation from ?. The sightline toward RX.J1236.0+2641, at 147 kpc and  $41^\circ$  azimuth angle on receding side, contains Ly $\alpha$  absorbers

<sup>2</sup> WebPlotDigitizer; <http://arohatgi.info/WebPlotDigitizer>



at velocities of 1009 and 1166  $\text{km s}^{-1}$  ( $\Delta v = -221, -64 \text{ km s}^{-1}$ ). Both are consistent with counter-rotating gas, although the line at 1166  $\text{km s}^{-1}$  is close to the allowed range of  $[-29, 146] \text{ km s}^{-1}$  for co-rotation. However, the presence of several satellite galaxies surely disrupts any possible extended disk rotation that would otherwise be detectable via sightline absorption.

### 5.5. UGC06446

UGC06446 is an Sd galaxy located at  $V_{\text{sys}} = 645 \text{ km s}^{-1}$  on the far northwest edge of the Ursa Major cluster of galaxies. We take the rotation curve and orientation information from (??). We detect  $\text{Ly}\alpha$  absorption toward two nearby QSO's, SDSSJ112448.30+531818.0 at 143 kpc and  $22^\circ$  azimuth ( $\text{Ly}\alpha$  at 664 and 1019  $\text{km s}^{-1}$ ) and RX\_J1117.6+5301 at 417 kpc and  $52^\circ$  azimuth ( $\text{Ly}\alpha$  at 685  $\text{km s}^{-1}$ ), are both located toward the southwest, receding side of UGC06446. The absorbers residing at 664  $\text{km s}^{-1}$  and 685  $\text{km s}^{-1}$  both agree well with the expected velocities for co-rotating gas ( $[-15, 61]$ ,  $[25, 35]$ ). The 1019  $\text{km s}^{-1}$  absorber toward SDSSJ112448.30+531818.0 is most likely associated with NGC3718, which is much closer in velocity ( $V_{\text{sys}} = 993 \text{ km s}^{-1}$ ). Interestingly, we note that the more distant absorber toward RX\_J1117.6+5301 has a larger  $\Delta v$  value ( $\Delta v = 40 \text{ km s}^{-1}$  compared to 19  $\text{km s}^{-1}$ ). If both absorbers were strictly following the expected NFW rotation curve, we would expect the opposite, with the more distant absorber velocity approaching  $V_{\text{sys}}$ .

### 5.6. UGC06399

UGC06399 is an edge-on Sm type galaxy located at  $V_{\text{sys}} = 791 \text{ km s}^{-1}$  on the far west edge of the Ursa Major cluster of galaxies. Again, we take the rotation curve and orientation information from (??). As noted by ?, this is the most isolated galaxy in the Ursa Major cluster. Unfortunately, the closest QSO is SBS1116+523, at 471 kpc away on the northwest, approaching side. Although we detect  $\text{Ly}\alpha$  absorption at 731  $\text{km s}^{-1}$ , this impact parameter is beyond our model radius of  $3R_{\text{vir}}$ . Nonetheless, the velocity of this absorber agrees well with the expected velocity range if we extend a static or NFW rotating disk out to  $4R_{\text{vir}}$ .

### 5.7. NGC3726

The third galaxy from the Ursa Major galaxy cluster we're including is NGC3726, a SAB(r)c type galaxy at  $V_{\text{sys}} = 866 \text{ km s}^{-1}$  on the southwestern edge of the cluster (?). Two QSO's, CSO1208 and RX\_J1142.7+4625, are located at 369 and 440 kpc, respectively, southeast of NGC3726. They lie very close to and on apposing sides of the minor axis, such that CSO1208 samples the receding side and RX\_J1142.7+4625 the approaching. Unfortunately, both are also closer to a small group of dwarf galaxies, including NGC3782 and MCG+08-21-092,  $\sim 100 \text{ km s}^{-1}$  blueward of NGC3726. We detect  $\text{Ly}\alpha$  at 731 and 874  $\text{km s}^{-1}$  toward CSO1208. The

731  $\text{km s}^{-1}$  line is most likely associated with the dwarf group, although the 874  $\text{km s}^{-1}$  line falls within the expected range ( $[-28, 21]$ ) for co-rotating gas.

CSO1208 at 369 kpc, 87 az ( $\text{Ly}\alpha$  at 731, 874 kms) RX\_J1142.7+4625 (PGC139665) at 440, 86 az (G130M but not in reduce) close to CSO1208

Both sightlines closer to some other little galaxies. NGC3877 nearby, has rot curve

### 5.8. NGC3067

The galaxy-QSO pair NGC3067-3C232 is a particularly well studied system. They are separated by only 11 kpc ( $74^\circ$  azimuth angle on NW side) and a LLS of  $N_{\text{HI}} = 110^{20} \text{ cm}^{-2}$  is detected toward 3C232 at 1421  $\text{km s}^{-1}$  ( $V_{\text{sys}} = 1476 \text{ km s}^{-1}$ ), which has been postulated as a high velocity cloud (HVC) orbiting NGC3067 (??). We obtained the rotation curve for NGC3067 from ? and the orientation from ?. While H measurements of this LLS fit a single component, we have fit 3 separate components at 1408, 1510, and 1641  $\text{km s}^{-1}$  to match the associated metal lines (namely, C IV, Si II, Si III, Si IV, Mg II, Fe II, and N I all show at least 2 separate components). This splitting has been analyzed in detail most recently by ? and ?, who find similar but slightly lower  $cz$  for all three absorbers.

3C232 is located directly north of NGC3067, thus placing it on the receding side. Our NFW model predicts a co-rotating cloud to have a velocity within the range  $[-139, 26] \text{ km s}^{-1}$ . Hence, the 1408 and 1510  $\text{km s}^{-1}$  lines can be described as co-rotating (assuming reasonable velocity uncertainties), while the 1641  $\text{km s}^{-1}$  line must be either a counter-rotating clouddlet or an outflow directed away from our line of sight.

SDSSJ095914.80+320357.0 at 128 kpc, 43 az (not ID'd, but probable line around 1500 kms) RX\_J1002.9+3240 at 359 kpc, 33 az (not in reduce yet)

We have also included the 5 galaxy-QSO systems analyzed by ?. We briefly summarize each of these systems here, and refer the reader to ? for a more complete discussion. **ALL? OF THESE GALAXIES WE PROJECT TO BE AT GREATER IMPACT PARAMETER THAN COTE DID. MOSTLY BASED ON BETTER RID DISTANCES THAN USED BY COTE**

### 5.9. NGC6140

NGC6140 is a small SB(s)cd pec type galaxy located at  $V_{\text{sys}} = 910 \text{ km s}^{-1}$  and 113 kpc from the sightline toward QSO MRK876. The approaching side is oriented toward the east, with MRK876 toward the northwest at an azimuth of  $21^\circ$  (although this is somewhat uncertain; the position angle for NGC6140 could be closer to  $60^\circ$  than our adopted value of  $94^\circ$  due to it being mostly face on, faint, and strongly barred). We detect

$\text{Ly}\alpha$  absorption toward MRK876 at  $939 \text{ km s}^{-1}$  ( $\Delta v = 29 \text{ km s}^{-1}$ ). This value implies co-rotation, but is just below the range predicted by our models ( $35 - 102 \text{ km s}^{-1}$ ).

#### 5.10. NGC4529

As an inclined, isolated galaxy with a QSO sightline located 159 kpc and only  $23^\circ$  off the major axis, NGC4529 represents an ideal test environment for this study. This sightline towards MRK771 contains a  $\text{Ly}\alpha$  absorber at  $2553 \text{ km s}^{-1}$ . With  $V_{sys} = 2536 \text{ km s}^{-1}$  and MRK771 located on the approaching side of NGC4529 (southwest side), this is a clearcut case of counter-rotating H I. As ? conclude, “there is simply no physical way to produce such a velocity with an extending co-rotating disk.”

#### 5.11. UGC04238

UGC04238 is an isolated and edge-on SBd type galaxy at  $V_{sys} = 1544 \text{ km s}^{-1}$ . We take the rotation curve and orientation from ?.  $\text{Ly}\alpha$  are detected at 1143, 1526, and  $1593 \text{ km s}^{-1}$  toward PG0804+761, located directly south at 148 kpc and  $59^\circ$  azimuth on the receding side of UGC04238. Only the absorber at  $1593 \text{ km s}^{-1}$ , the lowest  $W$  of the three, falls within the expected velocity range for co-rotation.

The following are from ?:

#### 5.12. NGC2770

NGC2770 is a large, edge-on Sc type galaxy at  $V_{sys} = 1947 \text{ km s}^{-1}$ . It is mostly isolated except for two nearby small dwarfs MCG+06-20-036NED02 and GALEXASCJ090946.88+330840.4 (both 25 kpc away, on opposite sides of NGC2770). We take the rotation curve and orientation from ?. There are five nearby QSOs. Toward the south on the approaching side they are the following: FBQJ0908+3246 at 204 kpc and  $59^\circ$  azimuth ( $\text{Ly}\alpha$  at 1915,  $1982 \text{ km s}^{-1}$ ), TON1009 at 267 kpc and  $41^\circ$  azimuth (**NEEDS TO BE IDENTIFIED**), and SDSSJ091127.30+325337.0 at 234 kpc and  $30^\circ$  azimuth ( $\text{Ly}\alpha$  at  $2063 \text{ km s}^{-1}$ ). Only the 1915  $\text{km s}^{-1}$  line can reasonably be explained by co-rotation.

Toward the northeast on the receding side they are the following: TON1015 at 218 kpc and  $61^\circ$  azimuth ( $\text{Ly}\alpha$  at 1833,  $1985 \text{ km s}^{-1}$ ), and SDSSJ091052.80+333008.0 at 239 kpc and  $66^\circ$  azimuth ( $\text{Ly}\alpha$  at 1824,  $1975 \text{ km s}^{-1}$ ). Two of these lines, at  $1985 \text{ km s}^{-1}$  and  $1975 \text{ km s}^{-1}$  are consistent with co-rotation. Interestingly, these two QSO’s appear to be sampling the same structures. The absorber pairs at 1833,  $1824 \text{ km s}^{-1}$  and 1985,  $1975 \text{ km s}^{-1}$  each have the equivalent  $EW$  and  $N_{H I}$  within errors, and remarkably similar appearing line-structure. Adopting a distance of 28.6 Mpc to this cloud, we calculate a linear separation between TON1015 and SDSSJ091052.80+333008.0 of 28 kpc. Hence, there appears to be two distinct clouds of at least 28 kpc in physical extent sandwiched around the system velocity of NGC2770. **MORE?**

#### 5.13. NGC3432

NGC3432 is an edge-on SB(s)m type galaxy at  $V_{sys} = 616 \text{ km s}^{-1}$ . It is interacting with the nearby dwarf galaxy UGC05983 located 11 kpc away and at  $V_{sys} = 765 \text{ km s}^{-1}$ . We take a rotation curve and orientation for NGC3432 from ?. The quasar CSO295 is located just 20 kpc away and just to the receding side of the minor axis. This is the second closest pair in our sample, after the 11 kpc separated NGC3067-3C232 system. We detect  $\text{Ly}\alpha$  toward CSO295 at 600 and  $662 \text{ km s}^{-1}$ . Both of these lines fall well within our expected range for co-rotation ( $[-37, 134] \text{ km s}^{-1}$  for an NFW halo). This would represent the lower-velocity cloud existing toward the near-edge of the halo and the higher velocity cloud lying very close to the plane of the stellar disk.

CSO295 at 20 kpc, 82 az ( $\text{Ly}\alpha$  600, 662),  $[-37, 134]$ NFW  
 RXJ1054.2+3511 at 290 kpc, 57 az (Looks like a feature around  $600 \text{ km s}^{-1}$  but no ID?)  
 MS1047.3+3518 at 326 kpc, 26 az (low SN, not ID’d, possible feature?)  
 SDSSJ110349.70+371527.0 (low res only)

#### 5.14. NGC3666

NGC3666 is a mostly isolated and edge-on SA(rs)c type galaxy at  $V_{sys} = 1060 \text{ km s}^{-1}$ . We take a rotation curve and orientation from ?. The following three QSOs are located toward on the approaching, northeast side: SDSSJ112439.50+113117.0 at 58 kpc, SDSSJ112632.90+120437.0 at 279 kpc, and SDSSJ112756.70+115427.0 at 320 kpc. There are several dwarf galaxies  $\sim 400 \text{ kpc}$  away, but all are more distant than these three QSOs. We detect  $\text{Ly}\alpha$  toward SDSSJ112439.50+113117.0 at 1047 and  $1099 \text{ km s}^{-1}$  ( $\Delta v = -13, 39 \text{ km s}^{-1}$ ).

SDSSJ112439.50+113117.0 at 58 kpc - Borthakur target ( $\text{Ly}\alpha$  at 1047,  $1099 \text{ km s}^{-1}$ ),  $[-87, 20]$ ,  $[-136, 20]$ NFW  
 SDSSJ112632.90+120437.0 at 279 kpc - Borthakur target (not in reduce yet)  
 SDSSJ112756.70+115427.0 at 320 kpc - (not in reduce yet)

NW side is receding

#### 5.15. NGC5951

NGC5951 is a large, edge-on SBc type galaxy at  $V_{sys} = 1780 \text{ km s}^{-1}$ . The pair of galaxies NGC5954 and NGC5953 are nearby ( $\sim 100 \text{ kpc}$ ), but the sightline toward 2E1530+1511 is closer ( $\rho = 55 \text{ kpc}$  straight east of NGC5951) and on the opposite side ( $85^\circ$  azimuth). We take the rotation curve and orientation for NGC5951 from ?.  $\text{Ly}\alpha$  appears toward 2E1530+1511 at 1795 and  $1953 \text{ km s}^{-1}$  ( $\Delta v = 15, 173 \text{ km s}^{-1}$ ). The absorber



at  $1795 \text{ km s}^{-1}$  aligns well with co-rotation, but the  $1953 \text{ km s}^{-1}$  absorber has a great velocity than  $V_{rot}$  for NGC5951. Given the  $V_{sys}$  for the nearby galaxies NGC5954 and NGC5953 are 1959 and  $1965 \text{ km s}^{-1}$ , this absorber is likely also linked with that system.

#### 5.16. NGC7817

NGC7817 is an edge-on SAbc type galaxy at  $V_{sys} = 2309 \text{ km s}^{-1}$ . We take a rotation curve and orientation from ?. The dwarf galaxy ESDOF538-02 is the closest neighbor, at 57 kpc to the east. A sightline toward MRK335 is located 343 kpc toward the southeast, lying directly on the projected minor axis of NGC7817. Because of this our model predicts a narrow range of velocities for co-rotation ( $[-26, -24] \text{ km s}^{-1}$ ).  $\text{Ly}\alpha$  absorbers are detected at  $1954$  and  $2274 \text{ km s}^{-1}$ , the latter of which falling a mere  $9 \text{ km s}^{-1}$  outside this predicted range. Within a very reasonable level of uncertainty this  $2274 \text{ km s}^{-1}$  absorber could easily be co-rotating with NGC7817. The absorption at  $1954 \text{ km s}^{-1}$  is likely not directly associated with NGC7817, given the large velocity difference.

#### 5.17. UGC08146

UGC08146 is an isolated and edge-on Sd type galaxy at  $V_{sys} = 670 \text{ km s}^{-1}$ . This galaxy (and the nearby QSO PG1259+593) are included in the ? sample also, but we have taken the rotation curve and orientation from ?. PG1259+593 is located 114 kpc northwest of UGC08146, at  $50^\circ$  azimuth on the receding side. While ? cite a single  $\text{Ly}\alpha$  component at  $679 \text{ km s}^{-1}$ , we find two components, at  $646$  and  $683 \text{ km s}^{-1}$ , in the higher signal-to-noise COS data now available for PG1259+593. Both of these components are consistent with co-rotating gas, assuming reasonable uncertainty in  $V_{sys}$ ,  $V_{\text{Ly}\alpha}$ , and  $V_{rot}$ .

### 6. DISCUSSION

### 7. SUMMARY

### 8. SIGHTLINES NEEDED

The following targets need to be added:

SBS1304+594  
SDSSJ112632.90+120437.0  
SDSSJ112756.70+115427.0  
RX\_J1054.2+3511 - check for feature around  $600 \text{ km s}^{-1}$   
MS1047.3+3518  
TON1009  
SDSSJ095914.80+320357.0  
RX\_J1002.9+3240  
RX\_J1142.7+4625  
SDSSJ152053.59+571122.1  
SDSSJ104816.30+120735.0  
SDSSJ104709.80+130454.0  
SDSSJ104843.50+130605.0  
SDSSJ105220.60+101751.0  
SDSSJ104341.53+085558.2  
SDSSJ101622.60+470643.0

D. M. F. thanks Claire Murray for useful insights on our halo model, and Julie Davis for invaluable SALT data reduction pointers. This research has made use of the NASA/IPAC Extragalactic Database (NED) which is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration. Based on observations with the NASA/ESA *Hubble Space Telescope*, obtained at the Space Telescope Science Institute (STScI), which is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS 5-26555. **SALT ACKNOWLEDGEMENT.** Spectra were retrieved from the Barbara A. Mikulski Archive for Space Telescopes (MAST) at STScI. Over the course of this study, D.M.F. and B.P.W. were supported by grant AST-1108913, awarded by the US National Science Foundation, and by NASA grants *HST*-AR-12842.01-A, *HST*-AR-13893.01-A, and *HST*-GO-14240 (STScI).


*Facility:* HST (COS)

## APPENDIX


### A. ROTATION CURVES

Here we present rotation curves with finder charts indicating the slit position for each galaxy observed with SALT.


Galaxy	Target	$\rho$	Az.	$V_{sys}$	$V_{rot}$	$V_{Ly\alpha}$	$W_{Ly\alpha}$	Cyl. Model	NFW Model
		[kpc]	[Deg.]	[km s <sup>-1</sup> ]	[km s <sup>-1</sup> ]	[km s <sup>-1</sup> ]	[mÅ]	[km s <sup>-1</sup> ]	[km s <sup>-1</sup> ]
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)
CGCG039-137	RX_J1121.2+0326	99	71	6917.8	139	6975	678	[-36, 137]	[-37, 164]
CGCG039-137	SDSSJ112224.10+031802.0	491	24	6917.8	-139	6606	70	[-129, -115]	[-115, -99]
ESO343-G014	RBS1768	466	74	9138.9	205	9308	63	[-203, 10]	[-122, 31]
ESO343-G014	RBS1768	466	74	9138.9	205	9360	306	[-203, 10]	[-122, 31]
ESO343-G014	RBS1768	466	74	9138.9	205	9434	161	[-203, 10]	[-122, 31]
IC5325	RBS2000	314	64	1511.9	-125	1598	35	[-41, -20]	[-29, 1]
MCG-03-58-009	MRC2251-178	355	71	9014.9	171	9029	62	[-26, 137]	[-42, 83]
NGC1566	1H0419-577	303	10	1501.9	86	1075	249	[48, 76]	[-2, 31]
NGC1566	1H0419-577	303	10	1501.9	86	1123	269	[48, 76]	[-2, 31]
NGC1566	1H0419-577	303	10	1501.9	86	1188	240	[48, 76]	[-2, 31]
NGC1566	1H0419-577	303	10	1501.9	86	1264	91	[48, 76]	[-2, 31]
NGC1566	1H0419-577	303	10	1501.9	86	2020	9	[48, 76]	[-2, 31]
NGC1566	HE0429-5343	256	60	1501.9	-86	1167	79	[-53, -2]	[-22, 17]
NGC1566	HE0429-5343	256	60	1501.9	-86	1358	136	[-53, -2]	[-22, 17]
NGC1566	HE0435-5304	396	62	1501.9	-86	1512	220	[-44, -43]	[-16, -15]
NGC1566	HE0435-5304	396	62	1501.9	-86	1633	217	[-44, -43]	[-16, -15]
NGC1566	HE0435-5304	396	62	1501.9	-86	1690	204	[-44, -43]	[-16, -15]
NGC1566	RBS567	423	69	1501.9	-86	1664	517	[-46, -7]	[-22, 10]
NGC1566	HE0439-5254	459	65	1501.9	-86	1148	72	[-50, -22]	[-23, 1]
NGC1566	HE0439-5254	459	65	1501.9	-86	1649	501	[-50, -22]	[-23, 1]
NGC3513	H1101-232	60	67	1203.7	20	1182	635	[-19, 27]	[-19, 28]
NGC3633	SDSSJ112224.10+031802.0	413	50	2587.2	-157	-99	-99	[-158, -51]	[-64, 0]
NGC3633	SDSSJ112005.00+041323.0	468	78	2587.2	-157	2285	27	[-59, -55]	[-36, -28]
NGC3633	SDSSJ112005.00+041323.0	468	78	2587.2	-157	2578	116	[-59, -55]	[-36, -28]
NGC3633	RX_J1121.2+0326	184	58	2587.2	-157	2605	180	[-153, -14]	[-77, 10]
NGC4536	3C273.0	349	11	1866.9	139	1580	369	[87, 121]	[5, 41]
NGC4536	3C273.0	349	11	1866.9	139	2156	42	[87, 121]	[5, 41]
NGC4536	3C273.0	349	11	1866.9	139	2267	27	[87, 121]	[5, 41]
NGC4536	HE1228+0131	338	51	1866.9	139	1495	160	[18, 51]	[2, 32]
NGC4536	HE1228+0131	338	51	1866.9	139	1571	23	[18, 51]	[2, 32]
NGC4536	HE1228+0131	338	51	1866.9	139	1686	321	[18, 51]	[2, 32]
NGC4536	HE1228+0131	338	51	1866.9	139	1721	303	[18, 51]	[2, 32]
NGC4536	HE1228+0131	338	51	1866.9	139	1854	78	[18, 51]	[2, 32]
NGC4536	HE1228+0131	338	51	1866.9	139	2311	343	[18, 51]	[2, 32]
NGC4939	PG1302-102	254	61	3092.8	-275	3448	72	[-219, 14]	[-119, 36]
NGC5364	SDSSJ135726.27+043541.4	165	84	1238	55	1124	83	[-26, 108]	[-30, 68]
NGC5364	SDSSJ135726.27+043541.4	165	84	1238	55	1296	92	[-26, 108]	[-30, 68]
NGC5786	QSO1500-4140	453	1	2974.6	172	3141	177	[106, 160]	[19, 67]
UGC09760	SDSSJ151237.15+012846.0	123	90	2093.7	-46	2051	496	[-30, 30]	[-30, 86]
NGC3198	RX_J1017.5+4702	370	55	660	152	629	71	[-153, -21]	[-91, 6]
NGC3198	SDSSJ101622.60+470643.0	401	61	660	152	-99	-99	[-154, -17]	[-89, 8]
NGC4565	RX_J1236.0+2641	147	41	1230	253	1009	365	[-2, 246]	[-30, 144]
NGC4565	RX_J1236.0+2641	147	41	1230	253	1166	305	[-2, 246]	[-30, 144]
UGC04238	PG0804+761	148	59	1544	91.6	1143	183	[-3, 86]	[-6, 94]
UGC04238	PG0804+761	148	59	1544	91.6	1526	62	[-3, 86]	[-6, 94]
UGC04238	PG0804+761	148	59	1544	91.6	1593	32	[-3, 86]	[-6, 94]
NGC3351	SDSSJ104335.90+115129.0	31	43	778	198	717	823	[-99, 12]	[-68, 20]
NGC3351	SDSSJ104335.90+115129.0	31	43	778	198	882	621	[-99, 12]	[-68, 20]
NGC3351	SDSSJ104335.90+115129.0	31	43	778	198	1030	391	[-99, 12]	[-68, 20]
NGC4529	MRK771	159	23	2536	106.4	2553	240	[-103, -40]	[-105, -35]
NGC6140	MRK876	113	21	910	138.11	939	379	[40, 101]	[35, 102]
NGC5907	SBS1503+570	413	47	667	227.4	708	301	[31, 228]	[-24, 101]
NGC5907	SDSSJ152053.59+571122.1	286	63	667	227.4	-99	-99	[-18, 228]	[-42, 114]
NGC5907	RBS1503	478	63	667	227.4	-99	-99	[-228, -9]	[-96, 33]
UGC06446	RX_J1117.6+5301	417	52	645	79.4	685	342	[35, 47]	[24, 35]
UGC06446	SDSSJ112448.30+531818.0	143	22	645	79.4	664	339	[-9, 65]	[-15, 61]
UGC06446	SDSSJ112448.30+531818.0	143	22	645	79.4	1019	71	[-9, 65]	[-15, 61]
UGC06399	SBS1116+523	471	13	791	88.2	731	259	[-91, -54]	[-80, -42]
NGC3726	CSO1208	369	88	866	167.2	731	470	[-27, 29]	[-28, 21]
NGC3726	CSO1208	369	88	866	167.2	874	506	[-27, 29]	[-28, 21]
NGC3726	RX_J1142.7+4625	440	86	866	167.2	818	375	[-34, -14]	[-30, -7]





**Figure 1.** A map of the locations of each absorber normalized with respect to the galaxy virial radius. Concentric rings indicate distances of 1, 2, 3 and 4  $R_{vir}$ . All galaxies are rotated to  $PA = 90$  or  $270$ , such that their major axis' are horizontal and their approaching side is on the left as indicated. The color and style of each point indicates the line-of-sight velocity compared to that of the rotation of the nearby galaxy. Green diamonds indicate co-rotation, red X's indicate anti-rotation, and purple circles indicate cases where either is possible due to a combination of orientation and velocity uncertainties. The size of each point is scaled to reflect the EW of the absorber. The labeled galaxy-QSO pairs are as follows: 1. NGC3067 : 3C232, 2. NGC3351 : SDSSJ104335.90 + 115129.0, 3. CGCG039 - 137 : RXJ1121.2 + 0326, 4. NGC3513 : H1101 - 232, 5. NGC1566 : RBS567, 6. NGC3726 : CSO1208, 7. NGC1566 : HE0439 - 5254, 8. UGC09760 : SDSSJ151237.15 + 012846.0, 9. NGC6140 : MRK876, 10. NGC3726 : RXJ1142.7 + 4625, 11. NGC4536 : 3C273.0, 12. NGC4565 : RXJ1236.0 + 2641, 13. NGC4536 : HE1228 + 0131, 14. UGC06446 : RXJ1117.6 + 5301, 15. UGC06446 : SDSSJ112448.30 + 531818.0, 16. ESO343 - G014 : RBS1768, 17. NGC5907 : SBS1503 + 570, 18. NGC2770 : SDSSJ091127.30 + 325337.0, 19. NGC1566 : 1H0419 - 577, 20. NGC2770 : SDSSJ091052.80 + 333008.0, 21. UGC06399 : SBS1116 + 523, 22. NGC2770 : TON1015, 23. NGC4529 : MRK771, 24. NGC2770 : FBQSJ0908+3246, 25. NGC1566 : HE0435 - 5304, 26. UGC04238 : PG0804+761, 27. NGC3633 : RXJ1121.2+0326, 28. NGC5786 : QSO1500 - 4140, 29. NGC1566 : HE0429 - 5343, 30. NGC3633 : SDSSJ112005.00 + 041323.0, 31. NGC4536 : LBQS1230 - 0015, 32. NGC5364 : SDSSJ135726.27 + 043541.4, 33. NGC4939 : PG1302 - 102, 34. NGC3198 : RXJ1017.5+4702, 35. CGCG039 - 137 : SDSSJ112224.10 + 031802.0, 36. MCG - 03 - 58 - 009 : MRC2251 - 178, 37. IC5325 : RBS2000, 38. NGC3198 : SDSSJ101622.60 + 470643.0, 39. NGC5907 : SDSSJ152053.59 + 571122.1, 40. NGC5907 : RBS1503, 41. NGC3067 : SDSSJ095914.80 + 320357.0, 42. NGC3067 : RXJ1002.9 + 3240, 43. NGC2770 : TON1009





**Figure A1.** a) Rotation curve of CGCG039-137. The solid green line indicates the weighted mean velocity over the corresponding x-axis region, and the shaded green indicates the  $1\sigma$  error in the mean. b) SALT finder chart for CGCG039-137 showing the position of the slit in red.




**Figure A2.** a) Rotation curve of IC5325. The solid green line indicates the weighted mean velocity over the corresponding x-axis region, and the shaded green indicates the  $1\sigma$  error in the mean. b) SALT finder chart for IC5325 showing the position of the slit in red.




**Figure A3.** a) Rotation curve of MCG-03-58-009. The solid green line indicates the weighted mean velocity over the corresponding x-axis region, and the shaded green indicates the  $1\sigma$  error in the mean. b) SALT finder chart for MCG-03-58-009 showing the position of the slit in red.



**Figure A4.** a) Rotation curve of NGC1566. The solid green line indicates the weighted mean velocity over the corresponding x-axis region, and the shaded green indicates the  $1\sigma$  error in the mean. b) SALT finder chart for NGC1566 showing the position of the slit in red.





**Figure A5.** a) Rotation curve of NGC3513. The solid green line indicates the weighted mean velocity over the corresponding x-axis region, and the shaded green indicates the  $1\sigma$  error in the mean. b) SALT finder chart for NGC3513 showing the position of the slit in red.





**Figure A6.** a) Rotation curve of NGC4536. The solid green line indicates the weighted mean velocity over the corresponding x-axis region, and the shaded green indicates the  $1\sigma$  error in the mean. b) SALT finder chart for NGC3633 showing the position of the slit in red.







**Figure A7.** a) Rotation curve of NGC4536. The solid green line indicates the weighted mean velocity over the corresponding x-axis region, and the shaded green indicates the  $1\sigma$  error in the mean. b) SALT finder chart for NGC4536 showing the position of the slit in red.





**Figure A8.** a) Rotation curve of NGC4939. The solid green line indicates the weighted mean velocity over the corresponding x-axis region, and the shaded green indicates the  $1\sigma$  error in the mean. b) SALT finder chart for NGC4939 showing the position of the slit in red.



**Figure A9.** a) Rotation curve of NGC5364. The solid green line indicates the weighted mean velocity over the corresponding x-axis region, and the shaded green indicates the  $1\sigma$  error in the mean. b) SALT finder chart for NGC5364 showing the position of the slit in red.



**Figure A10.** a) Rotation curve of NGC5786. The solid green line indicates the weighted mean velocity over the corresponding x-axis region, and the shaded green indicates the  $1\sigma$  error in the mean. b) SALT finder chart for NGC5786 showing the position of the slit in red.



**Figure A11.** a) Rotation curve of NGC5364. The solid green line indicates the weighted mean velocity over the corresponding x-axis region, and the shaded green indicates the  $1\sigma$  error in the mean. b) SALT finder chart for NGC5364 showing the position of the slit in red.