

MATCHING LY α ABSORPTION TO NEARBY GALAXY HALOS WITH A LIKELIHOOD-BASED METHOD*

DAVID M. FRENCH, BART P. WAKKER
 Department of Astronomy, University of Wisconsin - Madison
Draft version October 4, 2016

ABSTRACT

We present initial results from an ongoing large-scale study of the circumgalactic medium in the nearby Universe ($cz \leq 10,000$ km/s), using archival Cosmic Origins Spectrograph (COS) spectra of background QSOs. This initial sample contains 33 sight lines chosen for their proximity to large galaxies ($D \geq 25$ kpc) and high signal-to-noise ratio ($S/N \geq 10$), yielding 48 Ly α systems which we have paired with individual galaxies. We introduce a likelihood parameter to facilitate the matching of galaxies to absorption lines in a reproducible manner. We find the usual anti-correlation between Ly α equivalent width (EW) and impact parameter (ρ) when we normalize by galaxy virial radius (R_{vir}). Galaxies associated with a Ly α absorber are found to be more highly-inclined than the average distribution of galaxies in the survey volume at a $> 99\%$ confidence level (3σ). Contrary to suggestions in other recent papers, we do not see obvious correlations with azimuth angle.

Subject headings: IGM, CGM, galaxies

1. INTRODUCTION

It is well known that galaxies must continue to accrete gas throughout their lifetimes in order to sustain their observed levels of star formation (e.g. Erb 2008, Putman et al. 2009b). This additional gas must come from the diffuse intergalactic medium (IGM), where the majority of the baryons in the universe reside (Penton et al. 2002, 2004; Lehner et al. 2007; Danforth & Shull 2008; Shull et al. 2012). How exactly this IGM gas eventually falls into the halos and disks of galaxies is still highly uncertain, as observational constraints are hard to come by. Because of the diffuse nature of IGM gas it is most readily and sensitively detected as absorption in the spectra of background active galactic nuclei (AGN). The advent of the sensitive UV spectrographs STIS and COS on the Hubble Space Telescope (HST) has provided a wealth of information on the properties and distribution of both the ions of heavy elements as well as the Lyman series of neutral H I gas around galaxies.

Individual concentrations of gas along a given sightline imprint absorption lines on the spectrum in the direction of the QSO. The metal lines trace the star formation history within the intervening gas, and neutral hydrogen lines (Ly α) indicate both the location and velocities of outflowing gas, as well as the presence of fuel for future star formation. Numerous studies using these observations have shown that many Ly α absorbers trace individual galaxy halos (e.g. Lanzetta et al 1995, Tripp et al. 1998, Chen et al. 1998, 2001a, Wakker & Savage 2009, Steidel et al. 2010, Prochaska et al. 2011, Thom et al 2012, Tumlinson et al. 2011 & 2013, Stocke et al. 2013 & 2014, Danforth et al. 2014, Liang et al 2014).

Some recent studies find that about half of Ly α absorbers lie within galaxy haloes, at impact parameters

$\rho < 350$ kpc (Côté et al. 2005, Prochaska et al. 2006, Wakker & Savage 2009). In addition, Wakker & Savage (2009) find that an absorber lies within 400 kpc and 400 km/s for 90% of galaxies brighter than $0.1L_*$, and all galaxies have a Ly α absorber within 1.5 Mpc. Higher redshift studies, such as Rudie et al. (2012) at $2 < z < 3$, find evidence for an elevated density of absorbers up to 2 Mpc from galaxies. Wakker & Savage (2009) also discovered a correlation between Ly α absorption linewidth and impact parameter ρ , observing that the broadest lines ($FWHM > 150$ km/s) are only seen within 350 kpc of a galaxy, while at $\rho > 1$ Mpc, only lines with $FWHM < 75$ km/s occur. This suggests that the temperature and/or turbulence of gas increases in the presence of galaxies.

In addition, studying the enrichment of galaxy halos is necessary for constraining outflow models and informing stellar feedback prescriptions. Directly measuring the velocity field and column densities of absorbers as a function of impact parameter and orientation around galaxies would provide the clearest evidence of inflow or outflow activity, but results are still uncertain. Kacprzak et al. (2011) claim to find that Mg II equivalent widths correlate with galaxy inclination, but Mathes et al. (2014) find no such correlation for Ly α and O VI absorbers. Furthermore, we should expect outflowing gas to be more highly enriched and trace the metallicity of the associated galaxy, with inflowing gas instead appearing only in H I. Both Stocke et al. (2013) and Liang & Chen (2014) find an “edge” to heavy ion absorption at $\sim 0.5R_{vir}$, but with Ly α covering fractions of $\sim 0.75 - 1$ continuing out to R_{vir} . However, Mathes et al. (2014) measures O VI absorption out to $\sim 3R_{vir}$, and Savage et al. (2014) find that more than half of O VI absorption occurs beyond $1 R_{vir}$ from the nearest galaxy.

Recent results from Kacprzak et al. (2011 & 2012) suggest that absorbing systems have a preferred orientation with respect to the major and minor axes of the galaxies they are associated with. This could be evidence of inflows and outflows, or an effect of the global structure of galaxy halos, but the statistics are not yet good enough to provide consistent answers. A larger-scale study of in-

* This research has made use of the NASA/IPAC Extragalactic Database (NED) which is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration. Based on observations with the NASA/ESA *Hubble Space Telescope*, obtained at the Space Telescope Institute, which is operated by AURA, Inc., under NASA contract NAS 5-26555.

clination and azimuthal angles vs. absorber properties is needed in order to elucidate the distribution of absorbing systems around galaxies. This is most easily done for the largest galaxies in the nearby universe, where it is possible to obtain inclinations and unambiguous absorber associations.

Previous studies have suffered from small sample sizes (e.g. Mathes et al. 2014 use 14 galaxies, Stocke et al. 2013 use 11, Werk et al. 2014 use 44), and incompleteness due to their higher mean redshifts (e.g. the Mathes et al. 2014 sample is $0.12 < z < 0.67$, and Werk et al. 2014 are complete to $\sim L^*$ at $z \sim 0.2$). To address these shortcomings, we are conducting a large survey of the properties of intergalactic gas in the nearby universe, where we have good and relatively complete information on both faint and bright galaxies, in order to reveal how the IGM and galaxies affect each other. We are taking advantage of the over 500 archived QSO and Seyfert spectra taken by the Cosmic Origins Spectrograph (COS) and Space Telescope Imaging Spectrograph (STIS) on the Hubble Space Telescope (HST), combined with the wealth of information available for the $\sim 100,000$ galaxies with $cz < 10,000$ km/s found in the NASA Extragalactic Database (NED) to probe the environment of absorbing gas systems in the nearby universe. This approach allows for an unbiased understanding of the distribution of the gas around galaxies, which requires looking for both detections and non-detections of gas, both near as well as far away from galaxies.

This paper presents initial results from our pilot study of 33 sight lines, chosen for their proximity to large galaxies and high signal-to-noise spectra. This paper is organized as follows: in Section 2 we present the data and analysis techniques, in Section 3 we present the results, and in Section 4 we discuss possible interpretations of our results.

2. DATA AND ANALYSIS

2.1. Galaxy Data

Achieving the goal of this study relies on knowing the locations and properties of all galaxies near detected Ly α absorption lines. To facilitate this, we have constructed a database of all $z \leq 0.033$ ($cz \leq 10,000$ km/s) galaxies with published data available through the NASA Extragalactic Database (NED). A full description of this catalog will be presented in French & Wakker 2017 (in prep). Here we summarize its important aspects.

The galaxy dataset contains over 108,000 entries, and includes data from SDSS, 2MASS, 2dF, 6dF, RC3, and many other, smaller surveys. Our criteria for including a galaxy in this dataset is only an accurate, spectroscopic redshift which places the galaxy in the $cz \leq 10,000$ km/s velocity range. This restriction leads to a completeness limit of $B \lesssim 18.7$ mag, or $\sim 0.2L_*$, at $cz = 10,000$ km/s, and progressively better towards lower velocities (see Figure 1). This limit will vary depending on which major surveys include a particular region of the sky. The major contributor is whether or not SDSS data is available, which begins around $cz = 5,000$ km/s. Figure 1 is split into 4 velocity bins to illustrate this. Our data is complete down to $\sim 0.1L_*$ in the first bin, $0 \leq cz \leq 2,500$ km/s. At slightly higher velocity, $2500 \leq cz \leq 6000$ km/s, the completeness falls to barely better than $\sim 1.0L_*$ as




Figure 1. Distribution of L/L_* values for all galaxies in the dataset. Black vertical lines highlight 1, 0.5, 0.1, 0.05 and 0.01 L_* . The turnoff around $0.1L_*$ shows that on average, the dataset is mostly complete to $0.2L_*$.

we move past the near and well studied galaxies, but have yet to reach the footprint of deep all sky surveys. SDSS data becomes available in the last two bins, spanning $6000 \leq cz \leq 10,000$ km/s, and correspondingly completeness remains high down to the SDSS limits of $B \lesssim 18.7$ mag, or $\sim 0.2L_*$ at $cz = 10,000$ km/s.

Additionally, we have homogenized the galaxy data beyond the steps taken by NED by normalizing all measurements of galaxy inclination, position angle, and diameter to 2MASS K -band values. Most galaxies in NED have measures of inclination, position angle and diameter available in several different bands, so in order to make meaningful comparisons it is necessary to automatically choose one band for all measurements. We chose 2MASS values for this because it was an all-sky survey, and represents the largest fraction of available galaxy data. Physical galaxy diameters are derived from 2MASS K_s “total” angular diameter measurements and galaxy distances. 2MASS K_s “total” diameter estimates are surface brightness extrapolation measurements and are derived as

$$r_{tot} = r' + a(\ln(148))^b, \quad (1)$$

where r_{tot} is defined as the point where the surface brightness extends to 5 disk scale lengths, r' is the starting point radius ($> 5'' - 10''$ beyond the nucleus, or core influence), and a and b are Sersic exponential function scale length parameters ($f = f_0 \exp(-r/a)^{(1/b)}$, see Jarret et al. 2003 for a full description). Approximately 50% of all the galaxies have this 2MASS K_s “total” diameter. Of the remainder, 20% have SDSS diameters, 27% have no published diameter, and 3% have diameters from other surveys. We convert values in these other bands to 2MASS K_s “total” diameters via a simple least squares linear fit when necessary.

We used B -band magnitudes to estimate each galaxy’s luminosity in units of L_* as follows:

$$\frac{L}{L_*} = 10^{-0.4(M_B - M_{B_*})}. \quad (2)$$

We adopt the CfA galaxy luminosity function by Marzke et al. (1994), which sets $B_* = -19.57$. Direct B

Target	R.A.	Dec.	z	Program	Grating	Obs ID	Obs Date	T_{exp}^* [ks]	S/N* [1238]
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)
1H0717+714	7.0 21.0 53.3	71.0 20.0 36.0	0.5003	12025	G130M	LBG812	11-12-27	6.0	37
1H0717+714	7.0 21.0 53.3	71.0 20.0 36.0	0.5003	12025	G160M	LBG812	11-12-27	8.3	31
2dFGRS_S393Z082	2.0 45.0 0.8	-30.0 7.0 23.0	0.3392	12988	G130M	LC1040	13-05-27,28	17.7	10
						LC1045			
FBQJ1431+2442	14.0 31.0 25.8	24.0 42.0 20.0	0.4069	13342	G130M	LC8903	15-03-29	16.5	17
				12603		LBS314	13-03-08		
H1101-232	11.0 3.0 37.7	-23.0 29.0 31.0	0.1860	12025	G130M	LBG804	11-07-05	13.3	16
H1101-232	11.0 3.0 37.7	-23.0 29.0 31.0	0.1860	12025	G160M	LBG805	11-07-06	13.3	10
HE0241-3043	2.0 43.0 37.7	-30.0 30.0 48.0	0.6693	12988	G130M	LC1070	13-06-21	7.0	14
LBQS1230-0015	12.0 33.0 4.1	-0.0 31.0 34.0	0.4709	11598	G130M	LB5N15	10-08-01	10.3	13
				12486		LBP250	12-04-26		
LBQS1230-0015	12.0 33.0 4.1	-0.0 31.0 34.0	0.4709	11598	G160M	LB5N15	10-08-01	5.9	7
MRC2251-178	22.0 54.0 5.9	-17.0 34.0 55.0	0.0661	12029	G130M	LBGB03	11-09-29	5.5	42
MRC2251-178	22.0 54.0 5.9	-17.0 34.0 55.0	0.0661	12029	G160M	LBGB03	11-09-29	7.1	30
MRK290	15.0 35.0 52.3	57.0 54.0 9.0	0.0296	11524	G130M	LB4Q02	09-10-28	3.9	38
MRK290	15.0 35.0 52.3	57.0 54.0 9.0	0.0296	11524	G160M	LB4Q02	09-10-28	4.8	18
MRK876	16.0 13.0 57.2	65.0 43.0 10.0	0.1290	11524	G130M	LB4Q03	10-04-08,09	12.6	65
				11686		LB4F05			
MRK876	16.0 13.0 57.2	65.0 43.0 10.0	0.1290	11686	G160M	LB4F05	10-04-09,10	11.8	44
MRK1014	1.0 59.0 50.2	0.0 23.0 41.0	0.1630	12569	G130M	LBP404	12-01-25	1.8	17
PG0832+251	8.0 35.0 35.9	24.0 59.0 41.0	0.3310	12025	G130M	LBG808	12-04-19	6.1	14
PG0832+251	8.0 35.0 35.9	24.0 59.0 41.0	0.3310	12025	G160M	LBG808	12-04-19	6.8	9
PG0003+158	0.0 5.0 59.3	16.0 9.0 49.0	0.4509	12038	G130M	LBGL17	11-10-22	10.4	25
PG0003+158	0.0 5.0 59.3	16.0 9.0 49.0	0.4509	12038	G160M	LBGL27	11-10-22	10.9	20
PG1001+054	10.0 4.0 20.1	5.0 13.0 1.0	0.1610	13347	G130M	LCCV02	14-06-19	5.2	14
				13423		LC9W02	14-04-04		
PG1001+054	10.0 4.0 20.1	5.0 13.0 1.0	0.1610	13347	G160M	LCCV02	14-06-18,19	3.8	10
PG1302-102	13.0 5.0 33.0	-10.0 33.0 20.0	0.2784	12038	G130M	LBGL04	11-08-16	6.0	27
PG1302-102	13.0 5.0 33.0	-10.0 33.0 20.0	0.2784	12038	G160M	LBGL04	11-08-16	6.9	34
RBS1768	21.0 38.0 49.7	-38.0 28.0 40.0	0.1830	12936	G130M	LC1201	13-06-25	7.0	24
RBS1768	21.0 38.0 49.7	-38.0 28.0 40.0	0.1830	12936	G160M	LC1201	13-06-25	3.8	11
RX_J0714.5+7408	7.0 14.0 36.2	74.0 8.0 11.0	0.3710	12275	G130M	LBH402	11-03-18	8.3	18
RX_J1017.5+4702	10.0 17.0 30.9	47.0 2.0 25.0	0.3354	13314	G130M	LC9M04	14-01-29	8.7	12
RX_J1117.6+5301	11.0 17.0 40.5	53.0 1.0 50.0	0.1587	14240	G130M	LCWM05	16-04-13	4.9	11
RX_J1236.0+2641	12.0 36.0 4.1	26.0 41.0 36.0	0.2092	12248	G130M	LBH087	12-01-29	4.2	11
RX_J1236.0+2641	12.0 36.0 4.1	26.0 41.0 36.0	0.2092	12248	G160M	LBH087	12-01-29	6.8	6
RX_J1330.8+3119	13.0 30.0 53.2	31.0 19.0 32.0	0.2423	12248	G130M	LBHO85	11-07-11	4.3	11
RX_J1330.8+3119	13.0 30.0 53.2	31.0 19.0 32.0	0.2423	12248	G160M	LBHO85	11-07-11	6.8	11
RX_J1356.4+2515	13.0 56.0 25.6	25.0 15.0 23.0	0.1640	12248	G130M	LBH057	12-02-03	2.3	10
RX_J1356.4+2515	13.0 56.0 25.6	25.0 15.0 23.0	0.1640	12248	G160M	LBH057	12-02-03	2.8	6
RX_J1503.2+6810	15.0 3.0 16.5	68.0 10.0 6.0	0.1140	12276	G130M	LBI609	10-12-31	1.9	11
RX_J1544.5+2827	15.0 44.0 30.5	28.0 27.0 56.0	0.2314	13423	G130M	LC9W08	14-02-25	2.1	10
RX_J2043.1+0324	20.0 43.0 6.2	3.0 24.0 50.0	0.2710	13840	G130M	LCJW02	14-10-23	7.8	15
RX_J2139.7+0246	21.0 39.0 44.2	2.0 46.0 5.0	0.2600	13840	G130M	LCJW03	14-10-27	7.9	16
SBS0957+599	10.0 1.0 2.6	59.0 44.0 15.0	0.7475	12248	G130M	LBHO65	11-03-18,19	3.3	12
SBS0957+599	10.0 1.0 2.6	59.0 44.0 15.0	0.7475	12248	G160M	LBHO65	11-03-19	5.2	11
SDSSJ021218.32-073719.8	2.0 12.0 18.3	-7.0 37.0 20.0	0.1739	12248	G130M	LBHO83	11-06-26	6.5	12
						LBHO92	11-08-21		
SDSSJ021218.32-073719.8	2.0 12.0 18.3	-7.0 37.0 20.0	0.1739	12248	G160M	LBHO83	11-06-26	6.8	7
						LBHO92	11-08-21		
SDSSJ080838.80+051440.0	8.0 8.0 38.8	5.0 14.0 40.0	0.3606	12603	G130M	LBS330	12-03-17	4.7	10
SDSSJ091728.60+271951.0	9.0 17.0 28.6	27.0 19.0 51.0	0.0756	14071	G130M	LCX202	15-11-30	15.5	11
						LCX2Z2	16-02-06		
SDSSJ112224.10+031802.0	11.0 22.0 24.1	3.0 18.0 2.0	0.4753	12603	G130M	LBS318	13-03-29	7.6	13
SDSSJ130524.30+035731.0	13.0 5.0 24.3	3.0 57.0 31.0	0.5457	12603	G130M	LBS321	12-06-25,26	7.6	13
SDSSJ135726.27+043541.4	13.0 57.0 26.2	4.0 35.0 41.0	1.2345	12264	G130M	LBJ005	11-06-22	14.1	21
SDSSJ135726.27+043541.4	13.0 57.0 26.2	4.0 35.0 41.0	1.2345	12264	G160M	LBJ006	11-06-14	28.2	13
						LBJ007	11-06-26		
SDSSJ140428.30+335342.0	14.0 4.0 28.3	33.0 53.0 42.0	0.5500	12603	G130M	LBS320	13-03-03	7.7	10
TON1009	9.0 9.0 6.1	32.0 36.0 31.0	0.8103	12603	G130M	LBS328	12-04-22	4.7	12

Table 1

COS targets in this sample. *Total exposure time and S/N ratio is given for multi-orbit exposures unless different gratings are used.

band measurements are available for $\sim 30\%$ of galaxies, and most of the rest have SDSS g and r magnitudes, which can be converted to B via $B = g + 0.39(g - r) + 0.21$ (Jester et al. 2005). Finally, we also compute an estimate of the virial radius of each galaxy as $\log R_{\text{vir}} = 0.69 \log D + 1.24$. This follows the parametrization of Stocke et al. (2013) relating a galaxy’s luminosity to its virial radius, and the Wakker & Savage (2009) empirical relation between diameter and luminosity (see Wakker et al. 2015 and references therein for further details). Errors are propagated from the original published magnitude errors.

This homogeneous galaxy data table allows us to draw direct comparisons between the properties of the absorbers and the properties, separations, and environments of nearby galaxies with unprecedented completeness. The full dataset will be publicly released and discussed in further detail in a forthcoming paper (French et al. 2017, in prep).

2.2. Spectra

This initial pilot study contains 33 sightlines to bright QSOs observed with COS. We chose sightlines by first sorting the galaxy data table described above by galaxy diameter. This sorted list is then correlated with the full list of publicly available sightlines, and only those systems with impact parameter less than 500 Mpc and galaxy diameter, D , greater than 25 kpc are kept. Finally, we select the top 33 sightlines with $S/N > 10$, as sorted by galaxy diameter.

All COS spectra for the target sightlines were obtained through the Barbara A. Mikulski Archive for Space Telescopes (MAST), and processed with CALCOS v3.0 or later. We combined individual exposures by the method of Wakker et al. (2015), which corrects the COS wavelength scale by cross-correlating all ISM and IGM lines in each exposure. This method addresses the up to ± 40 km/s misalignments produced by CALCOS, and produces a corrected error array based on Poisson noise, which better matches the measured errors than the errors delivered in the x1d files. We then combine multiple exposures by aligning Galactic absorption lines with 21-cm spectra, and adding up the total counts in each pixel before converting to flux using the original, average flux-count ratio at each wavelength.

3. RESULTS

We have identified 48 Ly α absorption lines in the spectra of our initial 33 QSO sample which can be unambiguously associated with a single nearby galaxy of diameter $D \geq 25$ kpc. In order to be considered for a pairing, a galaxy and absorption feature must appear within 400 km/s in velocity and 500 kpc in physical impact parameter from each other. When multiple galaxies pass these criteria for a particular line, we are left with two options. 1) one galaxy is obviously far larger and closer in physical and velocity space to the line, and may have several satellite galaxies, or 2) no single galaxy is obviously dominant, and we do not include this line in further analysis.

To facilitate this decision, we calculate the likelihood, \mathcal{L} , of every possible galaxy-absorber pairing as follows:

$$\mathcal{L} = A e^{-(\frac{\rho}{R_{\text{eff}}})^2} e^{-(\frac{\Delta v}{200})^2}. \quad (3)$$




Figure 2. a) An example of Ly α lines found in a sightline towards MRK290 at 3090 and 3192 km/s. b) A map of *all* galaxies within a 500 kpc impact parameter of target MRK290 sightline and with velocity (cz) within 400 km/s of absorption detected at 3090 and 3192 km/s (central black star). The galaxy NGC5987 ($v = 3010$ km/s, inclination = 65°) can be unambiguously paired with the Ly α absorption features at $v = 3090, 3192$ km/s because it is the largest and closest galaxy in both physical and velocity space to the absorption feature.

Here ρ is the physical impact parameter, Δv the velocity difference between the absorber and the galaxy ($\Delta v = v_{\text{galaxy}} - v_{\text{absorber}}$), and A is a factor included to increase the likelihood in the case that $\rho \leq R_{\text{eff}}$ (in which case $A = 2$, otherwise $A = 1$).

We compute \mathcal{L} for two different values of R_{eff} : R_{vir} , the virial radius of the galaxy, and $d^{1.5}$, the major diameter of the galaxy to the power of 1.5. \mathcal{L} computed with R_{vir} is liable to select satellite galaxies instead of the larger hosts, so including a version with $d^{1.5}$ serves as a two-tiered selection system. An absorber-galaxy system separated by 200 km/s in velocity and $1R_{\text{vir}}$ would have $\mathcal{L} = 0.27$. In order for an absorber to be marked as “associated” with a particular galaxy, we require that its \mathcal{L} must be a factor of 5 larger than the next best possible association, and $\mathcal{L} \geq 0.01$ for at least one of $\mathcal{L}_{R_{\text{vir}}}$ or $\mathcal{L}_{d^{1.5}}$. We visually inspect systems with only one \mathcal{L} meet-

Target	Galaxy	\mathcal{L}	R_{vir} [kpc]	ρ (kpc) [kpc]	v_{galaxy} [km/s]	Δv [km/s]	Inc. [deg]	Az. [deg]	$v_{Ly\alpha}$ [km/s]	$W_{Ly\alpha}$ [mÅ]
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)
1H0717+714	UGC03804	0.24	173	207	2887	17	55	7	2870	343±6
1H0717+714	UGC03804	0.21	173	207	2887	-69	55	7	2956	39±4
2dFGRS.S393Z082	NGC1097	1.9*	304	112	1271	32	58	27	1239	570±21
H1101-232	MCG-04-26-019	0.33	173	179	3623	43	68	26	3580	573±12
HE0241-3043	NGC1097	1.6*	304	219	1271	50	58	77	1221	83±12
HE0241-3043	NGC1097	1.6*	304	219	1271	-39	58	77	1310	184±15
LBQS1230-0015	NGC4517	1.6*	208	110	1128	1	90	90	1127	473±16
MRC2251-178	MCG-03-58-009	1.4*	319	320	9030	-21	61	39	9051	60±4
MRK1014	NGC0768	0.042*	231	486	7021	-59	64	85	7080	117±11
MRK290	NGC5987	0.77*	322	486	3010	-95	67	12	3105	511±5
MRK290	NGC5987	0.37*	322	486	3010	-197	67	12	3207	319±4
MRK876	UGC10294	0.063	165	274	3504	26	51	7	3478	280±3
PG0003+158	NGC7814	0.081	171	197	1050	217	68	47	833	131±15
PG0832+251	KUG0833+252	0.041	165	294	6964	-16	62	55	6980	133±14
PG0832+251	KUG0833+252	0.01	165	294	6964	-237	62	55	7201	48±10
PG1001+054	UGC05432	0.14	164	217	3995	-97	36	78	4092	222±10
PG1302-102	NGC4939	0.05*	235	265	3110	-338	48	61	3448	71±5
RBS1768	RFGC3781	0.056*	253	464	9162	-198	90	74	9360	364±4
RBS1768	RFGC3781	0.024*	253	464	9162	-272	90	74	9434	160±5
RX_J0714.5+7408	UGC03717	0.13*	202	271	4188	114	63	83	4074	58±7
RX_J0714.5+7408	UGC03717	0.15*	202	271	4188	-76	63	83	4264	410±9
RX_J1017.5+4702	NGC3198	0.02	191	378	663	34	73	55	629	60±17
RX_J1117.6+5301	NGC3631	0.32	187	198	1156	25	16	47	1131	356±20
RX_J1117.6+5301	NGC3631	0.25	187	198	1156	-103	16	47	1259	57±17
RX_J1236.0+2641	NGC4559	0.27	165	188	807	12	64	31	795	295±37
RX_J1236.0+2641	NGC4565	0.54*	292	159	1230	218	90	39	1012	337±32
RX_J1236.0+2641	NGC4565	1.7*	292	159	1230	42	90	39	1188	288±24
RX_J1330.8+3119	UGC08492	0.081*	204	335	7414	13	16	41	7401	330±15
RX_J1356.4+2515	CGCG132-055	0.35*	206	190	8671	196	36	25	8475	126±18
RX_J1503.2+6810	CGCG318-012	0.031*	250	325	9765	-357	52	1	10122	44±14
RX_J1544.5+2827	CGCG166-047	0.031	175	326	9646	4	43	61	9642	183±14
RX_J1544.5+2827	CGCG166-047	0.023	175	326	9646	-113	43	61	9759	169±12
RX_J2043.1+0324	NGC6954	0.037	166	301	4067	-13	56	66	4080	82±10
RX_J2139.7+0246	UGC11785	1.5	203	108	4074	-9	90	69	4083	490±7
RX_J2139.7+0246	UGC11785	1.2*	203	108	4074	-107	90	69	4181	529±7
SBS0957+599	MCG+10-14-058	1.4*	261	206	9501	32	75	19	9469	78±12
SDSSJ021218.32-073719.8	SDSSJ021315.79-073942.7	0.09	174	268	4800	44	52	10	4756	528±15
SDSSJ021218.32-073719.8	SDSSJ021315.79-073942.7	0.092	174	268	4800	-33	52	10	4833	500±17
SDSSJ080838.80+051440.0	UGC04239	0.87*	279	378	8763	23	45	38	8740	883±24
SDSSJ080838.80+051440.0	UGC04239	0.45*	279	378	8763	-164	45	38	8927	130±19
SDSSJ091728.60+271951.0	UGC04895	0.022*	204	408	7073	-68	61	32	7141	374±23
SDSSJ112224.10+031802.0	NGC3640	0.4	180	139	1251	202	38	22	1049	288±30
SDSSJ112224.10+031802.0	NGC3640	1.1	180	139	1251	-13	38	22	1264	424±27
SDSSJ130524.30+035731.0	UGC08186	1.3*	268	249	7006	-33	82	14	7039	480±14
SDSSJ135726.27+043541.4	NGC5364	0.74*	211	183	1241	117	57	84	1124	85±11
SDSSJ135726.27+043541.4	NGC5364	0.97*	211	183	1241	-55	57	84	1296	98±9
SDSSJ140428.30+335342.0	KUG1402+341	1.4	204	118	7919	35	72	63	7884	889±28
TON1009	NGC2770	0.19*	204	274	1947	-14	87	41	1961	350±21

Table 2

All associated systems. The largest \mathcal{L} value is given, with a (*) indicating that this corresponds to $\mathcal{L}_{d1.5}$, otherwise the quoted \mathcal{L} was computed with R_{vir} .

ing these criteria, and decide to reject or include it based on the complexity of the nearby galaxy environment.

Figures 2.2 and 2(b) show a clean example of a Ly α absorption line with a map of its galaxy environment, showing an unambiguous pairing between the absorption features at 3090, 3192 km/s toward MRK290 and galaxy NGC5987 ($\mathcal{L} = 0.37$). Unless explicitly stated, all following analysis concerns similarly unambiguous “associated” systems.

Additionally, we split the absorber-galaxy catalog

based on the velocity difference of the two, Δv . With this scheme, we refer to an absorber with a velocity *lower* than the associated galaxy as *blueshifted*, while an absorber with a velocity *higher* is referred to as *redshifted*. The rest of the results will be analyzed based upon this splitting. In all figures blue and red-shifted absorbers are represented as blue diamonds and red circles, respectively, and red diamonds correspond to systems where *both* a red and blue-shifted absorber is detected.




Figure 3. a) Equivalent width of each absorber as a function of impact parameter, ρ . b) Equivalent width as a function of ρ/R_{vir} . The anti-correlation is strongest when scaling ρ by the galaxy virial radius. Absorbers are separated into red and blue-shifted samples based on Δv . Bins of mean EW are overplotted in red-dashed, and blue-dotted lines for their respective samples.




Figure 4. a) Equivalent width of each absorber as a function of the virial radius of the associated galaxy. The blue-dotted and red-dashed lines shows the average EW in 50 kpc bins of impact parameter for the blueshifted and redshifted absorbers, respectively. b) Impact parameter to each absorber as a function of the virial radius of the associated galaxy. The blue-dotted and red-dashed lines shows the average impact parameter in 50 kpc bins of R_{vir} for the blueshifted and redshifted absorbers, respectively.

3.1. EW - ρ Anti-correlation

Numerous previous studies have suggested that Ly α equivalent width (EW) is anti-correlated with impact parameter (ρ) to the nearest galaxy. We find a weak correlation, as shown in Figure 3(a). However, we find a stronger anti-correlation when we normalize ρ by R_{vir} . Figure 3(b) shows this expected anti-correlation when plotting EW vs ρ/R_{vir} . A possible explanation for this trend is that larger galaxies host larger, more physically extended CGM halos. We would thus expect the absorber EW to also correlate positively with R_{vir} . Figure 4(a) shows EW as a function of R_{vir} , with the blue-dashed

and red-dotted lines show the average EW in bins of 50 kpc of R_{vir} , showing little evidence of a correlation. However, by similarly plotting ρ as a function of R_{vir} , we instead find some evidence that absorbers around larger galaxies tend to be found at higher impact parameters. While we expect the upper-left quadrant of this figure to be sparsely populated (our likelihood-based method would tend not to choose small galaxies at large distances), it is unclear to us why the lower-right quadrant (large galaxies with absorbers at low impact parameter) is also sparsely populated. The full-sized sample at the completion of our study should provide a clearer picture.




Figure 5. Equivalent width of each absorber as a function of the inclination angle of the associated galaxy. The dashed black line shows the mean EW of all absorbers in bins of 15° .




Figure 6. **Top:** Distribution of inclinations for all associated galaxies, split into red and blue shifted sets. **Bottom:** Distribution of inclinations of all observed galaxies in the $cz \leq 10,000$ km/s redshift range. The dashed line shows the inclination distribution for a truly random sample (i.e. no observational biases).

3.2. Inclination

In this section we examine the inclinations of the associated galaxies compared to the distributions of absorbers. We correct for the finite thickness of galaxies, which causes b/a to deviate from $\cos i$ at high inclinations, by computing galaxy inclinations with the following formula from Heidmann et al. (1972a):

$$\cos(i) = \sqrt{\frac{q^2 - q_0^2}{1 - q_0^2}}, \quad (4)$$

where $q = b/a$, the ratio of the minor to major axis, and q_0 is the intrinsic axis ratio, set to $q_0 = 0.2$ for all galaxies (e.g., Jones, Davies, and Trewella 1996). 6 of the 48 total absorbers are associated with E or S0 type galaxies, but we have chosen to keep the value of $q_0 = 0.2$ uniform throughout. The calculated values of $\cos(i)$ we use for these galaxies are thus conservative under-estimates of




Figure 7. Equivalent width as a function of the velocity separation between the galaxy and absorption line.

their true inclinations.

Figure 5 shows red and blueshifted absorbers' EW plotted against the inclinations of their associated galaxies. We note that there is a clear excess of absorbers near galaxies of high inclination, with 77% of redshifted and 73% of blueshifted absorbers being associated with galaxies of $i \geq 50$ deg, and only 3 absorbers associated with a galaxy of $i < 35$. The solid-black and dashed-grey lines show mean and 90th percentile histograms, respectively, in bins of 12 deg. There does not appear to be much evolution of EW across galaxy inclination, although a slight increase of mean EW is possibly present towards higher inclinations.

In total 75% of absorbers are associated with high inclined galaxies ($i \geq 50$ deg). Only 56% of all galaxies in the survey volume are highly inclined, indicating a preference for detecting absorption around inclined galaxies. Figure 6 shows the distribution of galaxy inclinations for both the red and blue-shifted associated galaxies and all galaxies within the survey volume. We tested the difference between the full distribution of inclination angles for all galaxies in our survey volume and the distribution for all associated galaxies (red + blue-shifted absorbers) using the Anderson-Darling (AD) and Kolmogorov-Smirnov (KS) statistical distribution tests, yielding p-values of $KS_p = 0.014$, and $AD_p = 0.00037$. Hence, there is less than a 1% probability (3σ) that our associated galaxies are sampled from the average distribution of galaxies within this survey volume.

It is worth noting here that the observed distribution of galaxy inclinations is *not* flat, as one might expect. The dashed line in Figure 6 shows the distribution of inclinations for a random, uniform sample (i.e., a uniform distribution of $q = b/a$ values between $0.2 - 1.0$). There could be a number of effects contributing to this. If our sample is magnitude limited and we assume galaxies are mostly optically thin but with an very-thin optically thick component (e.g., a dust lane), then mostly face-on and mostly edge-on galaxies would be underrepresented due to surface-brightness and dust obscuration effects (e.g., see Jones, Davies and Trewella 1996). It is possible that a similar effect is also responsible for the

over-abundance of Ly α detections around highly inclined galaxies. If we assume a disk or oblate spheroid shape and some covering fraction for the CGM, the probability of encountering a cloud near an inclined galaxy would increase with the increased path-length through the halo.

3.3. Velocity Difference (Δv)

We find evidence for an anti-correlation between absorber EW and the velocity difference between the galaxy and the associated absorption, Δv . The mean and maximal EW of absorption increases with decreasing Δv (see Figure 7). In total, 32/48 (67%) of absorbers are found within ± 100 km/s. This ± 100 km/s threshold also applies to absorber EW , with only 1 absorber of $EW \geq 400$ found with $\Delta v > 100$ km/s. Blueshifted absorbers are on average closer to their associated galaxy, with $\overline{\Delta v}_{blue} = 68 \pm 16$ km/s, compared to $\overline{\Delta v}_{red} = -108 \pm 20$ km/s for the redshifted sample, and correspondingly have higher average equivalent width, $\overline{EW}_{blue} = 329 \pm 52$ mÅ compared to $\overline{EW}_{red} = 245 \pm 34$ mÅ.

Additionally, of the 48 associated absorbers, 29 are matched with the same galaxy as another absorber (for a total of 14 unique galaxies in this subset). All but one of these cases involve two absorbers in the same sightline yet separated in velocity around a galaxy. 23/29 of these are oriented such that the higher EW absorber has the smaller Δv , and the 6 others are close in either velocity or EW . The one galaxy with 3 associated absorbers, NGC1097, shows this trend across two sightlines as well, with absorbers at $\Delta v = 32$ km/s and $EW = 570$ mÅ towards 2dFGRS_S393Z082, and $\Delta v = -39$ km/s and $EW = 184$ mÅ and $\Delta v = 50$ km/s and $EW = 83$ mÅ towards HE0241-3043.

3.4. Azimuth

In this section we examine properties of absorbers as a function of their azimuthal angle with respect to their associated galaxy. Azimuth is defined as the angle between the major axis of a galaxy and the vector connecting the absorption feature and the midpoint of the galaxy plane. Figure 8 illustrates this.

The mean azimuth angle for blueshifted absorbers is $43 \pm 5^\circ$, and $49 \pm 5^\circ$ for redshifted absorbers. Figure 9 shows the distribution of azimuth angles for both red and blue-shifted absorbers. Unlike the findings of Kacprzak et al. (2011, 2012), who find a bimodal distribution of Mg II absorbers around galaxies, our distributions of Ly α absorbers are generally consistent with a flat, or random distribution. There is possibly a slight overabundance of redshifted absorbers around 0° (minor-axis) and blueshifted absorbers between $20 - 50^\circ$ (just off major-axis), but we cannot assign this observation much significance yet given the small sample size. We additionally find no significant correlation between azimuth angle and EW or Δv . See figure 10 for a map of the locations of absorbers relative to their associated galaxies, split into three bins of inclination.

4. SUMMARY

We have measured 48 Ly α absorption lines in the spectra of 33 COS targets and matched each to a single, large ($D \geq 25$ kpc) galaxy. Table 3 presents a breakdown of




Figure 8. Azimuth is the angle, α , between the major axis of the galaxy, a , and a vector extending from the AGN target to the galaxy center. Image of NGC891 credit: Composite Image Data - Subaru Telescope (NAOJ), Hubble Legacy Archive, Michael Joner, David Laney (West Mountain Observatory, BYU); Processing - Robert Gendler.




Figure 9. The distributions of azimuth angles for red and blue-shifted samples, with the combined sample plotted in black. $Azimuth = 0$ corresponds to along the projected major axis of the galaxy, and $azimuth = 90$ is along the minor axis.

our results when separating absorber-galaxy pairs into red and blue-shifted samples. The following summarizes our findings:

- We introduce a likelihood parameter based on Gaussian profiles centered around ρ/R_{vir} and Δv to automate the matching of absorbers with associated galaxies.

Statistic	Blueshifted Absorbers	Redshifted Absorbers
Number	22	26
Mean EW [mÅ]	329 ± 52	245 ± 34
Median EW [mÅ]	292 ± 16	177 ± 10
Mean R_{vir} [kpc]	215 ± 10	224 ± 10
Mean ρ [kpc]	218 ± 17	298 ± 23
Mean Δv [kms $^{-1}$]	68 ± 16	-108 ± 20
Mean Inc. [deg]	58 ± 4	61 ± 4
Mean Az. [deg]	43 ± 5	49 ± 5

Table 3

Average properties of the associated galaxy sample split into red and blue-shifted bins based on Δv .




Figure 10. A map of where each absorber was detected with respect to the associated galaxies, separated into three bins of inclination (illustrated by the gray ellipse in the bottom left corner of each plot). **LEFT:** Absorbers associated with galaxies of inclination $0 \leq Inc \leq 40$, **CENTER:** $40 < Inc \leq 65$, and **RIGHT:** $65 \leq Inc$.

- EW anti-correlates most strongly with ρ when normalized by R_{vir} . It follows that EW weakly correlates and anti-correlates with R_{vir} and ρ , respectively.
- The mean and maximal EW of absorbers increases with decreasing Δv . The strongest absorbers are nearly all found within $\Delta v = \pm 100$ km/s of their associated galaxies.
- We find a slight dichotomy in the EW of absorption blue-ward vs red-ward of associated galaxies. Redshifted absorbers are weaker, with $\overline{EW} = 245 \pm 34$ mÅ compared to $EW = 329 \pm 52$ mÅ for blueshifted absorbers.
- Ly α absorbers are most associated with inclined galaxies. 73% of blueshifted and 77% of redshifted absorbers are associated with galaxies with $i \geq 50$ deg, whereas 56% of all galaxies in the survey volume have similarly high inclinations. The distributions of associated vs all galaxy inclinations differ at a greater than 99% confidence, or 3σ , level according to the Anderson-Darling distribution test.
- We find no strong evidence for azimuth preference for absorption - Ly α absorbers appear to be distributed uniformly around galaxies.

REFERENCES

- Carswell, B., Schaye, J., & Kim, T.-S. 2002, ApJ, 578, 43
- Cen, R. 2013, ApJ, 770, 139
- . 2014, ApJ, 789, L21
- Chen, H.-W., Lanzetta, K. M., Webb, J. K., & Barcons, X. 1998, ApJ, 498, 77
- . 2001, ApJ, 559, 654
- Corwin, Jr., H. G., Buta, R. J., & de Vaucouleurs, G. 1994, AJ, 108, 2128
- Côté, S., Wyse, R. F. G., Carignan, C., Freeman, K. C., & Broadhurst, T. 2005, ApJ, 618, 178
- Danforth, C. W., & Shull, J. M. 2005, ApJ, 624, 555
- . 2008, ApJ, 679, 194
- Danforth, C. W., Shull, J. M., Rosenberg, J. L., & Stocke, J. T. 2006, ApJ, 640, 716
- Danforth, C. W., Keeney, B. A., Tilton, E. M., et al. 2014, ArXiv e-prints
- Erb, D. K. 2008, ApJ, 674, 151
- Heidmann, J., Heidmann, N., & de Vaucouleurs, G. 1972, MmRAS, 75, 85
- Jones, H., Davies, J. I., & Trewheella, M. 1996, MNRAS, 283, 316
- Kacprzak, G. G., Churchill, C. W., Barton, E. J., & Cooke, J. 2011a, ApJ, 733, 105
- Kacprzak, G. G., Churchill, C. W., Evans, J. L., Murphy, M. T., & Steidel, C. C. 2011b, MNRAS, 416, 3118
- Kacprzak, G. G., Churchill, C. W., & Nielsen, N. M. 2012, ApJ, 760, L7
- Keeney, B. A., Stocke, J. T., Syphers, D., et al. 2013, in American Astronomical Society Meeting Abstracts, Vol. 222, American Astronomical Society Meeting Abstracts, 214.18
- Kereš, D., & Hernquist, L. 2009, ApJ, 700, L1
- Kim, T.-S., Bolton, J. S., Viel, M., Haehnelt, M. G., & Carswell, R. F. 2007, MNRAS, 382, 1657
- Lanzetta, K. M., Bowen, D. V., Tytler, D., & Webb, J. K. 1995, ApJ, 442, 538
- Lehner, N., Savage, B. D., Richter, P., et al. 2007, ApJ, 658, 680
- Liang, C. J., & Chen, H.-W. 2014, MNRAS, 445, 2061
- Lundgren, B. F., Brunner, R. J., York, D. G., et al. 2009, ApJ, 698, 819
- Mathes, N. L., Churchill, C. W., Kacprzak, G. G., et al. 2014, ApJ, 792, 128
- McLIn, K. M., Stocke, J. T., Weymann, R. J., Penton, S. V., & Shull, J. M. 2002, ApJ, 574, L115
- Meiksin, A. A. 2009, Reviews of Modern Physics, 81, 1405
- Oppenheimer, B. D., & Davé, R. 2009, MNRAS, 395, 1875
- Oppenheimer, B. D., Davé, R., Katz, N., Kollmeier, J. A., & Weinberg, D. H. 2012, MNRAS, 420, 829
- Penton, S. V., Stocke, J. T., & Shull, J. M. 2002, ApJ, 565, 720
- . 2004, ApJS, 152, 29
- Prochaska, J. X., Weiner, B., Chen, H.-W., Mulchaey, J., & Cooksey, K. 2011, ApJ, 740, 91
- Prochaska, J. X., Weiner, B. J., Chen, H.-W., & Mulchaey, J. S. 2006, ApJ, 643, 680
- Putman, M. E., Peek, J. E. G., & Heitsch, F. 2009a, ArXiv e-prints
- Putman, M. E., Henning, P., Bolatto, A., et al. 2009b, in Astronomy, Vol. 2010, astro2010: The Astronomy and Astrophysics Decadal Survey, 241
- Rudie, G. C., Steidel, C. C., & Pettini, M. 2012a, ApJ, 757, L30
- Rudie, G. C., Steidel, C. C., Shapley, A. E., & Pettini, M. 2013, ApJ, 769, 146
- Rudie, G. C., Steidel, C. C., Trainor, R. F., et al. 2012b, ApJ, 750, 67
- Savage, B. D., Kim, T.-S., Wakker, B. P., et al. 2014, ApJS, 212, 8

- Shull, J. M., Smith, B. D., & Danforth, C. W. 2012, *ApJ*, 759, 23
- Smith, B. D., Hallman, E. J., Shull, J. M., & O'Shea, B. W. 2011, *ApJ*, 731, 6
- Steidel, C. C., Erb, D. K., Shapley, A. E., et al. 2010, *ApJ*, 717, 289
- Stewart, K. R., Kaufmann, T., Bullock, J. S., et al. 2011, *ApJ*, 738, 39
- Stocke, J. T., Keeney, B. A., Danforth, C. W., et al. 2013, *ApJ*, 763, 148
- Tejos, N., Morris, S. L., Crighton, N. H. M., et al. 2012, *MNRAS*, 425, 245
- Tejos, N., Morris, S. L., Finn, C. W., et al. 2014, *MNRAS*, 437, 2017
- Thom, C., Tumlinson, J., Werk, J. K., et al. 2012, *ApJ*, 758, L41
- Tripp, T. M., Aracil, B., Bowen, D. V., & Jenkins, E. B. 2006a, *ApJ*, 643, L77
- Tripp, T. M., Bowen, D. V., Sembach, K. R., et al. 2006b, in *Astronomical Society of the Pacific Conference Series*, Vol. 348, *Astrophysics in the Far Ultraviolet: Five Years of Discovery with FUSE*, ed. G. Sonneborn, H. W. Moos, & B.-G. Andersson, 341
- Tripp, T. M., Lu, L., & Savage, B. D. 1998, *ApJ*, 508, 200
- Tripp, T. M., Sembach, K. R., Bowen, D. V., et al. 2008, *ApJS*, 177, 39
- Tully, R. B., Courtois, H. M., Dolphin, A. E., et al. 2013, *AJ*, 146, 86
- Tumlinson, J., Werk, J. K., Thom, C., et al. 2011, *ApJ*, 733, 111
- van de Voort, F., & Schaye, J. 2012, *MNRAS*, 423, 2991
- van de Voort, F., Schaye, J., Booth, C. M., Haas, M. R., & Dalla Vecchia, C. 2011, *MNRAS*, 414, 2458
- Wakker, B. P., Hernandez, A. K., French, D. M., et al. 2015, *ApJ*, 814, 40
- Wakker, B. P., & Savage, B. D. 2009, *ApJS*, 182, 378
- Werk, J. K., Prochaska, J. X., Tumlinson, J., et al. 2014, *ApJ*, 792, 8
- York, D. G., vanden Berk, D., Richards, G. T., et al. 2005, in *IAU Colloq. 199: Probing Galaxies through Quasar Absorption Lines*, ed. P. Williams, C.-G. Shu, & B. Menard, 58–64