Supplemental Table 2: Detected mutant loads in gDNA and cell samples of patients and their affected children after direct mutation site amplification by conventional PCR and after WGA followed by mutation site amplification

Sample type	Detected mutant load (%)						
	Patient A	**Daughter of Patient A**	**Patient B**	**Son of Patient B**	**Patient C**	**Daughter of Patient C**	**Patient D**
gDNA subjected to conventional PCR	55.0[1] 51.9[1] 56.3[2] 54.4[2]	100[3] 100[3]	ND[1]	97.1[1]	ND[1] ND[1] ND[1]	84.2[2] 85.0[2] 85.0[2]	6.0[1] 13.2[2]
gDNA subjected to WGA and conventional PCR	59.6[1] 60.8[1] 52.8[1] 49.2[1] 51.9[2] 50.6[2] 56.2[2] 65.8[2]	100[3] 100[3]	ND[1] ND[1]	98.4[1] 91.9[1] 96.6[1]	ND[1] ND[1]	84.5[2] 85.6[2] 82.2[2] 82.1[2] 84.9[2]	6.7[1] 14.7[2]
Cell samples subjected to conventional PCR	97.3[1,4] ND[1,4] 59.8[1,5] 91.1[1,5] 74.8[1,6] 19.6[1,6]		ND[1] ND[1] ND[1]	93.0[1,4] 100[1,4] 99.9[1,4]	ND[1] ND[2]	83.5[2,4] 83.9[2,4] 93.9[2,4] 90.5[2,4]	ND[1,4] 49.0[1,4] ND[1,4] ND[1,4] ND[1,4] ND[1,4] ND[2,4] ND[2,4] 39.1[2,4] ND[2,4] 63.8[2,4] ND[2,4]
Cell samples subjected to WGA and conventional PCR	ND[1,4] 100[1,4] 33.1[1,5] 60.2[1,5] 72.2[1,6] 95.7[1,6]		ND[1] ND[1] ND[1] ND[1] ND[1] ND[1]	100[1,4] 100[1,4] 99.9[1,4] 98.8[1,4] 100[1,4] 98.9[1,4]	ND[1] ND[1] ND[2] ND[2]	77.3[2,4] 81.7[2,4] 94.3[2,4] 75.2[2,4] 85.9[2,4] 77.3[2,4]	49.7[1,4] ND[1,4] ND[1,4] ND[1,4] ND[1,4] ND[1,4]

				39.5[1,4] 100[1,4] 98.0[1,4]			ND[2,4] 1.2[2,4] 1.3[2,4] ND[2,4] ND[2,4] 1.6[2,4]

Mean coverage of mutation loci was 75,420 reads, ranging from 12,376-421,777. [1]gDNA or lymphocyte(s) from peripheral blood. [2]gDNA or buccal cells from a cheek swab. [3]Source unknown. [4]Single cell. [5]Three cells. [6]Five cells. ND=not detected.