

An Introduction to String Theory

Mate Zoltan Farkas

July 15, 2020

Table of Contents

- 1 Introduction
 - Notation
- 2 The Relativistic String
 - The Relativistic Point Particle
 - Dynamics of a Relativistic String
 - The Nambu-Goto Action
 - The Polyakov Action
- 3 Covariant Quantization of the Solutions
- 4 Quantization in the Lightcone Gauge

► Metric:

- $\eta_{\mu\nu} = \text{diag}(-1, +1, \dots, +1)$

The Relativistic Point Particle – The Action


- ▶ Consider the action of a point particle (with fixed coordinates $X_\mu = (t, \vec{x})$ in a given frame):

$$S = -m \int dt \sqrt{1 - \dot{\vec{x}}\dot{\vec{x}}}$$

→ not Lorentz-invariant, due to mixture of spacial and temporal coordinates under a Lorentz-transformation Λ .

- ▶ Consider instead for a generalized coordinate τ along the line element:

$$S = -m \int d\tau \sqrt{-\frac{dX^\mu}{d\tau} \frac{dX^\nu}{d\tau} \eta_{\mu\nu}}$$



Ref: [1]

Remark: S is proportional to the integral over the worldline of the particle

The Relativistic Point Particle – The Action


- ▶ Consider the action of a point particle (with fixed coordinates $X_\mu = (t, \vec{x})$ in a given frame):

$$S = -m \int dt \sqrt{1 - \dot{\vec{x}}\dot{\vec{x}}}$$

→ not Lorentz-invariant, due to mixture of spacial and temporal coordinates under a Lorentz-transformation Λ .

- ▶ Consider instead for a generalized coordinate τ along the line element:

$$S = -m \int d\tau \sqrt{-\frac{dX^\mu}{d\tau} \frac{dX^\nu}{d\tau} \eta_{\mu\nu}}$$



Ref: [1]

Remark: S is proportional to the integral over the worldline of the particle

The Relativistic Point Particle – Action Symmetries

What are the symmetries of

$$S = -m \int d\tau \sqrt{-\frac{dX^\mu}{d\tau} \frac{dX^\nu}{d\tau} \eta_{\mu\nu}}$$

- Reparametrization invariance:

Let $\tilde{\tau} = \tilde{\tau}(\tau)$. Then:

$$S' = -m \int d\tilde{\tau} \sqrt{-\frac{dX^\mu}{d\tilde{\tau}} \frac{dX^\nu}{d\tilde{\tau}} \eta_{\mu\nu}} = S$$

→ gauge symmetry of the action → still D-1 dof!

- Poincaré invariance:

Let:

$$X'^\mu = \Lambda^\mu{}_\nu X^\nu + c^\mu$$

Then:

$$S' = S, \quad \text{as} \quad \Lambda^\mu{}_\rho \eta_{\mu\nu} \Lambda^\nu{}_\sigma = \eta_{\rho\sigma}$$

The Relativistic Point Particle – Action Symmetries

What are the symmetries of

$$S = -m \int d\tau \sqrt{-\frac{dX^\mu}{d\tau} \frac{dX^\nu}{d\tau} \eta_{\mu\nu}}$$

- Reparametrization invariance:

Let $\tilde{\tau} = \tilde{\tau}(\tau)$. Then:

$$S' = -m \int d\tilde{\tau} \sqrt{-\frac{dX^\mu}{d\tilde{\tau}} \frac{dX^\nu}{d\tilde{\tau}} \eta_{\mu\nu}} = S$$

→ gauge symmetry of the action → still D-1 dof!

- Poincaré invariance:

Let:

$$X'^\mu = \Lambda^\mu{}_\nu X^\nu + c^\mu$$

Then:

$$S' = S, \quad \text{as} \quad \Lambda^\mu{}_\rho \eta_{\mu\nu} \Lambda^\nu{}_\sigma = \eta_{\rho\sigma}$$

Dynamics of a Relativistic String – The Nambu-Goto Action

Action of the Relativistic String?

$$\text{particle} \quad \Leftrightarrow \quad \text{worldline} \quad \Leftrightarrow \quad S = -m \int d\tau \underbrace{\sqrt{-\frac{dX^\mu}{d\tau} \frac{dX^\nu}{d\tau} \eta_{\mu\nu}}}_{\text{line element } ds}$$

$$\text{closed string} \quad \Leftrightarrow \quad ? \quad \Leftrightarrow \quad ?$$

Boundary condition of closed strings living in D dimensions:

$$X^\mu(\tau, \sigma) = X^\mu(\tau, \sigma + 2\pi) \quad \text{for } \mu = 0, 1, \dots, D-1; \sigma \in [0; 2\pi)$$

Shorthand notation: $\sigma^\alpha = (\tau, \sigma)$ for $\alpha \in \{0; 1\}$

image of a string

Dynamics of a Relativistic String – The Nambu-Goto Action

Action of the Relativistic String?

$$\text{particle} \quad \Leftrightarrow \quad \text{worldline} \quad \Leftrightarrow \quad S = -m \int d\tau \sqrt{-\frac{dX^\mu}{d\tau} \frac{dX^\nu}{d\tau} \eta_{\mu\nu}}$$

$$\text{closed string} \quad \Leftrightarrow \quad \text{worldsheet} \quad \Leftrightarrow \quad ?$$

Boundary condition of closed strings living in D dimensions:

$$X^\mu(\tau, \sigma) = X^\mu(\tau, \sigma + 2\pi) \quad \text{for } \mu = 0, 1, \dots, D-1; \sigma \in [0; 2\pi)$$

Shorthand notation: $\sigma^\alpha = (\tau, \sigma)$ for $\alpha \in \{0; 1\}$

image of a string

Dynamics of a Relativistic String – The Nambu-Goto Action

Action of the Relativistic String?

$$\text{particle} \quad \Leftrightarrow \quad \text{worldline} \quad \Leftrightarrow \quad S = -m \int d\tau \sqrt{-\frac{dX^\mu}{d\tau} \frac{dX^\nu}{d\tau} \eta_{\mu\nu}}$$

closed string \Leftrightarrow worldsheet \Leftrightarrow Nambu-Goto/Dirac Action

Boundary condition of closed strings living in D dimensions:

$$X^\mu(\tau, \sigma) = X^\mu(\tau, \sigma + 2\pi) \quad \text{for } \mu = 0, 1, \dots, D-1; \sigma \in [0; 2\pi)$$

Shorthand notation: $\sigma^\alpha = (\tau, \sigma)$ for $\alpha \in \{0; 1\}$

image of a string

Dynamics of a Relativistic String – The Nambu-Goto Action

How to describe the surface area to construct the action?

$$S \propto \int_{\partial V} d^2x = \int_{\partial V} d^2\sigma |\det J|$$

For this, first remember that any metric is given by:

$$g_{\alpha\beta} = g(e_\alpha, e_\beta)$$

So the metric on the worldsheet (the pullback metric) can be written as:

$$\gamma_{\alpha\beta} = \underbrace{\frac{\partial X^\mu}{\partial \sigma^\alpha} \frac{\partial X^\nu}{\partial \sigma^\beta} \eta_{\mu\nu}}_{(J^T \eta J)_{\alpha\beta}}$$

Thus,

$$\begin{aligned} \det \gamma &= \det \eta \det^2 J = -\det^2 J \\ |\det J| &= \sqrt{-\det \gamma} = \sqrt{-\gamma} \end{aligned}$$

Dynamics of a Relativistic String – The Nambu-Goto Action

So write the action as:

$$S_{NG} = -\frac{1}{2\pi\alpha'} \int d^2\sigma \sqrt{-\gamma}$$

This action is invariant under,

- ▶ Poincaré transformations
- ▶ Reparametrization

and the equations of motion (EoM) are given by:

$$\partial_\alpha \left(\sqrt{-\gamma} \gamma^{\alpha\beta} \partial_\beta X^\mu \right) = 0$$

⇒ rather hard to quantize in this form! Alternatives?

Dynamics of a Relativistic String – The Polyakov Action

Way out: **The Polyakov Action:**

$$S_P = -\frac{1}{4\pi\alpha'} \int d^2\sigma \sqrt{-g} g^{\alpha\beta} \partial_\alpha X^\mu \partial_\beta X^\nu \eta_{\mu\nu}$$

The EoMs:

► for X^μ :

- same as for the Nambu-Goto action!

$$\partial_\alpha \left(\sqrt{-g} g^{\alpha\beta} \partial_\beta X^\mu \right) = 0$$

► for $g_{\alpha\beta}$:

$$g_{\alpha\beta} = 2 \frac{\partial_\alpha X^\mu \partial_\beta X^\nu \eta_{\mu\nu}}{g^{\rho\sigma} \partial_\rho X^\mu \partial_\sigma X^\nu \eta_{\mu\nu}} \equiv 2 \frac{\partial_\alpha X \partial_\beta X}{g^{\rho\sigma} \partial_\rho X \partial_\sigma X}$$

Gains of the Polyakov action?

Dynamics of a Relativistic String – The Polyakov Action

Symmetries of the Polyakov action

$$S_P = -\frac{1}{4\pi\alpha'} \int d^2\sigma \sqrt{-g} g^{\alpha\beta} \partial_\alpha X^\mu \partial_\beta X^\nu \eta_{\mu\nu}$$

- ▶ Poincaré invariance
- ▶ Reparametrization invariance
- ▶ Invariance under:

$$g'_{\alpha\beta} = \Omega^2(\sigma) g_{\alpha\beta}$$

⇒ **Weyl invariance**

However, using the conformal gauge and writing g as

$$g_{\alpha\beta} = e^{2\phi} \eta_{\alpha\beta}$$

can be undone by a Weyl transformation ($\phi = 0$). Thus,

$$g_{\alpha\beta} = \eta_{\alpha\beta}$$

Plot of the transformation in action

Never forget the Titles!

▶ first item

- subitem

- subsubitem

▶ second item

1. item 1

- 1.1 subitem 1

- 1.2 subitem 2

2. item 2

▶ third item

Never forget the Titles!

- ▶ first item
 - subitem
 - subsubitem
- ▶ second item
 - 1. item 1
 - 1.1 subitem 1
 - 1.2 subitem 2
 - 2. item 2
- ▶ third item

Never forget the Titles!

- ▶ first item
 - subitem
 - subsubitem
- ▶ second item
 - 1. item 1
 - 1.1 subitem 1
 - 1.2 subitem 2
 - 2. item 2
- ▶ third item

Never forget the Titles!

- ▶ first item
 - subitem
 - subsubitem
- ▶ second item
 - 1. item 1
 - 1.1 subitem 1
 - 1.2 subitem 2
 - 2. item 2
- ▶ third item

Never forget the Titles!

- ▶ first item
 - subitem
 - subsubitem
- ▶ second item
 - 1. item 1
 - 1.1 subitem 1
 - 1.2 subitem 2
 - 2. item 2
- ▶ third item

Covariant Quantization of the Solutions of the Nambu-Goto Action

$$S = \frac{1}{2\pi\alpha'} \int \sqrt{-\det g} \, \partial_\mu$$

Quantization of X^μ in the Lightcone Gauge

- [1] David Tong. “Lectures on String Theory”. In: *arXiv:0908.0333 [hep-th]* (Feb. 2012). arXiv: 0908.0333. URL: <http://arxiv.org/abs/0908.0333> (visited on 07/13/2020).