


Topology, symplectic topology, . . . What's next?

Ronen Brilleslijper & Oliver Fabert


0-dimensional objects

Morse theory on smooth
manifolds




0-dimensional objects

Morse theory on smooth manifolds




1-dimensional objects

Floer theory on symplectic manifolds




0-dimensional objects

Morse theory on smooth manifolds



1-dimensional objects

Floer theory on symplectic manifolds




2-dimensional objects

???


0-dimensional objects

Morse theory on smooth manifolds



1-dimensional objects

Floer theory on symplectic manifolds



2-dimensional objects

Holomorphic symplectic? Polysymplectic?

Introduction
●●

Polysymplectic geometry
○○○○

Dirac operators
○○○○○○

Holomorphic symplectic geometry
○○○○

Rigidity
○○○○○

Application
○○

End
○○

Physics motivation

Symplectic geometry

Mechanics

Physics motivation

Symplectic geometry
Mechanics

Polysymplectic geometry
Field theory

Physics motivation

Symplectic geometry

Mechanics

ODEs

Polysymplectic geometry

Field theory

Physics motivation

Symplectic geometry

Mechanics

ODEs

Polysymplectic geometry

Field theory

PDEs

Physics motivation

Two types of field theory

Minkowski

Euclidean

Physics motivation


Two types of field theory

Minkowski

Electromagnetism

Wave equation

Hyperbolic PDEs



Euclidean

Physics motivation


Two types of field theory

Minkowski

Electromagnetism

Wave equation

Hyperbolic PDEs




Euclidean

Electrostatics

Laplace equation

Elliptic PDEs



Physics motivation


Two types of field theory

Minkowski

Electromagnetism

Wave equation

Hyperbolic PDEs




Euclidean

Electrostatics

Laplace equation

Elliptic PDEs



Physics motivation


Two types of field theory

Minkowski

Electromagnetism

Wave equation

Hyperbolic PDEs




[BF23]

Euclidean

Electrostatics

Laplace equation

Elliptic PDEs



Definition

An \mathbb{R}^d -valued 2-form $\Omega = \sum_{i=1}^d \eta_i \otimes \partial_i$ on a manifold M is called a *polysymplectic form* if it is closed and non-degenerate

Definition

An \mathbb{R}^d -valued 2-form $\Omega = \sum_{i=1}^d \eta_i \otimes \partial_i$ on a manifold M is called a *polysymplectic form* if it is closed and non-degenerate in the sense that

$$\begin{aligned}\Omega^b : TM &\rightarrow \operatorname{Hom}(TM, \mathbb{R}^d) \\ X &\mapsto \Omega(X, \cdot)\end{aligned}$$

is injective.

Definition

An \mathbb{R}^d -valued 2-form $\Omega = \sum_{i=1}^d \eta_i \otimes \partial_i$ on a manifold M is called a *polysymplectic form* if it is closed and non-degenerate in the sense that

$$\begin{aligned}\Omega^b : TM &\rightarrow \operatorname{Hom}(TM, \mathbb{R}^d) \\ X &\mapsto \Omega(X, \cdot)\end{aligned}$$

is injective. Equivalently $\bigcap_i \ker \eta_i^b = 0$.

Definition

An \mathbb{R}^d -valued 2-form $\Omega = \sum_{i=1}^d \eta_i \otimes \partial_i$ on a manifold M is called a *polysymplectic form* if it is closed and non-degenerate in the sense that

$$\begin{aligned}\Omega^b : TM &\rightarrow \text{Hom}(TM, \mathbb{R}^d) \\ X &\mapsto \Omega(X, \cdot)\end{aligned}$$

is injective. Equivalently $\bigcap_i \ker \eta_i^b = 0$.

Given a function $H : M \rightarrow \mathbb{R}$ a map $Z : \mathbb{R}^d \rightarrow M$ is called a *solution* if

$$\sum_{i=1}^d \eta_i(\cdot, \partial_i Z) = dH.$$

Example

Let $d = 2$ and $M = \mathbb{R}^3 \ni (q, p_1, p_2)$ with

$$\eta_1 = dp_1 \wedge dq$$

$$\eta_2 = dp_2 \wedge dq$$

Example

Let $d = 2$ and $M = \mathbb{R}^3 \ni (q, p_1, p_2)$ with

$$\eta_1 = dp_1 \wedge dq$$

$$\eta_2 = dp_2 \wedge dq$$

Then $Z = (q, p_1, p_2) : \mathbb{R}^2 \rightarrow \mathbb{R}^3$ is a solution of $H(q, p_1, p_2) = \frac{1}{2}p_1^2 + \frac{1}{2}p_2^2 + V(q)$ if

$$\begin{cases} -\partial_1 p_1 - \partial_2 p_2 &= V'(q) \\ \partial_1 q &= p_1 \\ \partial_2 q &= p_2 \end{cases}$$

Example

Let $d = 2$ and $M = \mathbb{R}^3 \ni (q, p_1, p_2)$ with

$$\eta_1 = dp_1 \wedge dq$$

$$\eta_2 = dp_2 \wedge dq$$

Then $Z = (q, p_1, p_2) : \mathbb{R}^2 \rightarrow \mathbb{R}^3$ is a solution of $H(q, p_1, p_2) = \frac{1}{2}p_1^2 + \frac{1}{2}p_2^2 + V(q)$ if

$$\begin{cases} -\partial_1 p_1 - \partial_2 p_2 &= V'(q) \\ \partial_1 q &= p_1 \\ \partial_2 q &= p_2 \end{cases}$$

$$\iff -(\partial_1^2 + \partial_2^2)q = V'(q)$$

Example (*De Donder-Weyl equations*)

Let $d = 2$ and $M = \mathbb{R}^3 \ni (q, p_1, p_2)$ with

$$\eta_1 = dp_1 \wedge dq$$

$$\eta_2 = dp_2 \wedge dq$$

Then $Z = (q, p_1, p_2) : \mathbb{R}^2 \rightarrow \mathbb{R}^3$ is a solution of $H(q, p_1, p_2) = \frac{1}{2}p_1^2 + \frac{1}{2}p_2^2 + V(q)$ if

$$\begin{cases} -\partial_1 p_1 - \partial_2 p_2 &= V'(q) \\ \partial_1 q &= p_1 \\ \partial_2 q &= p_2 \end{cases}$$

$$\iff -(\partial_1^2 + \partial_2^2)q = V'(q)$$

Hamiltonians?

- $H : M \rightarrow \mathbb{R}$

Hamiltonians?

$$\bullet \quad H : M \rightarrow \mathbb{R} \quad \implies \quad \eta_1(\cdot, \partial_1 Z) + \eta_2(\cdot, \partial_2 Z) = dH$$

Hamiltonians?

- $H : M \rightarrow \mathbb{R} \quad \implies \quad \eta_1(\cdot, \partial_1 Z) + \eta_2(\cdot, \partial_2 Z) = dH$
- ??? \implies vector field

Hamiltonians?

- $H : M \rightarrow \mathbb{R} \quad \implies \quad \eta_1(\cdot, \partial_1 Z) + \eta_2(\cdot, \partial_2 Z) = dH$
- $F : M \rightarrow \mathbb{R}^2? \quad \implies \quad \text{vector field}$

Hamiltonians?

- $H : M \rightarrow \mathbb{R} \quad \implies \quad \eta_1(\cdot, \partial_1 Z) + \eta_2(\cdot, \partial_2 Z) = dH$
- $F : M \rightarrow \mathbb{R}^2 \quad \implies \quad \text{vector field}$

Definition

A function $F : M \rightarrow \mathbb{R}^2$ is called a *current* if there exists a vector field X_F on M such that

$$\Omega(X_F, \cdot) = dF$$

Hamiltonians?

- $H : M \rightarrow \mathbb{R} \quad \implies \quad \eta_1(\cdot, \partial_1 Z) + \eta_2(\cdot, \partial_2 Z) = dH$
- $F : M \rightarrow \mathbb{R}^2 \quad \implies \quad \text{vector field}$

Definition

A function $F : M \rightarrow \mathbb{R}^2$ is called a *current* if there exists a vector field X_F on M such that

$$\Omega(X_F, \cdot) = dF$$

The flow of these vector fields preserve Ω .

The good and the bad

- ✓ Geometric framework
for PDEs

The good and the bad

- ✓ Geometric framework for PDEs
- ✓ Hamiltonian and Lagrangian formalism

The good and the bad

- ✓ Geometric framework for PDEs
- ✓ Hamiltonian and Lagrangian formalism
- ✓ Action functional

The good and the bad

- ✓ Geometric framework for PDEs
- ✓ Hamiltonian and Lagrangian formalism
- ✓ Action functional

✗ No Darboux theorem

The good and the bad

- ✓ Geometric framework for PDEs
- ✓ Hamiltonian and Lagrangian formalism
- ✓ Action functional

- ✗ No Darboux theorem
- ✗ No holomorphic curve techniques

The good and the bad

- ✓ Geometric framework for PDEs
- ✓ Hamiltonian and Lagrangian formalism
- ✓ Action functional

- ✗ No Darboux theorem
- ✗ No holomorphic curve techniques
- ✗ Not elliptic

Laplace equation

$$-(\partial_1^2 + \partial_2^2)q = \nabla V(q)$$

$$q = (q_1, q_2) : \mathbb{R}^2 \rightarrow \mathbb{R}^{2n}$$

Laplace equation

$$-(\partial_1^2 + \partial_2^2)q = \nabla V(q)$$

$$q = (q_1, q_2) : \mathbb{R}^2 \rightarrow \mathbb{R}^{2n}$$

$$-4\partial_{\bar{t}}\partial_t q = \nabla V(q)$$

$$\partial_t = \frac{1}{2}(\partial_1 - i\partial_2)$$

Laplace equation

$$-(\partial_1^2 + \partial_2^2)q = \nabla V(q) \qquad q = (q_1, q_2) : \mathbb{R}^2 \rightarrow \mathbb{R}^{2n}$$

$$-4\partial_{\bar{t}}\partial_t q = \nabla V(q) \qquad \partial_t = \frac{1}{2}(\partial_1 - i\partial_2)$$

De Donder-Weyl: 4 momentum vectors $p_i^\alpha = \partial_\alpha q_i$

Laplace equation

$$-(\partial_1^2 + \partial_2^2)q = \nabla V(q) \qquad q = (q_1, q_2) : \mathbb{R}^2 \rightarrow \mathbb{R}^{2n}$$

$$-4\partial_{\bar{t}}\partial_t q = \nabla V(q) \qquad \partial_t = \frac{1}{2}(\partial_1 - i\partial_2)$$

De Donder-Weyl: 4 momentum vectors $p_i^\alpha = \partial_\alpha q_i$

Only need: $p = 2\partial_t q$

Laplace equation

$$-(\partial_1^2 + \partial_2^2)q = \nabla V(q) \quad q = (q_1, q_2) : \mathbb{R}^2 \rightarrow \mathbb{R}^{2n}$$

$$-4\partial_{\bar{t}}\partial_t q = \nabla V(q) \quad \partial_t = \frac{1}{2}(\partial_1 - i\partial_2)$$

De Donder-Weyl: 4 momentum vectors $p_i^\alpha = \partial_\alpha q_i$

Only need: $p = 2\partial_t q$

$$p_1 = \partial_1 q_1 + \partial_2 q_2$$

$$p_2 = \partial_1 q_2 - \partial_2 q_1$$

Laplace equation

$$-(\partial_1^2 + \partial_2^2)q = \nabla V(q) \quad q = (q_1, q_2) : \mathbb{R}^2 \rightarrow \mathbb{R}^{2n}$$

$$-4\partial_{\bar{t}}\partial_t q = \nabla V(q) \quad \partial_t = \frac{1}{2}(\partial_1 - i\partial_2)$$

De Donder-Weyl: 4 momentum vectors $p_i^\alpha = \partial_\alpha q_i$

Only need: $p = 2\partial_t q, -2\partial_{\bar{t}} p = \nabla V(q)$

$$p_1 = \partial_1 q_1 + \partial_2 q_2$$

$$p_2 = \partial_1 q_2 - \partial_2 q_1$$

$$\partial_{q_1} V = -\partial_1 p_1 + \partial_2 p_2$$

$$\partial_{q_2} V = -\partial_1 p_2 - \partial_2 p_1$$

Laplace equation

$$-(\partial_1^2 + \partial_2^2)q = \nabla V(q) \quad q = (q_1, q_2) : \mathbb{R}^2 \rightarrow \mathbb{R}^{2n}$$

$$-4\partial_{\bar{t}}\partial_t q = \nabla V(q) \quad \partial_t = \frac{1}{2}(\partial_1 - i\partial_2)$$

De Donder-Weyl: 4 momentum vectors $p_i^\alpha = \partial_\alpha q_i$

Only need: $p = 2\partial_t q$, $-2\partial_{\bar{t}} p = \nabla V(q)$

$$\not\partial Z := \begin{pmatrix} 0 & -2\partial_{\bar{t}} \\ 2\partial_t & 0 \end{pmatrix} Z = \nabla H(Z),$$

where

$$Z = (q, p) : \mathbb{R}^2 \rightarrow \mathbb{R}^{4n}$$

$$H(q, p) = \frac{1}{2}|p|^2 + V(q)$$

Polysymplectic formulation

$$\delta Z := \begin{pmatrix} 0 & -2\partial_{\bar{t}} \\ 2\partial_t & 0 \end{pmatrix} Z = \nabla H(Z) \quad Z = (q_1, q_2, p_1, p_2)$$

Polysymplectic formulation

$$\delta Z := \begin{pmatrix} 0 & -2\partial_{\bar{t}} \\ 2\partial_t & 0 \end{pmatrix} Z = \nabla H(Z) \quad Z = (q_1, q_2, p_1, p_2)$$

Define $\Omega = \omega_1 \otimes \partial_1 + \omega_2 \otimes \partial_2$ for

$$\omega_1 = dp_1 \wedge dq_1 + dp_2 \wedge dq_2 \quad \omega_2 = dp_1 \wedge dq_2 - dp_2 \wedge dq_1$$

Polysymplectic formulation

$$\partial Z := \begin{pmatrix} 0 & -2\partial_{\bar{t}} \\ 2\partial_t & 0 \end{pmatrix} Z = \nabla H(Z) \quad Z = (q_1, q_2, p_1, p_2)$$

Define $\Omega = \omega_1 \otimes \partial_1 + \omega_2 \otimes \partial_2$ for

$$\omega_1 = dp_1 \wedge dq_1 + dp_2 \wedge dq_2 \quad \omega_2 = dp_1 \wedge dq_2 - dp_2 \wedge dq_1$$

The Dirac equation is equivalent to

$$\omega_1(\cdot, \partial_1 Z) + \omega_2(\cdot, \partial_2 Z) = dH$$

Polysymplectic formulation

Define $\Omega = \omega_1 \otimes \partial_1 + \omega_2 \otimes \partial_2$ for

$$\omega_1 = dp_1 \wedge dq_1 + dp_2 \wedge dq_2 \quad \omega_2 = dp_1 \wedge dq_2 - dp_2 \wedge dq_1$$

Note

- Both ω_1 and ω_2 are symplectic forms related by $\omega_2 = -\omega_1(\cdot, I\cdot)$, where

$$I = \begin{pmatrix} i & 0 \\ 0 & i^* \end{pmatrix} = \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix}$$

on $\mathbb{R}^{4n} = \mathbb{R}^{2n} \times (\mathbb{R}^{2n})^*$.

Polysymplectic formulation

Define $\Omega = \omega_1 \otimes \partial_1 + \omega_2 \otimes \partial_2$ for

$$\omega_1 = dp_1 \wedge dq_1 + dp_2 \wedge dq_2 \quad \omega_2 = dp_1 \wedge dq_2 - dp_2 \wedge dq_1$$

Note

- Both ω_1 and ω_2 are symplectic forms related by $\omega_2 = -\omega_1(\cdot, I\cdot)$, where

$$I = \begin{pmatrix} i & 0 \\ 0 & i^* \end{pmatrix} = \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix}$$

on $\mathbb{R}^{4n} = \mathbb{R}^{2n} \times (\mathbb{R}^{2n})^*$.

- Let $\omega^{\mathbb{C}} = dp \wedge dq = \omega_1 + i\omega_2$, then

$$\Omega = \omega^{\mathbb{C}} \otimes \partial_t + \bar{\omega}^{\mathbb{C}} \otimes \partial_{\bar{t}}.$$

Definition (B.-F. '24)

Let (W, I) a complex manifold. An \mathbb{R}^2 -valued polysymplectic form $\Omega = \omega_1 \otimes \partial_1 + \omega_2 \otimes \partial_2$ on W is called *complex-regularized* if $\omega_2 = -\omega_1(\cdot, I\cdot)$.

Definition (B.-F. '24)

Let (W, I) a complex manifold. An \mathbb{R}^2 -valued polysymplectic form $\Omega = \omega_1 \otimes \partial_1 + \omega_2 \otimes \partial_2$ on W is called *complex-regularized* if $\omega_2 = -\omega_1(\cdot, I\cdot)$.

The triple (W, I, Ω) is called a *complex-regularized polysymplectic (CRPS) manifold*.

Example

Let (Q, i) a complex manifold and $W = T^*Q$ with induced complex structure I .

Example

Let (Q, i) a complex manifold and $W = T^*Q$ with induced complex structure I . Note $\pi : T^*Q \rightarrow Q$ is holomorphic.

Example

Let (Q, i) a complex manifold and $W = T^*Q$ with induced complex structure I . Note $\pi : T^*Q \rightarrow Q$ is holomorphic. For $Z = (q, p) \in W$ and $X \in T_Z W$ define

$$(\theta_1)_Z(X) = p \circ d\pi(X)$$

$$(\theta_2)_Z(X) = p \circ d\pi(IX).$$

Example

Let (Q, i) a complex manifold and $W = T^*Q$ with induced complex structure I . Note $\pi : T^*Q \rightarrow Q$ is holomorphic. For $Z = (q, p) \in W$ and $X \in T_Z W$ define

$$\begin{aligned}(\theta_1)_Z(X) &= p \circ d\pi(X) \\ (\theta_2)_Z(X) &= p \circ d\pi(IX).\end{aligned}$$

Then $\Omega = d\theta_1 \otimes \partial_1 - d\theta_2 \otimes \partial_2$ is CRPS on (W, I) .

First properties

Lemma

Let $(W, I, \Omega = \omega_1 \otimes \partial_1 + \omega_2 \otimes \partial_2)$ a CRPS manifold. Then both ω_1 and ω_2 are symplectic forms.

First properties

Lemma

Let $(W, I, \Omega = \omega_1 \otimes \partial_1 + \omega_2 \otimes \partial_2)$ a CRPS manifold. Then both ω_1 and ω_2 are symplectic forms.

Proof idea: Since $\omega_2 = -\omega_1(\cdot, I\cdot)$

$$\ker \omega_1^b = \ker \omega_2^b.$$

First properties

Lemma

Let $(W, I, \Omega = \omega_1 \otimes \partial_1 + \omega_2 \otimes \partial_2)$ a CRPS manifold. Then both ω_1 and ω_2 are symplectic forms.

Proof idea: Since $\omega_2 = -\omega_1(\cdot, I\cdot)$

$$\ker \omega_1^b = \ker \omega_2^b.$$

So Ω being non-degenerate implies $\ker \omega_1^b = \ker \omega_2^b = 0$. □

First properties

Lemma

Let $(W, I, \Omega = \omega_1 \otimes \partial_1 + \omega_2 \otimes \partial_2)$ a CRPS manifold. Then both ω_1 and ω_2 are symplectic forms.

Lemma

Let $\psi : (W, I, \Omega) \rightarrow (W', I', \Omega')$ a diffeomorphism between CRPS manifolds, such that

$$\psi^* \Omega' = \Omega.$$

Then ψ is holomorphic with respect to I and I' .

Quick recap

Definition

A holomorphic 2-form $\omega^{\mathbb{C}}$ on a complex manifold (W, I) is called a *holomorphic symplectic form* if it is closed and its restriction to $T^{(1,0)}W$ is non-degenerate.

Quick recap

Definition

A holomorphic 2-form $\omega^{\mathbb{C}}$ on a complex manifold (W, I) is called a *holomorphic symplectic form* if it is closed and its restriction to $T^{(1,0)}W$ is non-degenerate.

A holomorphic Hamiltonian system is a tuple $(W, I, \omega^{\mathbb{C}}, F)$, where $F : W \rightarrow \mathbb{C}$ is holomorphic.

Quick recap

Definition

A holomorphic 2-form $\omega^{\mathbb{C}}$ on a complex manifold (W, I) is called a *holomorphic symplectic form* if it is closed and its restriction to $T^{(1,0)}W$ is non-degenerate.

A holomorphic Hamiltonian system is a tuple $(W, I, \omega^{\mathbb{C}}, F)$, where $F : W \rightarrow \mathbb{C}$ is holomorphic. It induces a holomorphic vector field \mathcal{X}_F by

$$\omega^{\mathbb{C}}(\mathcal{X}_F, \cdot) = dF.$$

Relation to CRPS manifolds

Proposition

For a complex manifold (W, I) there is a bijection

$$\{\text{CRPS forms}\} \xleftrightarrow{1-1} \{\text{holomorphic symplectic forms}\}$$

Given by

$$\begin{aligned}\omega_1 \otimes \partial_1 + \omega_2 \otimes \partial_2 &\mapsto \omega_1 + i\omega_2 \\ \omega^{\mathbb{C}} \otimes \partial_t + \bar{\omega}^{\mathbb{C}} \otimes \partial_{\bar{t}} &\leftarrow \omega^{\mathbb{C}}\end{aligned}$$

Introduction
○○

Polysymplectic geometry
○○○○

Dirac operators
○○○○○○

Holomorphic symplectic geometry
○○●○

Rigidity
○○○○○

Application
○○

End
○○

CRPS geometry

Holomorphic symplectic geometry

CRPS geometry

- Real valued Hamiltonians

Holomorphic symplectic geometry

CRPS geometry

- Real valued Hamiltonians

Holomorphic symplectic geometry

- Holomorphic Hamiltonians

CRPS geometry

- Real valued Hamiltonians

Holomorphic symplectic geometry

- Holomorphic Hamiltonians

Recall: $F : (W, I, \Omega) \rightarrow \mathbb{R}^2$ is a current if there exists X_F such that $\Omega(X_F, \cdot) = dF$.

CRPS geometry

- Real valued Hamiltonians

Holomorphic symplectic geometry

- Holomorphic Hamiltonians

Recall: $F : (W, I, \Omega) \rightarrow \mathbb{R}^2$ is a current if there exists X_F such that $\Omega(X_F, \cdot) = dF$.

Lemma

Let $(W, I, \omega_1 \otimes \partial_1 + \omega_2 \otimes \partial_2)$ a CRPS manifold and $\omega^{\mathbb{C}} = \omega_1 + i\omega_2$. By identifying $\mathbb{R}^2 \cong \mathbb{C}$ and $TW \cong T^{(1,0)}W$ we get that $F : W \rightarrow \mathbb{R}^2 \cong \mathbb{C}$ is a current if and only if it is a holomorphic function.

CRPS geometry

- Real valued Hamiltonians

Holomorphic symplectic geometry

- Holomorphic Hamiltonians

Recall: $F : (W, I, \Omega) \rightarrow \mathbb{R}^2$ is a current if there exists X_F such that $\Omega(X_F, \cdot) = dF$.

Lemma

Let $(W, I, \omega_1 \otimes \partial_1 + \omega_2 \otimes \partial_2)$ a CRPS manifold and $\omega^{\mathbb{C}} = \omega_1 + i\omega_2$. By identifying $\mathbb{R}^2 \cong \mathbb{C}$ and $TW \cong T^{(1,0)}W$ we get that $F : W \rightarrow \mathbb{R}^2 \cong \mathbb{C}$ is a current if and only if it is a holomorphic function. In this case $X_F = \mathcal{X}_F$.

Darboux theorem

Corollary

Let (W, I, Ω) a CRPS manifold. Around every point in W there exist coordinates $\{q_1^\alpha, q_2^\alpha, p_1^\alpha, p_2^\alpha\}$ where $\alpha = 1, \dots, n$ such that

$$I \frac{\partial}{\partial q_1^\alpha} = \frac{\partial}{\partial q_2^\alpha} \qquad I \frac{\partial}{\partial p_1^\alpha} = -\frac{\partial}{\partial p_2^\alpha}$$

and

$$\begin{aligned} \omega_1 &= \sum_{\alpha} (dp_1^\alpha \wedge dq_1^\alpha + dp_2^\alpha \wedge dq_2^\alpha) \\ \omega_2 &= \sum_{\alpha} (dp_1^\alpha \wedge dq_2^\alpha - dp_2^\alpha \wedge dq_1^\alpha). \end{aligned}$$

Darboux theorem

Corollary

$$I \frac{\partial}{\partial q_1^\alpha} = \frac{\partial}{\partial q_2^\alpha}$$

$$I \frac{\partial}{\partial p_1^\alpha} = - \frac{\partial}{\partial p_2^\alpha}$$

and

$$\omega_1 = \sum_{\alpha} (dp_1^\alpha \wedge dq_1^\alpha + dp_2^\alpha \wedge dq_2^\alpha)$$

$$\omega_2 = \sum_{\alpha} (dp_1^\alpha \wedge dq_2^\alpha - dp_2^\alpha \wedge dq_1^\alpha).$$

Proof: Follows from the Darboux theorem for holomorphic symplectic manifolds (see thesis of Wagner for proof).



Action functional

Assume $\Omega = d\Theta$ on W is exact.

Action functional

Assume $\Omega = d\Theta$ on W is exact.

Let $d\mathcal{V} = dt_1 \wedge dt_2 = -\frac{1}{2i} dt \wedge d\bar{t}$ the volume form on \mathbb{T}^2 .

Action functional

Assume $\Omega = d\Theta$ on W is exact.

Let $d\mathcal{V} = dt_1 \wedge dt_2 = -\frac{1}{2i} dt \wedge d\bar{t}$ the volume form on \mathbb{T}^2 .

Contract $\Theta = \theta_1 \otimes \partial_1 + \theta_2 \otimes \partial_2$ with $d\mathcal{V}$ to give

$$\tilde{\Theta} = \theta_1 \wedge dt_2 - \theta_2 \wedge dt_1 \in \Lambda^2(W \times \mathbb{T}^2).$$

Action functional

Assume $\Omega = d\Theta$ on W is exact.

Let $d\mathcal{V} = dt_1 \wedge dt_2 = -\frac{1}{2i} dt \wedge d\bar{t}$ the volume form on \mathbb{T}^2 .

Contract $\Theta = \theta_1 \otimes \partial_1 + \theta_2 \otimes \partial_2$ with $d\mathcal{V}$ to give

$$\tilde{\Theta} = \theta_1 \wedge dt_2 - \theta_2 \wedge dt_1 \in \Lambda^2(W \times \mathbb{T}^2).$$

For $Z : \mathbb{T}^2 \rightarrow W$ define $\tilde{Z} = Z \times \text{id} : \mathbb{T}^2 \rightarrow W \times \mathbb{T}^2$ and

$$\begin{aligned} \mathcal{A}(Z) &= \int_{\mathbb{T}^2} \tilde{Z}^* \tilde{\Theta} \\ &= \int_{\mathbb{T}^2} (\theta_1(\partial_1 Z) + \theta_2(\partial_2 Z)) d\mathcal{V} \end{aligned}$$

Action functional

$$\mathcal{A}(Z) = \int_{\mathbb{T}^2} (\theta_1(\partial_1 Z) + \theta_2(\partial_2 Z)) \, d\mathcal{V}$$

Action functional

$$\mathcal{A}(Z) = \int_{\mathbb{T}^2} (\theta_1(\partial_1 Z) + \theta_2(\partial_2 Z)) \, d\mathcal{V}$$

- The action is real-valued so we may study its gradient lines.

Action functional

$$\mathcal{A}(Z) = \int_{\mathbb{T}^2} (\theta_1(\partial_1 Z) + \theta_2(\partial_2 Z)) \, d\mathcal{V}$$

- The action is real-valued so we may study its gradient lines.
- Critical points of

$$\mathcal{A}_H(Z) = \mathcal{A}(Z) - \int_{\mathbb{T}^2} H(Z) \, d\mathcal{V}$$

are multiperiodic solutions to the corresponding PDE.

Action functional

$$\mathcal{A}(Z) = \int_{\mathbb{T}^2} (\theta_1(\partial_1 Z) + \theta_2(\partial_2 Z)) \, d\mathcal{V}$$

- The action is real-valued so we may study its gradient lines.
- Critical points of

$$\mathcal{A}_H(Z) = \mathcal{A}(Z) - \int_{\mathbb{T}^2} H(Z) \, d\mathcal{V}$$

are multiperiodic solutions to the corresponding PDE.

- Pick a metric g such that $\omega_1 = g(\cdot, J\cdot)$ and $\omega_2 = g(\cdot, K\cdot)$ for almost complex structures J, K , then $K = IJ$.

Action functional

$$\mathcal{A}(Z) = \int_{\mathbb{T}^2} (\theta_1(\partial_1 Z) + \theta_2(\partial_2 Z)) \, d\mathcal{V}$$

- The action is real-valued so we may study its gradient lines.
- Critical points of

$$\mathcal{A}_H(Z) = \mathcal{A}(Z) - \int_{\mathbb{T}^2} H(Z) \, d\mathcal{V}$$

are multiperiodic solutions to the corresponding PDE.

- Pick a metric g such that $\omega_1 = g(\cdot, J\cdot)$ and $\omega_2 = g(\cdot, K\cdot)$ for almost complex structures J, K , then $K = IJ$.

$$\text{grad}_{L^2} \mathcal{A}(Z) = J\partial_1 Z + K\partial_2 Z$$

Action functional

$$\mathcal{A}(Z) = \int_{\mathbb{T}^2} (\theta_1(\partial_1 Z) + \theta_2(\partial_2 Z)) \, d\mathcal{V}$$

- The action is real-valued so we may study its gradient lines.
- Critical points of

$$\mathcal{A}_H(Z) = \mathcal{A}(Z) - \int_{\mathbb{T}^2} H(Z) \, d\mathcal{V}$$

are multiperiodic solutions to the corresponding PDE.

- Pick a metric g such that $\omega_1 = g(\cdot, J\cdot)$ and $\omega_2 = g(\cdot, K\cdot)$ for almost complex structures J, K , then $K = IJ$.

$$\text{grad}_{L^2} \mathcal{A}(Z) = J\partial_1 Z + K\partial_2 Z$$

Gradient lines are given by the Fueter equation

$$\partial_s Z + J\partial_1 Z + K\partial_2 Z = 0$$

Action functional

$$\mathcal{A}(Z) = \int_{\mathbb{T}^2} (\theta_1(\partial_1 Z) + \theta_2(\partial_2 Z)) \, d\mathcal{V}$$

- The action is real-valued so we may study its gradient lines.
- Critical points of

$$\mathcal{A}_H(Z) = \mathcal{A}(Z) - \int_{\mathbb{T}^2} H(Z) \, d\mathcal{V}$$

are multiperiodic solutions to the corresponding PDE.

- Pick a metric g such that $\omega_1 = g(\cdot, J\cdot)$ and $\omega_2 = g(\cdot, K\cdot)$ for almost complex structures J, K , then $K = IJ$.

$$\text{grad}_{L^2} \mathcal{A}(Z) = J\partial_1 Z + K\partial_2 Z$$

Gradient lines are given by the Fueter equation

$$I\partial_s Z + K\partial_1 Z - J\partial_2 Z = 0.$$

Arnold conjecture

Theorem (B.-F. '24)

Let $Q = \mathbb{T}^{2n}$ and $W = T^*Q$ with the standard CRPS form $\omega_1 \otimes \partial_1 + \omega_2 \otimes \partial_2$.

Arnold conjecture

Theorem (B.-F. '24)

Let $Q = \mathbb{T}^{2n}$ and $W = T^*Q$ with the standard CRPS form $\omega_1 \otimes \partial_1 + \omega_2 \otimes \partial_2$.

If $H : \mathbb{T}^2 \times W \rightarrow \mathbb{R}$ is given by $H(t, q, p) = \frac{1}{2}|p|^2 + h(t, q, p)$ for $h : \mathbb{T}^2 \times W \rightarrow \mathbb{R}$ smooth with finite C^2 -norm,

Arnold conjecture

Theorem (B.-F. '24)

Let $Q = \mathbb{T}^{2n}$ and $W = T^*Q$ with the standard CRPS form $\omega_1 \otimes \partial_1 + \omega_2 \otimes \partial_2$.

If $H : \mathbb{T}^2 \times W \rightarrow \mathbb{R}$ is given by $H(t, q, p) = \frac{1}{2}|p|^2 + h(t, q, p)$ for $h : \mathbb{T}^2 \times W \rightarrow \mathbb{R}$ smooth with finite C^2 -norm, then

$$dH = \omega_1(\cdot, \partial_1 Z) + \omega_2(\cdot, \partial_2 Z)$$

has at least $(2n + 1)$ solutions.

Arnold conjecture

Theorem (B.-F. '24)

Let $Q = \mathbb{T}^{2n}$ and $W = T^*Q$ with the standard CRPS form $\omega_1 \otimes \partial_1 + \omega_2 \otimes \partial_2$.

If $H : \mathbb{T}^2 \times W \rightarrow \mathbb{R}$ is given by $H(t, q, p) = \frac{1}{2}|p|^2 + h(t, q, p)$ for $h : \mathbb{T}^2 \times W \rightarrow \mathbb{R}$ smooth with finite C^2 -norm, then

$$dH = \omega_1(\cdot, \partial_1 Z) + \omega_2(\cdot, \partial_2 Z)$$

has at least $(2n + 1)$ solutions.

Corollary

When $V : Q \rightarrow \mathbb{R}$ has finite C^2 -norm then $-\Delta q = \nabla V(q)$ has at least $(2n + 1)$ solutions.

Overview of proof of Arnold conjecture

- Follows from studying Floer curves

$$\partial_s Z + J\partial_1 Z + K\partial_2 Z = \nabla H(Z).$$

Overview of proof of Arnold conjecture

- Follows from studying Floer curves

$$\partial_s Z + J\partial_1 Z + K\partial_2 Z = \nabla H(Z).$$

- C^0 -bounds for solutions and for Floer curves.

Overview of proof of Arnold conjecture

- Follows from studying Floer curves

$$\partial_s Z + J\partial_1 Z + K\partial_2 Z = \nabla H(Z).$$

- C^0 -bounds for solutions and for Floer curves.
- Moduli space of Floer curves is a compact 1-dimensional manifold for generic choice of h .

Overview of proof of Arnold conjecture

- Follows from studying Floer curves

$$\partial_s Z + J\partial_1 Z + K\partial_2 Z = \nabla H(Z).$$

- C^0 -bounds for solutions and for Floer curves.
- Moduli space of Floer curves is a compact 1-dimensional manifold for generic choice of h .
- Rest of the proof follows the line of reasoning of the cuplength results from Albers-Hein.

Non-squeezing

Let $(W, I, \Omega = \omega_1 \otimes \partial_1 + \omega_2 \otimes \partial_2)$ a CRPS manifold and $\psi : W \rightarrow W$ a diffeomorphism.

Non-squeezing

Let $(W, I, \Omega = \omega_1 \otimes \partial_1 + \omega_2 \otimes \partial_2)$ a CRPS manifold and $\psi : W \rightarrow W$ a diffeomorphism.

$$\psi^* \Omega = \Omega \quad \Longleftrightarrow \quad \psi^* \omega_1 = \omega_1 \text{ and } \psi^* \omega_2 = \omega_2$$

Non-squeezing

Let $(W, I, \Omega = \omega_1 \otimes \partial_1 + \omega_2 \otimes \partial_2)$ a CRPS manifold and $\psi : W \rightarrow W$ a diffeomorphism.

$$\psi^* \Omega = \Omega \quad \Longleftrightarrow \quad \psi^* \omega_1 = \omega_1 \text{ and } \psi^* \omega_2 = \omega_2$$

Theorem

Let $\psi : \mathbb{R}^{4n} \rightarrow \mathbb{R}^{4n}$ a diffeomorphism preserving Ω such that

$$\psi(B_r^{4n}) \subseteq B_R^2 \times \mathbb{R}^{4n-2}$$

where B_R^2 is the R -ball in the (q_i^α, p_j^α) -plane for some $i, j \in \{1, 2\}$ and $\alpha \in \{1, \dots, n\}$.

Then $r \leq R$.

Non-squeezing

Let $(W, I, \Omega = \omega_1 \otimes \partial_1 + \omega_2 \otimes \partial_2)$ a CRPS manifold and $\psi : W \rightarrow W$ a diffeomorphism.

$$\psi^* \Omega = \Omega \quad \Longleftrightarrow \quad \psi^* \omega_1 = \omega_1 \text{ and } \psi^* \omega_2 = \omega_2$$

Corollary

Let $\psi_\nu : W \rightarrow W$ a sequence of diffeomorphisms preserving Ω that converge to a diffeomorphism $\psi : W \rightarrow W$ in the C^0 -limit. Then $\psi^* \Omega = \Omega$.

Holomorphic Lagrangians \rightarrow harmonic maps

Question: Given a holomorphic symplectic manifold $(W^{4n}, \omega^{\mathbb{C}})$ and a closed complex manifold L^{2n} , is there a holomorphic embedding $\iota : L \hookrightarrow W$ such that $\iota^* \omega^{\mathbb{C}} = 0$.

Holomorphic Lagrangians \rightarrow harmonic maps

Question: Given a holomorphic symplectic manifold $(W^{4n}, \omega^{\mathbb{C}})$ and a closed complex manifold L^{2n} , is there a holomorphic embedding $\iota : L \hookrightarrow W$ such that $\iota^* \omega^{\mathbb{C}} = 0$.


Real symplectic manifolds

Holomorphic Lagrangians \rightarrow harmonic maps

Question: Given a holomorphic symplectic manifold $(W^{4n}, \omega^{\mathbb{C}})$ and a closed complex manifold L^{2n} , is there a holomorphic embedding $\iota : L \hookrightarrow W$ such that $\iota^* \omega^{\mathbb{C}} = 0$.

Real symplectic manifolds

Morse theory of geodesics on L




Holomorphic Lagrangians \rightarrow harmonic maps

Question: Given a holomorphic symplectic manifold $(W^{4n}, \omega^{\mathbb{C}})$ and a closed complex manifold L^{2n} , is there a holomorphic embedding $\iota : L \hookrightarrow W$ such that $\iota^* \omega^{\mathbb{C}} = 0$.

Holomorphic symplectic manifolds

Morse theory of geodesics on L




Holomorphic Lagrangians \rightarrow harmonic maps

Question: Given a holomorphic symplectic manifold $(W^{4n}, \omega^{\mathbb{C}})$ and a closed complex manifold L^{2n} , is there a holomorphic embedding $\iota : L \hookrightarrow W$ such that $\iota^* \omega^{\mathbb{C}} = 0$.

Holomorphic symplectic manifolds

Morse theory of **minimal spheres** on L



Harmonic maps \rightarrow holomorphic Lagrangians

Question: Given a Kähler manifold (Q, i, g) , does there exist a minimal surface $\Sigma \rightarrow Q$ with boundary $\partial\Sigma$ in some submanifold $\Gamma \subseteq Q$?

Harmonic maps \rightarrow holomorphic Lagrangians

Question: Given a Kähler manifold (Q, i, g) , does there exist a minimal surface $\Sigma \rightarrow Q$ with boundary $\partial\Sigma$ in some submanifold $\Gamma \subseteq Q$?

One-dimensional analogue

Harmonic maps \rightarrow holomorphic Lagrangians

Question: Given a Kähler manifold (Q, i, g) , does there exist a minimal surface $\Sigma \rightarrow Q$ with boundary $\partial\Sigma$ in some submanifold $\Gamma \subseteq Q$?

One-dimensional analogue

Existence of geodesics with boundary on $\Gamma \subseteq Q$

Harmonic maps \rightarrow holomorphic Lagrangians

Question: Given a Kähler manifold (Q, i, g) , does there exist a minimal surface $\Sigma \rightarrow Q$ with boundary $\partial\Sigma$ in some submanifold $\Gamma \subseteq Q$?

One-dimensional analogue

Existence of geodesics with boundary on $\Gamma \subseteq Q$

$$H(q, p) = \frac{1}{2}|p|_g^2 \text{ on } T^*Q$$

Harmonic maps \rightarrow holomorphic Lagrangians

Question: Given a Kähler manifold (Q, i, g) , does there exist a minimal surface $\Sigma \rightarrow Q$ with boundary $\partial\Sigma$ in some submanifold $\Gamma \subseteq Q$?

One-dimensional analogue

Existence of geodesics with boundary on $\Gamma \subseteq Q$

$$H(q, p) = \frac{1}{2}|p|_g^2 \text{ on } T^*Q$$

Boundary fixed to the conormal bundle $N^*\Gamma \subseteq T^*Q$

Harmonic maps \rightarrow holomorphic Lagrangians

Question: Given a Kähler manifold (Q, i, g) , does there exist a minimal surface $\Sigma \rightarrow Q$ with boundary $\partial\Sigma$ in some submanifold $\Gamma \subseteq Q$?

Minimal surface case

Existence of geodesics with boundary on $\Gamma \subseteq Q$

$$H(q, p) = \frac{1}{2}|p|_g^2 \text{ on } T^*Q$$

Boundary fixed to the conormal bundle $N^*\Gamma \subseteq T^*Q$

Harmonic maps \rightarrow holomorphic Lagrangians

Question: Given a Kähler manifold (Q, i, g) , does there exist a minimal surface $\Sigma \rightarrow Q$ with boundary $\partial\Sigma$ in some submanifold $\Gamma \subseteq Q$?

Minimal surface case

Existence of **harmonic maps** with boundary on $\Gamma \subseteq Q$

$$H(q, p) = \frac{1}{2}|p|_g^2 \text{ on } T^*Q$$

Boundary fixed to the conormal bundle $N^*\Gamma \subseteq T^*Q$

Harmonic maps \rightarrow holomorphic Lagrangians

Question: Given a Kähler manifold (Q, i, g) , does there exist a minimal surface $\Sigma \rightarrow Q$ with boundary $\partial\Sigma$ in some submanifold $\Gamma \subseteq Q$?




Minimal surface case

Existence of **harmonic maps** with boundary on $\Gamma \subseteq Q$

$$H(q, p) = \frac{1}{2}|p|_g^2 \text{ on } T^*Q$$

Boundary fixed to the conormal bundle $N^*\Gamma \subseteq T^*Q \rightarrow$
holomorphic Lagrangian if Γ is a complex submanifold

Bibliography

-  Ronen Brilleslijper and Oliver Fabert.
 Regularized polysymplectic geometry and first steps towards
 Floer theory for covariant field theories.
Journal of Geometry and Physics, 183:104703, 2023.
-  Ronen Brilleslijper and Oliver Fabert.
Generalizing symplectic topology from 1 to 2 dimensions.
arXiv preprint arXiv:2412.16223, 2024.
-  Oliver Fabert and Ronen Brilleslijper.
 From Euclidean field theory to hyperkähler Floer theory via
 regularized polysymplectic geometry.
Communications in Contemporary Mathematics, 2025.

Thank you!