

Chapter 7

Logistic Regression Models

This chapter introduces the modeling framework for categorical data in the simple situation where we have a categorical response variable, often binary, and one or more explanatory variables. A fitted model provides both statistical inference and prediction, accompanied by measures of uncertainty. Data visualization methods for discrete response data must often rely on smoothing techniques, including both direct, non-parametric smoothing and the implicit smoothing that results from a fitted parametric model. Diagnostic plots help us to detect influential observations which may distort our results.

{ch:logistic}

7.1 Introduction

{sec:logist-intro}

All models are wrong, but some are useful

George E. P. Box, (Box and Draper, 1987, p. 424)

Chapters 4–6 have been concerned primarily with simple, exploratory methods for studying the relations among categorical variables and with testing hypotheses about their associations through non-parametric tests and with overall goodness-of-fit statistics.

This chapter begins our study of model-based methods for the analysis of discrete data. These models differ from those we have examined earlier primarily in that they consider *explicitly* an assumed probability distribution for the observations, and make clear distinctions between the systematic component, which is explained by the model, and the random component, which is not. More importantly, the model-based approach allows a compact summary of categorical data in terms of a (hopefully) small number of parameters accompanied by measures of uncertainty (standard errors), and the ability to estimate predicted values over the range of explanatory variables.

This model-fitting approach has several advantages: (a) Inferences for the model parameters include both hypothesis tests and confidence intervals. (b) The former help us to assess which explanatory variables affect the outcome; the size of the estimated parameters and the widths of their confidence intervals help us to assess the strength and importance of these effects. (c) There are a variety of methods for model selection, designed to help determine a favorable trade-off between goodness-of-fit and parsimony. (d) Finally, the predicted values obtained from the model effectively smooth the discrete responses, allow predictions for unobserved values of the explanatory variables, and provide important means to interpret the fitted relationship graphically.

Figure 7.1 provides a visual overview of the steps for fitting and graphing with model-based methods in R. (a) A modeling function such as `glm()` is applied to an input data frame. The result




Figure 7.1: Overview of fitting and graphing for model-based methods in R.

{fig:governview}

is a **model object** containing all the information from the fitting process. (b) As is standard in R, `print()` and `summary()` methods give, respectively, basic and detailed printed output. (c) Many modeling functions have `plot()` methods that produce different types of summary and diagnostic plots. (d) For visualizing the fitted model, most model methods provide a `predict()` method that can be used to plot the fitted values from the model over the ranges of the predictors. Such plots can be customized by the addition of points (showing the observations), lines, confidence bands, and so forth.

In this chapter we consider models for a **binary response**, such as “success” or “failure”, or the number of “successes” in a fixed number of “trials”, where we might reasonably assume a binomial distribution for the random component. These methods extend readily to a **polytomous response** with more than two outcome categories, such as improvement in therapy, with categories “none,” “some” and “marked.”

These models can be seen as simple extensions of familiar ANOVA and regression models for quantitative data. They are also important special cases of a more general approach, the **generalized linear model** that subsumes a wide variety of families of techniques within a single, unified framework. However, rather than starting at the top with the fully general version, this chapter details the important special cases of models for discrete outcomes, beginning with binary responses.

This chapter proceeds as follows: in Section 7.2 we introduce the simple logistic regression model for a binary response and a single quantitative predictor. This model extends directly to models for grouped, binomial data (Section 7.2.4) and to models with any number of regressors (Section 7.3), which can be quantitative, discrete factors and more general forms. For interpreting and understanding the results of a fitted model, we emphasize plotting predicted probabilities and predicted log odds in various ways, for which effect plots (Section 7.3.3) are particularly useful for complex models. Individual observations sometimes exert great influence on a fitted model. Some measures of influence and diagnostic plots are illustrated in Section 7.5. In Section 7.6, we develop several approaches to modelling a multi-category (polytomous) response.

7.2 The logistic regression model

The logistic regression model describes the relationship between a discrete outcome variable, the “response”, and a set of explanatory variables. The response variable is often **dichotomous**, although extensions to the model permit multi-category, **polytomous** outcomes, discussed in Section 7.6. The explanatory variables may be continuous or (with factor variables) discrete.

For a binary response, Y , and a continuous explanatory variable, X , we may be interested in modeling the probability of a successful outcome, which we denote $\pi(x) \equiv \Pr(Y = 1 | X = x)$. That is, at a given value $X = x$, you can imagine that there is a binomial distribution of the responses, $\text{Bin}(\pi(x), n_x)$.

The simplest naive model, called the **linear probability model**, supposes that this probability, $\pi(x)$ varies linearly with the value of x ,

$$E(Y | x) = \pi(x) = \alpha + \beta x, \quad (7.1)$$

{eq:logit0}

{sec:logist-model}

where the notation $E(Y | x)$ indicates that the probability $\pi(x)$ represents the population conditional average of the 1s and 0s for all observations with a fixed value of x . For binary observations, this is simply the proportion of 1s.

Figure 7.2 illustrates the basic setup for modeling a binary outcome using the *Arthritis* data, and described more fully in Example 7.1–Example 7.3. The 0/1 observations are shown as (jittered) points. The predicted values under the linear probability model Eqn. (7.1) are shown as the red lines in both panels. As you can see, this model cannot be right, because it predicts a probability less than 0 for small values of Age, and would also predict probabilities greater than 1 for larger values of Age.




Figure 7.2: Arthritis treatment data, for the relationship of the binary response “Better” to Age, shown as jittered points. The left panel shows the predicted values and 95% confidence envelope under the linear probability model. The right panel shows the fitted logistic regression, together with the simple linear regression (red) and a non-parametric (loess) smoothed curve (green).

{fig:arthritis-age}

The linear probability model is also wrong because it assumes that the distribution of residuals, $Y_i - \hat{\pi}(x_i)$ is normal, with mean 0 and constant variance. However, because Y is dichotomous, the residuals are also dichotomous, and have variance $\pi(x_i)(1 - \pi(x_i))$, which is maximal for $\pi = 0.5$ and decreases as π goes toward 0 or 1.

One way around the difficulty of needing to constrain the predicted values to the interval $[0, 1]$ is to re-specify the model so that a *transformation* of π has a linear relation to x , and that transformation keeps $\hat{\pi}$ between 0 and 1 for all x . This idea, of modeling a transformation of the response that has desired statistical properties is one of the fundamental ones that led to the development of **generalized linear models**, which we treat more fully later in Chapter 9.

A particularly convenient choice of the transformation gives the **linear logistic regression model** (or **linear logit model**¹) which posits a linear relation between the **log odds** (or **logit**) of this probability and x ,

$$\text{logit}[\pi(x)] \equiv \log\left(\frac{\pi(x)}{1 - \pi(x)}\right) = \alpha + \beta x. \quad (7.2) \quad \{\text{eq:logit1}\}$$

When $\beta > 0$, $\pi(x)$ and the log odds increase as X increases; when $\beta < 0$ they decrease with X .

¹Some writers use the term *logit model* to refer to those using only categorical predictors; we use the terms logistic regression and logit regression interchangeably.

This model can also be expressed as a model for the probabilities $\pi(x)$ in terms of the *inverse* of the logit transformation used in Eqn. (7.2),

$$\pi(x) = \text{logit}^{-1}[\pi(x)] = \frac{1}{1 + \exp[-(\alpha + \beta x)]} \quad (7.3) \quad \{\text{eq:logit1a}\}$$

This transformation uses the cumulative distribution function of the logistic distribution, $\Lambda(p) = \frac{1}{1 + \exp(-p)}$, giving rise to the term *logistic regression*.²

From Eqn. (7.2) we see that the odds of a success response can be expressed as

$$\text{odds}(Y = 1) \equiv \frac{\pi(x)}{1 - \pi(x)} = \exp(\alpha + \beta x) = e^\alpha (e^\beta)^x, \quad (7.4) \quad \{\text{eq:logit2}\}$$

which is a multiplicative model for the odds. So, under the logistic model,

- β is the change in the log odds associated with a unit increase in x . The odds are multiplied by e^β for each unit increase in x .
- α is log odds at $x = 0$; e^α is the odds of a favorable response at this x -value (which may not have a reasonable interpretation if $X = 0$ is far from the range of the data).

It is easy to explore the relationships among probabilities, odds and log odds using **R** as we show below, using the function `fractions()` in **MASS** to print the odds corresponding to probability p as a fraction.

```
> library(MASS)
> p <- c(.05, .10, .25, .50, .75, .90, .95)
> data.frame(p,
+            odds=as.character(fractions(p/(1-p))),
+            logit=log(p/(1-p)))
```

	p	odds	logit
1	0.05	1/19	-2.9444
2	0.10	1/9	-2.1972
3	0.25	1/3	-1.0986
4	0.50	1	0.0000
5	0.75	3	1.0986
6	0.90	9	2.1972
7	0.95	19	2.9444

Thus, a probability of $\pi = 0.25$ represents an odds of 1 to 3, or 1/3, while a probability of $\pi = 0.75$ represents an odds of 3 to 1, or 3. The logits are symmetric around 0, so `logit(.25) = -logit(.75)`.

Another simple way to interpret the parameter β in the logistic regression model is to consider the relationship between the probability $\pi(x)$ and x . From Eqn. (7.3) it can be shown that the fitted curve (the blue line in Figure 7.2) has slope equal to $\beta\pi(1 - \pi)$. This has a maximum value of $\beta/4$ when $\pi = \frac{1}{2}$, so taking $\beta/4$ gives a quick estimate of the maximum effect of x on the probability scale.

In Figure 7.2 and other plots later in this chapter we try to show the binary responses (as jittered points or a rug plot) to help you appreciate how the fitted logistic curve arises from their distribution across the range a predictor. For didactic purposes this can be seen more readily by plotting the conditional distributions of $x|y = \{0, 1\}$ as a histogram, boxplot or density plot. The function

²Any other cumulative probability transformation serves the purpose of constraining the probabilities to the interval [0, 1]. The cumulative normal transformation $\pi(x) = \Phi(\alpha + \beta x)$ gives the *linear probit regression* model. We don't treat probit models here because: (a) The logistic and probit models give results so similar that it is hard to distinguish them in practice; (b) The logistic model is simpler to interpret as a linear model for the log odds or multiplicative model for the odds.

`logi.hist.plot()` in the `probio` package is a nice implementation of this idea (de la Cruz Rot, 2005). The call below produces Figure 7.3, and it is easy to see how increasing age produces a greater probability of a Better response.

```
> with(Arthritis,
+     logi.hist.plot(Age, Better, type="hist", counts=TRUE,
+                   ylabel="Probability (Better)", xlab="Age",
+                   col.cur="blue", col.hist="lightblue", col.box="lightblue")
+ )
```




Figure 7.3: Plot of the Arthritis treatment data, showing the conditional distributions of the 0/1 observations of the Better response by histograms and boxplots.

{fig:arth-logi-hist}

7.2.1 Fitting a logistic regression model

{sec:logist-fitting}

Logistic regression models are the special case of generalized linear models fit in R using `glm()` for a binary response using `family=binomial`. We first illustrate how simple models can be fit and interpreted.

{ex:arthrit6}

EXAMPLE 7.1: Arthritis treatment

In Chapter 4 we examined the data on treatment for rheumatoid arthritis in relation to two categorical predictors, sex of patient and treatment. In addition, the *Arthritis* data gives the age of each patient in this study, and we focus here on the relationship between Age and the outcome, Improved. This response variable has three categories (none, some, or marked improvement), but for now we consider whether the patient showed any improvement at all, defining the event *Better* to be some or marked improvement.

```
> data("Arthritis", package="vcd")
> Arthritis$Better <- as.numeric(Arthritis$Improved > "None")
```

The logistic regression model is fit using `glm()` as shown below, specifying `family=binomial` for a binary response.

```
> arth.logistic <- glm(Better ~ Age, data=Arthritis, family=binomial)
```

As usual for R modeling functions, the `print()` method for "glm" objects gives brief printed output, while the `summary()` method is more verbose, and includes standard errors and hypothesis tests for the model coefficients. To save some space, it is convenient to use the generic function `coeftest()` from the `lmtest` package. Then, we can use this instead of the more detailed `summary()`:

```
> library(lmtest)
> coeftest(arth.logistic)

z test of coefficients:

              Estimate Std. Error z value Pr(>|z|)
(Intercept)  -2.6421    1.0732   -2.46    0.014 *
Age           0.0492    0.0194    2.54    0.011 *
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

In the output above, the parameter estimates are $\alpha = -2.642$, and $\beta = 0.0492$. So, the estimated odds of a better response are multiplied by $e^\beta = \exp(0.0492) = 1.05$ for each one year increase in age. Equivalently, you can think of this as a 5% increase per year (using $100(e^\beta - 1)$ to convert). Over 10 years, the odds are multiplied by $\exp(10 \times 0.0492) = 1.64$, a 64% increase, a substantial effect in the range for these data. You can do these calculations in R using the `coef()` method for the "glm" object.

```
> exp(coef(arth.logistic))

(Intercept)      Age
  0.071214    1.050482

> exp(10*coef(arth.logistic)[2])

Age
1.6364
```

For comparison with the logistic model, we could fit the linear probability model Eqn. (7.1) using either `lm()` or `glm()` with the default `family=gaussian` argument.

```
> arth.lm <- glm(Better ~ Age, data=Arthritis)
> coef(arth.lm)

(Intercept)      Age
 -0.107170    0.011379
```

The coefficient for age can be interpreted to indicate that the probability of a better response increases by 0.011 for each one year increase in age. You can compare this with the $\beta/4$ rule of thumb, that gives $0.0492/4 = 0.0123$. Even though the linear probability model is inappropriate theoretically, you can see in Figure 7.2 (the black line) that it gives similar predicted probabilities to those of the logistic model between age 25–75, where most of the data points are located.

△

7.2.2 Model tests for simple logistic regression

There are two main types of hypothesis tests one might want to perform for a logistic regression model. We postpone general discussion of this topic until Section 7.3, but introduce the main ideas here using the analysis of the *Arthritis* data.

- The most basic test answers the question “How much better is the fitted model, $\text{logit}(\pi) = \alpha + \beta x$ than the null model $\text{logit}(\pi) = \alpha$ that includes only the regression intercept?” One answer to this question is given by the (Wald) test of the coefficient for age testing the hypothesis $H_0 : \beta = 0$ that appeared in the output from `summary(arth.logistic)` shown above. The more direct test compares the deviance of the fitted model to the deviance of the null model, and can be obtained using the `anova()` function:

```
> anova(arth.logistic, test="Chisq")

Analysis of Deviance Table

Model: binomial, link: logit

Response: Better

Terms added sequentially (first to last)

      Df Deviance Resid. Df Resid. Dev Pr(>Chi)
NULL                83      116
Age    1         7.29    82      109    0.007 **
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

- A second question is “How bad is this model, compared to a model (the *saturated model*) that fits the data perfectly?” This is a test of the size of the residual deviance, that is given by the function `LRstats()` in `vcdExtra`.

```
> LRstats(arth.logistic)

Likelihood summary table:
      AIC BIC LR Chisq Df Pr(>Chisq)
arth.logistic 113 118    109 82    0.024 *
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

The summary of these tests is that linear logistic model Eqn. (7.2) fits significantly better than the null model, but that model also shows significant lack of fit.

7.2.3 Plotting a binary response

{sec:logist-plotting}

It is often difficult to understand how a binary response can give rise to a smooth, continuous relation between the predicted response, usually the probability of an event, and a continuous explanatory variable. Beyond this, plots of the data together with fitted models help you to interpret what these models imply.

We illustrate two approaches below using the *Arthritis* data shown in Figure 7.2, first using R base graphics, and then with the `ggplot2` package that makes such graphs somewhat easier to do.

That plot, which was designed for didactic purposes, has the following features:

- It shows the *data*, that is, the 0/1 observations of the `Better` response in relation to age. To do this effectively and avoid over-plotting, the binary responses are jittered.
- It plots the predicted (fitted) logistic regression relationship on the scale of probability, together with a 95% confidence band.
- It also plots the predicted probabilities from the linear probability model.
- A smoothed, non-parametric regression curve for the binary observations is also added to the plot to give some indication of possible non-linearity in the relationship of `Better` to age.

{ex:arthrit7}

EXAMPLE 7.2: Arthritis treatment – Plotting logistic regression with base graphics

Here we explain how plots similar to Figure 7.2 can be constructed using R base graphics. We describe the steps needed to calculate predicted values and confidence bands and how to add these to a basic plot. These ideas are the basis for the higher-level and more convenient plotting methods illustrated later in this chapter. The steps detailed below give the plot shown in Figure 7.4.




Figure 7.4: A version of plot of the Arthritis treatment data (Figure 7.2) produced with R base graphics, showing logistic, linear regression and lowess fits.

First, we set up the basic plot of the jittered values of `Better` vs. `Age`, setting `xlim` to a larger range than that in the data, only to emphasize where the logistic and linear probability models diverge.

```
> plot(jitter(Better, .1) ~ Age, data=Arthritis,
+       xlim = c(15, 85), pch=16,
+       ylab="Probability (Better)")
```

The fitted logistic curve can be obtained using the `predict()` method for the "glm" object `arth.logistic`. For this example, we wanted to get fitted values for the range of `Age` from 15–85, which is specified in the `newdata` argument.³ The argument `type="response"` gives fitted values of the probabilities. (The default, `type="link"` would give predicted logits.) Standard errors of the fitted values are not calculated by default, so we set `se.fit=TRUE`.

```
> xvalues <- seq(15, 85, 5)
> pred.logistic <- predict(arth.logistic,
+                           newdata=data.frame(Age=xvalues),
+                           type="response", se.fit=TRUE)
```

³Omitting the `newdata` argument would give predicted values using the linear predictors in the data used for the fitted model. Some care needs to be taken if the predictor(s) contain missing values.

When `se.fit=TRUE`, the `predict()` function returns its result in a list, with components `fit` for the fitted values and `se.fit` for the standard errors. From these, we can calculate 95% pointwise prediction intervals using the standard normal approximation.

```
> upper <- pred.logistic$fit + 1.96 * pred.logistic$se.fit
> lower <- pred.logistic$fit - 1.96 * pred.logistic$se.fit
```

We can then plot the confidence band using `polygon()` and the fitted logistic curve using `lines()`. A graphics trick is used here to use a transparent color for the confidence band using `rgb(r, g, b, alpha)`, where `alpha` is the transparency value.

```
> polygon(c(xvalues, rev(xvalues)),
+         c(upper, rev(lower)),
+         col=rgb(0, 0, 1, .2), border=NA)
> lines(xvalues, pred.logistic$fit, lwd=4, col="blue")
```

This method, using `predict()` for calculations and `polygon()` and `lines()` for plotting can be used to display the predicted relationships and confidence bands under other models. Here, we simply used `abline()` to plot the fitted line for the linear probability model `arth.lm` and `lowess()` to calculate a smoothed, non-parametric curve.

```
> abline(arth.lm, lwd=2)
> lines(lowess(Arthritis$Age, Arthritis$Better, f=.9), col="red", lwd=2)
```

△

{ex:arthrit8}

EXAMPLE 7.3: Arthritis treatment – Plotting logistic regression with ggplot2

Model-based plots such as Figure 7.2 are relatively more straight-forward to produce using `ggplot2`. The basic steps here are to:

- set up the plot frame with `ggplot()` using `Age` and `Better` as (x, y) coordinates;
- use `geom_point()` to plot the observations, whose positions are jittered with `position_jitter()`;
- use `stat_smooth()` with `method = "glm"` and `family = binomial` to plot the predicted probability curve and confidence band. By default, `stat_smooth()` calculates and plots 95% confidence bands on the response (probability) scale.

```
> library(ggplot2)
> # basic logistic regression plot
> gg <- ggplot(Arthritis, aes(x=Age, y=Better)) +
+   xlim(5, 95) + theme_bw() +
+   geom_point(position = position_jitter(height = 0.02, width = 0)) +
+   stat_smooth(method = "glm", family = binomial, alpha = 0.1, fill="blue",
+               size=2.5, fullrange=TRUE)
```

Finally, we can add other smoothers to the plot, literally by using `+` to add these to the "ggplot" object.

```
> # add linear model and loess smoothers
> gg <- gg + stat_smooth(method = "lm", se=FALSE,
+                        size=1.2, color="black", fullrange=TRUE)
> gg <- gg + stat_smooth(method = "loess", se=FALSE,
+                        span=0.95, colour="red", size=1.2)
> gg # show the plot
```

△

7.2.4 Grouped binomial data

{sec:logist-grouped}

A related case occurs with grouped data, where rather than binary observations, $y_i \in \{0, 1\}$ in case form, the data is given in what is called *events/trials form* that records the number of successes, y_i that occurred in n_i trials associated with each setting of the explanatory variable(s) x_i .⁴ Case form, with binary observations is the special case where $n_i = 1$.

Data in events/trials form often arises from contingency table data with a binary response. For example in the *UCBAdmissions* data, the response variable Admit with levels "Admitted", "Rejected" could be treated in this way using the number of applicants as the number of trials.

As before, we can consider y_i/n_i to estimate the probability of success, π_i and the distribution of Y to be binomial, $\text{Bin}(\pi_i, n_i)$ at each x_i .

In practical applications, there are two main differences between the cases of ungrouped, case form data and grouped, event/trials form.

- In fitting models using `glm()`, the model formula, `response ~ terms`, can be given using a response consisting of a two-column matrix, whose columns contain the numbers of successes y_i and failures $n_i - y_i$. Alternatively, the response can be given as the proportion of successes, y_i/n_i , but then it is necessary to specify the number of trials as a weight.
- In plotting the fitted model on the scale of probability, you usually have to explicitly plot the fraction of successes, y_i/n_i .

{ex:nasa-temp}

EXAMPLE 7.4: Space shuttle disaster

In Example 1.2 and Example 1.10 we described the background behind the post-mortem examination of the evidence relating to the disastrous launch of the space shuttle *Challenger* on January 28, 1986. Here we consider a simple, but proper analysis of the data available at the time of launch. We also use this example to illustrate some details of the fitting and plotting of grouped binomial data. As well, we describe some of the possibilities for dealing with missing data.

The data set *SpaceShuttle* in *vcd* contains data on the failures of the O-rings in 24 NASA launches preceding the launch of *Challenger*, as given by Dalal *et al.* (1989) and Tufte (1997) also analysed by Lavine (1991).

Each launch used two booster rockets with a total of six O-rings, and the data set records as `nFailures` the number of these that were considered damaged after the rockets were recovered at sea. In one launch (flight # 4), the rocket was lost at sea, so the relevant response variables are missing.

In this example, we focus on the variable `nFailures` as a binomial with $n_i = 6$ trials. The missing data for flight 4 can be handled in several ways in the call to `glm()`

```
> data("SpaceShuttle", package="vcd")
> shuttle.mod <- glm(cbind(nFailures, 6 - nFailures) ~ Temperature,
+                   data = SpaceShuttle, na.action = na.exclude,
+                   family = binomial)
```

Alternatively, we can add an explicit `trials` variable, represent the response as the proportion `nFailures/trials`, and use `weight = trials` to indicate the total number of observations.

```
> SpaceShuttle$trials <- 6
> shuttle.modw <- glm(nFailures/trials ~ Temperature, weight = trials,
+                   data = SpaceShuttle, na.action = na.exclude,
+                   family = binomial)
```

⁴Alternatively, the data may record the number of successes, y_i , and number of failures, $n_i - y_i$.

These two approaches give identical results for all practical purposes:

```
> all.equal(coef(shuttle.mod), coef(shuttle.modw))
[1] TRUE
```

As before, we can test whether temperature significantly improves prediction of failure probability using `anova()`:

```
> # testing, vs. null model
> anova(shuttle.mod, test="Chisq")

Analysis of Deviance Table

Model: binomial, link: logit

Response: cbind(nFailures, 6 - nFailures)

Terms added sequentially (first to last)

              Df Deviance Resid. Df Resid. Dev Pr(>Chi)
NULL                22      24.2
Temperature  1         6.14      21      18.1    0.013 *
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

The code below gives a `ggplot2` version in Figure 7.5 of the plot we showed earlier in Example 1.2 (Figure 1.2). The relevant details here are:

- We specify `y = nFailures / trials` to calculate the failure probabilities.
- Points are jittered in the call to `geom_point()` to prevent overplotting.
- In the call to `geom_smooth()`, we need to use `weight = trials`, just as in the call to `glm()` above.
- `fullrange = TRUE` makes the fitted regression curve and confidence band extend across the entire plot

```
> library(ggplot2)
> ggplot(SpaceShuttle, aes(x = Temperature, y = nFailures / trials)) +
+   xlim(30, 81) + theme_bw() +
+   xlab("Temperature (F)") +
+   ylab("O-Ring Failure Probability") +
+   geom_point(position=position_jitter(width=0, height=0.01),
+             aes(size = 2)) +
+   theme(legend.position="none") +
+   geom_smooth(method = "glm", family = binomial, fill="blue",
+             aes(weight = trials), fullrange = TRUE, alpha=0.2, size=2)
```

△

7.3 Multiple logistic regression models

As is the case in classical regression, generalizing the simple logistic regression to an arbitrary number of explanatory variables is quite straightforward. We let $\mathbf{x}_i = (x_{i1}, x_{i2}, \dots, x_{ip})$ denote the vector of p explanatory variables for case or cluster i . Then the general logistic regression model can be expressed as

$$\begin{aligned} \text{logit}(\pi_i) \equiv \log \frac{\pi_i}{1 - \pi_i} &= \alpha + \mathbf{x}_i^\top \boldsymbol{\beta} \\ &= \alpha + \beta_1 x_{i1} + \beta_2 x_{i2} + \cdots + \beta_p x_{ip} . \end{aligned} \quad (7.5)$$

{sec:logist-mult}




Figure 7.5: Space shuttle data, with fitted logistic regression model

{fig:nasa-temp-ggplot}

Equivalently, we can represent this model in terms of probabilities as the logistic transformation of the *linear predictor*, $\eta_i = \alpha + \mathbf{x}_i^\top \boldsymbol{\beta}$,

$$\begin{aligned} \pi_i = \Lambda(\eta_i) &= \Lambda(\alpha + \mathbf{x}_i^\top \boldsymbol{\beta}) \\ &= \frac{1}{1 + \exp(\alpha + \beta_1 x_{i1} + \beta_2 x_{i2} + \cdots + \beta_p x_{ip})} . \end{aligned} \quad (7.6) \quad \text{{eq:logistm1}}$$

The x s can include any of the following sorts of regressors, as in the general linear model:

- **quantitative** variables (e.g., age, income)
- **polynomial** powers of quantitative variables (e.g., age, age², age³)
- **transformations** of quantitative variables (e.g., log salary)
- factors, represented as **dummy** variables for qualitative predictors (e.g., P_1, P_2, P_3 for four political party affiliations)
- **interaction** terms (e.g., sex \times age, or age \times income)

{ex:arthrit-mult}

EXAMPLE 7.5: Arthritis treatment

We continue with the analysis of the *Arthritis* data, fitting a model containing the main effects of Age, Sex and Treatment, with Better as the response. This model has the form

$$\text{logit}(\pi_i) = \alpha + \beta_1 x_{i1} + \beta_2 x_{i2} + \beta_3 x_{i3}$$

where x_1 is Age and x_2 and x_3 are the factors representing Sex and Treatment, respectively. Using the default (0/1) dummy coding that R uses (“treatment” contrasts against the lowest factor

level),⁵ they are defined as:

$$x_2 = \begin{cases} 0 & \text{if Female} \\ 1 & \text{if Male} \end{cases} \quad x_3 = \begin{cases} 0 & \text{if Placebo} \\ 1 & \text{if Treatment} \end{cases}$$

In this model,

- α doesn't have a sensible interpretation here, but formally it would be the log odds of improvement for a person at age $x_1 = 0$ in the baseline or reference group with $x_2 = 0$ and $x_3 = 0$ —females receiving the placebo. To make the intercept interpretable, we will fit the model centering age near the mean, by using $x_1 - 50$ as the first regressor.
- β_1 is the increment in log odds of improvement for each one-year increase in age.
- β_2 is the increment in log odds for male as compared to female. Therefore, e^{β_2} gives the odds of improvement for males relative to females.
- β_3 is the increment in log odds for being in the treated group. e^{β_3} gives the odds of improvement for the active treatment group relative to placebo.

We fit the model as follows. In `glm()` model formulas, “~” has a special meaning, so we use the identity function, `I(Age-50)` to center age.

```
> arth.logistic2 <- glm(Better ~ I(Age-50) + Sex + Treatment,
+                       data=Arthritis,
+                       family=binomial)
```

The parameters defined here are *incremental effects*. The intercept corresponds to a baseline group (50 year-old females given the placebo); the other parameters are incremental effects for the other groups compared to the baseline group. Thus, when α , β_1 , β_2 and β_3 have been estimated, the fitted logits and predicted odds at `Age==50` are:

Sex	Treatment	Logit	Odds Improved
Female	Placebo	α	e^α
Female	Treated	$\alpha + \beta_2$	$e^{\alpha+\beta_2}$
Male	Placebo	$\alpha + \beta_1$	$e^{\alpha+\beta_1}$
Male	Treated	$\alpha + \beta_1 + \beta_2$	$e^{\alpha+\beta_1+\beta_2}$

We first focus on the interpretation of the coefficients estimated for this model shown below.

```
> coefTest(arth.logistic2)

z test of coefficients:

              Estimate Std. Error z value Pr(>|z|)
(Intercept)   -0.5781    0.3674   -1.57    0.116
I(Age - 50)     0.0487    0.0207    2.36    0.018 *
SexMale        -1.4878    0.5948   -2.50    0.012 *
TreatmentTreated 1.7598    0.5365    3.28    0.001 **
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

⁵For factor variables with the default treatment contrasts, you can change the reference level using `relevel()`. In this example, you could make male the baseline category using `Arthritis$Sex <- relevel(Arthritis$Sex, ref = "Male")`.

To interpret these in terms of odds ratios and also find confidence intervals, just use `exp()` and `confint()`.

```
> exp(cbind(OddsRatio=coef(arth.logistic2),
+           confint(arth.logistic2)))
```

	OddsRatio	2.5 %	97.5 %
(Intercept)	0.5609	0.26475	1.1323
I(Age - 50)	1.0500	1.01000	1.0963
SexMale	0.2259	0.06524	0.6891
TreatmentTreated	5.8113	2.11870	17.7266

Here,

- $\alpha = -0.578$: At age 50, females given the placebo have an odds of improvement of $\exp(-0.578) = 0.56$.
- $\beta_1 = 0.0487$: Each year of age multiplies the odds of improvement by $\exp(0.0487) = 1.05$, or a 5% increase.
- $\beta_2 = -1.49$: Males are only $\exp(-1.49) = 0.26$ times as likely to show improvement relative to females. (Or, females are $\exp(1.49) = 4.437$ times more likely than males to improve.)
- $\beta_3 = 1.76$: People given the active treatment are $\exp(1.76) = 5.8$ times more likely to show improvement compared to those given the placebo.

As you can see, the interpretation of coefficients in multiple logistic models is straightforward, though a bit cumbersome. This becomes more difficult in larger models, particularly when there are interactions. In these cases, you can understand (and explain) a fitted model more easily through plots of predicted values, either on the scale of response probability or on the logit scale of the linear predictor. We describe these methods in Section 7.3.1–Section 7.3.3 below.

△

7.3.1 Conditional plots

c:logist-condplots}

The simplest kind of plots display the data together with a representation of the fitted relationship (predicted values, confidence bands) separately for subsets of the data defined by one or more of the predictors. Such plots can show the predicted values for the response variable on the ordinate against one chosen predictor on the abscissa, and can use multiple curves and multiple panels to represent other predictors.

However, these plots are *conditional plots*, meaning that the data shown in each panel and used in each fitted curve are limited to the subset of the observations defined by the curve and panel variables. As well, predictors that are not shown in a given plot are effectively ignored (or marginalized), as was the case in Figure 7.2 that showed only the effect of age in the *Arthritis* data.

{ex:arth-cond}

EXAMPLE 7.6: Arthritis treatment – conditional plots

For the *Arthritis* data, a basic conditional plot of Better vs. Age, showing the observations as jittered points (with `geom_point()`) and the fitted logistic curves (with `stat_smooth()` using `method="glm"`) can be produced with `ggplot2` as shown below, giving Figure 7.6.

```
> library(ggplot2)
> gg <- ggplot(Arthritis, aes(Age, Better, color=Treatment)) +
+   xlim(5, 95) + theme_bw() +
+   geom_point(position = position_jitter(height = 0.02, width = 0)) +
+   stat_smooth(method = "glm", family = binomial, alpha = 0.2,
+             aes(fill=Treatment), size=2.5, fullrange=TRUE)
> gg # show the plot
```




Figure 7.6: Conditional plot of Arthritis data showing separate points and fitted curves stratified by Treatment. A separate fitted curve is shown for the two treatment conditions, ignoring Sex.

{fig:arth-cond1}

In this call to `ggplot()`, specifying `color=Treatment` gives different point and line colors, but also automatically stratifies the fitted curves using the levels of this variable.

With such a plot, it is easy to add further stratifying variables in the data using *facets* to produce separate panels (functions `facet_wrap()` or `facet_grid()`, with different options to control the details). The following line further stratifies by Sex, producing Figure 7.7.

```
> gg + facet_wrap(~ Sex)
```




Figure 7.7: Conditional plot of Arthritis data, stratified by Treatment and Sex. The unusual patterns in the panel for Males signals a problem with this data.

{fig:arth-cond2}

However, you can see from this plot how this method breaks down when the sample size is small in some of the groups defined by the stratifying factors. The panel for males shows a paradoxical negative relation with age for the treated group and a step function for the placebo group. The explanation for this is shown in the two-way frequency table of the sex and treatment combinations:

```
> addmargins(xtabs(~Sex + Treatment, data=Arthritis), 2)
Treatment
```


Sex	Placebo	Treated	Sum
Female	32	27	59
Male	11	14	25

Less than 1/3 of the sample were males, and of these only 11 were in the placebo group. `glm()` cannot estimate the fitted relationship against Age here— the slope coefficient is infinite, and the fitted probabilities are all either 0 or 1.⁶

△

7.3.2 Full-model plots

```
c:logist-fullplots}
```

For a model with two or more explanatory variables, *full-model plots* display the fitted response surface for all predictors together, rather than stratified by conditioning variables. Such plots show the predicted values for the response variable on the ordinate against one chosen predictor on the abscissa, and can use multiple curves and multiple panels to represent other predictors.

A simple R trick⁷ makes this method far easier and more general than the naive plotting method used in Example 7.2. The trick is simply to combine the columns in the original data frame with the result of the `predict()` method for the fitted model and plot the calculated fit value, together with confidence bands (if you use `se.fit=TRUE`).

```
{ex:arth-full}
```

EXAMPLE 7.7: Arthritis treatment – full-model plots

```
> arth.fit2 <- cbind(Arthritis,
+                   predict(arth.logistic2, se.fit = TRUE))
```

The fitted values here are on the logit scale, which means that it takes one more trick to show the data points on the plot. We simply define a new variable, `obs` with convenient logit values corresponding the Better values of 0 and 1.

```
> arth.fit2$obs <- c(-4, 4)[1+arth.fit2$Better]
```

We can then plot the fitted logit against Age using `x=Age`, `y=fit` from the data frame containing the fitted values. The call to `ggplot()` below produces Figure 7.8. Here, we used `color=Treatment` to produce separate points, lines and confidence bands colored by Treatment. Confidence bands in the plot are constructed using `geom_ribbon()`.

```
> ggplot(arth.fit2, aes(x=Age, y=fit, color=Treatment)) +
+   geom_line(size = 2) + theme_bw() +
+   geom_ribbon(aes(ymin = fit - 1.96 * se.fit,
+                 ymax = fit + 1.96 * se.fit,
+                 fill = Treatment), alpha = 0.2,
+             color = "transparent") +
+   labs(x = "Age", y = "Log odds (Better)") +
+   geom_point(aes(y=obs),
+             position=position_jitter(height=0.25, width=0)) +
+   facet_wrap(~ Sex)
```

This plot method has several nice features:

- Plotting on the logit scale shows the additive, linear effects of all predictors.
- It provides a visual representation of the information contained in the table of coefficients. Note, however, that the choice to display Treatment within each panel makes it easier to judge the size of this effect, compared to the effect of Sex which must be judged across the panels.

⁶This is called *complete separation*, and occurs whenever the responses have no overlap on the predictor variable(s) used in fitting the logistic regression model.

⁷Thanks to Dennis Murphy for suggesting this method.




Figure 7.8: Full-model plot of Arthritis data, showing fitted logits by Treatment and Sex.

- It shows the data as points, and the fitted lines and confidence bands are restricted to the range of the data in each. You can easily see the reason for the unusual pattern in the conditional plot for Males shown in Figure 7.7.
- It generalizes directly to any fitted model, because the same plotting code can be used once the model predicted values have been calculated.
- Additional predictors, either factors or quantitative variables can easily be accommodated by including them in the `facet_wrap()` call. For example, if the patients were also categorized by education and this had been included in the model, `facet_wrap(~ Sex + Education)` would produce separate panels for the combinations of these two variables.

While plots on the logit scale have a simpler form, many people find it easier to think about such relationships in terms of probabilities, as we have done in earlier plots in this chapter. You can do the same for full-model plots with a simple extension of this method. All you need to do is to transform the fit and end points of the confidence band back to the scale of probabilities. The function `plogis()` does this for the logistic distribution.

```
> arth.fit2 <- within(arth.fit2, {
+   prob <- plogis(fit)
+   lower <- plogis(fit - 1.96 * se.fit)
+   upper <- plogis(fit + 1.96 * se.fit)
+ })
```

The plot step is then similar to what we used above (but with `prob`, `lower` and `upper`), producing Figure 7.9.

```
> ggplot(arth.fit2, aes(x=Age, y=prob, color=Treatment)) +
+   geom_line(size = 2) + theme_bw() +
+   geom_ribbon(aes(ymin = lower,
+                 ymax = upper,
+                 fill = Treatment), alpha = 0.2,
+             color = "transparent") +
+   labs(x = "Age", y = "Probability (Better)") +
+   geom_point(aes(y=Better),
+             position=position_jitter(height=0.02, width=0)) +
+   facet_wrap(~ Sex)
```




Figure 7.9: Full-model plot of Arthritis data, showing fitted probabilities by Treatment and Sex.

{fig:arth-full2}

7.3.3 Effect plots

{sec:logist-effplots}

For more than two variables, full-model plots of the fitted response surface can be cumbersome, particularly when the model contains interactions or when the main substantive interest is focused on a given main effect or interaction, controlling for all other explanatory variables. The method of *effect displays* (tables and graphs), developed by John Fox (1987, 2003) and implemented in the *effects* package, is a useful solution to these problems.

The idea of effect plots is quite simple but very general:⁸ consider a particular subset of predictors (*focal predictors*) we wish to visualize in a given linear model or generalized linear model. The essence is to calculate fitted values (and standard errors) for the model terms involving these variables and all low-order relatives (e.g., main effects that are marginal to an interaction), as these variables are allowed to vary over their range. All other variables are “controlled” by being fixed at typical values. For example a quantitative covariate could be fixed at its mean or median; a factor could be fixed at equal proportions of its levels or its proportions in the data. The result, when plotted, shows all effects of the focal predictors and their low-order relatives, but with all other variables not included controlled or adjusted for.

More formally, assume we have fit a model with a linear predictor $\eta_i = \alpha + \mathbf{x}_i^T \boldsymbol{\beta}$ (on the logit scale, for logistic regression). Letting $\beta_0 = \alpha$ and $\mathbf{x}_0 = \mathbf{1}$, we can rewrite this in matrix form as $\boldsymbol{\eta} = \mathbf{X}\boldsymbol{\beta}$ where \mathbf{X} is the model matrix constructed by the modeling function, such as `glm()`. Fitting the model gives the estimated coefficients \mathbf{b} and its estimated covariance matrix $\widehat{\mathbf{V}}(\mathbf{b})$.

The `Effect()` function constructs an analogous *score model matrix*, \mathbf{X}^* , where the focal variables have been varied over their range, and all other variables represented as constant, typical values. Using this as input (the `newdata` argument) to the `predict()` function then gives the fitted values $\boldsymbol{\eta}^* = \mathbf{X}^* \mathbf{b}$. Standard errors used for confidence intervals are calculated by `predict()` (when `se.fit=TRUE`) as the square roots of $\text{diag}(\mathbf{X}^* \widehat{\mathbf{V}}(\mathbf{b}) \mathbf{X}^{*T})$. Note that these ideas work not only for `glm()` models, but potentially for any modeling function that has a `predict()` and `vcov()` method.⁹

These results are calculated on the scale of the linear predictor $\boldsymbol{\eta}$ (logits, for logistic regression)

⁸Less general expression of these ideas include the use of *adjusted means* in analysis of covariance, and *least squares means* or *population marginal means* (Searle *et al.*, 1980) in analysis of variance; for example, see the *lsmeans* package for classical linear models.

⁹For example, the *effects* package presently provides methods for models fit by `lm()` (including multivariate linear response models), `glm()`, `gls()`, `multinomial(multinom())` in the *nnet* package) and proportional odds models (`polr()` in *MASS*), polytomous latent class models (*poLCA* package), as well as a variety of multi-level and mixed-effects linear models fit with `lmer()` from the *lme4* package, or with `lme()` from the *nlme* package.

when the `type` argument to `predict()` is `type="link"` or on the response scale (probabilities, here) when `type="response"`. The latter makes use of the inverse transformation, Eqn. (7.6).

There are two main calculation functions in the `effects` package:

- `Effect()` takes a character vector of the names of a subset of focal predictors and constructs the score matrix X^* by varying these over their ranges, while holding all other predictors constant at “typical” values. There are many options that control these calculations. For example, `xlevels` can be used to specify the values of the focal predictors; `typical` or `given.values` respectively can be used to specify either a function (mean, median) or a list of specific typical values used for the variables that are controlled. The result is an object of class “`eff`”, for which there are `print()`, `summary()` and (most importantly) `plot()` methods. See `help(Effect)` for a complete description.
- `allEffects()` takes a model object, and calculates the effects for each high-order term in the model (including their low-order) relatives. Similar optional arguments control the details of the computation. The result is an object of class “`efflist`”.

In addition, the plotting methods for “`eff`” and “`efflist`” objects offer numerous options to control the plot details, only a few of which are used in the examples below. For logistic regression models, they also solve the problem of the trade-off between plots on the logit scale, that have a simple representation in terms of additive effects, and plots on the probability scale that are usually simpler to understand. By default, the fitted model effects are plotted on the logit scale, but the response y axis is labeled with the corresponding probability values.

{ex:arthrit-eff}

EXAMPLE 7.8: Arthritis treatment

Here we illustrate the use of the `effects` package with the simple main effects model which was fit in Example 7.5. `allEffects()` is used to calculate the predicted probabilities of the `Better` response for `Age` and the two factors, `Sex` and `Treatment`.

```
> library(effects)
> arth.eff2 <- allEffects(arth.logistic2)
> names(arth.eff2)

[1] "I(Age-50)" "Sex"      "Treatment"
```

The result, `arth.eff2` is a list containing the fitted values (response probabilities, by default) for each of the model terms. No `xlevels` argument was specified, so by default the function calculated the effects for `Age` at a reasonable selection of equally-spaced values:

```
> arth.eff2[[1]]

Age effect
Age
      30      40      50      60      70
0.24289 0.34311 0.45959 0.58066 0.69274
```

The default plot method for the “`efflist`” object produces one plot for each high-order term, which are just the main effect in this model. The call below produces Figure 7.10.

```
> plot(arth.eff2, rows=1, cols=3)
```

You can also produce full-model plots quite easily by using all predictors in the model in a call to `Effect()`.




Figure 7.10: Plot of all effects in the main effects model for the Arthritis data

{fig:arth-effplot1}

```
> arth.full <- Effect(c("Age", "Treatment", "Sex"), arth.logistic2)
```

Then plotting the result, with some options, gives the plot shown in Figure 7.11.

```
> plot(arth.full, multiline=TRUE, ci.style="bands",
+       colors = c("red", "blue"), lwd=3,
+       ticks=list(at=c(.05, .1, .25, .5, .75, .9, .95)),
+       key.args=list(x=.52, y=.92), grid=TRUE)
```




Figure 7.11: Full-model plot of the effects of all predictors in the main effects model for the Arthritis data, plotted on the logit scale.

{fig:arth-effplot2}

Alternatively, we can plot these results directly on the scale of probabilities, as shown in Figure 7.12.

```
> plot(arth.full, multiline=TRUE, ci.style="bands", rescale.axis=FALSE,
+       colors = c("red", "blue"), lwd=3,
+       key.args=list(x=.52, y=.92), grid=TRUE)
```

△




Figure 7.12: Full-model plot of the effects of all predictors in the main effects model for the Arthritis data, plotted on the probability scale.

7.4 Case studies

The examples below take up some issues of data analysis, model building and visualization in the context of multiple logistic regression models. We focus on the combination of exploratory plots to see the data, modeling steps and graphs to interpret a given model.

EXAMPLE 7.9: Donner Party

In Chapter 1, Example 1.3, we described the background behind the sad story of the Donner Party, perhaps the most famous tragedy in the history of the westward settlement in the United States. In brief, the party was stranded on the eastern side of the Sierra Nevada mountains by heavy snow in late October, 1846, and by the time the last survivor was rescued in April, 1847, nearly half of the members had died from famine and exposure to extreme cold. Figure 1.3 showed that survival decreased strongly with age.

Here we consider a more detailed analysis of these data, which are contained in the data set *Donner* in *vcdExtra*. This data set lists 90 people in the Donner Party by name, together with age, sex, survived (0/1) and the date of death for those who died.¹⁰

```
> data("Donner", package="vcdExtra") # load the data
> library(car)                       # for some() and Anova()
> some(Donner, 8)
```

	family	age	sex	survived	death
Breen, Peter	Breen	3	Male	1	<NA>
Donner, Jacob	Donner	65	Male	0	1846-12-21
Foster, Jeremiah	MurFosPik	1	Male	0	1847-03-13
Graves, Nancy	Graves	9	Female	1	<NA>
McCutchen, Harriet	McCutchen	1	Female	0	1847-02-02
Reed, James	Reed	46	Male	1	<NA>
Reinhardt, Joseph	Other	30	Male	0	1846-12-21
Wolfinger, Doris	FosdWolf	20	Female	1	<NA>

The main purpose of this example is to try to understand, through graphs and models, how

¹⁰Most historical sources count the number in the Donner Party at 87 or 89. An exact accounting of the members of the Donner Party is difficult, because: (a) several people joined the party in mid-route, at Fort Bridger and in the Wasatch Mountains; (b) several rode ahead to search for supplies and one (Charles Stanton) brought two more with him (Luis and Salvador); (c) five people died before reaching the Sierra Nevada mountains. It incorporates updated information from Kristin Johnson's listing, <http://user.xmission.com/~octa/DonnerParty/Roster.htm>.




Figure 7.13: Spineplot of survival in the Donner Party by family.

```
> library(gpairs)
> library(vcd)
> gpairs(Donner[,c(4,2,3,1)],
+   diag.pars=list(fontsize=20, hist.color="gray"),
+   mosaic.pars=list(gp=shading_Friendly), outer.rot=c(45,45)
+   )
```




Figure 7.14: Generalized pairs plot for the Donner data

{fig:donner1-gpairs}

From an exploratory perspective, we now proceed to examine the relationship of survival to

age and sex, beginning with the kind of conditional plots we illustrated earlier (in Example 7.6). Figure 7.15 shows a plot of `survived`, converted back to a 0/1 variable as required by `ggplot()`, together with the binary responses as points and the fitted logistic regressions separately for males and females.

```
> ggplot(Donner, aes(age, as.numeric(survived=="yes"), color = sex)) +
+   theme_bw() + ylab("Survived") +
+   geom_point(position = position_jitter(height = 0.02, width = 0)) +
+   stat_smooth(method = "glm", family = binomial, formula = y ~ x,
+               alpha = 0.2, size=2, aes(fill = sex))
```




Figure 7.15: Conditional plot of the Donner data, showing the relationship of survival to age and sex. The smoothed curves and confidence bands show the result of fitting separate linear logistic regressions on age for males and females.

{fig:donner1-cond1}

It is easy to see that survival among women was greater than for men, perhaps narrowing the gap among the older people, but the data gets thin towards the upper range of age.

The curves plotted in Figure 7.15 assume a linear relationship between the log odds of survival and age (expressed as `formula = y ~ x` in the call to `stat_smooth()`). One simple way to check whether the relationship between survival and age is non-linear is to re-do this plot, but now allow a quadratic relationship with age, using `formula = y ~ poly(x, 2)`. The result is shown in the left panel of Figure 7.16.

```
> gg <- ggplot(Donner, aes(age, as.numeric(survived=="yes"), color = sex)) +
+   theme_bw() + ylab("Survived") +
+   geom_point(position = position_jitter(height = 0.02, width = 0))
>
> gg + stat_smooth(method = "glm", family = binomial, formula = y ~ poly(x, 2),
+                 alpha = 0.2, size=2, aes(fill = sex))
>
> gg + stat_smooth(method = "loess", span=0.9, alpha = 0.2, size=2,
+                 aes(fill = sex)) + coord_cartesian(ylim=c(-.05, 1.05))
```

This plot is quite surprising. It suggests quite different regimes relating to survival for men and women. Among men, survival probability decreases steadily with age, at least after age 20. For women, those in the age range 10–35 were very likely to have lived, while those over 40 were almost all predicted to perish.




Figure 7.16: Conditionals plot of the Donner data, showing the relationship of survival to age and sex. Left: The smoothed curves and confidence bands show the result of fitting separate quadratic logistic regressions on age for males and females. Right: Separate loess smooths are fit to the data for males and females

{fig:donner1-cond3}

Another simple technique is to fit a non-parametric loess smooth, as shown in the right panel of Figure 7.16.¹¹ The curve for females is similar to that of the quadratic fit in the left panel, but the curve for males suggests that survival also has a peak around the teenage years. One lesson to be drawn from these graphs is that a linear logistic regression, such as shown in Figure 7.16 may tell only part of the story, and, for a binary response it is not easy to discern whether the true relationship is linear. If it really is, all these graphs would look much more similar. As well, we usually obtain a more realistic smoothing of the data using full-model plots or effect plots.

The suggestions from these exploratory graphs can be used to define and test some models for survival in the Donner Party. The substantive questions of interest are:

- Is relationship the same for men and women? This is, is it necessary to allow for an interaction of age with sex, or separate fitted curves for men and women?
- Is the relationship between survival and age well-represented in a linear logistic regression model?

The first question is the easiest to deal with: we can simply fit a model allowing an interaction of age (or some function of age) and sex,

```
survived ~ age * sex
survived ~ f(age) * sex
```

and compare the goodness of fit with the analogous additive, main-effects models.

From a modeling perspective, there is a wide variety of approaches for testing for non-linear relationships. We only scratch the surface here, and only for a single quantitative predictor, x , such as age in this example. One simple approach, illustrated in Figure 7.16 is to allow a quadratic (or higher-power, e.g., cubic) function to describe the relationship between the log odds and x ,

$$\text{logit}(\pi_i) = \alpha + \beta_1 x_i + \beta_2 x_i^2$$

¹¹A technical problem with the use of the loess smoother for binary data is that it can produce fitted values outside the $[0, 1]$ interval, as happens in the right panel of this figure. Kernel smoothers, such as the `KernSmooth` package avoid this problem, but are not available through `ggplot2`.

$$\text{logit}(\pi_i) = \alpha + \beta_1 x_i + \beta_2 x_i^2 + \beta_3 x_i^3 \dots$$

In R, these model terms can be fit using `poly(x, 2)`, `poly(x, 3)` ..., which generate orthogonal polynomials for the powers of x . A simple way to test for non-linearity is a likelihood ratio test comparing the more complex model to the linear one. This method is often sufficient for a hypothesis test, and, if the relationship truly is linear, the fitted logits and probabilities will not differ greatly from what they would be under a linear model. A difficulty with this approach is that polynomial models are often unrealistic, particularly for data that approach an asymptote.

Another simple approach is to use a **regression spline**, that fits the relationship with x in terms of a set of piecewise polynomials, usually cubic, joined at a collection of points, called *knots* so that the overall fitted relationship is smooth and continuous. See Fox (2008, §17.2) for a cogent, brief description of these methods.

One particularly convenient method is a **natural spline**, implemented in the `splines` package in the `ns()` function. This method constrains the fitted cubic spline to be linear at lower and upper limits of x , and, for k knots, fits $df = k + 1$ parameters not counting the intercept. The k knots can be conveniently chosen as k cutpoints in the percentiles of the distribution of x . For example, with $k = 1$, the knot would be placed at the median, or 50th percentile; with $k = 3$, the knots would be placed at the quartiles of the distribution of x ; $k = 0$ corresponds to no knots, i.e., a simple linear regression.

In the `ns()` function, you can specify the locations of knots or the number of knots with the `knots` argument, but it is conceptually simpler to specify the number of degrees of freedom used in the spline fit. Thus, `ns(x, 2)` and `poly(x, 2)` both specify a term in x of the same complexity, the former a natural spline with $k = 1$ knot and the later a quadratic function in x .

We illustrate these ideas in the remainder of this example, fitting a 2×2 collection of models to the *Donner* data corresponding to: (a) whether or not age and sex effects are additive; (b) whether the effect is linear on the logit scale or non-linear (quadratic, here). A brief summary of each model is given using the `Anova()` in the `car` package, providing Type II tests of each effect. As usual, `summary()` would give more detailed output, including tests for individual coefficients. First, we fit the linear models, without and with an interaction term:

```
> donner.mod1 <- glm(survived ~ age + sex,
+                     data=Donner, family=binomial)
> Anova(donner.mod1)

Analysis of Deviance Table (Type II tests)

Response: survived
      LR Chisq Df Pr(>Chisq)
age      5.52  1    0.0188 *
sex      6.73  1    0.0095 **
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

> donner.mod2 <- glm(survived ~ age * sex,
+                     data=Donner, family=binomial)
> Anova(donner.mod2)

Analysis of Deviance Table (Type II tests)

Response: survived
      LR Chisq Df Pr(>Chisq)
age      5.52  1    0.0188 *
sex      6.73  1    0.0095 **
age:sex    0.40  1    0.5269
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

The main effects of age and sex are both significant here, but the interaction term, `age:sex` is not in model `donner.mod2`. Note that the terms tested by `Anova()` in `donner.mod1` are a redundant subset of those in `donner.mod2`.

Next, we fit non-linear models, representing the linear and non-linear trends in age by `poly(age, 2)`.¹² The `Anova()` results for terms in both models are contained in the output from `Anova(donner.mod4)`.

```
> donner.mod3 <- glm(survived ~ poly(age,2) + sex,
+                     data=Donner, family=binomial)
> donner.mod4 <- glm(survived ~ poly(age,2) * sex,
+                     data=Donner, family=binomial)
> Anova(donner.mod4)
```

Analysis of Deviance Table (Type II tests)

Response: survived

	LR	Chisq	Df	Pr(>Chisq)
poly(age, 2)	9.91	2	0.0070	**
sex	8.09	1	0.0044	**
poly(age, 2):sex	8.93	2	0.0115	*

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Now, in model `donner.mod4`, the interaction term `poly(age, 2):sex` is significant, indicating that the fitted quadratics for males and females differ in “shape,” meaning either their linear (slope) or quadratic (curvature) components.

These four models address the questions posed earlier. A compact summary of these models, giving the likelihood ratio tests of goodness of fit, together with AIC and BIC statistics are shown below, using the `LRstats()` method in `vcdExtra` for a list of “glm” models. **TODO: LRstats gives the wrong pvalues**

```
> library(vcdExtra)
> LRstats(donner.mod1, donner.mod2, donner.mod3, donner.mod4)
```

Likelihood summary table:

	AIC	BIC	LR	Chisq	Df	Pr(>Chisq)
donner.mod1	117	125	111.1	87	0.042	*
donner.mod2	119	129	110.7	86	0.038	*
donner.mod3	115	125	106.7	86	0.064	.
donner.mod4	110	125	97.8	84	0.144	

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

By AIC and BIC, `donner.mod4` is best, and it is also the only model with a non-significant LR χ^2 (residual deviance). Because these models comprise a 2×2 set of hypotheses, it is easier to compare models by extracting the LR statistics and arranging these in a table, together with the their row and column differences. The entries in the table below are calculated as follows.

```
> mods <- list(donner.mod1, donner.mod2, donner.mod3, donner.mod4)
> LR <- sapply(mods, function(x) x$residuals)
> LR <- matrix(LR, 2, 2)
> rownames(LR) <- c("additive", "non-add")
> colnames(LR) <- c("linear", "non-lin")
> LR <- cbind(LR, diff= LR[,1]-LR[,2])
> LR <- rbind(LR, diff= c(LR[1,1:2]-LR[2,1:2], NA))
```

¹²Alternatively, we could use the term `ns(age, 2)` or higher-degree polynomials or natural splines with more knots, but we don't do this here.

	linear	non-linear	$\Delta\chi^2$	p-value
additive	111.128	106.731	4.396	0.036
non-additive	110.727	97.799	12.928	0.000
$\Delta\chi^2$	0.400	8.932		
p-value	0.527	0.003		

Thus, the answer to our questions seems to be that: (a) there is evidence that the relationship of survival to age differs for men and women in the Donner Party; (b) these relationships are not well-described by a linear logistic regression.

For simplicity, we used a quadratic effect, `poly(age, 2)`, to test for non-linearity here. An alternative test of the same complexity could use a regression spline, `ns(age, 2)`, also with 2 degrees of freedom for the main effect and interaction, or allow more knots. To illustrate, we fit two natural spline models with 2 and 4 df, and compare these with the quadratic model (`donner.mod4`), all of which include the interaction of age and sex.

```
> library(splines)
> donner.mod5 <- glm(survived ~ ns(age, 2) * sex, data=Donner,
+                   family=binomial)
> Anova(donner.mod5)

Analysis of Deviance Table (Type II tests)

Response: survived
              LR Chisq Df Pr(>Chisq)
ns(age, 2)      9.28  2    0.0097 **
sex             7.98  1    0.0047 **
ns(age, 2):sex   8.71  2    0.0129 *
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

> donner.mod6 <- glm(survived ~ ns(age, 4) * sex, data=Donner,
+                   family=binomial)
> Anova(donner.mod6)

Analysis of Deviance Table (Type II tests)

Response: survived
              LR Chisq Df Pr(>Chisq)
ns(age, 4)     22.05  4    0.0002 ***
sex            10.49  1    0.0012 **
ns(age, 4):sex  8.54  4    0.0737 .
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

> LRstats(donner.mod4, donner.mod5, donner.mod6)

Likelihood summary table:
              AIC BIC LR Chisq Df Pr(>Chisq)
donner.mod4  110 125   97.8 84    0.14
donner.mod5  111 126   98.7 84    0.13
donner.mod6  106 131   86.1 80    0.30
```

With four more parameters, `donner.mod6` fits better and has a smaller AIC.

We conclude this example with an effect plot for the spline model `donner.mod6` shown in Figure 7.17. The complexity of the fitted relationships for men and women is intermediate between the two conditional plots shown in Figure 7.16. (However, note that the fitted effects are plotted on the logit scale in Figure 7.17 and labeled with the corresponding probabilities, whereas the conditional plots are plotted directly on the probability scale.)

```

> library(effects)
> donner.eff6 <- allEffects(donner.mod6, xlevels=list(age=seq(0, 50, 5)))
> plot(donner.eff6, ticks=list(at=c(0.001, 0.01, 0.05, 0.1, 0.25,
+                                0.5, 0.75, 0.9, 0.95, 0.99, 0.999)))

```



Figure 7.17: Effect plot for the Donner data

{fig:donner-effect}

This plot confirms that for women in the Donner Party, survival was greatest for those aged 10-30. Survival among men was overall much less and there is a hint of greater survival for men aged 10-15.

Of course, this statistical analysis does not provide explanations for these effects, and it ignores the personal details of the Donner Party members and the individual causes and circumstances of death, which are generally well-documented in the historical record (Johnson, 1996). See <http://user.xmission.com/~octa/DonnerParty/> for a comprehensive collection of historical sources.

Grayson (1990) attributes the greater survival of women of intermediate age to demographic arguments that women are overall better able to withstand conditions of famine and extreme cold, and high age-specific mortality rates among the youngest and oldest members of human societies. He also concludes (without much analysis) that members with larger social and kinship networks would be more likely to survive. \triangle

{ex:arrests}

EXAMPLE 7.10: Racial profiling: Arrests for marijuana possession

In the summer of 2002, the *Toronto Star* newspaper launched an investigation on the topic of possible racial profiling by the Toronto police service. Through freedom of information requests, they obtained a data base of over 600,000 arrest records on all potential charges in the period from 1996–2002, the largest data bases on crime arrests and disposition ever assembled in Canada. An initial presentation of this study was given in Example 1.4.

In order to examine the issue of racial profiling (different treatment as a function of race) they excluded all charges such as assault, robbery, speeding and driving under the influence, where the police have no discretion regarding the laying of a charge. They focused instead on a subset of arrests, where the police had various options.

Among these, for people arrested for a single charge of simple possession of a small amount of marijuana, police have the option of releasing the arrestee, with a summons (“Form 9”) to appear in

court (similar to a parking ticket), or else the person could be given harsher treatment—brought to a police station or held in jail for a bail hearing (“Show cause”). The main question for the *Toronto Star* was whether the subject’s skin color had any influence on the likelihood that the person would be released with a summons.¹³

Their results, published in a week-long series of articles in December 2002, concluded that there was strong evidence that black and white subjects were treated differently. For example, the analysis showed that blacks were 1.5 times more likely than whites to be given harsher treatment than release with a summons; if the subject was taken to the police station, a black was 1.6 times more likely to be held in jail for a bail hearing. An important part of the analysis and the public debate that ensued was to show that other variables that might account for these differences had been controlled or adjusted for.¹⁴

The data set *Arrests* in the *effects* package gives a simplified version of the *Star* database, containing records for 5226 cases of arrest on the charge of simple possession of marijuana analyzed by the newspaper. The response variable here is *released* (Yes/No) and the main predictor of interest is skin color of the person arrested, *colour* (Black/White).¹⁵ A random subset of the data set is shown below.

```
> library(effects)
> data("Arrests", package="effects")
> Arrests[sample(nrow(Arrests), 6),]
```

	released	colour	year	age	sex	employed	citizen	checks
3768	Yes	Black	2000	23	Male	No	Yes	4
4576	Yes	Black	2001	17	Male	Yes	Yes	0
3976	No	White	2002	20	Male	No	Yes	3
4629	Yes	White	2000	18	Male	Yes	Yes	1
2384	No	Black	2000	19	Male	Yes	Yes	3
869	Yes	White	2001	15	Male	Yes	Yes	1

Other available predictors, to be used as control variables included the year of the arrest, age and sex of the person, and binary indicators of whether the person was employed and a citizen of Canada. In addition, when someone is stopped by police, his/her name is checked in six police data bases that record previous arrests, convictions, whether on parole, etc. The variable *checks* records the number, 0–6, in which the person’s name appeared.

A variety of logistic models were fit to these data including all possible main effects and some two-way interactions. To allow for possible non-linear effects of year, this variable was treated as a factor rather than as a (linear) numeric variable, but the effects of age and checks were reasonably linear on the logit scale. A reasonable model included the interactions of colour with both year and age, as fit below:

```
> Arrests$year <- as.factor(Arrests$year)
> arrests.mod <- glm(released ~ employed + citizen + checks
+                   + colour*year + colour*age,
+                   family=binomial, data=Arrests)
```

For such models, significance tests for the model terms are best carried out using the *Anova()* function in the *car* package that uses Type II tests ...

¹³Another discretionary charge they investigated was police stops for non-moving violations under the Ontario *Highway Traffic Act*, such as being pulled over for a faulty muffler or having an expired license plate renewal sticker. A disproportionate rate of charges against blacks is sometimes referred to as “driving while black” (DWB). This investigation found that the number of blacks so charged, but particularly young black males, far out-weighted their representation in the population.

¹⁴The Toronto Police Service launched a class-action libel law suit against the *Toronto Star* and the first author of this book, who served as their statistical consultant, claiming damages of \$5,000 for every serving police officer in the city, a total of over 20 million dollars. The suit was thrown out of court, and the Toronto police took efforts to enhance training programs to combat the perception of racial profiling.

¹⁵The original data set also contained the categories Brown and Other, but these appeared with small frequencies.

```
> library(car)
> Anova(arrests.mod)

Analysis of Deviance Table (Type II tests)

Response: released
      LR Chisq Df Pr(>Chisq)
employed    72.7  1  < 2e-16 ***
citizen     25.8  1  3.8e-07 ***
checks     205.2  1  < 2e-16 ***
colour      19.6  1  9.7e-06 ***
year         6.1  5   0.29785
age          0.5  1   0.49827
colour:year  21.7  5   0.00059 ***
colour:age   13.9  1   0.00019 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

The difficulty in interpreting these results from tables of coefficients can be seen in the output below:

```
> coeftest(arrests.mod)

z test of coefficients:

      Estimate Std. Error z value Pr(>|z|)
(Intercept)    0.34443    0.31007    1.11 0.26665
employedYes     0.73506    0.08477    8.67 < 2e-16 ***
citizenYes      0.58598    0.11377    5.15 2.6e-07 ***
checks        -0.36664    0.02603   -14.08 < 2e-16 ***
colourWhite     1.21252    0.34978    3.47 0.00053 ***
year1998       -0.43118    0.26036   -1.66 0.09770 .
year1999       -0.09443    0.26154   -0.36 0.71805
year2000       -0.01090    0.25921   -0.04 0.96647
year2001        0.24306    0.26302    0.92 0.35541
year2002        0.21295    0.35328    0.60 0.54664
age             0.02873    0.00862    3.33 0.00086 ***
colourWhite:year1998 0.65196    0.31349    2.08 0.03756 *
colourWhite:year1999 0.15595    0.30704    0.51 0.61152
colourWhite:year2000 0.29575    0.30620    0.97 0.33411
colourWhite:year2001 -0.38054    0.30405   -1.25 0.21073
colourWhite:year2002 -0.61732    0.41926   -1.47 0.14091
colourWhite:age   -0.03737    0.01020   -3.66 0.00025 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

By direct calculation (e.g., using `exp(coef(arrests.mod))`) you can find that the odds of a quick release was $\exp(0.735) = 2.08$ times greater for someone employed, $\exp(0.586) = 1.80$ times more likely for a Canadian citizen and $\exp(1.21) = 3.36$ times more likely for a white than a black person. It is much more difficult to interpret the interaction terms.

The primary question for the newspaper concerned the overall difference between the treatment of blacks and whites—the main effect of `colour`. We plot this as shown below, giving the plot shown in Figure 7.18. This supports the claim by the *Star* because the 95% confidence limits for blacks and whites do not overlap, and all other relevant predictors that could account for this effect have been controlled or adjusted for.

```
> plot(Effect("colour", arrests.mod),
+       lwd=3, ci.style="bands", main="",
+       xlab = list("Skin color of arrestee", cex=1.25),
+       ylab = list("Probability(released)", cex=1.25)
+       )
```

Of course, one should be very wary of interpreting main effects when there are important interactions, and the story turned out to be far more nuanced than was reported in the newspaper.




Figure 7.18: Effect plot for the main effect of skin color in the Arrests data.

{fig:arrests-eff1}

In particular, the interactions of color with with age and year provided a more complete account. Effect plots for these interactions are shown in Figure 7.19.

```
> # colour x age interaction
> plot(Effect(c("colour", "age"), arrests.mod),
+       lwd=3, multiline=TRUE,
+       xlab=list("Age", cex=1.25),
+       ylab=list("Probability(released)", cex=1.25),
+       key.args=list(x=.05, y=.99, cex=1.2)
+     )
> # colour x year interaction
> plot(Effect(c("colour", "year"), arrests.mod),
+       lwd=3, multiline=TRUE,
+       xlab=list("Year", cex=1.25),
+       ylab=list("Probability(released)", cex=1.25),
+       key.args=list(x=.7, y=.99, cex=1.2)
+     )
```

From the left panel in Figure 7.19, it is immediately apparent that the effect of age was in opposite directions for blacks and whites: Young blacks were indeed treated more severely than young whites; however for older people, blacks were treated less harshly than whites, controlling for all other predictors.

The right panel of Figure 7.19 shows the changes over time in the treatment of blacks and whites. It can be seen that up to the year 2000 there was strong evidence for differential treatment on these charges, again controlling for other predictors. There was also evidence to support the claim by the police that in the year 2001 they began training of officers to reduce racial effects in treatment.

Finally, the `effects` package provides a convenience function, `allEffects()`, that calculates the effects for all high-order terms in a given model. The `plot()` method for the "efflist" object can be used to plot individual terms selectively from a graphic menu, or plot all terms together in one comprehensive display using `ask=FALSE`.




Figure 7.19: Effect plots for the interactions of color with age (left) and year (right) in the Arrests data.

```
> arrests.effects <- allEffects(arrests.mod,
+                               xlevels=list(age=seq(15, 45, 5)))
> plot(arrests.effects,
+       ylab="Probability(released)", ci.style="bands", ask=FALSE)
```




Figure 7.20: Effect plot for all high-order terms in the model for the Arrests data

{fig:arrests-all}

The result, shown in Figure 7.20 is a relatively compact and understandable summary of the `arrests.mod` model: (a) people were more likely to be released if they were employed and citizens. (b) each additional police check decreased the likelihood of release with a summons. (c) the effect of skin color varied with age and year of arrest, in ways that tell a far more nuanced story than reported in the newspaper.

Finally, another feature of this plot bears mention: by default, the scales for each effect plot are

determined separately for each effect, to maximize use of the plot region. However, you have to read the Y scale values to judge the relative sizes of these effects. An alternative plot, using the *same* scale in each subplot¹⁶ would show the relative sizes of these effects.

△

7.4.1 More complex models: Model selection and visualization

{sec:complex}

Models with more predictors or more complex terms (interactions, non-linear terms) present additional challenges for model fitting, summarization, and visualization and interpretation. A very complicated model, with many terms and interactions may fit the data at hand quite well. However, because goodness-of-fit is optimized in the sample, terms that appear significant are less likely to be important in a future sample, and we need to worry about inflation of Type I error rates that accompany multiple significance tests. As well, it becomes increasingly difficult to visualize and understand a fitted model as the model becomes increasingly complex.

On the other hand, a very simple model may omit important predictors, interactions, or non-linear relationships with the response and give an illusion of a comfortable interpretation.

{ex:icul}

TODO: Complete this brief introduction to model selection and define AIC/BIC,

EXAMPLE 7.11: Death in the ICU

In this example we examine briefly some aspects of logistic regression related to model selection and graphical display with a large collection of potential predictors, including both quantitative and discrete variables. We use data from a classic study by Lemeshow *et al.* (1988) of patients admitted to an intensive care unit at Baystate Medical Center in Springfield, Massachusetts. The major goal of this study was to develop a model to predict the probability of survival (until hospital discharge) of these patients and to study the risk factors associated with ICU mortality. The data, contained in the data set *ICU* in *vcdExtra*, gives the results for a sample of 200 patients that was presented in Hosmer *et al.* (2013) (and earlier editions).

The *ICU* contains 22 variables of which the first, *died* is a factor. Among the predictors, two variables (*race*, *coma*) were represented initially as 3-level factors, but then recoded to binary variables (*white*, *uncons*).

```
> data("ICU", package="vcdExtra")
> names(ICU)

 [1] "died"      "age"      "sex"      "race"      "service"
 [6] "cancer"    "renal"     "infect"   "cpr"       "systolic"
[11] "hrtrate"   "previcu"  "admit"    "fracture"  "po2"
[16] "ph"        "pco"      "bic"      "creatin"   "coma"
[21] "white"     "uncons"

> ICU <- ICU[,-c(4, 20)] # remove redundant race, coma
```

Removing the 3-level versions leaves 19 predictors, of which three (age, heart rate, systolic blood pressure) are quantitative and the remainder are either binary (service, cancer) or had previously been dichotomized ($ph < 7.25$).

As an initial step, and a basis for comparison, we fit the full model containing all 19 predictors.

```
> icu.full <- glm(died ~ ., data=ICU, family=binomial)
> summary(icu.full)
```

¹⁶With the *effects* package, you can set the *ylim* argument to equate the vertical range for all plots, but this should be done on the logit scale. For this plot, `ylim = plogis(c(0.5, 1))` would work.

```

Call:
glm(formula = died ~ ., family = binomial, data = ICU)

Deviance Residuals:
    Min       1Q   Median       3Q      Max
-1.8040  -0.5606  -0.2044  -0.0863   2.9773

Coefficients:
              Estimate Std. Error z value Pr(>|z|)
(Intercept)   -6.72670    2.38551   -2.82   0.0048 **
age             0.05639    0.01862    3.03   0.0025 **
sexMale        0.63973    0.53139    1.20   0.2286
serviceSurgical -0.67352    0.60190   -1.12   0.2631
cancerYes      3.10705    1.04585    2.97   0.0030 **
renalYes       -0.03571    0.80165   -0.04   0.9645
infectYes      -0.20493    0.55319   -0.37   0.7110
cprYes         1.05348    1.00661    1.05   0.2953
systolic       -0.01547    0.00850   -1.82   0.0686 .
hrtrate        -0.00277    0.00961   -0.29   0.7732
previcuYes     1.13194    0.67145    1.69   0.0918 .
admitEmergency 3.07958    1.08158    2.85   0.0044 **
fractureYes    1.41140    1.02971    1.37   0.1705
po2<=60        0.07382    0.85704    0.09   0.9314
ph<7.25        2.35408    1.20880    1.95   0.0515 .
pco>45         -3.01844    1.25345   -2.41   0.0160 *
bic<18         -0.70928    0.90978   -0.78   0.4356
creatin>2       0.29514    1.11693    0.26   0.7916
whiteNon-white 0.56573    0.92683    0.61   0.5416
unconsYes      5.23229    1.22630    4.27   2e-05 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

(Dispersion parameter for binomial family taken to be 1)

    Null deviance: 200.16  on 199  degrees of freedom
Residual deviance: 120.78  on 180  degrees of freedom
AIC: 160.8

Number of Fisher Scoring iterations: 6

```

You can see that a few predictors are individually significant, but many are not.

However, it is useful to carry out a simultaneous global test of $H_0 : \beta = 0$ that *all* regression coefficients are zero. If this test is not significant, it makes little sense to use selection methods to choose individually significant predictors. For convenience, we define a simple function, `LRtest()`, to calculate the likelihood ratio test from the model components.

```

> LRtest <- function(model)
+   c(LRchisq=(model$null.deviance - model$deviance),
+     df=(model$df.null - model$df.residual))
>
> (LR <- LRtest(icu.full))

LRchisq      df
 79.383  19.000

> (pvalue=1-pchisq(LR[1],LR[2]))

LRchisq
2.3754e-09

```

At this point, it is tempting to examine the output from `summary(icu.full)` shown above

and eliminate those predictors which fail significance at some specified level such as the conventional $\alpha = 0.05$. This is generally a bad idea for many reasons.¹⁷

A marginally better approach is to remove non-significant variables whose coefficients have signs that don't make sense from the substance of the problem. For example, in the full model, both `renal` (history of chronic renal failure) and `infect` (infection probable at ICU admission) have negative signs, meaning that their presence *decreases* the odds of death. We remove those variables using `update()`; as expected they make little difference.

```
> icu.full11 <- update(icu.full, . ~ . - renal - fracture)
> anova(icu.full11, icu.full, test="Chisq")

Analysis of Deviance Table

Model 1: died ~ age + sex + service + cancer + infect + cpr + systolic +
  hrtrate + previcu + admit + po2 + ph + pco + bic + creatin +
  white + uncons
Model 2: died ~ age + sex + service + cancer + renal + infect + cpr +
  systolic + hrtrate + previcu + admit + fracture + po2 + ph +
  pco + bic + creatin + white + uncons
   Resid. Df Resid. Dev Df Deviance Pr(>Chi)
1         182         122
2         180         121   2      1.7    0.43
```

Before proceeding to consider model selection, it is useful to get a better visual overview of the current model than is available from a table of coefficients and significance tests. Some very useful `print()`, `summary()` and `plot()` methods are available in the `rsm` package. Unfortunately, these require that the logistic model is fitted with `lrm()` in that package rather than with `glm()`. We pause here to refit the same model as `icu.full11` in order to show a plot of odds ratios for the terms in this model.

```
> library(rms)
> dd <- datadist(ICU[, -1])
> options(datadist="dd")
> icu.lrm1 <- lrm(died ~ ., data=ICU)
> icu.lrm1 <- update(icu.lrm1, . ~ . - renal - fracture)
```

The `summary()` method for "rms" objects produces a much more detailed descriptive summary of a fitted model, and the `plot()` method for that summary object gives a sensible plot of the odds ratios for the model terms together with confidence intervals, at levels (0.9, 0.95, 0.99) by default. The following lines produce Figure 7.21.

```
> sum.lrm1 <- summary(icu.lrm1)
> plot(sum.lrm1, log=TRUE, main="Odds ratio for 'died'", cex=1.25,
+       col = rgb(0.1, 0.1, 0.8, alpha = c(0.3, 0.5, 0.8)))
```

In this plot, continuous variables are shown at the top, followed by the discrete predictors. In each line, the range or levels of the predictors are given in the form $a : b$, such that the value a corresponds to the numerator of the odds ratio plotted. Confidence intervals that don't overlap the vertical line for odds ratio = 1 are significant, but this graph shows those at several confidence levels, allowing you to decide what is "significant" visually. As well, the widths of those intervals convey the precision of these estimates.

Among several stepwise selection methods in R for "glm" models, `stepAIC()` in the `MASS` package implements a reasonable collection of methods for forward, backward and stepwise selection using penalized AIC-like criteria that balance goodness of fit against parsimony. The method

¹⁷It ignores the facts of (a) an arbitrary cutoff value for significance, (b) the strong likelihood that chance features of the data or outliers influence the result, (c) problems of collinearity, etc. See Harrell (2001, §4.3) for a useful discussion of these issues.




Figure 7.21: Odds ratios for the terms in the model for the ICU data. Each line shows the odds ratio for a term, together with lines for 90, 95 and 99% confidence intervals in progressively darker shades.

g:icul-odds-ratios}

takes an argument, `scope`, which is a list of two model formulae; `upper` defines the largest (most complex) model to consider and `lower` defines the smallest (simplest) model, e.g., `lower = ~ 1` is the intercept-only model.

By default, the function produces verbose printed output showing the details of each step, but we suppress that here to save space. It returns the final model as its result, along with an `anova` component that summarises the deviance and AIC from each step.

```
> library(MASS)
> icu.step1 <- stepAIC(icu.full1, trace = FALSE)
> icu.step1$anova
```

Stepwise Model Path
Analysis of Deviance Table

Initial Model:
died ~ age + sex + service + cancer + infect + cpr + systolic +
hrtrate + previcu + admit + po2 + ph + pco + bic + creatin +
white + uncons

Final Model:
died ~ age + cancer + systolic + admit + ph + pco + uncons

	Step	Df	Deviance	Resid. Df	Resid. Dev	AIC
1				182	122.48	158.48
2	- po2	1	0.062446	183	122.54	156.54
3	- creatin	1	0.059080	184	122.60	154.60
4	- hrtrate	1	0.072371	185	122.67	152.67
5	- infect	1	0.122772	186	122.79	150.79
6	- white	1	0.334999	187	123.13	149.13
7	- service	1	0.671313	188	123.80	147.80
8	- bic	1	0.377521	189	124.18	146.18

9	- cpr	1	1.148260	190	125.33	145.33
10	- sex	1	1.543523	191	126.87	144.87
11	- previcu	1	1.569976	192	128.44	144.44

Alternatively, we can use the BIC criterion, by specifying $k=\log(n)$, which generally will select a smaller model when the sample size is reasonably large.

```
> icu.step2 <- stepAIC(icu.full, trace = FALSE, k=log(200))
> icu.step2$anova
```

Stepwise Model Path
Analysis of Deviance Table

Initial Model:
died ~ age + sex + service + cancer + renal + infect + cpr +
systolic + hrtrate + previcu + admit + fracture + po2 + ph +
pco + bic + creatin + white + unconc

Final Model:
died ~ age + cancer + admit + unconc

	Step	Df	Deviance	Resid. Df	Resid. Dev	AIC
1				180	120.78	226.74
2	- renal	1	0.0019881	181	120.78	221.45
3	- po2	1	0.0067968	182	120.79	216.16
4	- creatin	1	0.0621463	183	120.85	210.92
5	- hrtrate	1	0.0658870	184	120.92	205.69
6	- infect	1	0.2033221	185	121.12	200.59
7	- white	1	0.3673180	186	121.49	195.66
8	- bic	1	0.6002993	187	122.09	190.96
9	- service	1	0.7676303	188	122.85	186.43
10	- fracture	1	1.3245086	189	124.18	182.46
11	- cpr	1	1.1482598	190	125.33	178.31
12	- sex	1	1.5435228	191	126.87	174.55
13	- previcu	1	1.5699762	192	128.44	170.83
14	- ph	1	4.4412370	193	132.88	169.97
15	- pco	1	2.7302934	194	135.61	167.40
16	- systolic	1	3.5231028	195	139.13	165.63

This model differs from model `icu.step1` selected using AIC in the last three steps, that also removed `ph`, `pco` and `systolic`.

```
> coefTest(icu.step2)
```

z test of coefficients:

	Estimate	Std. Error	z value	Pr(> z)
(Intercept)	-6.8698	1.3188	-5.21	1.9e-07 ***
age	0.0372	0.0128	2.91	0.00360 **
cancerYes	2.0971	0.8385	2.50	0.01238 *
admitEmergency	3.1022	0.9186	3.38	0.00073 ***
unconcYes	3.7055	0.8765	4.23	2.4e-05 ***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

These two models are nested, so we can compare them directly using a likelihood ratio test from `anova()`.

```
> anova(icu.step2, icu.step1, test="Chisq")

Analysis of Deviance Table

Model 1: died ~ age + cancer + admit + uncons
Model 2: died ~ age + cancer + systolic + admit + ph + pco + uncons
  Resid. Df Resid. Dev Df Deviance Pr(>Chi)
1      195      139
2      192      128  3      10.7    0.013 *
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

The larger model is significantly better by this test, but the smaller model is simpler to interpret. We retain these both as “candidate models” to be explored further, but for ease in this example, we do so using the smaller model, `icu.step2`.

Another important step is to check for non-linearity of quantitative predictors such as age and interactions among the predictors. This is easy to do using `update()` and `anova()` as shown below. First, allow a non-linear term in age, and all two-way interactions of the binary predictors.

```
> icu.glm3 <- update(icu.step2, . ~ . -age + ns(age,3) + (cancer+admit+uncons)^2)
> anova(icu.step2, icu.glm3, test="Chisq")

Analysis of Deviance Table

Model 1: died ~ age + cancer + admit + uncons
Model 2: died ~ cancer + admit + uncons + ns(age, 3) + cancer:admit +
  cancer:uncons + admit:uncons
  Resid. Df Resid. Dev Df Deviance Pr(>Chi)
1      195      139
2      191      135  4      3.73    0.44
```

Next, we can check for interactions with age:

```
> icu.glm4 <- update(icu.step2, . ~ . + age*(cancer+admit+uncons))
> anova(icu.step2, icu.glm4, test="Chisq")

Analysis of Deviance Table

Model 1: died ~ age + cancer + admit + uncons
Model 2: died ~ age + cancer + admit + uncons + age:cancer + age:admit +
  age:uncons
  Resid. Df Resid. Dev Df Deviance Pr(>Chi)
1      195      139
2      192      134  3      5.37    0.15
```

None of these additional terms have much effect.

So, we will tentatively adopt the simple main effects model, `icu.step2`, and consider how to visualize in interpret this result. One interesting display is a *nomogram* that shows how values on the various predictors translate into a predicted value of the log odds, and the relative strengths of their effects on this prediction. This kind of plot is shown in Figure 7.22, produced using `nomogram()` in the `rms` package as follows. It only works with models fit using `lrm()`, so we have to refit this model.

```
> icu.lrm2 <- lrm(died ~ age + cancer + admit + uncons, data=ICU)
> plot(nomogram(icu.lrm2), cex.var=1.2, lplabel="Log odds death")
```

In this nomogram, each predictor is scaled according to the size of its effect on a common scale of 0–100 “points.” A representative observation is shown by the marked points, corresponding to




Figure 7.22: Nomogram for predicted values in the simple main effects model for the ICU data. Each predictor is scaled in relation to its effect on the outcome in terms of “points”, 0–100. Adding the points for a given case gives total points that have a direct translation to log odds. The marked points show the prediction for someone of age 60, admitted to the emergency ward and unconscious.

{fig:icu-nomogram}

a person of age 60, without cancer, who was admitted to emergency and was unconscious at that time. Adding the points associated with each variable value gives the result shown on the scale of total points. For this observation, the result is $50 + 0 + 84 + 100 = 234$, for which the scale of log odds at the bottom gives a predicted logit of 2.2, or a predicted probability of death of $1/(1 + \exp(-2.2)) = 0.90$.

This leaves us with the problem of how to visualize the fitted model compactly and comprehensively. Full-model plots and effect plots, as we have used them, are somewhat unwieldy with four or more predictors if we want to view all effects simultaneously because it is more difficult to make comparisons across multiple panels (particularly if the vertical scales differ).

One way to reduce the visual complexity of such graphs is to combine some predictors that would otherwise be shown in separate panels into a recoding that can be shown as multiple curves for their combinations in fewer panels. In general, this can be done by combining some predictors interactively; for example with sex and education as factors, their combinations, `M:Hi`, `M:Lo`, etc. could be used to define a new variable, `group` used as the curves in one plot, rather than separate panels.

In this case, because age is continuous, it makes sense to plot fitted values against age. With cancer, admit and uncons as binary factors associated with risk of death, it is also sensible to combine them all into a single variable, `risks`, indicating which one or more risk factors are present for each case. We first convert each variable to an abbreviation for the risk, if present, or “”, and paste these together.

```
> # combine categorical risk factors to a single string
> risks <- ICU[, c("cancer", "admit", "uncons")]
> risks[,1] <- ifelse(risks[,1]=="Yes", "Cancer", "")
> risks[,2] <- ifelse(risks[,2]=="Emergency", "Emerg", "")
> risks[,3] <- ifelse(risks[,3]=="Yes", "Uncons", "")
> risks <- apply(risks, 1, paste, collapse="")
```

```
> risks[risks==""] <- "(none)"
```

```
> table(risks)
```

```
risks
(none)      Cancer CancerEmerg      Emerg EmergUncons
      37         15          5        128         14
Uncons      1
```

The frequency counts of the risk combinations show that admission to the emergency ward alone was most frequent, and only one patient had unconsciousness as the only risk.

As done before, we can then get the fitted logit values for the chosen model, and combine these with the data and the `risks` variable.

```
> icu.glm2 <- glm(died ~ age + cancer + admit + uncons,
+               data=ICU, family=binomial)
> icu.fit <- cbind(ICU, predict(icu.glm2, se=TRUE), risks)
```

In the plot step below, we use `geom_ribbon()` to plot a one standard error confidence band around the fitted logits. `color=risks` gives separate curves for each level of the risks factor.

```
> gg <- ggplot( icu.fit, aes(x=age, y=fit, color=risks)) +
+   geom_line(size = 1.2) + theme_bw() +
+   geom_ribbon(aes(ymin = fit - se.fit,
+                 ymax = fit + se.fit,
+                 fill = risks),
+             alpha = 0.2,
+             color = "transparent") +
+   theme_bw() +
+   labs(x = "Age", y = "Log odds (died)") +
+   geom_point(size=2)
```

By default, `ggplot()` uses a legend to display the labels for the curve variable, but the graph is more readable using `directlabels`, giving the plot shown in Figure 7.23.

```
> library(directlabels)
> direct.label(gg+xlim(10,100), last.points)
```

From this graph, it is apparent that the log odds of mortality increases with age in all cases. Relative to the line labeled `(none)`, mortality is higher when any of these risk factors are present, particularly when the patient is admitted to Emergency; it is highest when the patient is also unconscious at admission. The vertical gaps between lines that share a common risk (e.g., `Cancer`, `CancerEmerg`) indicate the additional increment from one more risk. Finally, the plotted points show the number and age distribution of these various combinations.

Before concluding that this model provides an adequate description of the data, we should examine whether any individual cases are unduly influencing the predicted results, and more importantly, the choice of variables in the model. We examine this question in Section 7.5 where we return to these data (Example 7.13).

△

7.5 Influence and diagnostic plots

In ordinary least squares (OLS) regression, measures of *influence* (leverage, Cook's D, DFBETAs, etc.) and associated plots help you to determine whether individual cases (or cells in grouped

{sec:logist-infl}




Figure 7.23: Fitted log odds of death in the ICU data. Each line shows the relationship with age, for patients having various combinations of risk factors.

{fig:icul-fit-plot}

data) have undue impact on the fitted regression model and the coefficients of individual predictors. Analogs of most of these measures have been suggested for logistic regression and generalized linear models. Pregibon (1981) provided the theoretical basis for these methods, exploiting the relationship between logistic models and weighted least squares. Some additional problems occur in practical applications to logistic regression because the response is discrete, and because the leave-one-out diagnostics are more difficult to compute, but the ideas are essentially the same.

7.5.1 Residuals and leverage

As in ordinary least squares regression, the influence (actual impact) of an observation in logistic models depends multiplicatively on its residual (disagreement between y_i and \hat{y}_i) and its leverage (how unusual \mathbf{x}_i is in the space of the explanatory variables). A conceptual formula is

$$\text{Influence} = \text{Leverage} \times \text{Residual}$$

This multiplicative definition implies that a case is influential to the extent that it is both poorly fit *and* has unusual values of the predictors.

7.5.1.1 Residuals

In logistic regression, the simple raw residual is just $e_i \equiv y_i - \hat{p}_i$, where $\hat{p}_i = 1/[1 + \exp(-\mathbf{x}_i^T \mathbf{b})]$.

The Pearson and deviance residuals are more useful for identifying poorly fitted observations, and are components of overall goodness-of-fit statistics. The (raw) **Pearson residual** is defined as

$$r_i \equiv \frac{e_i}{\sqrt{\hat{p}_i(1 - \hat{p}_i)}} \quad (7.7)$$

and the Pearson chi-square is therefore $\chi^2 = \sum r_i^2$. The **deviance residual** is

$$g_i \equiv \pm 2[y_i \log \hat{p}_i + (1 - y_i) \log(1 - \hat{p}_i)]^{1/2} \quad (7.8)$$

where the sign of g_i is the same as that of e_i . Likewise, the sum of squares of the deviance residuals gives the overall deviance, $G^2 = -2 \log \mathcal{L}(\mathbf{b}) = \sum g_i^2$.

When y_i is a binomial count based on n_i trials (grouped data), the Pearson residuals Eqn. (7.7) then become

$$r_i \equiv \frac{y_i - n_i \hat{p}_i}{\sqrt{n_i \hat{p}_i (1 - \hat{p}_i)}}$$

with similar modifications made to Eqn. (7.8).

In R, `residuals()` is the generic function for obtaining (raw) residuals from a model fitted with `glm()` (or `lm()`). However **standardized residuals**, given by `rstandard()`, and **studentized residuals**, provided by `rstudent()` are often more useful because they rescale the residuals to have unit variance. They use, respectively, an overall estimate, $\hat{\sigma}^2$ of error variance, and the leave-one-out estimate, $\hat{\sigma}_{(-i)}^2$, omitting the i th observation; the studentized version is usually to be preferred in model diagnostics because it also accounts for the impact of the observation on residual variance.

7.5.1.2 Leverage

Leverage measures the *potential* impact of an individual case on the results, which is directly proportional to how far an individual case is from the centroid in the space of the predictors. Leverage is defined as the diagonal elements, h_{ii} , of the “Hat” matrix, \mathbf{H} ,

$$\mathbf{H} = \mathbf{X}^* (\mathbf{X}^{*\top} \mathbf{X}^*)^{-1} \mathbf{X}^{*\top}$$

where $\mathbf{X}^* = \mathbf{V}^{1/2} \mathbf{X}$, and $\mathbf{V} = \text{diag}[\hat{p}(1 - \hat{p})]$. As in OLS, leverage values are between 0 and 1, and a leverage value, $h_{ii} > \{2 \text{ or } 3\}k/n$ is considered “large”; here, $k = p + 1$ is the number of coefficients including the intercept and n is the number of cases. In OLS, however, the hat values depend only on the X s, whereas in logistic regression, they also depend on the dependent variable values and the fitted probabilities (through \mathbf{V}). As a result, an observation may be extremely unusual on the predictors, yet not have a large hat value, if the fitted probability is near 0 or 1. The function `hatvalues()` calculates these values for a fitted “glm” model object.

7.5.2 Influence diagnostics

{sec:logist-inflldiag}

Influence measures assess the effect that deleting an observation has on the regression parameters, fitted values, or the goodness-of-fit statistics. In OLS, these measures can be computed exactly from a single regression. In logistic regression, the exact effect of deletion requires refitting the model with each observation deleted in turn, a time-intensive computation. Consequently, Pregibon (1981) showed how analogous deletion diagnostics may be approximated by performing one additional step of the iterative procedure. Most modern implementations of these methods for generalized linear models follow Williams (1987).

The simplest measure of influence of observation i is the standardized change in the coefficient for each variable due to omitting that observation, termed **DFBETAs**. From the relation (Pregibon, 1981, p. 716)

$$\mathbf{b} - \mathbf{b}_{(-i)} = (\mathbf{X}^\top \mathbf{V} \mathbf{X})^{-1} \mathbf{x}_i (y_i - \hat{p}_i) / (1 - h_{ii}) ,$$

the estimated standardized change in the coefficient for variable j is

$$\text{DFBETA}_{ij} \equiv \frac{b_{(-i)j} - b_j}{\hat{\sigma}(b_j)} , \quad (7.9) \quad \{\text{eq:dfbeta}\}$$

where $\hat{\sigma}(b_j)$ is the estimated standard error of b_j . With k regressors, there are $k + 1$ sets of DFBETAs, which makes their examination burdensome. Graphical displays ease this burden, as do various summary measures considered below.

The most widely used summary of the overall influence of observation i on the estimated regression coefficients is **Cook's distance**, which measures the average squared distance between \mathbf{b} for all the data and $\mathbf{b}_{(-i)}$ estimated without observation i . It is defined as

$$C_i \equiv (\mathbf{b} - \mathbf{b}_{(-i)})^\top \mathbf{X}^\top \mathbf{V} \mathbf{X} (\mathbf{b} - \mathbf{b}_{(-i)}) / k \hat{\sigma}^2 .$$

However, Pregibon (1981) showed that C_i could be calculated simply as

$$C_i = \frac{r_i^2 h_{ii}}{k(1 - h_{ii})^2} , \quad (7.10)$$

where $r_i = y_i - \hat{p}_i / \sqrt{v_{ii}(1 - h_{ii})}$ is the i th standardized Pearson residual and v_{ii} is the i th diagonal element of \mathbf{V} . Rules of thumb for noticeably “large” values of Cook's D are only rough indicators, and designed so that only “noteworthy” observations are nominated as unusually influential. One common cutoff for an observation to be treated as influential is $C_i > 1$. Others refer the values of C_i to a χ_k^2 or $F_{k, n-k}$ distribution.

Another commonly used summary statistic of overall influence is the **DFFITs** statistic, a standardized measure of the difference between the predicted value \hat{y}_i using all the data and the predicted value $\hat{y}_{(-i)}$ calculated omitting the i th observation.

$$\text{DFFITs}_i = \frac{\hat{y}_i - \hat{y}_{(-i)}}{\hat{\sigma}_{(-i)} \sqrt{h_{ii}}} ,$$

where $\hat{\sigma}_{(-i)}$ is the estimated standard error with the i th observation deleted. For computation, DFFITS can be expressed in terms of the standardized Pearson residual and leverage as

$$\text{DFFITs}_i = r_i \sqrt{\frac{h_{ii}}{(1 - h_{ii})} \frac{v_{ii}}{v_{(-ii)}}} . \quad (7.11)$$

From Eqn. (7.10) and Eqn. (7.11) it can be shown that Cook's distance is nearly the square of DFFITS divided by k ,

$$C_i = \frac{v_{(-ii)}^2}{v_{ii}^2} \frac{\text{DFFITs}_i^2}{k} . \quad (7.12)$$

Noteworthy values of DFFITS are often nominated by the rule-of-thumb $\text{DFFITs}_i > 2$ or $3\sqrt{k/n - k}$.

In R, these influence measures are calculated for a fitted “glm” model using `cooks.distance()` and `dffits()`. The `dfbeta()` function calculates and returns the matrix of all standardized changes in the model coefficients (Eqn. (7.9)) due to omitting each observation in turn.¹⁸ A convenience function, `influence.measures()` gives a tabular display showing the DFBETA_{ij} for each model variable, DFFITS, Cook's distances and the diagonal elements of the hat matrix. Cases which are influential with respect to any of these measures are marked with an asterisk.¹⁹

Beyond printed output of these numerical summaries, plots of these measures can shed light on potential problems due to influential or other noteworthy cases. By highlighting them, such plots provide the opportunity to determine if and how any of these affect your conclusions, or to take some corrective action.

A basic collection of diagnostic plots is provided by the `plot()` method for a “glm” model object. The `car` package contains a variety of other functions for model diagnostic plots. We illustrate some of these in the examples below.

¹⁸**TODO:** Not quite true: `dfbeta()` doesn't have a “glm” method. Either omit this or write a `dfbeta.glm()` method that gives the same results as `influence.measures()`.

¹⁹See `help(influence.measures)` for the description of all of these functions for residuals, leverage and influence diagnostics in generalized linear models.

EXAMPLE 7.12: Donner Party

This example re-visits the data on the Donner Party examined in Example 7.9. For illustrative purposes, we consider the influence measures and diagnostic plots for one specific model, the model `donner.mod3`, that included a quadratic effect of age and a main effect of sex, but no interaction.

The simplest overview of the adequacy of a fitted model is provided by the `plot()` method for a "glm" (or "lm") object. This function can produce up to six different plots that can be plotted individually or selected (using the argument `which`) and composed into a single overview figure using `par(mfrow=c(rows, cols))` as shown below.

It is useful to see the entire collection because, by default, only four are plotted (`which=c(1:3, 5)`) and this selection (sometimes called the *regression quartet* of diagnostic plots) is tuned more to classical linear models for quantitative data. Important feature of these plots are that (a) plot annotations are added to each showing trends or expected behaviour under the assumptions of a fitted model; (b) noteworthy observations are labeled individually.

```
> caption = list("(1) Residuals vs Fitted",
+               "(2) Normal Q-Q",
+               "(3) Scale-Location",
+               "(4) Cook's distance",
+               "(5) Residuals vs Leverage",
+               expression("(6) Cook's dist vs Leverage "
+                           * h[ii] / (1 - h[ii])))
> op <- par(mfrow=c(3,2), mar=c(4,4,2,1)+.1, cex.lab=1.2, cex=1)
> plot(donner.mod3, which=1:6, caption=caption)
> par(op)
```

The six plots, corresponding to the values of `which`, shown in Figure 7.24 for the `donner.mod3` model are:

1. a plot of residuals against fitted values. In a classical linear model, this plot should appear unstructured (random around the zero line), but for logistic regression there will always be two sequences of points, corresponding to the 0/1 observations.
2. a normal Q-Q plot of ordered residuals vs. the corresponding quantiles of the gaussian distribution. In a classical linear model, all points should follow the dotted reference line, but this will rarely hold for logistic regression models.
3. a Scale-Location plot of $\sqrt{|\text{residuals}|}$ against fitted values, with a loess smoothed curve showing the trend for variance of the residual to change with the predicted value. This is useful to detect non-constant residual variance in classical models, but in logistic regression, you will almost always see a U-shaped pattern corresponding to the fact that the variance around the fitted value is a function of $\sqrt{\hat{p}_i(1 - \hat{p}_i)}$.
4. an index plot of Cook's distances versus observation numbers, with the largest `id.n` values labeled.
5. a plot of residuals against leverages, showing contours of Cook's distances. Among all of these plots, this is probably the most useful for assessment of influence in both classical and generalized linear models. The function `influencePlot()` in `car` provides a better version of this plot, using the size of a bubble symbol to also show Cook's distance directly.
6. a plot of Cook's distances against leverage/(1-leverage). In this plot contours of standardized residuals that are equal in magnitude are lines through the origin, and labeled with their absolute values. Consequently, more influential observations appear toward the top.

In all these plots, three observations are labeled as noteworthy, by one criterion or another with a default number given by `id.n=3`. Plotting just the residual-leverage graph (`which=5`) with some additional annotations to show the conventional cutoff values gives Figure 7.25.




Figure 7.24: Diagnostic plots for a glm object, using the fitted model `donner.mod3` for the Donner Party data. Each plot shows some additional annotations or smoothed curves and labels observations considered noteworthy in terms of influence.

{fig:donner2-plot}

```

> op <- par(mar=c(5,4,4,2)+.1)
> plot(donner.mod3, which=5, cex.id=1, cook.levels=c(0.25, 0.5), id.n=3)
> abline(h=c(-2, 2), col="gray")
> k <- length(coef(donner.mod3))
> n <- nrow(Donner)
> abline(v=c(2, 3)*k/n, col="gray")
> text(x=c(2, 3)*k/n, y=-2.3, c("2k/n", "3k/n"))
> par(op)

```




Figure 7.25: Residual vs. Leverage plot for the Donner data model. Horizontal and vertical reference lines show typical cutoff values for noteworthy residuals and leverage.

{fig:donner2-plot5}

Details of all the diagnostic measures for a given model including the DFBETAs for individual coefficients can be obtained using `influence.measures`. This can be useful for custom plots not provided elsewhere (see Example 7.13).

```

> infl <- influence.measures(donner.mod3)
> names(infl)

```

```
[1] "infmat" "is.inf" "call"
```

The `summary()` method for the "infl" object prints those observations considered noteworthy on one or more of these statistics, as indicated by a "*" next to the value.

```
> summary(infl)
```

```

Potentially influential observations of
glm(formula = survived ~ poly(age, 2) + sex, family = binomial,      data = Donner) :

```

	dfb.1_	dfb.p(,2)1	dfb.p(,2)2	dfb.sxM1	dffit	cov.r	cook.d	hat
Breen, Patrick	0.08	0.65	0.56	0.23	0.69_*	0.93	0.32	0.09
Donner, Elizabeth	-0.26	-0.34	-0.22	0.12	-0.40	1.15_*	0.03	0.14_*
Graves, Elizabeth C.	-0.24	-0.37	-0.26	0.10	-0.42	1.20_*	0.03	0.16_*

The function `influencePlot()` in the `car` package gives a similar plot, but uses the size

(area) of the plotting symbol to also show the value of Cook's D as shown in Figure 7.26. Like other diagnostic plots in `car`, it is considerably more general than illustrated here, because it allows for different `id.methods` to label noteworthy points, including `id.method="identify"` for interactive point identification by clicking with the mouse. The `id.n` argument works differently than with `plot()`, because it selects the most extreme `id.n` observations on *each* of the studentized residual, hat value and Cook's D, and labels all of these.

```
> library(car)
> res <- influencePlot(donner.mod3, id.col="blue", scale=8, id.n=2)
> text(x=c(2, 3)*k/n, y=-1.8, c("2k/n", "3k/n"))
```




Figure 7.26: Influence plot (residual vs. leverage) for the Donner data model, showing Cook's D as the size of the bubble symbol. Horizontal and vertical reference lines show typical cutoff values for noteworthy residuals and leverage.

g:donner2-inflplot}

Conveniently, `influencePlot()` returns a data frame containing the influence statistics for the points identified in the plot (`res` in the call above). We can combine this with the data values to help learn why these points are considered influential.

```
> # show data together with diagnostics for influential cases
> idx <- which(rownames(Donner) %in% rownames(res))
> cbind(Donner[idx,2:4], res)
```

	age	sex	survived	StudRes	Hat	CookD
Breen, Patrick	51	Male	yes	2.501	0.09148	0.5688
Donner, Elizabeth	45	Female	no	-1.114	0.13541	0.1846
Graves, Elizabeth C.	47	Female	no	-1.019	0.16322	0.1849
Reed, James	46	Male	yes	2.098	0.08162	0.3790

We can see that Patrick Breen and James Reed²⁰ are unusual because they were both older men

²⁰Breen and Reed, both born in Ireland, were the leaders of their family groups. Among others, both kept detailed diaries of their experiences, from which most of the historical record derives. Reed was also the leader of two relief parties sent out to find rescue or supplies over the high Sierra mountains, so it is all the more remarkable that he survived.

who survived, and have large positive residuals; Breen is the most influential by Cook's D, but this value is not excessively large. The two women were among the older women who died. They are selected here because they have the largest hat values, meaning they are unusual in terms of the distribution of age and sex, but they are not particularly influential in terms of Cook's D.

A related graphical display is the collection of index plots provided by `influenceIndexPlot()` in `car`, which plots various influence diagnostics against the observation numbers in the data. The `id.n` argument here works to label that number of the most extreme observations *individually* for each measure plotted. The following call produces Figure 7.27.

```
> influenceIndexPlot(donner.mod3, vars=c("Cook", "Studentized", "hat"),
+                   id.n=4)
```




Figure 7.27: Index plots of influence measures for the Donner data model. The four most extreme observations on each measure are labeled.

{fig:donner2-indexinfl

In our opinion, *separate* index plots are often less useful than combined plots such as the leverage-influence plot that shows residuals, leverage and Cook's D together. However, the `car` version in Figure 7.27 does that too, and allows us to consider how unusual the labeled observations are both individually and in combination.

△

{ex:icu2}

EXAMPLE 7.13: Death in the ICU

In Example 7.11 we examined several models to account for death in the *ICU* data set. We continue this analysis here, with a focus on the simple main effects model, `icu.glm2`, for which the fitted logits were shown in Figure 7.23. For ease of reference, we restate that model here:

```
> icu.glm2 <- glm(died ~ age + cancer + admit + uncons,
+                data=ICU, family=binomial)
```

The plot of residual vs. leverage for this model is shown in Figure 7.28.

```
> library(car)
> res <- influencePlot(icu.glm2, id.col="red", scale=8, id.cex=1.5, id.n=3)
```




Figure 7.28: Influence plot for the main effects model for the ICU data

Details for the cases identified in the figure are shown below, again using `rownames(res)` to select the relevant observations from the *ICU* data.

```
> idx <- which(rownames(ICU) %in% rownames(res))
> cbind(ICU[idx, c("died", "age", "cancer", "admit", "uncons")], res)
```

	died	age	cancer	admit	uncons	StudRes	Hat	CookD
84	No	59	No	Emergency	Yes	-2.258	0.06781	0.3626
371	No	46	Yes	Emergency	No	-1.277	0.16408	0.2210
766	No	31	Yes	Emergency	No	-1.028	0.17062	0.1719
881	No	89	No	Emergency	Yes	-2.718	0.03081	0.4106
127	Yes	19	No	Emergency	No	2.565	0.01679	0.2724
208	Yes	70	No	Elective	Yes	1.662	0.29537	0.4568
380	Yes	20	No	Emergency	No	2.548	0.01672	0.2668

None of the cases are particularly influential on the model coefficients overall: the largest Cook's D is only 0.45 for case 208. This observation also has the largest hat value. It is unusual on the predictors in this sample: a 70 year old man without cancer, admitted on an elective basis, who nonetheless died. However, this case is also highly unusual in having been unconscious on admission for an elective procedure, and signals that there might have been a coding error or other anomaly for this observation.

Another noteworthy observation identified here is case 881, an 89 year old male, admitted unconscious as an emergency; this case is poorly predicted because he survived. Similarly, two other cases (127, 380) with large studentized residuals are poorly predicted because they died, although they were young, did not have cancer, and conscious at admission. However, these cases have relatively small Cook's D values. From this evidence we might conclude that, case 208 bears further scrutiny, but none of these cases greatly affects the model, its coefficients, or interpretation.

For comparison with Figure 7.28, the related index plot of these measures is shown in Figure 7.29.

```
> influenceIndexPlot(icu.glm2, vars=c("Cook", "Studentized", "hat"), id.n=4)
```




Figure 7.29: Index plots of influence measures for the ICU data model. The four most extreme observations on each measure are labeled.

{fig:icu2-infl-index}

Cook's D and DFFITS are *overall* measures of the total influence that cases have on the regression coefficients and fitted values respectively. It might be that some cases have a large impact on some individual regression coefficients, but don't appear particularly unusual in these aggregate measures.

One way to study this is to make plots of the $DFBETA_{ij}$ statistics. Such plots are not available (as far as we know) in R packages, but it is not hard to construct them from the result returned by `influence.measures()`. To do this, we select the appropriate columns from the `infmt` component returned by that function.

```
> infl <- influence.measures(icu.glm2)
> dfbetas <- data.frame(infl$infmt[,2:5])
> colnames(dfbetas) <- c("dfb.age", "dfb.cancer", "dfb.admit", "dfb.uncons")
> head(dfbetas)
```

	dfb.age	dfb.cancer	dfb.admit	dfb.uncons
8	0.047340	0.013418	0.004067	0.009254
12	0.018988	0.018412	-0.004174	0.018106
14	-0.001051	0.014882	0.026278	0.005555
28	0.031562	0.018424	-0.001511	0.016640
32	-0.164084	0.003788	-0.036505	0.023488
38	-0.021525	0.016539	-0.011937	0.020803

To illustrate this idea, plotting an individual column of `dfbetas` using `type = "h"` gives an index plot against the observation number. This is shown in Figure 7.30 for the impact on the coefficient for age. The lines and points are colored blue or red according to whether the patient lived or died. Observations for which the $|DFBETA_{age}| > 0.2$ (an arbitrary value) are labeled.

```

> cols=ifelse (ICU$died=="Yes", "red", "blue")
> op <- par(mar=c(5,5,1,1)+.1)
> plot(dfbetas[,1], type = "h", col=cols,
+      xlab="Observation index",
+      ylab=expression(Delta * beta[Age]),
+      cex.lab=1.3)
> points(dfbetas[,1], col=cols)
> # label some points
> big <- abs(dfbetas[,1]) > .25
> idx <- 1:nrow(dfbetas)
> text(idx[big], dfbetas[big,1], label=rownames(dfbetas)[big],
+      cex=0.9, pos=ifelse(dfbetas[big,1]>0, 3, 1),
+      xpd=TRUE)
> abline(h=c(-.25, 0, .25), col="gray")
> par(op)

```




Figure 7.30: Index plot for DFBETA (Age) in the ICU data model. The observations are colored blue or red according to whether the patient lived or died.

{fig:icu2-dbage}

None of the labeled points here are a cause for concern, since the standardized DFBETAs are all relatively small. However, the plot shows that patients who died have generally larger impacts on this coefficient.

An alternative to individual index plots is a scatterplot matrix, that shows the pairwise changes in the regression coefficients for the various predictors. Here we use `scatterplotMatrix()` from `car` that offers features for additional plot annotations, including identifying the most unusual points in each pairwise plot. In each off-diagonal panel, a 95% data ellipse and linear regression line helps to show the marginal relationship between the two measures and highlight why the labeled points are atypical in each plot.²¹

```

> scatterplotMatrix(dfbetas, smooth=FALSE, id.n=2,
+   ellipse=TRUE, levels=0.95, robust=FALSE,
+   diagonal="histogram",
+   groups=ICU$died, col=c("blue", "red"))

```

²¹ This plot uses the `id.method="mahal"` method to label the most extreme observations according to the Mahalanobis distance of each point from the centroid in the plot.




Figure 7.31: Scatterplot matrix for DFBETAs from the model for the ICU data. Those who lived or died are shown with blue circles and red triangles, respectively. The diagonal panels show histograms of each variable.

fig:icu2-dbscatmat}



7.5.3 Other diagnostic plots*

The graphical methods described in this section are relatively straight-forward indicators of the adequacy of a particular model, with a specified set of predictors, each expressed in a given way. More sophisticated methods have also been proposed, which focus on the need to include a particular predictor and whether its relationship is linear. These include the *component-plus-residual plot*, the *added-variable plot*, and the *constructed variable plot*, which are all analogous to techniques developed in OLS.

7.5.3.1 Component-plus-residual plots

The *component-plus-residual plot* (also called a *partial residual plot*) proposed originally by Larsen and McCleary (1972) is designed to show whether a given quantitative predictor, x_j , included linearly in the model, actually shows a nonlinear relation, requiring transformation. The essential idea is to move the linear term for x_j back into the residual, by calculating the *partial residuals*,

$$r_j^* = r + \beta_j x_j$$

Then, a plot of r_j^* against x_j will have the same slope, β_j , as the full model including it among other predictors. However, any non-linear trend will be shown in the pattern of the points, usually aided by a smoothed non-parametric curve.

As adapted to logistic regression by Landwehr *et al.* (1984), the partial residual for variable x_j is defined as

$$r_j^* = V^{-1}r + \beta_j x_j$$

The partial residual plot is then a plot of r_j^* against x_j , possibly with the addition of a smoothed lowess curve (Fowlkes, 1987) and a linear regression line to aid interpretation. The linear regression of the partial residuals on x_j has the same slope, β_j , as in the full model.

If x_j affects the binary response linearly, the plot should be approximately linear with a slope approximately equal to β_j . A nonlinear plot suggests that x_j needs to be transformed, and the shape of the relation gives a rough guide to the required transformation. For example, a parabolic shape would suggest a term in x_j^2 . These plots complement the conditional data plots described earlier (Section 7.3.1), and are most useful when there several quantitative predictors, so that it is more convenient and sensible to examine their relationships individually.

The *car* package implements these plots in the `crPlots()` and `crPlot()` functions. They also work for models with factor predictors (using parallel boxplots for the factor levels) but not for those with interaction terms.

EXAMPLE 7.14: Donner Party

In Example 7.12, we fit several models for the Donner Party data, and we recall two here to illustrate component-plus-residual plots. Both assert additive effects of age and sex, but the model `donner.mod3` allows a quadratic effect of age.

```
> donner.mod1 <- glm(survived ~ age + sex, data=Donner, family=binomial)
> donner.mod3 <- glm(survived ~ poly(age,2) + sex, data=Donner, family=binomial)
```

Had we not made exploratory plots earlier (Example 7.12), and naively fit only the linear model in age, `donner.mod1`, we could use `crPlots()` to check for a non-linear relationship of survival with age as follows, giving Figure 7.32.




Figure 7.33: Component-plus-residual plot for the non-linear additive model, `donner.mod3`

{fig:donner-cr2}

$\mathbf{y} \sim \mathbf{X}^{(-i)}$ with one weighted least squares regression of x_i on $\mathbf{X}^{(-i)}$ to find the residual part, x_i^* , of x not predicted by the other regressors.

Let \mathbf{r} be the vector of Pearson residuals from the initial logistic fit of \mathbf{y} on the variables in $\mathbf{X}^{(-i)}$, and let \mathbf{H} and $\mathbf{V} = \text{diag}[\hat{p}(1 - \hat{p})]$ be the hat matrix and \mathbf{V} matrix from this analysis. Then, the added-variable plot is a scatterplot of the residuals \mathbf{r} against the x_i -residuals,

$$x_i^* = (\mathbf{I} - \mathbf{H})\mathbf{V}^{1/2}\mathbf{x}.$$

There are several important uses of added-variable plots:

First, *marginal* plots of the response variable \mathbf{y} against the predictor variables x_i can conceal or misrepresent the relationships in a model including several predictors together due to correlations or associations among the predictors. This problem is compounded by the fact that graphical methods for discrete responses (boxplots, mosaic plots) cannot easily show influential observations or non-linear relationships. Added-variable plots solve this problem by plotting the residuals, $y_i^* = y_i | \text{others}_i$, which are less discrete than the marginal responses in \mathbf{y} .

Second, the numerical measures and graphical methods for detecting influential observations described earlier in this section are based on the idea of *single-case deletion*, comparing coefficients or fitted values for the full data, with those that result from deleting each case in turn. Yet, it is well-known (Lawrance, 1995), that sets of two (or more) observations can have *joint influence*, that greatly exceeds their individual influential. Similarly, the influence of one discrepant point can be offset by another influential point in an opposite direction, a phenomenon called *masking*. The main cases of joint influence are illustrated in Figure 7.34. Added-variable plots, showing the partial regression for one predictor controlling all others can make such cases visually apparent.

Finally, given a tentative model using predictors \mathbf{x} , the added-variable plot for another regressor, z can provide a useful visual assessment of its additional contribution. An overall test could be based on the difference in G^2 for the enlarged model $\text{logit}(\mathbf{p}) = \mathbf{X}\boldsymbol{\beta} + \gamma z$, compared to the reduced model $\text{logit}(\mathbf{p}) = \mathbf{X}\boldsymbol{\beta}$. But the added-variable plot shows whether the evidence for including z is spread throughout the sample or confined to a small subset of observations. The regressor z may be a new explanatory variable, or a higher-order term for variable(s) already in the model.

The `car` package implements these plots with the function `avPlot()` for a single term and `avPlots()` for all terms in a linear or generalized linear model, as shown in the example(s) below. See <http://www.datavis.ca/gallery/animation/duncanAV/> for an animated




Figure 7.34: Jointly influential points in regression models. In each panel, the thick black line shows the regression of y on x using all the data points. The solid purple line shows the regression deleting *both* the red and blue points and the broken and dotted lines show the regression retaining only the point in its color in addition to the constant gray points. (a) Two points whose joint influence enhance each other; (b) two points where the influence of one is masked by that of the other; (c) two points whose combined influence greatly exceeds the effect of either one individually.

{fig:joint}

graphic showing the transition between a marginal plot of the relationship of y to x and the added-variable plot of y^* to x^* for the case of multiple linear regression with a quantitative response.

{ex:donner4}

EXAMPLE 7.15: Donner Party

The simple additive model `donner.mod1` for the Donner Party data can be used to illustrate some features of added-variable plots. In the call to `avPlots()` below, we use color the plotting symbol to distinguish those who survived vs. died, shape to distinguish men from women.

```
> col <- ifelse(Donner$survived=="yes", "blue", "red")
> pch <- ifelse(Donner$sex=="Male", 16, 17)
> avPlots(donner.mod1, id.n=2, col=col, pch=pch, col.lines="darkgreen")
```

These plots have the following properties:

1. The slope in the simple regression of y_i^* on x_i^* is the same as the partial coefficient β_i in the full multiple regression model including both predictors here (or all predictors in general).
2. The residuals from this regression line are the same as the residuals in the full model.
3. Because the response, `survived`, is binary, the vertical axis y_{age}^* in the left panel for `age` is the part of the logit for survival that cannot be predicted from `sex`. Similarly, the vertical axis in the right panel is the part of survival that cannot be predicted from `age`. This property allows the clusters of points corresponding to discrete variables to be seen more readily, particularly if they are distinguished by visual attributes such as color and shape, as in Figure 7.35.

△

{ex:icu3}

EXAMPLE 7.16: Death in the ICU

We illustrate some of the uses of added-variable plots using the main effects model, `icu.glm2`, predicting death in the ICU from the variables `age`, `cancer`, `admit` and `uncons`.

To see why marginal plots of the discrete response against each predictor are often unrevealing for the purpose of model assessment, consider the collection of plots in Figure 7.36 showing the default plots (spineplots) for the factor response, `died` against each predictor. These show the marginal distribution of each predictor by the widths of the bars, and highlight the proportion who




Figure 7.35: Added-variable plots for age (left) and sex (right) in the Donner Party main effects model. Those who survived are shown in blue; those who died in red. Men are plotted with filled circles; women with filled triangles.

{fig:donner4-avp}

died by color. Such plots are useful for some purposes, but not for assessing the adequacy of the fitted model.

```
> op <- par(mfrow=c(2,2), mar=c(4,4,1,2.5)+.1, cex.lab=1.4)
> plot(died ~ age, data=ICU, col=c("lightblue", "red"))
> plot(died ~ cancer, data=ICU, col=c("lightblue", "red"))
> plot(died ~ admit, data=ICU, col=c("lightblue", "red"))
> plot(died ~ uncons, data=ICU, col=c("lightblue", "red"))
> par(op)
```

The added-variable plot for this model is shown in Figure 7.37. In each plot, the solid red line shows the partial slope, β_j for the focal predictor, controlling for all others.

```
> pch <- ifelse(ICU$died=="No", 1, 2)
> avPlots(icu.glm2, id.n=2, pch=pch, cex.lab=1.3)
```

The labeled points in each panel use the default `id.method` for `avPlots()`, selecting those with either large absolute model residuals or extreme x_i^* residuals, given all other predictors. Cases 127 and 881, identified earlier as influential stand out in all these plots.

Next, we illustrate the use of added-variable plots for checking the effect of influential observations on the decision to include an additional predictor in some given model. In the analysis of the *ICU* data using model selection methods, the variable `systolic` (systolic blood pressure at admission) was nominated by several different procedures. Here we take a closer look at the evidence for inclusion of this variable in a predictive model. We fit a new model adding `systolic` to the others and test the improvement with a likelihood ratio test:

```
> icu.glm2a <- glm(died ~ age + cancer + admit + uncons + systolic,
+                 data=ICU, family=binomial)
> anova(icu.glm2, icu.glm2a, test="Chisq")
```




Figure 7.36: Marginal plots of the response `died` against each of the predictors in the model `icu.glm2` for the *ICU* data

Analysis of Deviance Table

```
Model 1: died ~ age + cancer + admit + uncons
Model 2: died ~ age + cancer + admit + uncons + systolic
  Resid. Df Resid. Dev Df Deviance Pr(>Chi)
1      195      139
2      194      136  1      3.52    0.061 .
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

So, the addition of systolic blood pressure is nearly significant at the conventional $\alpha = 0.05$ level. The added-variable plot for this variable in Figure 7.38 shows the strength of evidence for its contribution, above and beyond the other variables in the model, as well as the partial leverage and influence of individual points.

```
> avPlot(icu.glm2a, "systolic", id.n=3, pch=pch)
```

In this plot, cases 331 and 921 have high partial leverage, but they are not influential. Case 84, however, has high leverage and a large residual, so is possibly influential on the evidence for inclusion of `systolic` in the model. Note also that the partial regression line in this plot nicely separates nearly all the patients who died from those who survived.

△

TODO: This chapter is quite long. Could start a new chapter here, but that would require re-numbering file names, or introduce Ch07a.

7.6 Polytomous response models

Polytomous response data arise when the outcome variable, Y , takes on $m > 2$ discrete values. For example, (a) patients may record that their improvement after treatment is “none,” “some” or

{sec:logist-poly}




Figure 7.37: Added-variable plots for the predictors in the model for the ICU data. Those who died and survived are shown by triangles (\triangle) and circles (\circ) respectively.

{fig:icu3-avp1}

“marked;” (b) high school students may choose a general, vocational or academic program; (c) women’s labor force participation may be recorded in a survey as not working outside the home, working part-time, or working full-time; (d) Canadian voters may express a preference for the Conservative, Liberal, NDP, Green party. These response categories may be considered ordered or simply nominal.

In this situation, there are several different ways to model the response probabilities. Let $\pi_{ij} \equiv \pi_j(x_i)$ be the probability of response j for case or group i , given the predictors x_i . Because $\sum_j \pi_{ij} = 1$, only $m - 1$ of these probabilities are independent. The essential idea here is to construct a model for the polytomous (or multinomial) response composed of $m - 1$ logit comparisons among the response categories in a similar way to how factors are treated in the predictor variables.

The simplest approach uses the *proportional odds model*, described in Section 7.6.1. This model applies only when the response is ordinal (as in improvement after therapy) *and* an additional assumption (the proportional odds assumption) holds. This model can be fit using `polr()` in the MASS package, `lrm()` in the rms package, and `vglm()` in VGAM.

However, if the response is purely nominal (e.g., vote Conservative, Liberal, NDP, Green), or if the proportional odds assumption is untenable, another particularly simple strategy is to fit separate models to a set of $m - 1$ *nested dichotomies* derived from the polytomous response (described in Section 7.6.3). This method allows you to resolve the differences among the m response categories into independent statistical questions (similar to orthogonal contrasts in ANOVA). For example, for




Figure 7.38: added-variable plot for the effect of adding systolic blood pressure to the main effects model for the ICU data.

{fig:icu3-avp2}

women's labor force participation, it might be substantively interesting to contrast not working vs. (part-time and full-time) and then part-time vs. full-time for women who are working. You fit such nested dichotomies by running the $m - 1$ binary logit models and combining the statistical results.

The most general approach is the **generalized logit model**, also called the **multinomial logit model**. This model fits *simultaneously* the $m - 1$ simple logit models against a baseline or reference category, for example, the last category, m . With a 3-category response, there are two generalized logits, $L_{i1} = \log(\pi_{i1}/\pi_{i3})$ and $L_{i2} = \log(\pi_{i2}/\pi_{i3})$, contrasting response categories 1 and 2 against category 3. In this approach, it doesn't matter which response category is chosen as the baseline, because all pairwise comparisons can be recovered from whatever is estimated. This model is conveniently fitted using `multinom()` in `nnet`.

7.6.1 Ordinal Response: Proportional Odds Model

{sec:ordinal}

For an ordered response Y , with categories $j = 1, 2, \dots, m$, the ordinal nature of the response can be taken into account by forming logits based on the $m - 1$ adjacent category cutpoints between successive categories. That is, if the cumulative probabilities are

$$\Pr(Y \leq j | \mathbf{x}) = \pi_1(\mathbf{x}) + \pi_2(\mathbf{x}) + \dots + \pi_j(\mathbf{x}) ,$$

then the **cumulative logit** for category j is defined as

$$L_j \equiv \text{logit}[\Pr(Y \leq j | \mathbf{x})] = \log \frac{\Pr(Y \leq j | \mathbf{x})}{\Pr(Y > j | \mathbf{x})} = \log \frac{\Pr(Y \leq j | \mathbf{x})}{1 - \Pr(Y \leq j | \mathbf{x})} \quad (7.13) \quad \text{{eq:cumlogit}}$$

for $j = 1, 2, \dots, m - 1$.

In our running example of responses to arthritis treatment, the actual response variable is `Improved`, with ordered levels "None" < "Some" < "Marked". In this case, the cumulative logits would




Figure 7.39: Proportional odds model for an ordinal response. The model assumes equal slopes for the cumulative response logits. Left: logit scale; right: probability scale.

{fig:podds}

be defined as

$$L_1 = \log \frac{\pi_1(\mathbf{x})}{\pi_2(\mathbf{x}) + \pi_3(\mathbf{x})} = \text{logit}(\text{None vs. [Some or Marked]})$$

$$L_2 = \log \frac{\pi_1(\mathbf{x}) + \pi_2(\mathbf{x})}{\pi_3(\mathbf{x})} = \text{logit}(\text{[None or Some] vs. Marked}),$$

where \mathbf{x} represents the predictors (sex, treatment and age).

The **proportional odds model** (PO) (McCullagh, 1980) proposes a simple and parsimonious account of these effects, where the predictors in (\mathbf{x}) are constrained to have the same slopes for all cumulative logits,

{eq:propodds}

$$L_j = \alpha_j + \mathbf{x}^T \boldsymbol{\beta} \quad j = 1, \dots, m-1. \quad (7.14)$$

That is, the effect of the predictor x_i is the same, β_i , for all cumulative logits. The cumulative logits differ only in their intercepts. In this formulation, the $\{\alpha_j\}$ increase with j , because $\Pr(Y \leq j | \mathbf{x})$ increases with j for fixed \mathbf{x} .²² Figure 7.39 portrays the PO model for a single quantitative predictor x with $m = 4$ response categories.

The name “proportional odds” stems from the fact that under Eqn. (7.14), for fixed \mathbf{x} , the cumulative log odds (logits) for categories j and j' are constant, $(\alpha_j - \alpha_{j'})$, so the odds, $\exp(\alpha_j - \alpha_{j'})$ have a constant ratio, or are proportional. Similarly, the ratio of the cumulative odds of making a response $Y \leq j$ at values of the predictors $\mathbf{x} = \mathbf{x}_1$ are $\exp((\mathbf{x}_1 - \mathbf{x}_2)^T \boldsymbol{\beta})$ times the odds of this response at $\mathbf{x} = \mathbf{x}_2$, so the log cumulative odds ratio is proportional to the difference between \mathbf{x}_1 and \mathbf{x}_2 .

7.6.1.1 Latent variable interpretation

For a binary response, an alternative motivation for logistic regression regards the relation of the observed Y as arising from a continuous, unobserved, (latent) response variable, ξ representing the propensity for a “success” (1) rather than “failure” (0). The latent response is assumed to be linearly related to the predictors \mathbf{x} according to

{eq:latent}

$$\xi_i = \alpha + \mathbf{x}_i^T \boldsymbol{\beta} + \epsilon_i = \alpha + \beta_1 x_{i1} + \dots + \beta_p x_{ip} + \epsilon_i \quad (7.15)$$

²²Some authors and some software describe the PO model in terms of $\text{logit}[\Pr(Y > j | \mathbf{x})]$, so the signs and order of the intercepts, α_j are reversed.




Figure 7.40: Latent variable representation of the proportional odds model for $m = 4$ response categories and a single quantitative predictor, x . *Source:* Adapted from Fox (2008, Fig 14.10), using code provided by John Fox.

{fig:latent}

However, we can only observe $Y_i = 1$ when ξ_i passes some threshold, that with some convenient scaling can be taken as $\xi_i > 0 \implies Y_i = 1$.²³

The latent variable motivation extends directly to an ordinal response under the PO model. We now assume that there is a set of $m - 1$ thresholds, $\alpha_1 < \alpha_2 < \dots < \alpha_{m-1}$ for the latent variable ξ_i in Eqn. (7.15) and we observe

$$Y_i = j \quad \text{if} \quad \alpha_{j-1} < \xi_i \leq \alpha_j ,$$

with appropriate modifications to the inequalities at the end points.

This is illustrated in Figure 7.40 for a response with $m = 4$ ordered categories and a single quantitative predictor, x . The observable response Y categories are shown on the right vertical axis, and the corresponding latent continuous variable ξ on the left axis together with the thresholds $\alpha_1, \alpha_2, \alpha_3$. The (conditional) logistic distribution of ξ is shown at two values of x , and the shaded areas under the curve give the conditional probabilities $\Pr(Y = 4 | x_i)$ for the two values x_1 and x_2 .

7.6.1.2 Fitting the proportional odds model

As mentioned earlier, there are a number of different R packages that provide facilities for fitting the PO model. These have somewhat different capabilities for reporting results, testing hypotheses and plotting, so we generally use `polr()` in the MASS package, except where other packages offer greater convenience.

Unless the response variable has numeric values, it is important to ensure that it has been defined

²³The latent variable derivation of logistic regression (and the related probit model) was fundamental in the history of statistical methods for discrete response outcomes. An early example was Thurstone's (1927) *Law of comparative judgment* designed to account for psychological preference by assuming an underlying latent continuum of "hedonic values." Similarly, the probit model arose from dose-response studies in toxicology (Bliss, 1934, Finney, 1947) where the number killed by some chemical agent was related to its' type, dose or concentration. The idea of a latent variable was also at the heart of the development of factor analysis **TODO: citation?** and latent class analysis (Lazarsfeld, 1950, 1954) was developed to treat the problem of classifying individuals into discrete latent classes from fallible measurements. See Bollen (2002) for a useful overview of latent variable models in the social sciences.

as an *ordered* factor (using `ordered()`). In the *Arthritis* data, the response, Improved was setup this way, as we can check by printing some of the values.²⁴

```
> head(Arthritis$Improved, 8)

[1] Some    None    None    Marked Marked Marked None    Marked
Levels: None < Some < Marked
```

We fit the main effects model for the ordinal response using `polr()` as shown below. We also specify `Hess=TRUE` to have the function return the observed information matrix (called the Hessian), that is used in other operations to calculate standard errors.

```
> arth.polr <- polr(Improved ~ Sex + Treatment + Age,
+                  data=Arthritis, Hess=TRUE)
> summary(arth.polr)

Call:
polr(formula = Improved ~ Sex + Treatment + Age, data = Arthritis,
      Hess = TRUE)

Coefficients:
              Value Std. Error t value
SexMale      -1.2517   0.5464  -2.29
TreatmentTreated  1.7453   0.4759   3.67
Age           0.0382   0.0184   2.07

Intercepts:
              Value Std. Error t value
None|Some     2.532   1.057    2.395
Some|Marked   3.431   1.091    3.144

Residual Deviance: 145.46
AIC: 155.46
```

The output from the `summary()` method, shown above, gives the estimated coefficients (β) and intercepts (α_j) labeled by the cutpoint on the ordinal response. It provides standard errors and *t*-values ($\beta_i/SE(\beta_i)$), but no significance tests or *p*-values.

```
> library(car)
> Anova(arth.polr)

Analysis of Deviance Table (Type II tests)

Response: Improved
              LR Chisq Df Pr(>Chisq)
Sex              5.69  1  0.01708 *
Treatment       14.71  1  0.00013 ***
Age              4.57  1  0.03251 *
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

7.6.1.3 Testing the proportional odds assumption

The simplicity of the PO model is achieved only when the proportional odds model holds for a given data set. In essence, a test of this assumption involves a contrast between the PO model and a generalized logit NPO model that allows different effects (slopes) of the predictors across the

²⁴As an unordered factor, the levels would be treated as ordered alphabetically, i.e., Marked, None, Some.

response categories:

$$\text{(eq:po)} \quad \text{PO: } L_j = \alpha_j + \mathbf{x}^T \boldsymbol{\beta} \quad j = 1, \dots, m-1 \quad (7.16)$$

$$\text{(eq:npo)} \quad \text{NPO: } L_j = \alpha_j + \mathbf{x}^T \boldsymbol{\beta}_j \quad j = 1, \dots, m-1 \quad (7.17)$$

The most general test involves fitting both models and testing the difference in the residual deviance by a likelihood ratio test or using some other measure (such as AIC) for model comparison. The PO model (Eqn. (7.16)) has $(m-1) + p$ parameters, while the NPO model (Eqn. (7.17)) has $(m-1)(1+p) = m(1+p)$ parameters, which may be difficult to fit if this is large relative to the number of observations. An intermediate model, the **partial proportional odds model** (Peterson and Harrell, 1990) allows one subset of predictors, \mathbf{x}_{po} , to satisfy the proportional odds assumption (equal slopes), while the remaining predictors \mathbf{x}_{npo} have slopes varying with the response level:

$$\text{(eq:ppo)} \quad \text{PPO: } L_j = \alpha_j + \mathbf{x}_{po}^T \boldsymbol{\beta} + \mathbf{x}_{npo}^T \boldsymbol{\beta}_j \quad j = 1, \dots, m-1 . \quad (7.18)$$

In R, the PO and NPO models can be readily contrasted by fitting them both using `vglm()` in the **VGAM** package. This defines the cumulative family of models and allows a `parallel` option. With `parallel=TRUE`, this is equivalent to the `polr()` model, except that the signs of the coefficients are reversed.

```
> library(VGAM)
> arth.po <- vglm(Improved ~ Sex + Treatment + Age, data=Arthritis,
+               family = cumulative(parallel=TRUE))
> arth.po

Call:
vglm(formula = Improved ~ Sex + Treatment + Age, family = cumulative(parallel = TRUE),
     data = Arthritis)

Coefficients:
  (Intercept):1      (Intercept):2      SexMale
    2.531990         3.430988         1.251671
TreatmentTreated      Age
   -1.745304      -0.038163

Degrees of Freedom: 168 Total; 163 Residual
Residual deviance: 145.46
Log-likelihood: -72.729
```

The more general NPO model can be fit using `parallel=FALSE`.

```
> arth.npo <- vglm(Improved ~ Sex + Treatment + Age, data=Arthritis,
+               family = cumulative(parallel=FALSE))
> arth.npo

Call:
vglm(formula = Improved ~ Sex + Treatment + Age, family = cumulative(parallel = FALSE),
     data = Arthritis)

Coefficients:
  (Intercept):1      (Intercept):2      SexMale:1
    2.618539         3.431175         1.509827
SexMale:2 TreatmentTreated:1 TreatmentTreated:2
    0.866434      -1.836929      -1.704011
      Age:1          Age:2
   -0.040866      -0.037294

Degrees of Freedom: 168 Total; 160 Residual
Residual deviance: 143.57
Log-likelihood: -71.787
```

The VGAM package defines a `coef()` method that can print the coefficients in a more readable matrix form giving the category cutpoints:

```
> coef(arth.po, matrix=TRUE)

              logit(P[Y<=1]) logit(P[Y<=2])
(Intercept)      2.531990      3.430988
SexMale          1.251671      1.251671
TreatmentTreated -1.745304     -1.745304
Age              -0.038163     -0.038163

> coef(arth.npo, matrix=TRUE)

              logit(P[Y<=1]) logit(P[Y<=2])
(Intercept)      2.618539      3.431175
SexMale          1.509827      0.866434
TreatmentTreated -1.836929     -1.704011
Age              -0.040866     -0.037294
```

In most cases, nested models can be tested using an `anova()` method, but the VGAM package has not implemented this for "vglm" objects. Instead, it provides an analogous function, `lrtest()`:

```
> VGAM::lrtest(arth.npo, arth.po)

Likelihood ratio test

Model 1: Improved ~ Sex + Treatment + Age
Model 2: Improved ~ Sex + Treatment + Age
#Df LogLik Df Chisq Pr(>Chisq)
1 160 -71.8
2 163 -72.7 3 1.88 0.6
```

The LR test can be also calculated as “manually” shown below using the difference in residual deviance for the two models.

```
> tab <- cbind(
+   Deviance = c(deviance(arth.npo), deviance(arth.po)),
+   df = c(df.residual(arth.npo), df.residual(arth.po))
+ )
> tab <- rbind(tab, diff(tab))
> rownames(tab) <- c("GenLogit", "PropOdds", "LR test")
> tab <- cbind(tab, pvalue=1-pchisq(tab[,1], tab[,2]))
> tab

              Deviance  df  pvalue
GenLogit 143.5741 160 0.81966
PropOdds 145.4579 163 0.83435
LR test   1.8838   3 0.59686
```

The `vglm()` can also fit partial proportional odds models, by specifying a formula giving the terms for which the PO assumption should be taken as TRUE or FALSE. Here we illustrate this using `parallel=FALSE ~ Sex`, to fit separate slopes for males and females, but parallel lines for the other predictors. The same model would be fit using `parallel=TRUE ~ Treatment + Age`.

```
> arth.ppo <- vglm(Improved ~ Sex + Treatment + Age, data=Arthritis,
+   family = cumulative(parallel=FALSE ~ Sex))
> coef(arth.ppo, matrix=TRUE)

              logit(P[Y<=1]) logit(P[Y<=2])
```

(Intercept)	2.542452	3.615561
SexMale	1.483336	0.867362
TreatmentTreated	-1.775742	-1.775742
Age	-0.039622	-0.039622

7.6.1.4 Graphical assessment of proportional odds

There are several graphical methods for visual assessment of the proportional odds assumption. These are all *marginal* methods, in that they treat the predictors one at a time. However, that provides one means to determine if a partial proportional odds model might be more appropriate. Harrell's 2001, Ch. 13-14 *Regression Modeling Strategies* and the corresponding `rms` package provide an authoritative treatment and methods in R.

One simple idea is to plot the conditional mean or expected value $E(X | Y)$ of a given predictor, X , at each level of the ordered response Y . If the response behaves ordinaly in relation to X , these means should be strictly increasing or decreasing with Y . For comparison, one can also plot the estimated conditional means $\hat{E}(X | Y = j)$ under the fitted PO model X as the only predictor. If the PO assumption holds for this X , the model-mean curve should be close to the data mean curve.

```
> library(rms)
> arth.po2 <- lrm(Improved ~ Sex + Treatment + Age, data=Arthritis)
> arth.po2
```

Logistic Regression Model

```
lrm(formula = Improved ~ Sex + Treatment + Age, data = Arthritis)
```

		Model Likelihood		Discrimination		Rank Discrim.	
		Ratio Test		Indexes		Indexes	
Obs	84	LR chi2	24.46	R2	0.291	C	0.750
None	42	d.f.	3	g	1.335	Dxy	0.500
Some	14	Pr(> chi2)	<0.0001	gr	3.801	gamma	0.503
Marked	28			gp	0.280	tau-a	0.309
max deriv	1e-07			Brier	0.187		

	Coef	S.E.	Wald Z	Pr(> Z)
y>=Some	-2.5320	1.0570	-2.40	0.0166
y>=Marked	-3.4310	1.0911	-3.14	0.0017
Sex=Male	-1.2517	0.5464	-2.29	0.0220
Treatment=Treated	1.7453	0.4759	3.67	0.0002
Age	0.0382	0.0184	2.07	0.0382

The plot of conditional X means is produced using the `plot.xmean.ordinaly()` as shown below. It produces one marginal panel for each predictor in the model. For categorical predictors, it plots only the overall most frequent category. The resulting plot is shown in Figure 7.41.

```
> op <- par(mfrow=c(1,3))
> plot.xmean.ordinaly(Improved ~ Sex + Treatment + Age, data=Arthritis,
+                       lwd=2, pch=16, subn=FALSE)
> par(op)
```

In Figure 7.41, there is some evidence that the effect of `Sex` is non-monotonic and the means differ from their model-implied values under the PO assumption. The effect of `Treatment` looks good by this method, and the effect of `Age` hints that the upper two categories may not be well-distinguished as an ordinal response.

Of course, this example has only a modest total sample size, and this method only examines the marginal effects of the predictors. Nevertheless, it is a useful supplement to the statistical methods described earlier.




Figure 7.41: Visual assessment ordinality and the proportional odds assumption for predictors in the Arthritis data. Solid lines connect the stratified means of X given Y . Dashed lines show the estimated expected value of X given $Y=j$ if the proportional odds model holds for X .

{fig:arth-rmsplot}

7.6.2 Visualizing results for the proportional odds model

{sec:vis-propodds}

Results from the PO model (and other models for polytomous responses) can be graphed using the same ideas and methods shown earlier for a binary or binomial response. In particular, full-model plots (described earlier in Section 7.3.2) and effect plots (Section 7.3.3) are still very helpful.

But now there is the additional complication that the response variable has $m > 2$ levels and so needs to be represented by $m - 1$ curves or panels in addition to those related to the predictor variables.

7.6.2.1 Full-model plots

{sec:po-fullplots}

For full-model plots, we continue the idea of appending the fitted response probabilities (or log-its) to the data frame and plotting these in relation to the predictors. The `predict()` method returns the highest probability category label by default (with `type="class"`), so to get the fitted probabilities you have to ask for `type="probs"`, as shown below.

```
> arth.fitp <- cbind(Arthritis,
+                   predict(arth.polr, type="probs"))
> head(arth.fitp)
```

	ID	Treatment	Sex	Age	Improved	Better	None	Some	Marked
1	57	Treated	Male	27	Some	1	0.73262	0.13806	0.12932
2	46	Treated	Male	29	None	0	0.71740	0.14443	0.13816
3	77	Treated	Male	30	None	0	0.70960	0.14763	0.14277
4	17	Treated	Male	32	Marked	1	0.69363	0.15400	0.15237
5	36	Treated	Male	46	Marked	1	0.57025	0.19504	0.23471
6	23	Treated	Male	58	Marked	1	0.45634	0.21713	0.32653

For plotting, it is most convenient to reshape these from wide to long format using `melt()` in the `reshape2` package. The response category is named `Level`.

```
> library(reshape2)
> plotdat <- melt(arth.fitp,
+               id.vars = c("Sex", "Treatment", "Age", "Improved"),
+               measure.vars=c("None", "Some", "Marked"),
+               variable.name = "Level",
+               value.name = "Probability")
> ## view first few rows
> head(plotdat)
```

	Sex	Treatment	Age	Improved	Level	Probability
1	Male	Treated	27	Some	None	0.73262
2	Male	Treated	29	None	None	0.71740
3	Male	Treated	30	None	None	0.70960
4	Male	Treated	32	Marked	None	0.69363
5	Male	Treated	46	Marked	None	0.57025
6	Male	Treated	58	Marked	None	0.45634

We can now plot Probability against Age, using Level to assign different colors to the lines for the response categories. `facet_grid()` is used to split the plot into separate panels by Sex and Treatment. In this example, the `directlabels` package is also used replace the default legend created by `ggplot()` with category labels on the curves themselves, which is easier to read.

```
> library(ggplot2)
> library(directlabels)
> gg <- ggplot(plotdat, aes(x = Age, y = Probability, colour = Level)) +
+   geom_line(size=2.5) + theme_bw() + xlim(10,80) +
+   geom_point(color="black", size=1.5) +
+   facet_grid(Sex ~ Treatment,
+             labeller = function(x, y) sprintf("%s = %s", x, y)
+             )
> direct.label(gg)
```




Figure 7.42: Predicted probabilities for the proportional odds model fit to the Arthritis data

{fig:arth-polr1}

Although we now have three response curves in each panel, this plot is relatively easy to understand: (a) In each panel, the probability of no improvement decreases with age, while that for marked improvement increases. (b) It is easy to compare the placebo and treated groups in each row, showing that no improvement decreases, while marked improvement increases with the active treatment. (On the other hand, this layout makes it harder to compare panels vertically for males and females in each condition.) (c) The points show where the observations are located in each panel; so, we can see that the data is quite thin for males given the placebo.²⁵

7.6.2.2 Effect plots

For PO models fit using `polr()`, the **effects** package provides two different styles for plotting a given effect. By default, curves are plotted in separate panels for the different response levels of a given effect, together with confidence bands for predicted probabilities. This form provides confidence bands and rug plots for the observations, but the default vertical arrangement of the panels makes it harder to compare the trends for the different response levels. The alternative *stacked* format shows the changes in response level more directly, but doesn't provide confidence bands.

Figure 7.43 shows these two styles for the main effect of Age in the proportional odds model, `arth.polr` fit earlier.

```
> plot(Effect("Age", arth.polr))
> plot(Effect("Age", arth.polr), style='stacked',
+       key.args=list(x=.55, y=.9))
```




Figure 7.43: Effect plots for the effect of Age in the proportional odds model for the Arthritis data. Left: responses shown in separate panels. Right: responses shown in stacked format

Even though this model includes only main effects, you can still plot the higher-order effects

²⁵One way to improve (pun intended) this graph would be to show the points on the lines only for the actual level of Improve for each observation.

for more focal predictors in a coherent display. Figure 7.44 shows the predicted probabilities for all three predictors together. Again, visual comparison is easier horizontally for placebo versus treated groups, but you can also see that the prevalence of marked improvement is greater for females than for males.

```
> plot(Effect(c("Treatment", "Sex", "Age"), arth.polr),
+       style="stacked", key.arg=list(x=.8, y=.9))
```




Figure 7.44: Effect plot for the effects of Treatment, Sex and Age in the Arthritis data.

{fig:arth-po-eff2}

Finally, the latent variable interpretation of the PO model provides for simpler plots on the logit scale. Figure 7.45 shows this plot for the effects of Treatment and Age (collapsed over Sex) produced with the argument `latent=TRUE` to `Effect()`. In this plot, there is a single line in each panel for the effect (slope) of Age on the log odds. The dashed horizontal lines give the thresholds between the adjacent response categories corresponding to the intercepts.

```
> plot(Effect(c("Treatment", "Age"), arth.polr, latent=TRUE), lwd=3)
```

7.6.3 Nested dichotomies

{sec:nested}

The method of *nested dichotomies* provides another simple way to analyse a polytomous response in the framework of logistic regression (or other generalized linear models). This method does not require an ordinal response or special software. Instead, it uses the familiar binary logistic model and fits $m - 1$ separate models for each of a hierarchically nested set of comparisons among the response categories.

Taken together, this set of models for the dichotomies comprises a complete model for the polytomous response. As well, these models are statistically independent, so test statistics such as G^2 or Wald tests can be added to give overall tests for the full polytomy.




Figure 7.45: Latent variable effect plot for the effects of Treatment and Age in the Arthritis data.

{fig:arth-po-eff3}

For example, the response categories $Y = \{1,2,3,4\}$ could be divided first as $\{1,2\}$ vs. $\{3,4\}$, as shown in the left side of Figure 7.46. Then these two dichotomies could be divided as $\{1\}$ vs. $\{2\}$, and $\{3\}$ vs. $\{4\}$. Alternatively, these response categories could be divided as shown in the right side of Figure 7.46: first, $\{1\}$ vs. $\{2,3,4\}$, then $\{2\}$ vs. $\{3,4\}$, and finally $\{3\}$ vs. $\{4\}$.




Figure 7.46: Nested dichotomies. The boxes show two different ways a four-category response can be represented as three nested dichotomies. Adapted from Fox (2008).


{fig:nested2}

Such models make the most sense when there are substantive reasons for considering the response categories in terms of such dichotomies. Two examples are shown in Figure 7.47.

- For the *Arthritis* data, it is sensible to consider one dichotomy (“better”), with logit L_1 , between the categories of “None” compared to “Some” or “Marked”. A second dichotomy, with logit L_2 , would then distinguish between the some and marked response categories.
- For a second case where patients are classified into $m = 4$ psychiatric diagnostic categories, the first dichotomy, with logit L_1 distinguishes those considered normal from all others given a clinical diagnosis. Two other dichotomies are defined to further divide the non-normal categories.

Then, consider the separate logit models for these $m - 1$ dichotomies, with different intercepts α_j and slopes β_j for each dichotomy,

$$L_1 = \alpha_1 + \mathbf{x}^\top \beta_1$$



{fig:nested1}

Figure 7.47: Examples of nested dichotomies and the corresponding logits

$$\begin{aligned}
 L_2 &= \alpha_2 + \mathbf{x}^\top \beta_2 \\
 \vdots &= \vdots \\
 L_{m-1} &= \alpha_{m-1} + \mathbf{x}^\top \beta_{m-1}
 \end{aligned}$$

{ex:wlfpart1}

EXAMPLE 7.17: Women's labor force participation

The data set *Womenlwf* in the *car* package gives the result of a 1977 Canadian survey. It contains data for 263 married women of age 21–30 who indicated their working status (outside the home) as not working, working part time or working full time, together with their husband's income and a binary indicator of whether they had one or more young children in their household. (Another variable, region of Canada, had no effects in these analyses, and is not examined here.) This example follows Fox and Weisberg (2011, §5.8).

```

> library(car) # for data and Anova()
> data("Womenlwf", package="car")
> some(Womenlwf)

   partic hincome children  region
6   not.work      7  present Ontario
25  not.work     23  present Ontario
36  not.work     19  absent  Ontario
83  fulltime     17  present Ontario
138 not.work     13  present Ontario
166 not.work      9  present Atlantic
168 fulltime     13  absent  Ontario
173 not.work      7  present Ontario
229 parttime     23  present  Quebec
233 fulltime     15  absent  Quebec

```

In this example, it makes sense to consider a first dichotomy (*working*) between women who are not working, vs. those who are (full time or part time). A second dichotomy (*fulltime*) contrasts full time work vs. part time work, among those women who are working at least part time. These two binary variables are created in the data frame using the `recode()` function from the *car* package.

```
> # create dichotomies
> Womenlf <- within(Womenlf, {
+   working <- recode(partic, " 'not.work' = 'no'; else = 'yes' ")
+   fulltime <- recode(partic,
+     " 'fulltime' = 'yes'; 'parttime' = 'no'; 'not.work' = NA" )})
> some(Womenlf)
```

	partic	hincome	children	region	fulltime	working
1	not.work	15	present	Ontario	<NA>	no
10	not.work	23	present	Ontario	<NA>	no
11	not.work	23	present	Ontario	<NA>	no
24	fulltime	11	absent	Ontario	yes	yes
67	not.work	15	present	Ontario	<NA>	no
98	fulltime	15	absent	Ontario	yes	yes
122	not.work	23	present	Atlantic	<NA>	no
167	not.work	15	present	Ontario	<NA>	no
208	fulltime	11	absent	Quebec	yes	yes
241	not.work	13	present	Quebec	<NA>	no

The tables below show how the response `partic` relates to the recoded binary variables, `working` and `fulltime`. Note that the `fulltime` variable is recoded to `NA` for women who are not working.

```
> with(Womenlf, table(partic, working))
```

	working	
partic	no	yes
fulltime	0	66
not.work	155	0
parttime	0	42

```
> with(Womenlf, table(partic, fulltime, useNA="ifany"))
```

	fulltime		
partic	no	yes	<NA>
fulltime	0	66	0
not.work	0	0	155
parttime	42	0	0

We proceed to fit two separate binary logistic regression models for the derived dichotomous variables. For the `working` dichotomy, we get the following results:

```
> mod.working <- glm(working ~ hincome + children, family=binomial,
+   data=Womenlf)
> summary(mod.working)
```

Call:

```
glm(formula = working ~ hincome + children, family = binomial,
    data = Womenlf)
```

Deviance Residuals:

Min	1Q	Median	3Q	Max
-1.677	-0.865	-0.777	0.929	1.997

Coefficients:

	Estimate	Std. Error	z value	Pr(> z)
(Intercept)	1.3358	0.3838	3.48	0.0005 ***
hincome	-0.0423	0.0198	-2.14	0.0324 *
childrenpresent	-1.5756	0.2923	-5.39	7e-08 ***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```
(Dispersion parameter for binomial family taken to be 1)
```

```
Null deviance: 356.15 on 262 degrees of freedom
Residual deviance: 319.73 on 260 degrees of freedom
AIC: 325.7
```

```
Number of Fisher Scoring iterations: 4
```

And, similarly for the fulltime dichotomy:

```
> mod.fulltime <- glm(fulltime ~ hincome + children, family=binomial,
+                      data=Womenlwf)
> summary(mod.fulltime)
```

Call:

```
glm(formula = fulltime ~ hincome + children, family = binomial,
    data = Womenlwf)
```

Deviance Residuals:

```
Min      1Q  Median      3Q      Max
-2.405  -0.868   0.395   0.621   1.764
```

Coefficients:

```
Estimate Std. Error z value Pr(>|z|)
(Intercept) 3.4778 0.7671 4.53 5.8e-06 ***
hincome -0.1073 0.0392 -2.74 0.0061 **
childrenpresent -2.6515 0.5411 -4.90 9.6e-07 ***
---
```

```
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
(Dispersion parameter for binomial family taken to be 1)
```

```
Null deviance: 144.34 on 107 degrees of freedom
Residual deviance: 104.49 on 105 degrees of freedom
(155 observations deleted due to missingness)
AIC: 110.5
```

```
Number of Fisher Scoring iterations: 5
```

Although these were fit separately, we can view this as a combined model for the three-level response, with the following coefficients:

```
> cbind(working=coef(mod.working), fulltime=coef(mod.fulltime))
```

```
      working fulltime
(Intercept)  1.335830  3.47777
hincome      -0.042308 -0.10727
childrenpresent -1.575648 -2.65146
```

Writing these out as equations for the logits, we have:

$$L_1 = \log \frac{\Pr(\text{working})}{\Pr(\text{notworking})} = 1.336 - 0.042 \text{ hincome} - 1.576 \text{ children} \quad (7.19) \quad \{\text{eq:wlf-logits}\}$$

$$L_2 = \log \frac{\Pr(\text{fulltime})}{\Pr(\text{parttime})} = 3.478 - 0.1072 \text{ hincome} - 2.652 \text{ children} \quad (7.20)$$

For both dichotomies, increasing income of the husband and the presence of young children decrease the log odds of a greater level of work. However, for those women who are working the effects of husband's income and children are greater on the choice between full time and part time work than they are for all women on the choice between working and not working.

As we mentioned above, the use of nested dichotomies implies that the models fit to the separate dichotomies are statistically independent. Thus, we can additively combine χ^2 statistics and degrees of freedom to give overall tests for the polytomous response.

For example, here we define a function, `LRtest()` to calculate the likelihood ratio test of the hypothesis $H_0 : \beta = 0$ for all predictors simultaneously. We then use this to display these tests for each sub-model, as well as the combined model based on the sums of the test statistic and degrees of freedom.

```
> LRtest <- function(model)
+   c(LRchisq=(model$null.deviance - model$deviance),
+     df=(model$df.null - model$df.residual))
> tab <- rbind(working=LRtest(mod.working),
+             fulltime=LRtest(mod.fulltime))
> tab <- rbind(tab, All = colSums(tab))
> tab <- cbind(tab, pvalue = 1- pchisq(tab[,1], tab[,2]))
> tab
```

	LRchisq	df	pvalue
working	36.418	2	1.2355e-08
fulltime	39.847	2	2.2252e-09
All	76.265	4	1.1102e-15

Similarly, you can carry out tests of individual predictors, $H_0 : \beta_i = 0$ for the polytomy by adding the separate χ^2 s from `Anova()`.

```
> Anova(mod.working)

Analysis of Deviance Table (Type II tests)

Response: working
      LR Chisq Df Pr(>Chisq)
hincome   4.82637  1  0.028028 *
children 31.32288  1 2.1849e-08 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

> Anova(mod.fulltime)

Analysis of Deviance Table (Type II tests)

Response: fulltime
      LR Chisq Df Pr(>Chisq)
hincome   8.9813  1  0.0027275 **
children 32.1363  1 1.4373e-08 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

For example, the test for husband's income gives $\chi^2 = 4.826 + 8.981 = 13.807$ with 2 df.

As before, you can plot the fitted values from such models, either on the logit scale (for the separate logit equations) or in terms of probabilities for the various responses. The general idea is the same: obtain the fitted values from `predict()` using data frame containing the values of the predictors. However, now we have to combine these for each of the sub-models.

We calculate these values below, on both the logit scale and the response scale of probabilities. The `newdata` argument to `predict()` is constructed as the combinations of values for `hincome` and `children`.²⁶

²⁶Alternatively, using the predictor values in the *Women1f* data would give the fitted values for the cases in the data, and allow a more data-centric plot as shown in Figure 7.42.

```

> predictors <- expand.grid(hincome=1:50,
+                           children=c('absent', 'present'))
> fit <- data.frame(predictors,
+                   p.working = predict(mod.working, predictors, type='response'),
+                   p.fulltime = predict(mod.fulltime, predictors, type='response'),
+                   l.working = predict(mod.working, predictors, type='link'),
+                   l.fulltime = predict(mod.fulltime, predictors, type='link'))
> print(some(fit, 5), digits=3)

```

	hincome	children	p.working	p.fulltime	l.working	l.fulltime
12	12	absent	0.696	0.899	0.828	2.191
17	17	absent	0.649	0.839	0.617	1.654
40	40	absent	0.412	0.307	-0.357	-0.813
65	15	present	0.294	0.314	-0.874	-0.783
94	44	present	0.109	0.020	-2.101	-3.893

One wrinkle here is that the probabilities for working full time and part time are conditional on working. We calculate the unconditional probabilities as shown below and choose to display the probability of *not* working as the complement of working.

```

> fit <- within(fit, {
+   full <- p.working * p.fulltime
+   part <- p.working * (1 - p.fulltime)
+   not <- 1 - p.working
+ })

```

Plotting these fitted values using `ggplot2` would require reshaping the `fit` data frame from wide to long format. Instead, we use R base graphics to produce plots of the probabilities and log odds. This method doesn't automatically give plots in separate panels, so a `for-loop` is used to generate panels for the levels of `children`. We set up an empty plot frame (`type="n"`) for each panel and then use `lines()` to plot the fitted probabilities. Using `par(mfrow=c(1,2))` places these plots in two side-by-side panels in a single display. The lines below give the plot shown in Figure 7.48.

```

> op <- par(mfrow=c(1,2), mar=c(5,4,4,1)+.1)
> Hinc <- 1:max(fit$hincome)
> for (kids in c("absent", "present")) {
+   dat <- subset(fit, children==kids)
+   plot(range(Hinc), c(0,1), type="n", cex.lab=1.25,
+         xlab="Husband's Income", ylab="Fitted Probability",
+         main = paste("Children", kids))
+   lines(Hinc, dat$not, lwd=3, col="black", lty=1)
+   lines(Hinc, dat$part, lwd=3, col="blue", lty=2)
+   lines(Hinc, dat$full, lwd=3, col="red", lty=3)
+   if (kids=="absent") {
+     legend("topright", lty=1:3, lwd=3, col=c("black", "blue", "red"),
+           legend=c('not working', 'part-time', 'full-time'))
+   }
+ }
> par(op)

```

We can see how that the decision not to work outside the home increases strongly with husband's income, and is higher when there are children present. As well, among working women, the decision to work full time as opposed to part time decreases strongly with husband's income, and is less likely with young children.

Similarly, we plot the fitted logits for the two dichotomies in `l.working` and `l.fulltime` as shown below, giving Figure 7.49.




Figure 7.48: Fitted probabilities from the models for nested dichotomies fit to the data on women's labor force participation.

{fig:wlf-fitted-prob}

```
> op <- par(mfrow=c(1,2), mar=c(5,4,1,1)+.1)
> for ( kids in c("absent", "present") ) {
+   dat <- subset(fit, children==kids)
+   plot( range(Hinc), c(-4,5), type="n", cex.lab=1.25,
+         xlab="Husband's Income", ylab='Fitted log odds')
+   lines(Hinc, dat$l.working, lwd=3, col="black", lty=1)
+   lines(Hinc, dat$l.fulltime, lwd=3, col="blue", lty=2)
+   text(25, 4.5, paste("Children", kids), cex=1.4)
+   if (kids=="absent") {
+     legend("bottomleft", lty=1:3, lwd=3, col=c("black", "blue"),
+           legend=c('working', 'full-time'))
+   }
+ }
> par(op)
```

This is essentially a graph of the fitted equations for L_1 and L_2 shown in Eqn. (7.19). It shows how the choice of full time work as opposed to part time depends more strongly on husband's income among women who are working than does the choice of working at all among all women. It also illustrates why the proportional odds assumption would not be reasonable for this data: that would require equal slopes for the two lines within each panel.

△

7.6.4 Generalized logit model

{sec:genlogit}

The generalized logit (or multinomial logit) approach models the probabilities of the m response categories directly as a set of $m - 1$ logits. These compare each of the first $m - 1$ categories to the last category, which serves as the baseline.²⁷ The logits for any other pair of categories can be retrieved from the $m - 1$ fitted ones.

When there are p predictors, x_1, x_2, \dots, x_p , which may be quantitative or categorical, the gen-

²⁷When the response is a factor, any category can be selected as the baseline level using `relevel()`.




Figure 7.49: Fitted log odds from the models for nested dichotomies fit to the data on women's labor force participation.

generalized logit model expresses the logits as

$$\begin{aligned} L_{jm} \equiv \log \frac{\pi_{ij}}{\pi_{im}} &= \beta_{0j} + \beta_{1j} x_{i1} + \beta_{2j} x_{i2} + \cdots + \beta_{kj} x_{ip} \quad j = 1, \dots, m-1 \\ &= \mathbf{x}_i^\top \boldsymbol{\beta}_j \end{aligned} \quad (7.21)$$

Thus, there is one set of fitted coefficients, $\boldsymbol{\beta}_j$ for each response category except the last. Each coefficient, β_{hj} , gives the effect, for a unit change in the predictor x_h , on the log odds that an observation had a response in category $Y = j$, as opposed to category $Y = m$.

The probabilities themselves can be expressed as

$$\begin{aligned} \pi_{ij} &= \frac{\exp(\mathbf{x}_i^\top \boldsymbol{\beta}_j)}{1 + \sum_{\ell=1}^{m-1} \exp(\mathbf{x}_i^\top \boldsymbol{\beta}_\ell)} \quad j = 1, 2, \dots, m-1 \\ \pi_{im} &= 1 - \sum_{i=1}^{m-1} \pi_{ij} \quad \text{for } Y = m \end{aligned}$$

Parameters in the $m-1$ equations Eqn. (7.21) can be used to determine the probabilities or the predicted log odds for any pair of response categories by subtraction. For instance, for an arbitrary pair of categories, a and b , and two predictors, x_1 and x_2 ,

$$\begin{aligned} L_{ab} &= \log \frac{\pi_{ia}/\pi_{im}}{\pi_{ib}/\pi_{im}} \\ &= \log \frac{\pi_{ia}}{\pi_{im}} - \log \frac{\pi_{ib}}{\pi_{im}} \\ &= (\beta_{0a} - \beta_{0b}) + (\beta_{1a} - \beta_{1b})x_{i1} + (\beta_{2a} - \beta_{2b})x_{i2} \end{aligned}$$

For example, the coefficient for x_{i1} in L_{ab} is just $(\beta_{1a} - \beta_{1b})$. Similarly, the predicted logit for any pair of categories can be calculated as

$$\hat{L}_{ab} = \hat{L}_{am} - \hat{L}_{bm}.$$

The generalized logit model can be fit most conveniently in R using the function `multinom()`

in the `nnet` package and the `effects` package has a set of methods for "multinom" models. These models can also be fit using `VGAM` and the `mlogit` package.

{ex:wlfpart2}

EXAMPLE 7.18: Women's labor force participation

To illustrate this method, we fit the generalized logit model to the women's labor force participation data as explained below. The response, `partic` is a character factor, and, by default `multinom()` treats these in alphabetical order and uses the *first* level as the baseline category.

```
> levels(Womenlwf$partic)
[1] "fulltime" "not.work" "parttime"
```

Although the multinomial model does not depend on the baseline category, it makes interpretation easier to choose "not.work" as the reference level, which we do with `relevel()`.²⁸

```
> # choose not working as baseline category
> Womenlwf$partic <- relevel(Womenlwf$partic, ref="not.work")
```

We fit the main effects model for husband's income and children as follows. As we did with `polr()` (Section 7.6.1), specifying `Hess=TRUE` saves the Hessian and facilitates calculation of standard errors and hypothesis tests.

```
> library(nnet)
> wlf.multinom <- multinom(partic ~ hincome + children,
+                           data=Womenlwf, Hess=TRUE)

# weights: 12 (6 variable)
initial value 288.935032
iter 10 value 211.454772
final value 211.440963
converged
```

The `summary()` method for "multinom" objects doesn't calculate test statistics for the estimated coefficients by default. The option `Wald=TRUE` produces Wald z -test statistics, calculated as $z = \beta/SE(\beta)$.

```
> summary(wlf.multinom, Wald=TRUE)

Call:
multinom(formula = partic ~ hincome + children, data = Womenlwf,
          Hess = TRUE)

Coefficients:
      (Intercept)      hincome childrenpresent
fulltime      1.9828 -0.0972321      -2.558605
parttime     -1.4323  0.0068938       0.021456

Std. Errors:
      (Intercept)      hincome childrenpresent
fulltime      0.48418 0.028096      0.36220
parttime      0.59246 0.023455      0.46904

Value/SE (Wald statistics):
      (Intercept)      hincome childrenpresent
fulltime      4.0953 -3.46071      -7.064070
parttime     -2.4176  0.29392       0.045744

Residual Deviance: 422.88
AIC: 434.88
```

²⁸Alternatively, we could declare `partic` an *ordered* factor, using `ordered()`.

Notice that the coefficients, their standard errors and the Wald test z values are printed in separate tables. The first line in each table pertains to the logit comparing full time work with the not working reference level; the second line compares part time work against not working.

For those who like p -values for significance tests, you can calculate these from the results returned by the `summary()` method in the `Wald.ratios` component, using the standard normal asymptotic approximation:

```
> stats <- summary(wlf.multinom, Wald=TRUE)
> z <- stats$Wald.ratios
> p <- 2 * (1 - pnorm(abs(z)))
> zapsmall(p)
```

	(Intercept)	hincome	childrenpresent
fulltime	0.00004	0.00054	0.00000
parttime	0.01562	0.76882	0.96351

The interpretation of these tests is that both husband's income and presence of children have highly significant effects on the comparison of working full time as opposed to not working, while neither of these predictors are significant for the comparison of working part time vs. not working.

So far, we have assumed that the effects of husband's income and presence of young children are additive on the log odds scale. We can test this assumption by allowing an interaction of those effects and testing it for significance.

```
> wlf.multinom2 <- multinom(partic ~ hincome * children,
+                             data=Womenlf, Hess=TRUE)

# weights: 15 (8 variable)
initial value 288.935032
iter 10 value 210.797079
final value 210.714841
converged

> Anova(wlf.multinom2)

Analysis of Deviance Table (Type II tests)

Response: partic
          LR Chisq Df Pr(>Chisq)
hincome      15.2  2  0.00051 ***
children     63.6  2  1.6e-14 ***
hincome:children  1.5  2  0.48378
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

The test for the interaction term, `hincome:children` is not significant, so we can abandon this model.

Full model plots of the fitted values can be plotted as shown earlier in Example 7.17: obtain the fitted values over a grid of the predictors and plot these.

```
> predictors <- expand.grid(hincome=1:50,
+                           children=c('absent', 'present'))
> fit <- data.frame(predictors,
+                   predict(wlf.multinom, predictors, type='probs'))
+                   )
```

Plotting these fitted values gives the plot shown in Figure 7.50.

```

> op <- par(mfrow=c(1,2), mar=c(5,4,4,1)+.1)
> Hinc <- 1:max(fit$hincome)
> for ( kids in c("absent", "present") ) {
+   dat <- subset(fit, children==kids)
+   plot( range(Hinc), c(0,1), type="n", cex.lab=1.25,
+         xlab="Husband's Income", ylab='Fitted Probability',
+         main = paste("Children", kids))
+   lines(Hinc, dat$not.work, lwd=3, col="black", lty=1)
+   lines(Hinc, dat$parttime, lwd=3, col="blue", lty=2)
+   lines(Hinc, dat$fulltime, lwd=3, col="red", lty=3)
+   if (kids=="absent") {
+     legend("topright", lty=1:3, lwd=3, col=c("black", "blue", "red"),
+           legend=c('not working', 'part-time', 'full-time'))
+   }
+ }
> par(op)

```




Figure 7.50: Fitted probabilities from the generalized logit model fit to the data on women's labor force participation.

fig:wlf-multi-prob)

The results shown in this plot are roughly similar to those obtained from the nested dichotomy models, graphed in Figure 7.48. However, the predicted probabilities of not working under the generalized logit model rise more steeply with husband's income for women with no children and level off sooner for women with young children.

The `effects` package has special methods for "multinom" models. It treats the response levels in the order given by `levels()`, so before plotting we use `ordered()` to arrange levels in their natural order. The `update()` method provides a simple way to get a new fitted model; in the call, the model formula `. ~ .` means to fit the same model as before, i.e., `partic ~ hincome + children`.

```

> levels(Womenlwf$partic)

[1] "not.work" "fulltime" "parttime"

> Womenlwf$partic <- ordered(Womenlwf$partic,
+                             levels=c('not.work', 'parttime', 'fulltime'))
> wlf.multinom <- update(wlf.multinom, . ~ .)

```

```
# weights: 12 (6 variable)
initial value 288.935032
iter 10 value 211.454772
final value 211.440963
converged
```

As illustrated earlier, you can use `plot(allEffects(model), ...)` to plot all the high-order terms in the model, either with separate curves for each response level (`style="lines"`) or as cumulative filled polygons (`style="stacked"`). Here, we simply plot the effects for the combinations of husband's income and children in stacked style, giving a plot (Figure 7.51) that is analogous to the full-model plot shown in Figure 7.50.

```
> plot(Effect(c("hincome", "children"), wlf.multinom),
+ style="stacked", key.args=list(x=.05, y=.9))
```




Figure 7.51: Effect plot for the probabilities of not working and working part time and full time from the generalized logit model fit to the women's labor force data.

{fig:wlf-multi-effect}

△

7.7 Chapter summary

{sec:ch07-summary}

- Model-based methods for categorical data provide confidence intervals for parameters and predicted values for observed and unobserved values of the explanatory variables. Graphical displays of predicted values help us to interpret the fitted relations by smoothing a discrete response.
- The logistic regression model (Section 7.2) describes the relationship between a categorical response variable, usually dichotomous, and a set of one or more quantitative or discrete explanatory variables (Section 7.3) It is conceptually convenient to specify this model as a linear model predicting the log odds (or logit) of the probability of a success from the explanatory variables.
- The relationship between a discrete response and a quantitative predictor may be explored

graphically by plotting the binary observations against the predictor with some smoothed curve(s), either parametric or non-parametric, possibly stratified by other predictors.

- For both quantitative and discrete predictors, the results of a logistic regression are most easily interpreted from full-model plots of the fitted values against the predictors, either on the scale of predicted probabilities or log odds (Section 7.3.2). In these plots, confidence intervals provide a visual indication of the precision of the predicted results.
- When there are multiple predictors and/or higher-order interaction terms, effect plots (Section 7.3.3) provide an important method for constructing simplified displays, focusing on the higher-order terms in a given model.
- Influence diagnostics (Section 7.5) assess the impact of individual cases or groups on the fitted model, predicted values, and the coefficients of individual predictors. Among other displays, plots of residuals against leverage showing Cook's D are often most useful.
- Other diagnostic plots (Section 7.5.3) include component-plus-residual plots, that are useful for detecting non-linear relationships for a quantitative predictor, and added-variable plots, that show the partial relations of the response to a given predictor, controlling or adjusting for all other predictors.
- Polytomous responses may be handled in several ways as extensions of binary logistic regression (Section 7.6): (a) The *proportional odds model* (Section 7.6.1) is simple and convenient, but its validity depends on an assumption of equal slopes for adjacent-category logits. (b) *Nested dichotomies* (Section 7.6.3) among the response categories give a set of models which may be regarded as a single, combined model for the polytomous response. (c) *Generalized logit models* (Section 7.6.4) may be used to construct models comparing any pair of categories.

7.8 Further reading

7.9 Lab exercises

Exercise 7.1 Arbutnot's data on the sex ratio of births in London was examined in Example 3.1. Use a binomial logistic regression model to assess whether the proportion of male births varied with the variables *Year*, *Plague* and *Mortality* in the *Arbutnot* data set. Produce effect plots for the terms in this model. What do you conclude?

Exercise 7.2 For the Donner Party data in *Donner*, examine Grayson's 1990 claim that survival in the Donner Party was also mediated by the size of the family unit. This takes some care, because the *family* variable in the *Donner* data is a simplified grouping based on the person's name and known alliances among families from the historical record. Use the following code to compute a *family.size* variable from each individual's last name:

```
> data("Donner", package="vcdExtra")
> Donner$survived <- factor(Donner$survived, labels=c("no", "yes"))
> # use last name for family
> lname <- strsplit(rownames(Donner), ",")
> lname <- sapply(lname, function(x) x[[1]])
> Donner$family.size <- as.vector(table(lname)[lname])
```

- (a) Choose one of the models (*donner.mod4*, *donner.mod6*) from Example 7.9 that include the interaction of age and sex and non-linear terms in age. Fit a new model that adds a main effect of *family.size*. What do you conclude about Grayson's claim?

- (b) Produce an effect plot for this model.
- (c) Continue, by examining whether the effect of family size can be taken as linear, or whether a non-linear term should be added.

{lab:7.3}

Exercise 7.3 Use component+residual plots (Section 7.5.3) to examine the additive model for the ICU data given by

```
> icu.glm2 <- glm(died ~ age + cancer + admit + uncons,
+               data=ICU, family=binomial)
```

- (a) What do you conclude about the linearity of the (partial) relationship between age and death in this model?
- (b) An alternative strategy is to allow some non-linear relation for age in the model using a quadratic (or cubic) term like `poly(age, 2)` (or `poly(age, 3)`) in the model formula. Do these models provide evidence for a non-linear effect of age on death in the ICU?

{lab:7.4}

Exercise 7.4 Explore the use of other marginal and conditional plots to display the relationships among the variables predicting death in the ICU in the model `icu.glm2`. For example, you might begin with a marginal `gpairs()` plot showing all bivariate marginal relations, something like this:

```
> library(gpairs)
> gpairs(ICU[,c("died", "age", "cancer", "admit", "uncons")],
+       diag.pars=list(fontsize=16, hist.color="lightgray"),
+       mosaic.pars=list(gp=shading_Friendly,
+                       gp_args=list(interpolate=1:4)))
```

{lab:7.5}

Exercise 7.5 For the women's labor force participation data (*Womenlf*) the response variable, `partic`, can be treated as ordinal by using

```
> Womenlf$partic <- ordered(Womenlf$partic,
+                          levels=c('not.work', 'parttime', 'fulltime'))
```

Use the methods in Section 7.6.1 to test whether the proportional odds model holds for these data.

{lab:7.6}

Exercise 7.6 The data set *birthwt* in the MASS package gives data on 189 babies born at Baystate Medical Center, Springfield, MA during 1986. The quantitative response is `bwt` (birth weight in grams), and this is also recorded as `low`, a binary variable corresponding to `bwt < 2500` (2.5 Kg). The goal is to study how this varies with the available predictor variables. The variables are all recorded as numeric, so in R it may be helpful to convert some of these into factors and possibly collapse some low frequency categories. The code below is just an example of how you might do this for some variables.

```
> data(birthwt, package="MASS")
> birthwt <- within(birthwt, {
+   race <- factor(race, labels = c("white", "black", "other"))
+   ptd <- factor(ptl > 0) # premature labors
+   ftv <- factor(ftv)     # physician visits
+   levels(ftv)[-1:2] <- "2+"
+   smoke <- factor(smoke>0)
+   ht <- factor(ht>0)
+   ui <- factor(ui>0)
+ })
```

- (a) Make some exploratory plots showing how low birth weight varies with each of the available predictors. In some cases, it will probably be helpful to add some sort of smoothed summary curves or lines.
- (b) Fit several logistic regression models predicting low birth weight from these predictors, with the goal of explaining this phenomenon adequately, yet simply.
- (c) Use some graphical displays to convey your findings.

{lab:7.7}

Exercise 7.7 The data set *housing* in the *MASS* package gives a $3 \times 3 \times 4 \times 2$ table in frequency form relating (a) satisfaction (*Sat*) of residents with their housing (High, Medium, Low), (b) perceived degree of influence (*Infl*) they have on the management of the property (High, Medium, Low), (c) Type of rental (Tower, Atrium, Apartment, Terrace), and (d) contact (*Cont*) residents have with other residents (Low, High). Consider satisfaction as the ordinal response variable.

- (a) Fit the proportional odds model with additive (main) effects of housing type, influence in management and contact with neighbors to this data. (Hint: Using `polr()`, with the data in frequency form, you need to use the `weights` argument to supply the *Freq* variable.)
- (b) Investigate whether any of the two-factor interactions among *Infl*, *Type* and *Cont* add substantially to goodness of fit of this model. (Hint: use `stepAIC()`, with the scope formula `~ .^2` and `direction="forward"`.)
- (c) For your chosen model from the previous step, use the methods of Section 7.6.2 to plot the probabilities of the categories of satisfaction.
- (d) Write a brief summary these analyses, interpreting *how* satisfaction with housing depends on the predictor variables.

{lab:7.8}

Exercise 7.8 The data *TV* on television viewing was analyzed using correspondence analysis in Example 6.4, ignoring the variable *Time* and extended in Exercise 6.7. Treating *Network* as a three-level response variable, fit a generalized logit model (Section 7.6.4) to explain the variation in viewing in relation to *Day* and *Time*. The *TV* data is a three-way table, so you will need to convert it to a frequency data frame first.

```
> data("TV", package="vcdExtra")
> TV.df <- as.data.frame.table(TV)
```

- (a) Fit the main-effects model, `Network ~ Day + Time` with `multinom()`. Note that you will have to supply the `weights` argument because each row of `TV.df` represents the number of viewers in the *Freq* variable.
- (b) Prepare an effects plot for the fitted probabilities in this model.
- (c) Interpret these results in comparison to the correspondence analysis analysis in Example 6.4.

{lab:7.9}

Exercise 7.9 Refer to Exercise 5.8 for a description of the *Accident* data. The interest here is to model the probability that an accident resulted in death rather than injury from the predictors *age*, *mode* and *gender*. With `glm()`, and the data in the form of a frequency table, you can use the argument `weight=Freq` to take cell frequency into account.

- (a) Fit the main effects model, `result=="Died" ~ age + mode + gender`. Use `car::Anova()` to assess the model terms.
- (b) Fit the model that allows all two-way interactions. Use `anova()` to test whether this model is significantly better than the main effects model.
- (c) Fit the model that also allows the three-way interaction of all factors. Does this offer any improvement over the two-way model?
- (d) Interpret the results of the analysis using effect plots for the two-way model, separately for each of the model terms. Describe verbally the nature of the *age*gender* effect. Which mode of transportation leads to greatest risk of death?

```
lab:logisfit(wbenvn)}

```

Exercise 7.10 Refer to Exercise ?? for a description of the *Vietnam* data set in *vcdExtra*. The goal here is to fit models for the polytomous response variable in relation to year and sex.

- Fit the proportional odds model to these data, allowing an interaction of year and sex.
- Is there evidence that the proportional odds assumption does not hold for this data set? Use the methods described in Section 7.6.1 to assess this.
- Fit the multinomial logistic model, also allowing an interaction. Use `car::Anova()` to assess the model terms.
- Produce an effect plot for this model and describe the nature of the interaction.
- Fit the simpler multinomial model in which there is no effect of year for females and the effect of year is linear for males (on the logit scale). Test whether this model is significantly worse than the general multinomial model with interaction.

TODO: Testing how to detach possibly conflicting packages

```
> .locals$ch07 <- setdiff(ls(), .globals)
> #.locals$ch07
> remove(list=.locals$ch07[ sapply(.locals$ch07, function(n) { !is.function(get(n)) }) ])
> detach(package:gpairs)
> detach(package:rms)
> detach(package:VGAM)
> .pkgs$ch07 <- setdiff(.packages(), .pkgs$ch07)
> .pkgs$ch07

[1] "nnet"          "reshape2"      "Hmisc"         "stats4"
[5] "proto"         "SparseM"       "gridExtra"     "Formula"
[9] "survival"      "splines"       "car"           "effects"
[13] "lmtest"        "zoo"           "ggplot2"       "directlabels"
[17] "quadprog"      "Lahman"        "lattice"       "MASS"
[21] "colorspace"    "vcdExtra"      "gnm"           "vcd"
[25] "grid"
```


References

- Bliss, C. I. (1934). The method of probits. *Science*, 79(2037), 38–39.
- Bollen, K. A. (2002). Latent variables in psychology and the social sciences. *Annual Review of Psychology*, 53(1), 605–634.
- Box, G. E. P. and Draper, N. R. (1987). *Empirical Model Building and Response Surfaces*. New York, NY: John Wiley & Sons.
- Cook, R. D. and Weisberg, S. (1999). *Applied Regression Including Computing and Graphics*. New York: Wiley.
- Dalal, S., Fowlkes, E. B., and Hoadley, B. (1989). Risk analysis of the space shuttle: Pre-Challenger prediction of failure. *Journal of the American Statistical Association*, 84(408), 945–957.
- de la Cruz Rot, M. (2005). Improving the presentation of results of logistic regression with r. *Bulletin of the Ecological Society of America*, 86, 41–48.
- Finney, D. J. (1947). *Probit analysis*. Cambridge England: University Press.
- Fowlkes, E. B. (1987). Some diagnostics for binary logistic regression via smoothing. *Biometrika*, 74(3), 503–5152.
- Fox, J. (1987). Effect displays for generalized linear models. In C. C. Clogg, ed., *Sociological Methodology, 1987*, (pp. 347–361). San Francisco: Jossey-Bass.
- Fox, J. (2003). Effect displays in R for generalized linear models. *Journal of Statistical Software*, 8(15), 1–27.
- Fox, J. (2008). *Applied Regression Analysis and Generalized Linear Models*. Thousand Oaks, CA: Sage, 2nd edn.
- Fox, J. and Weisberg, S. (2011). *An R Companion to Applied Regression*. Thousand Oaks CA: Sage, 2nd edn.
- Grayson, D. K. (1990). Donner party deaths: A demographic assessment. *Journal of Anthropological Research*, 46(3), 223–242.
- Harrell, Jr, F. E. (2001). *Regression Modeling Strategies: With Applications to Linear Models, Logistic Regression, and Survival Analysis*. Graduate Texts in Mathematics. New York: Springer.
- Hosmer, Jr, D. W., Lemeshow, S., and Sturdivant, R. X. (2013). *Applied Logistic Regression*. New York: John Wiley and Sons, 3rd edn.
- Johnson, K. (1996). *Unfortunate Emigrants: Narratives of the Donner Party*. Logan, UT: Utah State University Press.

- Landwehr, J. M., Pregibon, D., and Shoemaker, A. C. (1984). Graphical methods for assessing logistic regression models. *Journal of the American Statistical Association*, 79, 61–71.
- Larsen, W. A. and McCleary, S. J. (1972). The use of partial residual plots in regression analysis. *Technometrics*, 14, 781–790.
- Lavine, M. (1991). Problems in extrapolation illustrated with space shuttle O-ring data. *Journal of the American Statistical Association*, 86, 912–922.
- Lawrance, A. J. (1995). Deletion influence and masking in regression. *Journal of the Royal Statistical Society. Series B (Methodological)*, 57(1), 181–189.
- Lazarsfeld, P. F. (1950). The logical and mathematical foundation of latent structure analysis. In S. A. Stouffer, L. Guttman, E. A. Suchman, P. F. Lazarsfeld, S. A. Star, and J. A. Clausen, eds., *Studies in Social Psychology in World War II, vol. IV, Measurement and Prediction*, (pp. 362–412). Princeton, NJ: Princeton University Press.
- Lazarsfeld, P. F. (1954). A conceptual introduction to latent structure analysis. In P. F. Lazarsfeld, ed., *Mathematical Thinking in the Social Sciences*, (pp. 349–387). Glencoe, IL: Free Press.
- Lemeshow, S., Avrunin, D., and Pastides, J. S. (1988). Predicting the outcome of intensive care unit patients. *Journal of the American Statistical Association*, 83, 348–356.
- McCullagh, P. (1980). Regression models for ordinal data. *Journal of the Royal Statistical Society*, 42, 109–142.
- Peterson, B. and Harrell, Jr, F. E. (1990). Partial proportional odds models for ordinal response variables. *Applied Statistics*, 39, 205–217.
- Pregibon, D. (1981). Logistic regression diagnostics. *Annals of Statistics*, 9, 705–724.
- Searle, S. R., Speed, F. M., and Milliken, G. A. (1980). Population marginal means in the linear model: An alternative to least squares means. *The American Statistician*, 34(4), 216–221.
- Thurstone, L. L. (1927). A law of comparative judgment. *Psychological Review*, 34, 278–286.
- Tufte, E. R. (1997). *Visual Explanations: Images and Quantities, Evidence and Narrative*. Cheshire, CT: Graphics Press.
- Wang, P. C. (1985). Adding a variable in generalized linear models. *Technometrics*, 27, 273–276.
- Williams, D. A. (1987). Generalized linear model diagnostics using the deviance and single case deletions. *Applied Statistics*, 36, 181–191.