

Calculation of momentum velocity on collision of two masses

Friedemann Zintel

October 15, 2016

In order to calculate the momentum between two colliding masses it is necessary to determine the partial velocity vector of a moving mass in direction to the second mass. Let's assume a mass m_1 moves with velocity \vec{v} and would hit mass m_2 . For simplicity m_2 remains stationary and the shapes of both masses are spherical. Even when the direction of \vec{v} does not point directly to m_2 , it behaves as if m_1 hits m_2 with a (partial) velocity v_{m_2} which points in the direction to m_2 . We keep the calculation in two dimensional space. It should be easy to extend it to three dimensions.




Figure 1: collision

We need to calculate v_{m_2} . We define $\vec{d} = \vec{m}_2 - \vec{m}_1$, where \vec{m}_1 and \vec{m}_2 are the positional vectors for m_1 resp. m_2 . We require that the masses have a positive expansion ($r_1 > 0$, $r_2 > 0$), thus $\vec{d} \neq \vec{0}$ and $d_x \neq 0 \vee d_y \neq 0$ (collision takes place if $|\vec{d}| = r_1 + r_2$). Let \vec{s} be a (the) vector with $v_{m_2} = \vec{v} + \vec{s}$. Then \vec{s} must be orthogonal to v_{m_2} and thus to \vec{d} . That is because v_{m_2} is a partial vector of \vec{v} and points in the same direction as \vec{d} which resides on the momentum line. See figure 1.

Be $\lambda \in \mathbb{R}$, then the following equations hold:

$$\begin{aligned} v_{m_2} &= \vec{v} + \vec{s} = \lambda \vec{d} \\ \vec{s} \cdot \vec{d} &= 0 \end{aligned}$$

We end up with three equations and three unknown variables (s_x, s_y, λ) , which can easily be solved:

$$v_x + s_x = \lambda d_x \quad (1)$$

$$v_y + s_y = \lambda d_y \quad (2)$$

$$s_x d_x + s_y d_y = 0 \quad (3)$$

case: $d_x \neq 0$

$$(3) \Leftrightarrow s_x = -\frac{s_y d_y}{d_x} \quad (4)$$

$$(2) \Leftrightarrow s_y = \lambda d_y - v_y \quad (5)$$

$$\Rightarrow s_x = -\frac{(\lambda d_y - v_y) d_y}{d_x} \quad (6)$$

$$\text{in (1): } v_x - \frac{(\lambda d_y - v_y) d_y}{d_x} = \lambda d_x \quad (7)$$

$$\Leftrightarrow v_x - \frac{\lambda d_y^2}{d_x} + \frac{v_y d_y}{d_x} = \lambda d_x \quad (8)$$

$$\Leftrightarrow v_x + \frac{v_y d_y}{d_x} = \lambda \left(d_x + \frac{d_y^2}{d_x} \right) \quad (9)$$

$$\Leftrightarrow \frac{v_x d_x}{d_x} + \frac{v_y d_y}{d_x} = \lambda \left(\frac{d_x^2 + d_y^2}{d_x} \right) \quad (10)$$

$$\Leftrightarrow \frac{(v_x d_x + v_y d_y) d_x}{d_x (d_x^2 + d_y^2)} = \lambda \quad (11)$$

$$\Leftrightarrow \lambda = \frac{v_x d_x + v_y d_y}{d_x^2 + d_y^2} = \frac{\vec{v} \vec{d}}{\vec{d}^2} \quad (12)$$

case: $d_y \neq 0$

Calculation is analogue. We end up with the same result for λ .

As a result we get:

$$v_{\vec{m}_2} = \frac{\vec{v} \vec{d}}{\vec{d}^2} \vec{d}$$

□