HOORVIA - MEDICATION SCHEDULE CHART: PRINTABLE DAILY DOSE TRACKER

Quick Reference Cheat Sheet · US Letter Printable

Home, Family & Health · Sheet #1

Medication Example	Usual Timing	Food Guidance	Scheduling Tip
Multivitamin	Morning with breakfast	With food	Fat-soluble vitamins (A, D, E, K) absorb better with a meal
Iron supplement	Morning, 1 hour before food	Empty stomach	Vitamin C boosts absorption; keep away from coffee and calcium
Calcium supplement	Evening or split doses	With food	Absorbs about 500 mg at a time; never take together with iron
Short-acting statin (e.g. simvastatin)	Evening or bedtime	Either	Cholesterol synthesis peaks overnight, so evening dosing works best
Blood pressure medication	Same time daily, usually morning	Either	Consistency matters more than the exact hour
Levothyroxine (thyroid)	30-60 min before breakfast	Empty stomach	Take alone; coffee, calcium, and iron block absorption
Metformin (diabetes)	With meals	With food	Food cuts stomach upset; match doses to your meals
Antibiotics (e.g. amoxicillin)	Evenly spaced, every 8-12 hours	Either	Finish the full course; set alarms for overnight doses
Drowsy antihistamine	Evening or bedtime	Either	Works best at night, when sleepiness is a feature
Probiotic	With or just after a meal	With food	Food buffers stomach acid so more bacteria survive
Fish oil / omega-3	With a fat-containing meal	With food	Taking it at dinner reduces fishy aftertaste
Melatonin	30-60 min before bed	Empty stomach is fine	Best for short-term sleep resets, not as a nightly habit