

Neutrinos via Charm Decays in Astrophysical Sources

by

FRITZ ALI AGILDERE

fritz.agildere@udo.edu

Abstract

Acknowledgements

Abbreviations

Contents

1	Introduction	1
1.1	Motivation	1
1.2	Organisation	1
2	Methods	2
3	Results	3
4	Discussion	4
	Appendix	5

1 Introduction

1.1 Motivation

1.2 Organisation

2 Methods

$$\mu = \frac{BR^3}{2} \quad (2.1)$$

$$\dot{E} = -\frac{2\mu^2\omega^4}{3c^3} \sin^2\chi \quad (2.2)$$

$$\dot{E} = -\frac{\mu^2\omega^4}{c^3} (1 + \sin^2\chi) \quad (2.3)$$

$$\tau \equiv -\frac{E}{\dot{E}} \quad (2.4)$$

$$I = \frac{2MR^2}{5} \quad (2.5)$$

$$E = \frac{I\omega^2}{2} \quad (2.6)$$

$$\dot{E} = \omega\dot{\omega} = K\omega^4 \quad (2.7)$$

$$I\dot{\omega} = K\omega^3 \quad (2.8)$$

$$I\omega^{-3}d\omega = Kdt \quad (2.9)$$

$$\omega^{-2} = \omega_0^{-2} - \frac{2Kt}{I} = \omega_0^{-2} \left(1 + \frac{t}{\tau}\right) \quad (2.10)$$

$$\tau = -\frac{I}{2K\omega^2} \quad (2.11)$$

$$\omega = \frac{\omega_0}{\sqrt{1 + \frac{t}{\tau}}} \quad (2.12)$$

3 Results

4 Discussion

Appendix