# Neutrinos via Charm Decays in Astrophysical Sources

by

Fritz Ali Agildere

fritz.agildere@udo.edu

## Abstract

## Abbreviations

## Contents

1	Introduction	1
2	Background 2.1 Particle Physics 2.2 Multimessenger Astronomy 2.3 Astrophysical Sources	2 2 2 2
3	Results	3
4	Conclusion & Outlook	4
Ac	cknowledgements	5
Αp	ppendix	6
Bil	bliography	7

### 1 Introduction

αβγδηζξ χφφθθμνρ

```
first zeroth test.^{1-3}
second zeroth test.<sup>1,2</sup>
third zeroth \mathsf{test}^{1,3}
test.4
first test [1-3]
second test [1, 2]
third test [1, 3]
[5]
[4]
[6]
(2.998 \pm 0.001) \times 10^{10} \,\mathrm{cm\,s}^{-1}
\pm \times + - \cdot 6.626 \times 10^{-27} \text{ erg s}
4.803 \times 10^{-10} \text{ esu}
10^{14} \, \mathrm{G}
1.4 \times 10^0 \ 1.4 \times 10^0
                                                                   \sum\nolimits_{n=1}^{\infty}\frac{1}{n^2}=\frac{\pi^2}{6}
                                                                                                                                                             (1)
\sqrt{1-\beta^2}
```

### 2 Background

#### 2.1 Particle Physics

The interaction and classification of elementary particles is generally described by the *Standard Model* (SM) of particle physics. This view includes three of the four known fundamental forces, namely electromagnetism as well as the weak and strong interactions. Each of these sectors is characterized mathematically in the context of *Quantum Field Theory* (QFT), which combines aspects from classical field theory, special relativity and quantum mechanics to deliver some of the most precise predictions available today. Further gauge theoretical considerations as well as the symmetries encoded in the unitary  $U(1) \times SU(2) \times SU(3)$  group give rise to a self consistent model of physics on microscopic scales.

Despite its impressive predictive power, there are some important observations which this approach has so far been incapable of explaining. Perhaps most importantly, gravitational interactions as the fourth fundamental force are not incorporated at all. Due to the historical successes in quantizing classical theories, it is widely believed that some form of quantum gravity is the most likely candidate for explaining the emergence of macroscopic phenomena as described by *General Relativity* (GR), another extremely accurate model of the universe at large. However, in the energy regimes where such a hypothetical graviton is expected to become relevant, the formulation of the SM breaks down.

#### 2.2 Multimessenger Astronomy

#### 2.3 Astrophysical Sources

Magnetars

Active Galactic Nuclei

# 3 Results

# 4 Conclusion & Outlook

# Acknowledgements

# Appendix

### Bibliography

- 1. J. A. Carpio, K. Murase, M. H. Reno, I. Sarcevic, A. Stasto, *Physical Review D* **102**, ISSN: 2470-0029, DOI: 10.1103/PhysRevD.102.103001, arXiv: 2007.07945 [astro-ph.HE] (2020).
- 2. C. Alvarez, A. Carramiñana, *Astronomy & Astrophysics* **414**, 651–658, ISSN: 1432-0746, DOI: 10.1051/0004-6361:20031627, arXiv: astro-ph/0311267 (2004).
- 3. J. D. Jackson, Classical Electrodynamics (Wiley, New York, NY, Third Edition, 1999), ISBN: 9780471309321.
- 4. J. Li, A. Spitkovsky, A. Tchekhovskoy, *The Astrophysical Journal* **746**, 60, DOI: 10.1088/0004-637X/746/1/60, arXiv: 1107.0979 [astro-ph.HE] (2012).
- 5. C. Thompson, R. C. Duncan, *The Astrophysical Journal* **408**, 194, DOI: 10.1086/172580 (1993).
- 6. B. Haskell, A. Sedrakian, in *Astrophysics and Space Science Library* (Springer International Publishing, 2018), Chapter 8, 401–454, ISBN: 9783319976167, ISSN: 2214-7985, arXiv: 1709.10340 [astro-ph.HE].

# Figures

# Tables