



Neutrinos via Charm Decays in Astrophysical Sources


TALK IN PARTIAL FULFILLMENT OF THE REQUIREMENTS FOR THE DEGREE OF BACHELOR OF SCIENCE

Fritz Ali Agildere

September 3, 2024


Technical University of Dortmund & Ruhr University Bochum

Multimessenger Astrophysics




X. L. Qian et al.
Nucl. Sci. Tech. (2021)

Cosmic Ray Spectrum & Cutoff



K. A. Olive et al.
Chin. Phys. C (2014)


Plausible Cosmic Ray Origins



A. V. Olinto
AIP Conf. Proc. (2000)

Elementary Particles & Hadrons


Standard Model of Elementary Particles





Standard Model
Wikimedia Commons (2019)


Relevant Hadron Properties

h	q_h	τ_h / s	m_h / GeV	s_h / GeV^2	$B_{h \rightarrow \nu}$
p	uud	∞	0.94	15.98	0.000
π^+	$u\bar{d}$	26.0×10^{-9}	0.14	10.23	1.000
K^+	$u\bar{s}$	12.4×10^{-9}	0.49	12.62	0.636
D_0	$c\bar{u}$	0.4×10^{-12}	1.86	0.45	0.067
D^+	$c\bar{d}$	1.0×10^{-12}	1.87	0.40	0.176
D_s^+	$c\bar{s}$	0.5×10^{-12}	1.97	0.71	0.065
Λ_c^+	udc	0.2×10^{-12}	2.29	1.61	0.045











Young Neutron Stars




R. Takeshige
Magnetar (2020)

Dipolar Pulsar Model



E. Aliu et al.
Science (2008)

Particle Acceleration



P. Goldreich, W. H. Julian *ApJ* (1969)



Magnetar Spindown

- Electromagnetic Luminosity

Magnetar Spindown

■ Electromagnetic Luminosity

$$L = \frac{B_{ns}^2 R_{ns}^6 \omega_0^4 (1 + \sin^2 \chi)}{4c^3 (1 + t/t_{sd})^2}$$

Magnetar Spindown

■ Electromagnetic Luminosity

$$L = \frac{B_{ns}^2 R_{ns}^6 \omega_0^4 (1 + \sin^2 \chi)}{4c^3 (1 + t/t_{sd})^2}$$

Magnetar Spindown

■ Electromagnetic Luminosity

$$L = \frac{B_{ns}^2 R_{ns}^6 \omega_0^4 (1 + \sin^2 \chi)}{4c^3 (1 + t/t_{sd})^2}$$

Magnetar Spindown

■ Electromagnetic Luminosity

$$L = \frac{B_{ns}^2 R_{ns}^6 \omega_0^4 (1 + \sin^2 \chi)}{4c^3 (1 + t/t_{sd})^2}$$

Magnetar Spindown

■ Electromagnetic Luminosity

$$L = \frac{B_{ns}^2 R_{ns}^6 \omega_0^4 (1 + \sin^2 \chi)}{4c^3 (1 + t/t_{sd})^2}$$

Magnetar Spindown

- Electromagnetic Luminosity

$$L = \frac{B_{ns}^2 R_{ns}^6 \omega_0^4 (1 + \sin^2 \chi)}{4c^3 (1 + t/t_{sd})^2}$$

- Characteristic Timescale

Magnetar Spindown

■ Electromagnetic Luminosity

$$L = \frac{B_{ns}^2 R_{ns}^6 \omega_0^4 (1 + \sin^2 \chi)}{4c^3 (1 + t/t_{sd})^2}$$

■ Characteristic Timescale

$$t_{sd} = \frac{2I_{ns} c^3}{B_{ns}^2 R_{ns}^6 \omega_0^2 (1 + \sin^2 \chi)}$$

Magnetar Spindown

- Electromagnetic Luminosity

$$L = \frac{B_{ns}^2 R_{ns}^6 \omega_0^4 (1 + \sin^2 \chi)}{4c^3 (1 + t/t_{sd})^2}$$

- Characteristic Timescale

$$t_{sd} = \frac{2I_{ns} c^3}{B_{ns}^2 R_{ns}^6 \omega_0^2 (1 + \sin^2 \chi)}$$

- Monotonic Energy

Magnetar Spindown

■ Electromagnetic Luminosity

$$L = \frac{B_{ns}^2 R_{ns}^6 \omega_0^4 (1 + \sin^2 \chi)}{4c^3 (1 + t/t_{sd})^2}$$

■ Characteristic Timescale

$$t_{sd} = \frac{2I_{ns} c^3}{B_{ns}^2 R_{ns}^6 \omega_0^2 (1 + \sin^2 \chi)}$$

■ Monotonic Energy

$$E^M = \frac{\eta e B_{ns} R_{ns}^3 \omega_0^2}{2c^2 (1 + t/t_{sd})}$$

Magnetar Spindown

■ Electromagnetic Luminosity

$$L = \frac{B_{ns}^2 R_{ns}^6 \omega_0^4 (1 + \sin^2 \chi)}{4c^3 (1 + t/t_{sd})^2}$$

■ Characteristic Timescale

$$t_{sd} = \frac{2I_{ns} c^3}{B_{ns}^2 R_{ns}^6 \omega_0^2 (1 + \sin^2 \chi)}$$

■ Monotonic Energy

$$E^M = \frac{\eta e B_{ns} R_{ns}^3 \omega_0^2}{2c^2 (1 + t/t_{sd})}$$

Magnetar Spindown

■ Electromagnetic Luminosity

$$L = \frac{B_{ns}^2 R_{ns}^6 \omega_0^4 (1 + \sin^2 \chi)}{4c^3 (1 + t/t_{sd})^2}$$

■ Characteristic Timescale

$$t_{sd} = \frac{2I_{ns} c^3}{B_{ns}^2 R_{ns}^6 \omega_0^2 (1 + \sin^2 \chi)}$$

■ Monotonic Energy

$$E^M = \frac{\eta e B_{ns} R_{ns}^3 \omega_0^2}{2c^2 (1 + t/t_{sd})}$$

Magnetar Spindown

■ Electromagnetic Luminosity

$$L = \frac{B_{ns}^2 R_{ns}^6 \omega_0^4 (1 + \sin^2 \chi)}{4c^3 (1 + t/t_{sd})^2}$$


■ Characteristic Timescale

$$t_{sd} = \frac{2I_{ns} c^3}{B_{ns}^2 R_{ns}^6 \omega_0^2 (1 + \sin^2 \chi)}$$


■ Monotonic Energy


$$E^M = \frac{\eta e B_{ns} R_{ns}^3 \omega_0^2}{2c^2 (1 + t/t_{sd})}$$


Active Galactic Nuclei



V. Beckmann, C. Shrader
Active Galactic Nuclei (2012)









Injected Primary Spectrum

- Monochromatic Proton Energy

Injected Primary Spectrum

■ Monochromatic Proton Energy

$$\frac{d\dot{N}_p}{dE_p} = \frac{B_{ns} R_{ns}^3 \omega_0^2}{ec(1 + t/t_{sd})} \delta(E_p - E^M)$$

Injected Primary Spectrum

■ Monochromatic Proton Energy

$$\frac{d\dot{N}_p}{dE_p} = \frac{B_{ns} R_{ns}^3 \omega_0^2}{ec(1 + t/t_{sd})} \delta(E_p - E^M)$$

Injected Primary Spectrum

■ Monochromatic Proton Energy

$$\frac{d\dot{N}_p}{dE_p} = \frac{B_{ns} R_{ns}^3 \omega_0^2}{ec(1 + t/t_{sd})} \delta(E_p - E^M)$$

Injected Primary Spectrum

■ Monochromatic Proton Energy

$$\frac{d\dot{N}_p}{dE_p} = \frac{B_{ns} R_{ns}^3 \omega_0^2}{ec(1 + t/t_{sd})} \delta(E_p - E^M)$$

Injected Primary Spectrum

■ Monochromatic Proton Energy

$$\frac{d\dot{N}_p}{dE_p} = \frac{B_{ns} R_{ns}^3 \omega_0^2}{ec(1 + t/t_{sd})} \delta(E_p - E^M)$$

$$\frac{dN_p}{dE_p} \propto E_p^{-1}$$

Injected Primary Spectrum

■ Monochromatic Proton Energy

$$\frac{d\dot{N}_p}{dE_p} = \frac{B_{ns} R_{ns}^3 \omega_0^2}{ec(1 + t/t_{sd})} \delta(E_p - E^M)$$

$$\frac{dN_p}{dE_p} \propto E_p^{-1}$$

■ Stochastic Acceleration Power Law

Injected Primary Spectrum


■ Monochromatic Proton Energy


$$\frac{d\dot{N}_p}{dE_p} = \frac{B_{ns} R_{ns}^3 \omega_0^2}{ec(1 + t/t_{sd})} \delta(E_p - E^M)$$


$$\frac{dN_p}{dE_p} \propto E_p^{-1}$$

■ Stochastic Acceleration Power Law

$$\frac{dN_p}{dE_p} \propto E_p^{-2}$$







Magnetar Environment

- Supernova Shell Radius

Magnetar Environment

■ Supernova Shell Radius

$$R_{ej} = \beta_{ej} ct$$

Magnetar Environment

- Supernova Shell Radius

$$R_{ej} = \beta_{ej} ct$$

- Ejecta Number Density

Magnetar Environment

- Supernova Shell Radius

$$R_{ej} = \beta_{ej} ct$$

- Ejecta Number Density

$$n_{ej} = \frac{3M_{ej}}{4\pi R_{ej}^3 m_p}$$

Magnetar Environment

- Supernova Shell Radius

$$R_{ej} = \beta_{ej} ct$$

- Ejecta Number Density

$$n_{ej} = \frac{3M_{ej}}{4\pi R_{ej}^3 m_p}$$

Magnetar Environment

- Supernova Shell Radius

$$R_{ej} = \beta_{ej} ct$$

- Ejecta Number Density

$$n_{ej} = \frac{3M_{ej}}{4\pi R_{ej}^3 m_p}$$

Magnetar Environment

- Supernova Shell Radius

$$R_{ej} = \beta_{ej} ct$$

- Ejecta Number Density

$$n_{ej} = \frac{3M_{ej}}{4\pi R_{ej}^3 m_p}$$

- Numerical Setup

Magnetar Environment

■ Supernova Shell Radius

$$R_{ej} = \beta_{ej} ct$$

■ Ejecta Number Density

$$n_{ej} = \frac{3M_{ej}}{4\pi R_{ej}^3 m_p}$$

■ Numerical Setup

$$\begin{array}{llll} I_{ns} = 10^{45} \text{ g cm}^2 & R_{ns} = 10^6 \text{ cm} & B_{ns} = 10^{14.5} \text{ G} & t_{sd} = 10^{3.5} \text{ s} \\ M_{ej} = 10 M_{\odot} & \beta_{ej} = 0.1 & \eta = 0.1 & \end{array}$$

Accretion Disk

- Thin Limit Scale Height

Accretion Disk

■ Thin Limit Scale Height

$$h^2 = \frac{u_{sn}^2 R_{dc}^3}{GM_{bh}}$$

Accretion Disk

- Thin Limit Scale Height

$$h^2 = \frac{u_{sn}^2 R_{dc}^3}{GM_{bh}}$$

- Gaussian Density Profile

Accretion Disk

- Thin Limit Scale Height

$$h^2 = \frac{u_{sn}^2 R_{dc}^3}{GM_{bh}}$$

- Gaussian Density Profile

$$\rho \propto \exp\left(-\frac{GM_{bh} z^2}{2u_{sn}^2 R_{dc}^3}\right)$$

Accretion Disk

- Thin Limit Scale Height

$$h^2 = \frac{u_{sn}^2 R_{dc}^3}{GM_{bh}}$$

- Gaussian Density Profile

$$\rho \propto \exp\left(-\frac{GM_{bh} z^2}{2u_{sn}^2 R_{dc}^3}\right)$$

Accretion Disk

- Thin Limit Scale Height

$$h^2 = \frac{u_{sn}^2 R_{dc}^3}{GM_{bh}}$$

- Gaussian Density Profile

$$\rho \propto \exp\left(-\frac{GM_{bh} z^2}{2u_{sn}^2 R_{dc}^3}\right)$$

Accretion Disk

- Thin Limit Scale Height

$$h^2 = \frac{u_{sn}^2 R_{dc}^3}{GM_{bh}}$$

- Gaussian Density Profile

$$\rho \propto \exp\left(-\frac{GM_{bh} z^2}{2u_{sn}^2 R_{dc}^3}\right)$$

- Parameter Choice

Accretion Disk

■ Thin Limit Scale Height


$$h^2 = \frac{u_{sn}^2 R_{dc}^3}{GM_{bh}}$$


■ Gaussian Density Profile

$$\rho \propto \exp\left(-\frac{GM_{bh} z^2}{2u_{sn}^2 R_{dc}^3}\right)$$

■ Parameter Choice

$$n_{dc} = 10^{14} \text{ cm}^{-3} \qquad h = 10^{15} \text{ cm}$$







Direct Hadron Spectrum

- Single Interaction Distribution

Direct Hadron Spectrum

■ Single Interaction Distribution

$$F_{pp \rightarrow h} = E_p^{-1} \sigma_{pp}^{-1} \frac{d\sigma}{dx_E}$$

Direct Hadron Spectrum

- Single Interaction Distribution

$$F_{pp \rightarrow h} = E_p^{-1} \sigma_{pp}^{-1} \frac{d\sigma}{dx_E}$$

- Proton Number Convolution


Direct Hadron Spectrum


■ Single Interaction Distribution


$$F_{pp \rightarrow h} = E_p^{-1} \sigma_{pp}^{-1} \frac{d\sigma}{dx_E}$$

■ Proton Number Convolution

$$\frac{dN_h}{dE_h} = \int_{E_h}^{\infty} dE_p \frac{dN_p}{dE_p} F_{pp \rightarrow h}$$









Unweighted Energy Differential Cross Section

- Fragmentation Function

Unweighted Energy Differential Cross Section

■ Fragmentation Function

$$D_c^h = \frac{N_h z (1 - z)^2}{((1 - z)^2 + \epsilon_h z)^2}$$

Unweighted Energy Differential Cross Section

■ Fragmentation Function

$$D_c^h = \frac{N_h z(1-z)^2}{((1-z)^2 + \epsilon_h z)^2}$$

■ Charmed Hadron Spectra

Unweighted Energy Differential Cross Section

■ Fragmentation Function

$$D_c^h = \frac{N_h z(1-z)^2}{((1-z)^2 + \epsilon_h z)^2}$$

■ Charmed Hadron Spectra

$$\frac{d\sigma}{dx_E} = \left(1 - \frac{m_h}{x_E E_p}\right)^{1/2} \int_{x_E}^1 dz z^{-1} \frac{d\sigma}{dx_c} D_c^h$$

Unweighted Energy Differential Cross Section

■ Fragmentation Function

$$D_c^h = \frac{N_h z(1-z)^2}{((1-z)^2 + \epsilon_h z)^2}$$

■ Charmed Hadron Spectra

$$\frac{d\sigma}{dx_E} = \left(1 - \frac{m_h}{x_E E_p}\right)^{1/2} \int_{x_E}^1 dz z^{-1} \frac{d\sigma}{dx_c} D_c^h$$

Unweighted Energy Differential Cross Section

■ Fragmentation Function

$$D_c^h = \frac{N_h z(1-z)^2}{((1-z)^2 + \epsilon_h z)^2}$$

■ Charmed Hadron Spectra

$$\frac{d\sigma}{dx_E} = \left(1 - \frac{m_h}{x_E E_p}\right)^{1/2} \int_{x_E}^1 dz z^{-1} \frac{d\sigma}{dx_c} D_c^h$$

Unweighted Energy Differential Cross Section

■ Fragmentation Function

$$D_c^h = \frac{N_h z(1-z)^2}{((1-z)^2 + \epsilon_h z)^2}$$

■ Charmed Hadron Spectra

$$\frac{d\sigma}{dx_E} = \left(1 - \frac{m_h}{x_E E_p}\right)^{1/2} \int_{x_E}^1 dz z^{-1} \frac{d\sigma}{dx_c} D_c^h$$

■ Energy Fractions

Unweighted Energy Differential Cross Section

■ Fragmentation Function

$$D_c^h = \frac{N_h z(1-z)^2}{((1-z)^2 + \epsilon_h z)^2}$$

■ Charmed Hadron Spectra

$$\frac{d\sigma}{dx_E} = \left(1 - \frac{m_h}{x_E E_p}\right)^{1/2} \int_{x_E}^1 dz z^{-1} \frac{d\sigma}{dx_c} D_c^h$$


■ Energy Fractions

$$x_E = E_h/E_p$$

$$x_c = E_c/E_p$$

$$z = E_h/E_c$$

Weighted Energy Differential Cross Section




Derived Parametrizations

S. R. Kelner et al.
Phys. Rev. D (2006)

V. P. Gonçalves et al.
JHEP (2007)


Hadron Inelastic Scattering Cross Section




Combined Parametrizations

V. I. Belousov et al.
Phys. Atom. Nuclei (2016)

D. A. Fagundes, M. J. Menon
Nucl. Phys. A (2012)





Hadronic Cooling & Decay

■ Characteristic Timescales

Hadronic Cooling & Decay

- Characteristic Timescales

$$t_{cool} = (\kappa_{hp} \sigma_{hp} n c)^{-1}$$

Hadronic Cooling & Decay

■ Characteristic Timescales

$$t_{cool} = (\kappa_{hp} \sigma_{hp} n c)^{-1}$$

$$t_{dec} = \tau_h E_h / m_h$$

Hadronic Cooling & Decay

■ Characteristic Timescales

$$t_{cool} = (\kappa_{hp} \sigma_{hp} n c)^{-1}$$

$$t_{dec} = \tau_h E_h / m_h$$

■ Cooling Factor

Hadronic Cooling & Decay

■ Characteristic Timescales

$$t_{cool} = (\kappa_{hp} \sigma_{hp} n c)^{-1}$$

$$t_{dec} = \tau_h E_h / m_h$$

■ Cooling Factor

$$C = 1 - \exp\left(-\frac{t_{cool}}{t_{dec}}\right)$$

Hadronic Cooling & Decay

■ Characteristic Timescales

$$t_{cool} = (\kappa_{hp} \sigma_{hp} n c)^{-1}$$

$$t_{dec} = \tau_h E_h / m_h$$

■ Cooling Factor

$$C = 1 - \exp\left(-\frac{t_{cool}}{t_{dec}}\right)$$

■ Optical Depth

Hadronic Cooling & Decay

■ Characteristic Timescales

$$t_{cool} = (\kappa_{hp} \sigma_{hp} n c)^{-1}$$


$$t_{dec} = \tau_h E_h / m_h$$


■ Cooling Factor

$$C = 1 - \exp\left(-\frac{t_{cool}}{t_{dec}}\right)$$

■ Optical Depth

$$O = \kappa_{pp} \sigma_{pp} n d$$







Produced Hadron Spectrum

■ Modification Factors

Produced Hadron Spectrum

■ Modification Factors

$$\frac{dN_h}{dE_h} = CO \int_{E_h}^{\infty} dE_p \frac{dN_p}{dE_p} F_{pp \rightarrow h}$$

Produced Hadron Spectrum

■ Modification Factors

$$\frac{dN_h}{dE_h} = CO \int_{E_h}^{\infty} dE_p \frac{dN_p}{dE_p} F_{pp \rightarrow h}$$

Produced Hadron Spectrum

■ Modification Factors

$$\frac{dN_h}{dE_h} = CO \int_{E_h}^{\infty} dE_p \frac{dN_p}{dE_p} F_{pp \rightarrow h}$$

Produced Hadron Spectrum

■ Modification Factors

$$\frac{dN_h}{dE_h} = CO \int_{E_h}^{\infty} dE_p \frac{dN_p}{dE_p} F_{pp \rightarrow h}$$

■ Temporal Evolution


Produced Hadron Spectrum


■ Modification Factors

$$\frac{dN_h}{dE_h} = CO \int_{E_h}^{\infty} dE_p \frac{dN_p}{dE_p} F_{pp \rightarrow h}$$

■ Temporal Evolution

$$CO \propto t^{-3} (1 - e^{-t^3})$$







Resulting Neutrino Spectrum

- Complex Three Body Decay



Resulting Neutrino Spectrum

- Complex Three Body Decay
- Simple Two Body Decay

Resulting Neutrino Spectrum

- Complex Three Body Decay
- Simple Two Body Decay

$$\lambda_h = m_\mu^2 / m_h^2$$

Resulting Neutrino Spectrum

- Complex Three Body Decay
- Simple Two Body Decay

$$\lambda_h = m_\mu^2 / m_h^2$$

$$F_{h \rightarrow \nu} = B_{h \rightarrow \nu} E_h^{-1} (1 - \lambda_h)^{-1}$$

Resulting Neutrino Spectrum

- Complex Three Body Decay
- Simple Two Body Decay

$$\lambda_h = m_\mu^2 / m_h^2$$

$$F_{h \rightarrow \nu} = B_{h \rightarrow \nu} E_h^{-1} (1 - \lambda_h)^{-1}$$

- Hadron Number Convolution

Resulting Neutrino Spectrum


- Complex Three Body Decay
- Simple Two Body Decay


$$\lambda_h = m_\mu^2 / m_h^2$$

$$F_{h \rightarrow \nu} = B_{h \rightarrow \nu} E_h^{-1} (1 - \lambda_h)^{-1}$$


- Hadron Number Convolution

$$\frac{dN_\nu}{dE_\nu} = \int_{E_\nu}^{\infty} dE_h \frac{dN_h}{dE_h} F_{h \rightarrow \nu}$$









Normalized Magnetar Charm Contributions



Normalized Magnetar Neutrino Flux & Fluence





Published Estimated Magnetar Neutrinos




J. A. Carpio et al. *Phys. Rev. D* (2020)


Modified Magnetar Neutrino Flux & Fluence




Modified Magnetar Charm Contributions





Normalized AGN Charm Contributions




Normalized AGN Neutrino Fluence








Simulated Magnetar Formation





K. Hayashi *MPI Grav. Phys.* (2024)

Accretion Disk Structure




V. V. Zhuravlev et al. *ApJ* (2014)







Monte Carlo Event Generation



Data Courtesy
M. Schroller

Differential Cross Sections Comparison




Derived Parametrization S. R. Kelner et al. *Phys. Rev. D* (2006)

Event Data (SIBYLL 2.3c) Courtesy M. Schroll




Summary & Conclusion

Messenger Types & Detection




P. Mészáros et al.
Nat. Rev. Phys. (2019)

Source Regions & Events





P. Mészáros et al.
Nat. Rev. Phys. (2019)

Resistive Pulsar Magnetospheres



J. Li et al. *ApJ* (2012)

Accretion Disk & Corona



A. Müller Heidelberg University (2004)