



MASTER IN ASTROPHYSICS
AND SPACE SCIENCE



TOR VERGATA
UNIVERSITÀ DEGLI STUDI DI ROMA

Optimisation & Monte Carlo Sampling

INTRODUCTION TO NUMERICAL METHODS FOR ASTROPHYSICS

Fritz Ali Agildere

November 27, 2025

Overview

Probability Refresher

Definitions & Notation

Random Numbers

Optimisation

Common Methods

Problems

Markov Chain Monte Carlo

Popular Algorithms

Tuning & Diagnosing

Alternatives

Overview

Probability Refresher

Definitions & Notation

Random Numbers

Optimisation

Common Methods

Problems

Markov Chain Monte Carlo

Popular Algorithms

Tuning & Diagnosing

Alternatives

Frequentist View

- *With Prior Knowledge*: out of all n different but equally likely possible outcomes of an event, there are k that have a property A which is therefore measured with probability

$$P(A) = \frac{k}{n}$$

- *Without Prior Knowledge*: out of n independent measurements the outcome A occurs k times, defining

$$P(A) = \lim_{n \rightarrow \infty} \frac{k}{n}$$

Bayesian View

- $P(A|B)$ quantifies the plausibility of an assumption A under known information given by B
- *Kolmogorov Axioms* underpin the rules for calculating with probabilities
- $P(A, B) = P(A) P(B|A)$ quantifies the probability of measuring A and B simultaneously, and from assuming $P(A, B) = P(B, A)$ we find

Bayes' Theorem

$$P(A|B) = \frac{P(B|A) P(A)}{P(B)}$$

Continuous Distributions

■ *Cumulative Distribution Function:* $F(x) = P(X \leq x)$

■ *Probability Density Function:* $f(x) = \frac{d}{dx} F(x)$

$$P(a \leq X \leq b) = \int_a^b f(x) dx = F(b) - F(a)$$

■ *Normalization:* $1 = \int_{-\infty}^{+\infty} f(x) dx$ $\lim_{x \rightarrow -\infty} F(x) = 0$ $\lim_{x \rightarrow +\infty} F(x) = 1$

Expectation Values

$$E[h(x)] = \int_{-\infty}^{+\infty} h(x) f(x) dx$$

■ *Mean* or first raw moment: $\mu = E[X] = \int_{-\infty}^{+\infty} x f(x) dx$

■ *Variance* or second raw moment: $\sigma^2 = E[(X - E[X])^2] = \int_{-\infty}^{+\infty} (x - E[x])^2 f(x) dx$

Overview

Probability Refresher

Definitions & Notation

Random Numbers

Optimisation

Common Methods

Problems

Markov Chain Monte Carlo

Popular Algorithms

Tuning & Diagnosing

Alternatives

Drawing Numbers

$$x \sim U(a, b)$$

$$y \sim N(\mu, \sigma)$$

■ *Uniform Distribution:* $U(a, b) = f(x|a, b) = \begin{cases} \frac{1}{b-a} & a \leq x \leq b \\ 0 & \text{otherwise} \end{cases}$

■ *Gaussian Distribution:* $N(\mu, \sigma) = f(y|\mu, \sigma) = \frac{1}{\sqrt{2\pi}\sigma} \exp\left(-\frac{1}{2}\left(\frac{y-\mu}{\sigma}\right)^2\right)$

Number Generation

- *Pseudo Random Numbers: $x_{j+1} = g(x_1, \dots, x_j)$*
 - deterministic & reproducible
 - efficient & transparent
 - almost indistinguishable with long periods of recurrence
- *Equal Distribution Generators:*
 - *Linear Congruential*
 - *Linear Feedback Shift*
 - *Permutated Congruential*
- *Arbitrary Distributions:* obtained from uniformly distributed numbers by various methods

Overview

Probability Refresher

Definitions & Notation

Random Numbers

Optimisation

Common Methods

Problems

Markov Chain Monte Carlo

Popular Algorithms

Tuning & Diagnosing

Alternatives

Motivation

- *Model Optimisation*: finding the best possible model parameters
 - maximising objective function (*Fitting*)
 - minimising loss function (*Machine Learning*)
- *Example*: modified blackbody spectrum for galactic dust emission (M. Juvela, A&A 2023)

$$I_{\nu}(T, \beta) \equiv \frac{\nu^{3+\beta}}{\exp\left(\frac{h\nu}{kT}\right) - 1}$$

Example

$$I_{\nu}(T, \beta) = \frac{\nu^{3+\beta}}{\exp\left(\frac{h\nu}{kT}\right) - 1}$$

- *Objective:* optimise $\theta \equiv (T, \beta)$ such that the weighted squared error is minimal
- *Assumptions:*
 - independent measurements generated by $I_{\nu}(\theta)$ with normally distributed uncertainty
 - infinitesimally small error in ν so that we only need σ_{ν} in I_{ν} direction
 - suppose we have no prior information on the parameter distribution

Example

- *Likelihood:*

$$-\log L(I_v | v, \sigma_v, \theta) \propto \chi^2(\theta) = \sum_{i=1}^N \frac{(I_{v,i} - I_v(\theta))^2}{\sigma_{v,i}^2}$$


- *Posterior:*

$$\log P(\theta | I_v, v, \sigma_v) \propto -\chi^2(\theta)$$

- *Objective:* minimise the $\chi^2(\theta)$ measure to maximise the posterior probability

Measurements

How do we proceed from this dataset to maximise the likelihood for our nonlinear model?



Overview

Probability Refresher

Definitions & Notation

Random Numbers

Optimisation

Common Methods

Problems

Markov Chain Monte Carlo

Popular Algorithms

Tuning & Diagnosing


Alternatives

Parameter Grid

- *Idea:* in our $D = 2$ dimensional parameter space, we can create an $N \times N$ grid of (T, β) points and evaluate the likelihood at each one
- *Preparation:* for this as well as some of the following methods, it is a good idea to normalize the variation of each individual parameter
 - from the literature we expect $(T, \beta) \in [10, 25] \times [1.5, 2.0]$ as constraints
 - resize these intervals by defining $T \mapsto \tilde{T} = T / 30$ with $I_v(T, \beta) \rightarrow I_v(\tilde{T}, \beta)$
- *Problem:* as D increases we have N^D gridpoints, and with more complex parameter distributions, this procedure quickly becomes intractable due to computational limits

Parameter Grid

Can we think of more elegant ways to explore even very high dimensional parameter spaces?



General Case

- *Optimisation Algorithms*: intelligently move through parameter space towards optimum
- *Common Attributes*:
 1. choose initialisation point
 2. evaluate objective function or its derivatives to estimate local curvature
 3. propose new step in parameter space
 4. decide whether proposal is a good step and update position accordingly
 5. repeat until some predefined cutoff is satisfied
- *Interactive Visualisations*: check out Ben Frederickson¹ and Andy Casey² to see some nice examples

Nelder-Mead

- *Requirements*: only objective function, no information on gradients
- *Algorithm*:
 1. initialize simplex of $d + 1$ points in d dimensional parameter space
 2. rank points in order of values $f(x_1) \leq \dots \leq f(x_d) \leq f(x_{d+1})$
 3. compute centroid x_0 of $x_1, \dots, x_d$
 4. perform a bunch of conditional evaluations between x_0 and x_{d+1}
 5. reflect, expand, contract or shrink based on results to move points downhill and repeat
- *Notes*:
 - extrapolates function to slowly move geometric blob through parameter space
 - zeroth order method since no derivatives are used

Random Descent

- *Requirements:* only objective function, no information on gradients
- *Algorithm:*
 1. start from arbitrary initial value x_0
 2. draw a proposed shift from some chosen probability distribution $X \sim g(x)$
 3. propose updated position $\hat{x}_{k+1} = x_k + X$
 4. accept proposal $x_{k+1} = \hat{x}_{k+1}$ only if $f(\hat{x}_{k+1}) < f(x_{k+1})$ and repeat
- *Notes:*
 - traverses parameter space via random walk, can be very slow
 - zeroth order method since no derivatives are used
 - improve by considering last step and optimizing step range

Gradient Descent

- *Requirements:* first derivatives of the objective function, can be estimated numerically
- *Algorithm:*
 1. initialize at some x_0
 2. update current point with $x_{k+1} = x_k - \alpha \nabla f(x_k)$
 3. iterate until convergence $|x_{k+1}/x_k - 1| < \epsilon$ or maximum step count is reached
- *Notes:*
 - moves along direction of steepest descent at each point with fixed learning rate α
 - first order method since only the gradient is used
 - improve by performing line search for α_k at every step

Backtracking Line Search

- *Requirements:* first derivatives of the objective function, can be estimated numerically
- *Algorithm:*
 1. start from some fixed large α_0 for each step k of the gradient descent
 2. set $\alpha_{j+1} = \tau \alpha_j$ until $f(x_k) - f(x_k - \alpha_j \nabla f(x_k)) \geq \kappa \alpha_j (\nabla f(x_k))^2$ where $\tau, \kappa \in (0, 1)$
 3. use last value to set learning rate $\alpha_k = \alpha_j$ for next step
- *Notes:*
 - ensures $f(x_{k+1}) < f(x_k)$ such that the function steadily decreases with increasing k
 - favours secant between x_k and x_{k+1} steeper or almost as steep as tangent $\nabla f(x_k)$
 - robust convergence guarantees, larger step size for flat areas avoids saddle points
 - first order method since only the gradient is used
 - typical settings are $\tau = \kappa = 0.5$ (L. Armijo, *Pac. J. Math.* 1966)

Thank You!