

Characterization of Galaxies via Optical and Infrared Surveys

Fritz Ali Agildere

fritz.agildere@udo.edu

Department of Astronomy, Faculty of Mathematics, University of Belgrade

May 31, 2026

Abstract

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Ut purus elit, vestibulum ut, placerat ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetur id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.

1 Introduction

With ongoing and upcoming surveys providing ever larger amounts of data, the task of object classification is becoming increasingly relevant. In this work, we will summarize and compare select methods for characterizing narrow emission line galaxies and the extraction of their physical properties. For all steps in the following sections, we use data from the latest SDSS data release and match those to spectra obtained by WISE for the same objects (Wright et al. 2010, Kollmeier et al. 2026). Information to reproduce these results are included in the appendix.

2 Optical & Infrared Classification

Generally, the broad labeling of galaxies as Active Galactic Nuclei (AGN) or Star Formation (SF) dominated types relies on reconstructing the excitation and ionization mechanisms from emission lines, and determining whether these match the thermal continuum of stellar populations or the strongly nonthermal radiation produced in AGN powered by Super Massive Black Holes (SMBH) in their centers. Approaches analyzing the continuum emission provide complementary insights, and by probing different wavelengths we can peer through obscuring dust and gas to unveil the true underlying physics. Now, we will explore some famous examples.

2.1 BPT Diagram

By plotting the ratio of emission line fluxes for [OIII] at 5007 Å to Hβ at 4863 Å versus [NII] at 6583 Å to Hα at 6565 Å in their work, Baldwin, Phillips, and Terlevich draw a diagram that sorts galaxies into a distinctive winged shape (Baldwin et al. 1981). Due to lacking the energies to ionize heavier atoms, SF types tend towards the lower left of the image, while powerful AGN fall into the upper right tip. By choosing emission lines close to one another, this framework automatically cancels out reddening differences as well as calibration errors. Depicted in Figure 1 is our entire dataset as a scatter plot that has been shaded based on the density




Figure 1: Standard BPT diagram and class separation curves (Kewley et al. 2001, Kauffmann et al. 2003).

of nearby points. This convention is adopted for all such diagrams in this work for better interpretability of the data. Together with this, standard theoretical (Kewley et al. 2001) and empirical (Kauffmann et al. 2003) curves to discriminate between AGN and SF classes are shown.




Figure 2: Colored BPT diagram with derived class labels.

We label points between the lines as COMP for composite types and display the classes in Figure 2 encoded by color. Additionally, we have access to the already processed SDSS labels which we can similarly plot in the BPT diagram. This is done in Figure 3 for the AGN and SF as well as starburst SB subclasses. Comparing to the previous plot gives a first visual impression of their cross method agreement, which for the AGN wing seems acceptable, but will be evaluated more rigorously in the coming paragraphs.




Figure 3: Colored BPT diagram based on SDSS labels.

2.2 WHAN Diagram

Next up is the WHAN diagram, which plots the equivalent width $W_{H\alpha}$ of the $H\alpha$ line against the ratio of [NII] to $H\alpha$ fluxes (Cid Fernandes et al. 2011). This width quantifies line strengths in terms of the underlying continuum and is thereby independent of measurement effects. The concept behind a classification using the measure is that based on the prominence of lines above the background radiation, we can estimate the relative photon budget of the exciting or ionizing radiation field. In true AGN and SF galaxies, this value is high, whereas retired galaxies with old stellar populations have less pronounced emission lines. This is a distinction which the BPT classifier fails to take into account and can therefore improve our label confidence.




Figure 4: Raw data in WHAN diagram with separators.




Figure 5: Colored WHAN diagram with derived labels.

With the raw scattered data, Figure 4 includes discrimination lines. According to these, any galaxy with $W_{H\alpha} < 3.0 \text{ \AA}$ is classified as rGAL or retired. Points above this value with $\log_{10}([NII] / H\alpha) > -0.4$ are labeled AGN or else pSF for pure star formation. Within the active class, we also include the distinction for strong sAGN if $W_{H\alpha} > 6.0 \text{ \AA}$ and weak wAGN otherwise. There is further a category defined via $W_{H\alpha} < 0.5 \text{ \AA}$ and $W_{[NII]} < 0.5 \text{ \AA}$ for passive, actually lineless galaxies, but based on our selection, none fall into this class. Figure 5 shows the resulting categories, while Figure 6 presents the more restrictive SDSS classes, which show a smaller AGN region than what WHAN includes.




Figure 6: Colored WHAN diagram based on SDSS labels.

2.3 Category Mixing

In addition to the predefined SDSS label, we can check the mixing between BPT and WHAN classes. For this, Figure 7 displays WHAN colors in the BPT diagram, and Figure 8 the opposite. Here we can see how BPT conditions do not account for weak equivalent widths, making WHAN a more nuanced model in this regard. On the other hand, WHAN labels many objects AGN that sit far down in the BPT scale, indicating that information on the ionization potential of the source gets lost.




Figure 7: Colored BPT diagram based on WHAN labels.




Figure 8: Colored WHAN diagram based on BPT labels.

2.4 WISE Colors

For sources shrouded by dust, optical classifier such as BPT and WHAN can fail. Instead, infrared data from missions such as WISE with filters labeled W1 at $3.4\mu\text{m}$ and W2 at $4.6\mu\text{m}$ as well as W3 at $12\mu\text{m}$ center wavelengths can be used. In these scenarios, galaxies dominated by their stars appear in the more energetic bands corresponding to stellar atmospheres, while an AGN torus absorbs shorter wavelengths, leading to smaller frequencies characterizing their spectrum. In the typical WISE colors plot, we draw the differences of the respective magnitudes, equivalent to the logarithm of the flux ratio and therefore independent of true luminosity. In this diagram, we can define a region where most AGN fall, referred to as the wedge from here on (Mateos et al. 2012). Our data is visualized and classified according to this wedge in Figure 9 with schematic classes from (Wright et al. 2010) taken as a backdrop. Notably, there are only few objects being labeled as AGN by this method. For the actual statistics and a more robust comparison of the classifiers, the following subsection calculates the pairwise binary cross correlation and gives the raw numbers for our matched dataset.

2.5 Evaluating Diagnostics

Listed in Table 1 are the numbers for our processed dataset. To better compare the performance between the different classifiers, we compute the so called Matthews Correlation Coefficient (MCC) for each pair. This metric measures how the matching quality compares to a pure chance baseline. A value of 1 means perfect agreement, while 0 implies no advantage over randomly assigning an answer, and negative values signify worse than random anticorrelation between methods. It also corrects for unbalanced population sizes, as is the case for our sample in which AGN are far fewer than other types of galaxies. Figures 10 and 11 show our results for two cases of BPT classifier, one strict which restricts positives to AGN objects only, and one relaxed that also includes COMP labels.

Table 1: List of overall and relative positive AGN labels by method. In total, 10 000 unique SDSS objects were used, of which 9799 had matches in the WISE database with a $3''$ tolerance.

Method	Total	Fraction
SDSS	585	5.9 %
BPT (strict)	920	9.2 %
BPT (relaxed)	2491	24.9 %
WHAN	3464	34.6 %
WISE	88	0.9 %
SDSS & WISE	37	0.4 %
BPT (strict) & WISE	47	0.5 %
BPT (relaxed) & WISE	64	0.6 %
WHAN & WISE	58	0.6 %

We recognize that generally, optical methods agree with amongst each other to a higher degree than with infrared based results. On the other hand, agreement between WISE labels is barely better than random, especially for the WHAN results. For the strict BPT case, it matches best with SDSS classes, while the relaxed version shows better agreement with WHAN labels.

SDSS	1.000	0.680	0.301	0.145
BPT	0.680	1.000	0.271	0.144
WHAN	0.301	0.271	1.000	0.062
WISE	0.145	0.144	0.062	1.000
	SDSS	BPT	WHAN	WISE

Figure 10: Matrix of MCC results with strict BPT method.




Figure 9: Shaded scatter plot of data in WISE color diagram. Included is the wedge discriminator (Mateos et al. 2012) as well as the background adapted from (Wright et al. 2010) with shortened object type labels.

Based on this, we will proceed in the BPT framework for the following section and deliver SDSS and WHAN as well as WISE results for comparison in the appendix. There are more powerful statistical diagnostics that enable a detailed view of method performance, but these are omitted in this work due to its limited scope. Furthermore, sophisticated classifiers are likely able to achieve physically more sound results, though this comes at the cost of higher complexity.

3 Physical Properties

Other than classifying objects, we can also extract estimates of their physical properties from spectral data. The following subsections will outline this process for gas density and velocity, as well as derive the central Black Hole (BH) mass for our binned dataset.

3.1 Gas Density

First, we use the ratio of the two [NII] lines at 6716 Å and 6731 Å in our data. Since these both originate from a closely spaced doublet, their relative occurrence is independent of gas temperature. Instead, at low densities the ratio is fixed by their degeneracies as statistical weights, whereas at high densities, collisional deexcitation becomes relevant, which does not emit photons. Due to the lower wavelength state having a longer lifetime, it gets affected more strongly by this mechanism, dropping the ratio and allowing us to reconstruct the local electron density as a proxy for the gas (Sanders et al. 2016, Joh et al. 2021).

3.2 Gas Velocity

3.3 Black Hole Mass

3.4 Data Analysis

SDSS	1.000	0.433	0.301	0.145
BPT	0.433	1.000	0.631	0.104
WHAN	0.301	0.631	1.000	0.062
WISE	0.145	0.104	0.062	1.000
	SDSS	BPT	WHAN	WISE

Figure 11: Matrix of MCC results with relaxed BPT method.




Figure 12: Binned BPT data showing electron density.




Figure 15




Figure 13




Figure 16




Figure 14




Figure 17




Figure 18




Figure 19




Figure 20




Figure 21




Figure 22




Figure 23




Figure 24




Figure 25




Figure 26




Figure 27

4 Conclusion

References

- Baldwin, J. A. et al. *Publications of the Astronomical Society of the Pacific* (1981)
Cid Fernandes, R. et al. *Monthly Notices of the Royal Astronomical Society* (2011)
Joh, K. et al. *Publications of the Astronomical Society of Japan* (2021)
Kauffmann, G. et al. *Monthly Notices of the Royal Astronomical Society* (2003)
Kewley, L. J. et al. *The Astrophysical Journal* (2001)
Kollmeier, J. A. et al. *The Astronomical Journal* (2026)
Mateos, S. et al. *Monthly Notices of the Royal Astronomical Society* (2012)
Sanders, R. L. et al. *The Astrophysical Journal* (2016)
Wright, E. L. et al. *The Astronomical Journal* (2010)

Appendix

```

1 SELECT TOP 10000
2     g.sii_6717_flux, g.sii_6731_flux, g.nii_6584_flux, g.oi_6300_flux, g.oiii_5007_flux,
3     g.h_alpha_flux, g.h_beta_flux,
4     g.h_alpha_eqw, g.nii_6584_eqw, g.oiii_sigma,
5     w.w1mpro, w.w2mpro, w.w3mpro,
6     s.z, s.class, s.subclass, s.bestobjid
7 FROM SpecObj AS s
8 JOIN galSpecLine AS g ON s.specobjid = g.specobjid
9 JOIN PhotoTag AS p ON s.bestobjid = p.objid
10 OUTER APPLY ( SELECT TOP 1 x.wise_cntr
11                FROM wise_xmatch AS x
12                WHERE p.objid = x.sdss_objid AND x.match_dist < 3
13                ORDER BY x.match_dist ASC
14            ) AS y
15 LEFT JOIN wise_allsky AS w ON y.wise_cntr = w.cntr
16 WHERE s.class = 'GALAXY'
17 AND s.subclass != 'BROADLINE'
18 AND s.subclass != 'AGN BROADLINE'
19 AND s.subclass != 'STARFORMING BROADLINE'
20 AND s.subclass != 'STARBURST BROADLINE'
21 AND s.snmedian_r > 5.0
22 AND s.z < 0.35
23 AND g.sii_6717_eqw < 0.0
24 AND g.sii_6731_eqw < 0.0
25 AND g.nii_6584_eqw < 0.0
26 AND g.oi_6300_eqw < 0.0
27 AND g.oiii_5007_eqw < 0.0
28 AND g.h_alpha_eqw < 0.0
29 AND g.h_beta_eqw < 0.0
30 AND (2.355 * g.sigma_forbidden) < 500
31 AND (2.355 * g.sigma_balmer) < 500
32 AND g.sii_6717_flux > 3 * g.sii_6717_flux_err
33 AND g.sii_6731_flux > 3 * g.sii_6731_flux_err
34 AND g.nii_6584_flux > 3 * g.nii_6584_flux_err
35 AND g.oi_6300_flux > 3 * g.oi_6300_flux_err
36 AND g.oiii_5007_flux > 3 * g.oiii_5007_flux_err
37 AND g.h_alpha_flux > 3 * g.h_alpha_flux_err
38 AND g.h_beta_flux > 3 * g.h_beta_flux_err
39 ORDER BY s.specobjid ASC

```