

X-Ray Emitting Plasma in Galaxies and Galaxy Clusters

Fritz Ali Agildere

fritz.agildere@udo.edu

Department of Astronomy, Faculty of Mathematics, University of Belgrade
Astronomical Observatory of Belgrade

May 28, 2026

Abstract

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Ut purus elit, vestibulum ut, placerat ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetur id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.

1 Introduction

In studying the evolving nature of our universe on the largest scales, galaxy clusters represent the most expansive clearly defined structures in which matter approaches virialization. Due to this fact, we can use them as laboratories to study the cosmic growth rate and characterize density fluctuations, as well as reconstruct the dynamical history of merging systems and the effects of feedback from active members. A tenuous plasma called the Intracluster Medium (ICM) permeates these massive objects, with temperatures in the 10^7 K range and densities from 10^{-5} to 10^{-1} cm^{-3} between their outer regions and inner cores. In this hot state, the ICM radiates at several keV photon energies, making spectroscopy in the soft X-ray band a primary method to resolve the distribution of baryonic matter inside clusters (Böhringer & Werner 2010).

1.1 Models

At the broadest level, galaxy clusters are modeled as dark matter halos fully governed by gravitational collapse. They are thought to originate from primordial density fluctuations and grow via cluster mergers as well as by accreting matter from cosmic filaments. Their total mass is dominated by the dark component defined through its purely gravitational interactions, the density of which is well described using a Navarro-Frenk-White (NFW) profile. Most baryonic matter is not contained in the stellar mass of the member galaxies themselves but in the ICM gas, which follows a similar shape to the overall matter distribution in the outer parts where it usually makes up a fraction of roughly $f_g \sim 15\%$ depending on redshift, but flattens towards the central region due to counteracting pressures. Containing this material requires a sufficiently deep gravitational potential well that is provided by the high dark matter concentration, which strongly peaks in the center as collisional effects are negligible. Shown in Figure 1 are exemplary radial density profiles with realistic parameter choices for the observed galaxy cluster categories as explained in the following subsections (Pratt et al. 2019).




Figure 1: Typical mass density profiles for NCC and CC galaxy clusters are plotted as proxies for all baryonic matter. Shown as well is a standard NFW total density curve that is dominated by a collisionless dark matter component.

Adopting this view, radiative cooling of the higher density inner regions through X-ray emission generally leads to a drop in thermal pressure and a resulting influx of subsonic materials from the surrounding medium, called a cooling flow. This assumption predicts massive rates of molecular condensation near the central galaxies which should produce significant starburst activity. However, observations by the *Chandra* (Weisskopf et al. 2000) and *XMM-Newton* (Jansen et al. 2001) missions proved that this signature is not present in reality. Instead, the ICM cools towards the core but does not reach the neutral gas threshold. The leading hypothesis for the culprit of this deviation from theoretical expectations is the injection of energy by Active Galactic Nucleus (AGN) feedback phases regulated by a balancing of heating and fueling, although this description carries with it its own set of open questions, namely the mechanisms for isotropization of highly directional jets as well as stabilization and dissipation of the AGN bubbles (Böhringer & Werner 2010, Rosati et al. 2002, Pratt et al. 2019).




Figure 2: Selection of galaxy clusters used as examples in this work. Displayed here are composite overlays of second generation *Digitized Sky Survey* red band images and contours for *ROSAT All Sky Survey (RASS)* broad band count rates (Truemper 1982, Lasker 1994). Both optical and X-ray data are stretched using a simple 0.8 exponent power law. Each frame has a radius of $15'$ for its field of view. Refer to Figure 9 for raw *ROSAT* counts underlying the contours.

1.2 Classes

According to the spherical dark matter halo assumption, gas properties should follow selfsimilar laws along the radial distance to the cluster center and only scale with the total mass and redshift. Comparing to observations such as those presented in Figures 7 and 8 shows that this holds quite well for large radii but breaks down closer to the central regions. It turns out that based on these differences, galaxy clusters can be sorted into two categories. In order to do so, it is useful to introduce the cooling time. It derives from the total thermal energy $E_t \propto nT$ and volumetric cooling rate $\dot{E}_c \propto n^2 \Lambda_c(T)$ where the cooling function for bremsstrahlung scales as $\Lambda_c(T) \propto T^{1/2}$ to define a proportionality

$$t_c = \frac{E_t}{\dot{E}_c} \propto T^{1/2} n^{-1}$$

for the ICM properties. If this timescale is small $t_c < 1$ Gyr in its center, we call it a Cooling Core (CC) cluster. These systems radiate away heat efficiently, leading to a rapid drop in gas temperature and a corresponding peak in density. With these characteristics, they also have an almost critically low entropy floor, close to the catastrophic cooling flow collapse scenario and only barely stabilized by galactic winds from star formation and AGN feedback. Generally, the population

of CC clusters comprises dynamically relaxed systems with a high degree of order. In the opposite case, cooling is highly inefficient with $t_c > 10$ Gyr and frequently exceeding the Hubble time. Such clusters are labeled Noncool Core (NCC) and tend to possess isothermal or monotonically increasing temperatures and flat density profiles towards their centers. Their inner entropy floors are large and likely originate from merging of subclusters. Heating occurs through supersonic shocks, turbulence, contact discontinuities and sloshing of the central material.

From this follows that analysis of the thermal structure in principle allows us to infer the recent dynamical history of clusters. Additionally, measuring the total mass M inside some characteristic radius r makes it possible to map the distribution of large scale density inhomogeneities and their evolution throughout cosmological time. After determining plasma properties and by assuming hydrostatic equilibrium, we can rearrange the well known formula

$$dP/dr = g\rho = GM\rho/r^2$$

to solve for mass. A common choice for a universal distance normalization is the overdensity radius R_ℓ which constrains the sphere around the cluster center inside which the average mass density is a factor ℓ larger than the surrounding critical

density of the universe at that redshift. This way, contained masses are labeled M_ℓ and average quantities T_ℓ or P_ℓ as well as astrophysical entropy K_ℓ with $K \propto Tn^{-2/3}$ can be defined for comparison between systems of vastly different sizes. Typical values are $\ell = 200$ since R_{200} is close to the theoretical virialization radius, as well as $\ell = 500$ and R_{500} as the boundary up to which current X-ray telescopes can produce reliable measurements (Böhringer & Werner 2010, Pratt et al. 2019).

1.3 Examples

To have a consistent set of objects to compare theoretical predictions to real observations for this work, we select six clusters. Their optical appearance as well as X-ray surface brightness contours are depicted in Figure 2 with Table 1 listing their characterizing properties.

Table 1: Parameters for our galaxy cluster sample. Values given are those determined for the *X-COP* systems (Eckert et al. 2017).

Name	Class	z	R_{500} / kpc	$M_{500} / M_\odot$
A85	CC	0.0555	1200	5.6×10^{14}
A426	CC	0.0179	1300	5.9×10^{14}
A1644	NCC	0.0473	2100	3.5×10^{14}
A1656	NCC	0.0231	1300	6.0×10^{14}
A2029	CC	0.0766	1400	8.6×10^{14}
A3158	NCC	0.0590	1100	4.3×10^{14}

The Perseus or Abell 426 as well as the massive Abell 2029 clusters are archetypical examples of relaxed CC systems supported by regulated feedback loops, although there are still remnants of ancient disruptions detectable. Abell 85 also falls in the CC category but shows more prominent central sloshing as well as active accretion from intersecting cosmic filaments. The Coma or Abell 1656 cluster on the other hand is a prototype for NCC systems that have been disrupted by a recent massive merger event. Similarly, high resolution observations classify Abell 3158 as NCC despite its smooth X-ray surface. Lastly, Abell 1644 is clearly an NCC cluster as is readily apparent from its bimodal halo structure (Böhringer & Werner 2010, Eckert et al. 2017).

2 Radiation Mechanisms

Nulla malesuada porttitor diam. Donec felis erat, congue non, volutpat at, tincidunt tristique, libero. Vivamus viverra fermentum felis. Donec nonummy pellentesque ante. Phasellus adipiscing semper elit. Proin fermentum massa ac quam. Sed diam turpis, molestie vitae, placerat a, molestie nec, leo. Maecenas lacinia. Nam ipsum ligula, eleifend at, accumsan nec, suscipit a, ipsum. Morbi blandit ligula feugiat magna. Nunc eleifend consequat lorem. Sed lacinia nulla vitae enim. Pellentesque tincidunt purus vel magna. Integer non enim. Praesent euismod nunc eu purus. Donec bibendum quam in tellus. Nullam cursus pulvinar lectus. Donec et mi. Nam vulputate metus eu enim. Vestibulum pellentesque felis eu massa.

Quisque ullamcorper placerat ipsum. Cras nibh. Morbi vel justo vitae lacus tincidunt ultrices. Lorem ipsum dolor sit

amet, consectetur adipiscing elit. In hac habitasse platea dictumst. Integer tempus convallis augue. Etiam facilisis. Nunc elementum fermentum wisi. Aenean placerat. Ut imperdiet, enim sed gravida sollicitudin, felis odio placerat quam, ac pulvinar elit purus eget enim. Nunc vitae tortor. Proin tempus nibh sit amet nisl. Vivamus quis tortor vitae risus porta vehicula.

Fusce mauris. Vestibulum luctus nibh at lectus. Sed bibendum, nulla a faucibus semper, leo velit ultricies tellus, ac venenatis arcu wisi vel nisl. Vestibulum diam. Aliquam pellentesque, augue quis sagittis posuere, turpis lacus congue quam, in hendrerit risus eros eget felis. Maecenas eget erat in sapien mattis porttitor. Vestibulum porttitor. Nulla facilisi. Sed a turpis eu lacus commodo facilisis. Morbi fringilla, wisi in dignissim interdum, justo lectus sagittis dui, et vehicula libero dui cursus dui. Mauris tempor ligula sed lacus. Duis cursus enim ut augue. Cras ac magna. Cras nulla. Nulla egestas. Curabitur a leo. Quisque egestas wisi eget nunc. Nam feugiat lacus vel est. Curabitur consectetur.




Figure 3: Normalized bremsstrahlung continua for different temperatures over a range of magnitudes. Fitting the bend placement allows for the inference of local temperature.




Figure 4: Schematic view of the thermal Sunyaev-Zeldovich effect on the Cosmic Microwave Background (CMB) Planck spectrum with exaggerated $Y \sim 0.2$ Compton parameter in the nonrelativistic limit.

3 Instrumentation

Suspendisse vel felis. Ut lorem lorem, interdum eu, tincidunt sit amet, laoreet vitae, arcu. Aenean faucibus pede eu ante. Praesent enim elit, rutrum at, molestie non, nonummy vel, nisl. Ut lectus eros, malesuada sit amet, fermentum eu, sodales cursus, magna. Donec eu purus. Quisque vehicula, urna sed ultricies auctor, pede lorem egestas dui, et convallis elit erat sed nulla. Donec luctus. Curabitur et nunc. Aliquam dolor odio, commodo pretium, ultricies non, pharetra in, velit. Integer arcu est, nonummy in, fermentum faucibus, egestas vel, odio.

Sed commodo posuere pede. Mauris ut est. Ut quis purus. Sed ac odio. Sed vehicula hendrerit sem. Duis non odio. Morbi ut dui. Sed accumsan risus eget odio. In hac habitasse platea dictumst. Pellentesque non elit. Fusce sed justo eu urna porta tincidunt. Mauris felis odio, sollicitudin sed, volutpat a, ornare ac, erat. Morbi quis dolor. Donec pellentesque, erat ac sagittis semper, nunc dui lobortis purus, quis congue purus metus ultricies tellus. Proin et quam. Class aptent taciti sociosqu ad litora torquent per conubia nostra, per inceptos hymenaeos. Praesent sapien turpis, fermentum vel, eleifend faucibus, vehicula eu, lacus.

4 Spectral Characteristics

Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Donec odio elit, dictum in, hendrerit sit amet, egestas sed, leo. Praesent feugiat sapien aliquet odio. Integer vitae justo. Aliquam vestibulum fringilla lorem. Sed neque lectus, consectetur at, consectetur sed, eleifend ac, lectus. Nulla facilisi. Pellentesque eget lectus. Proin eu metus. Sed porttitor. In hac habitasse platea dictumst. Suspendisse eu lectus. Ut mi mi, lacinia sit amet, placerat et, mollis vitae, dui. Sed ante tellus, tristique ut, iaculis eu, malesuada ac, dui. Mauris nibh leo, facilisis non, adipiscing quis, ultrices a, dui.




Figure 5: Collection of *Chandra* binned spectra for various galaxy clusters (Weisskopf et al. 2000). Note the presence of emission lines, as well as possible hints of excesses in the relative count rates for both the soft and hard X-ray regimes.

4.1 Continuum

Morbi luctus, wisi viverra faucibus pretium, nibh est placerat odio, nec commodo wisi enim eget quam. Quisque libero justo, consectetur a, feugiat vitae, porttitor eu, libero. Suspendisse sed mauris vitae elit sollicitudin malesuada. Maecenas ultricies eros sit amet ante. Ut venenatis velit. Maecenas sed mi eget dui varius euismod. Phasellus aliquet volutpat odio. Vestibulum ante ipsum primis in faucibus orci luctus et ultrices posuere cubilia Curae; Pellentesque sit amet pede ac sem eleifend consectetur. Nullam elementum, urna vel imperdiet sodales, elit ipsum pharetra ligula, ac pretium ante justo a nulla. Curabitur tristique arcu eu metus. Vestibulum lectus. Proin mauris. Proin eu nunc eu urna hendrerit faucibus. Aliquam auctor, pede consequat laoreet varius, eros tellus scelerisque quam, pellentesque hendrerit ipsum dolor sed augue. Nulla nec lacus.

4.2 Line Emission

Suspendisse vitae elit. Aliquam arcu neque, ornare in, ullamcorper quis, commodo eu, libero. Fusce sagittis erat at erat tristique mollis. Maecenas sapien libero, molestie et, lobortis in, sodales eget, dui. Morbi ultrices rutrum lorem. Nam elementum ullamcorper leo. Morbi dui. Aliquam sagittis. Nunc placerat. Pellentesque tristique sodales est. Maecenas imperdiet lacinia velit. Cras non urna. Morbi eros pede, suscipit ac, varius vel, egestas non, eros. Praesent malesuada, diam id pretium elementum, eros sem dictum tortor, vel consectetur odio sem sed wisi.

5 Plasma Diagnostics

Etiam euismod. Fusce facilisis lacinia dui. Suspendisse potenti. In mi erat, cursus id, nonummy sed, ullamcorper eget, sapien. Praesent pretium, magna in eleifend egestas, pede pede pretium lorem, quis consectetur tortor sapien facilisis magna. Mauris quis magna varius nulla scelerisque imperdiet. Aliquam non quam. Aliquam porttitor quam a lacus. Praesent vel arcu ut tortor cursus volutpat. In vitae pede quis diam bibendum placerat. Fusce elementum convallis neque. Sed dolor orci, scelerisque ac, dapibus nec, ultricies ut, mi. Duis nec dui quis leo sagittis commodo.

Aliquam lectus. Vivamus leo. Quisque ornare tellus ullamcorper nulla. Mauris porttitor pharetra tortor. Sed fringilla justo sed mauris. Mauris tellus. Sed non leo. Nullam elementum, magna in cursus sodales, augue est scelerisque sapien, venenatis congue nulla arcu et pede. Ut suscipit enim vel sapien. Donec congue. Maecenas urna mi, suscipit in, placerat ut, vestibulum ut, massa. Fusce ultrices nulla et nisl.




Figure 6: Theoretical galaxy cluster radial profiles assuming universal selfsimilar scaling relations. *Top Left:* Electron number density (Vikhlinin et al. 2006). *Top Right:* Gas pressure (Arnaud et al. 2010). *Bottom Left:* Gas temperature (Vikhlinin et al. 2006). *Bottom Right:* Gas entropy (Cavagnolo et al. 2009). Compare to Figure 7 for real observations.




Figure 7: Sample of galaxy cluster radial profiles from *Chandra* observations (Weisskopf et al. 2000). *Top Left:* Electron number density. *Top Right:* Gas pressure. *Bottom Left:* Gas temperature. *Bottom Right:* Gas entropy. Compare to Figure 6 for a visualization of theoretical and empirical fits with realistic parameter choices.

6 Conclusion

nec, tempor congue, nulla. Duis viverra gravida mauris. Cras tincidunt. Curabitur eros ligula, varius ut, pulvinar in, cursus faucibus, augue.

Etiam ac leo a risus tristique nonummy. Donec dignissim tincidunt nulla. Vestibulum rhoncus molestie odio. Sed lobortis, justo et pretium lobortis, mauris turpis condimentum augue, nec ultricies nibh arcu pretium enim. Nunc purus neque, placerat id, imperdiet sed, pellentesque nec, nisl. Vestibulum imperdiet neque non sem accumsan laoreet. In hac habitasse platea dictumst. Etiam condimentum facilisis libero. Suspendisse in elit quis nisl aliquam dapibus. Pellentesque auctor sapien. Sed egestas sapien nec lectus. Pellentesque vel dui vel neque bibendum viverra. Aliquam porttitor nisl nec pede. Proin mattis libero vel turpis. Donec rutrum mauris et libero. Proin euismod porta felis. Nam lobortis, metus quis elementum commodo, nunc lectus elementum mauris, eget vulputate ligula tellus eu neque. Vivamus eu dolor.

Nulla in ipsum. Praesent eros nulla, congue vitae, euismod ut, commodo a, wisi. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Aenean nonummy magna non leo. Sed felis erat, ullamcorper in, dictum non, ultricies ut, lectus. Proin vel arcu a odio lobortis euismod. Vestibulum ante ipsum primis in faucibus orci luctus et ultrices posuere cubilia Curae; Proin ut est. Aliquam odio. Pellentesque massa turpis, cursus eu, euismod

References

- Arnaud, M. et al. *Astronomy and Astrophysics* (2010)
Böhringer, H. & Werner, N. *The Astronomy and Astrophysics Review* (2010)
Cavagnolo, K. W. et al. *The Astrophysical Journal Supplement* (2009)
Eckert, D. et al. *Astronomische Nachrichten* (2017)
Jansen, F. et al. *Astronomy and Astrophysics* (2001)
Lasker, B. M. *Bulletin of the American Astronomical Society* (1994)
Pratt, G. W. et al. *Space Science Reviews* (2019)
Rosati, P. et al. *Annual Review of Astronomy and Astrophysics* (2002)
Truemper, J. *Advances in Space Research* (1982)
Vikhlinin, A. et al. *The Astrophysical Journal* (2006)
Weisskopf, M. C. et al. *Proceedings of the Society of Photo Optical Instrumentation Engineers* (2000)

Appendix

A Cluster Profiles

B Reference Images




Figure 8: Plots of extracted radial profile projections for available *X-COP* galaxy clusters. *Upper Left:* Electron number density. *Upper Middle:* Gas pressure. *Upper Right:* Gas temperature. *Lower Right:* Gas entropy. *Lower Middle:* Gas mass fraction. *Lower Left:* Metal abundance. Consult the text for definitions of characteristic quantities and scaling. Compare as well to reconstructed distributions in Figure 7 which focus on the inner regions instead of the cluster outskirts.




Figure 9: Depicted are *RASS* counts used for X-ray contours shown in Figure 2 after applying a stretch and Gaussian blur (Truemper 1982). The sampled clusters are selected based on brightness and morphology, as well as data accessibility in the *Archive of Chandra Cluster Entropy Profile Tables (ACCEPT)* and *XMM Cluster Outskirts Project (X-COP)* catalogs (Cavagnolo et al. 2009, Eckert et al. 2017).