

X-Ray Emitting Plasma in Galaxies and Galaxy Clusters

Fritz Ali Agildere

fritz.agildere@udo.edu

*Department of Astronomy, Faculty of Mathematics, University of Belgrade
Astronomical Observatory of Belgrade*

May 29, 2026

Abstract

Galaxy clusters serve as valuable laboratories for studying cosmic growth on the largest scales. Permeated by thin, hot plasma, these structures are primarily observable through high resolution X-ray spectroscopy. This report examines fundamental models of central cooling and noncooling clusters, detailing how spectral analysis maps their distinct radial profiles. By isolating continuous radiation mechanisms and discrete emission lines, we can accurately determine local temperatures and elemental abundances. Furthermore, a review of upcoming X-ray instrumentation demonstrates how evolving microcalorimeter capabilities enable precise plasma diagnostics. Fitting advanced plasma models to these spectra successfully reconstructs thermodynamic properties, revealing the complex interplay between gravitational collapse, radiative cooling, and feedback mechanisms.

1 Introduction

In studying the evolving nature of our universe on the largest scales, galaxy clusters represent the most expansive clearly defined structures in which matter approaches virialization. Due to this fact, we can use them as laboratories to study the cosmic growth rate and characterize density fluctuations, as well as reconstruct the dynamical history of merging systems and the effects of feedback from active members. A tenuous plasma called the Intracluster Medium (ICM) permeates these massive objects, with temperatures in the 10^7 K range and densities from 10^{-5} to 10^{-1} cm^{-3} between their outer regions and inner cores. In this hot state, the ICM radiates at several keV photon energies, making spectroscopy in the soft X-ray band a primary method to resolve the distribution of baryonic matter inside clusters (Sarazin 1986).

1.1 Models

At the broadest level, galaxy clusters are modeled as dark matter halos fully governed by gravitational collapse. They are thought to originate from primordial density fluctuations and grow via cluster mergers as well as by accreting matter from cosmic filaments. Their total mass is dominated by the dark component defined through its purely gravitational interactions, the density of which is well described using a Navarro-Frenk-White (NFW) profile. Most baryonic matter is not contained in the stellar mass of the member galaxies themselves but in the ICM gas, which follows a similar shape to the overall matter distribution in the outer parts where it usually makes up a fraction of roughly $f_g \sim 15\%$ depending on redshift, but flattens towards the central region due to counteracting pressures. Containing this material requires a sufficiently deep gravitational potential well that is provided by the high dark matter concentration, which strongly peaks in the center as collisional effects are negligible. Shown in Figure 1 are exemplary radial density profiles with realistic parameter choices for the observed galaxy cluster categories as explained in the following subsections (Pratt et al. 2019).




Figure 1: Typical mass density profiles for NCC and CC galaxy clusters are plotted as proxies for all baryonic matter. Shown as well is a standard NFW total density curve that is dominated by a collisionless dark matter component.

Adopting this view, radiative cooling of the higher density inner regions through X-ray emission generally leads to a drop in thermal pressure and a resulting influx of subsonic materials from the surrounding medium, called a cooling flow. This assumption predicts massive rates of molecular condensation near the central galaxies which should produce significant starburst activity. However, observations by the *Chandra* (Weisskopf et al. 2000) and *XMM-Newton* (Jansen et al. 2001) missions proved that this signature is not present in reality. Instead, the ICM cools towards the core but does not reach the neutral gas threshold. The leading hypothesis for the culprit of this deviation from theoretical expectations is the injection of energy by Active Galactic Nucleus (AGN) feedback phases regulated by a balancing of heating and fueling, although this description carries with it its own set of open questions, namely the mechanisms for isotropization of highly directional jets as well as stabilization and dissipation of the AGN bubbles (Böhringer & Werner 2010, Rosati et al. 2002, Pratt et al. 2019).




Figure 2: Selection of galaxy clusters used as examples in this work. Displayed here are composite overlays of second generation *Digitized Sky Survey* red band images and contours for *ROSAT All Sky Survey (RASS)* broad band count rates (Truemper 1982, Lasker 1994). Both optical and X-ray data are stretched using a simple 0.8 exponent power law. Each frame has a radius of $15'$ for its field of view. Refer to Figure 9 for raw *ROSAT* counts underlying the contours.

1.2 Classes

According to the spherical dark matter halo assumption, gas properties should follow selfsimilar laws along the radial distance to the cluster center and only scale with the total mass and redshift. Comparing to observations such as those presented in Figures 7 and 8 shows that this holds quite well for large radii but breaks down closer to the central regions. It turns out that based on these differences, galaxy clusters can be sorted into two categories. In order to do so, it is useful to introduce the cooling time. It derives from the total thermal energy $E_t \propto nT$ and volumetric cooling rate $\dot{E}_c \propto n^2 \Lambda_c(T)$ where the cooling function for bremsstrahlung scales as $\Lambda_c(T) \propto T^{1/2}$ to define a proportionality

$$t_c = \frac{E_t}{\dot{E}_c} \propto T^{1/2} n^{-1}$$

for the ICM properties. If this timescale is small $t_c < 1$ Gyr in its center, we call it a Cooling Core (CC) cluster. These systems radiate away heat efficiently, leading to a rapid drop in gas temperature and a corresponding peak in density. With these characteristics, they also have an almost critically low entropy floor, close to the catastrophic cooling flow collapse scenario and only barely stabilized by galactic winds from star formation and AGN feedback. Generally, the population

of CC clusters comprises dynamically relaxed systems with a high degree of order. In the opposite case, cooling is highly inefficient with $t_c > 10$ Gyr and frequently exceeding the Hubble time. Such clusters are labeled Noncool Core (NCC) and tend to possess isothermal or monotonically increasing temperatures and flat density profiles towards their centers. Their inner entropy floors are large and likely originate from merging of subclusters. Heating occurs through supersonic shocks, turbulence, contact discontinuities and sloshing of the central material.

From this follows that analysis of the thermal structure in principle allows us to infer the recent dynamical history of clusters. Additionally, measuring the total mass M inside some characteristic radius r makes it possible to map the distribution of large scale density inhomogeneities and their evolution throughout cosmological time. After determining plasma properties and by assuming hydrostatic equilibrium, we can rearrange the well known formula

$$dP/dr = g\rho = GM\rho/r^2$$

to solve for mass. A common choice for a universal distance normalization is the overdensity radius R_ℓ which constrains the sphere around the cluster center inside which the average mass density is a factor ℓ larger than the surrounding critical

density of the universe at that redshift. This way, contained masses are labeled M_ℓ and average quantities T_ℓ or P_ℓ as well as astrophysical entropy K_ℓ with $K \propto Tn^{-2/3}$ can be defined for comparison between systems of vastly different sizes. Typical values are $\ell = 200$ since R_{200} is close to the theoretical virialization radius, as well as $\ell = 500$ and R_{500} as the boundary up to which current X-ray telescopes can produce reliable measurements (Böhringer & Werner 2010, Pratt et al. 2019, Voit 2005).

1.3 Examples

To have a consistent set of objects to compare theoretical predictions to real observations for this work, we select six clusters. Their optical appearance as well as X-ray surface brightness contours are depicted in Figure 2 with Table 1 listing their characterizing properties.

Table 1: Parameters for our galaxy cluster sample. Values given are those determined for the X-COP systems (Eckert et al. 2017).

Name	Class	z	R_{500} / kpc	$M_{500} / M_\odot$
A85	CC	0.0555	1200	5.6×10^{14}
A426	CC	0.0179	1300	5.9×10^{14}
A1644	NCC	0.0473	2100	3.5×10^{14}
A1656	NCC	0.0231	1300	6.0×10^{14}
A2029	CC	0.0766	1400	8.6×10^{14}
A3158	NCC	0.0590	1100	4.3×10^{14}

The Perseus or Abell 426 as well as the massive Abell 2029 clusters are archetypical examples of relaxed CC systems supported by regulated feedback loops, although there are still remnants of ancient disruptions detectable. Abell 85 also falls in the CC category but shows more prominent central sloshing as well as active accretion from intersecting cosmic filaments. The Coma or Abell 1656 cluster on the other hand is a prototype for NCC systems that have been disrupted by a recent massive merger event. Similarly, high resolution observations classify Abell 3158 as NCC despite its smooth X-ray surface. Lastly, Abell 1644 is clearly an NCC cluster as is readily apparent from its bimodal halo structure (Böhringer & Werner 2010, Eckert et al. 2017).

2 Radiation Mechanisms

While we can assume Local Thermodynamic Equilibrium (LTE) for thick plasmas such as those inside stars, where excitation obeys the Boltzmann distribution and ionization the Saha equation with radiation following a Planck law, the ICM is thin except for some resonant scattering effects. This means that photons escape easily, leaving the balance of ionization and recombination to be regulated entirely through the plasma particles themselves, in which electrons retain their Maxwellian velocity distribution. We refer to such systems as being in Collisional Ionization Equilibrium (CIE) which have certain characteristic radiation properties (Kaastra et al. 2008, Sazonov et al. 2002).

2.1 Free-Free Emission

When free electrons get deflected by the Coulomb field of an ion, they lose kinetic energy via the emission of photons. Since the amplitude and occurrence rate are dependent on the state of the thermalized plasma population, we call this mechanism thermal braking radiation or bremsstrahlung. Its emissivity is given by the equation

$$\epsilon \propto n_e n_i Z^2 T^{-1/2} g_{ff}(T, E) e^{-E/kT}$$

where the Gaunt factor $g_{ff}(T, E)$ is an averaged quantum mechanical correction of order unity. The spectra produced by bremsstrahlung present a characteristic broad continuum with an exponential rolloff once photon energies exceed the order of the thermal electron energy. This cutoff from the Gaunt factor dominated flat slope can therefore be used to determine the local temperature of an emitting region, as is shown in Figure 3 for different cases.




Figure 3: Normalized bremsstrahlung continua for different temperatures over a range of magnitudes. Fitting the bend placement allows for the inference of local temperature.

Since the ICM is very hot with thermal energies of the order 10 keV it is almost entirely dominated by the bremsstrahlung spectrum. However, systems with shallower gravitational potentials and lower temperatures than clusters are often still in CIE and exhibit a less prominent continuum, leading to other radiation components being more relevant. Examples for this are galaxy groups as well as giant elliptical galaxies (Böhringer & Werner 2010).

2.2 Bound-Free Component

If a free electron gets captured into a bound state by an ion, it emits photons with a frequency corresponding to the sum of its previous kinetic energy and the ionization potential of the specific atomic shell. This results in a spectrum defined by a sharp recombination edge at the minimum required energy for ionization of the excitation level and a smooth tail toward higher energies. Since lighter elements like hydrogen and helium are fully stripped of electrons, their contributions to the ICM radiation are negligible, although heavier metals like iron can show up as subtle steps in the bremsstrahlung continuum. At lower temperatures, this mechanism becomes more visible as ions retain more bound electrons.

2.3 Bound-Bound Features

Emission lines result from collisional or radiative excitation and the subsequent decay of an electron between two bound energy states. Due to the quantization of atomic shells, the emitted photons have a very narrow energy range. Realistic line shapes are not perfectly discrete because of the intrinsic Heisenberg uncertainty and can be further broadened by the thermal Doppler effect or deformation from collisions. From convolving Lorentzian and Gaussian curves, they obtain a Voigt profile depending on the plasma motion. Since in the ICM most atomic nuclei are stripped bare with no bound electrons left, few if any spectral lines are present in their bremsstrahlung continua. For the analysis of galaxy clusters through X-ray spectroscopy, the most important ones are the Fe K complex centered at 6.6 keV and the Fe L complex around 1.1 keV with Ne lines in the same range, as well as O lines near 0.6 keV for reasons discussed in the following sections (Rybicki & Lightman 1979, Kaastra et al. 2008, Böhringer & Werner 2010).

2.4 Sunyaev-Zeldovich Effect

Although it is observed in the microwave and submillimeter bands of the electromagnetic spectrum instead of at X-ray energies, the thermal Sunyaev-Zeldovich (SZ) effect offers a highly relevant complimentary approach for mapping ICM properties. It describes how Cosmic Microwave Background (CMB) photons are boosted towards higher energies through inverse Compton scattering with energetic ICM electrons. This produces a shadow below and an enhancement above a specific frequency versus the nearly homogenous CMB blackbody spectrum. Since the energy density at larger z is higher by a factor exactly equal to the dimming magnitude during travel of the signal, the SZ effect does not depend on redshift. As it involves one body, its surface brightness scales linearly with $P \propto nT$ as opposed to the square dependence in $S \propto n^2 T^{1/2}$ for bremsstrahlung from the interaction of two charges (Birkinshaw 1999, Böhringer & Werner 2010, Rybicki & Lightman 1979). Combining measurements of the ICM emission itself in X-rays and SZ cluster maps from *Planck* data allows us to build high quality models of their gas structure (Aghanim et al. 2020). Visualized in Figure 4 is a comparison to the CMB spectrum.




Figure 4: Schematic view of the thermal SZ effect on the CMB with exaggerated $Y \sim 0.2$ Compton parameter.

3 Instrumentation

By conducting the first sensitive all sky X-ray survey and building catalogs of galaxy clusters, the *Röntgen Satellite* (*ROSAT*) mission paved the way for following instruments (Truemper 1982). Currently, the two most established active telescopes are *Chandra* and *XMM-Newton* (Weisskopf et al. 2000, Jansen et al. 2001). While the National Aeronautics and Space Administration (NASA) operated *Chandra* can achieve very high spatial resolutions with its Advanced CCD Imaging Spectrometer (ACIS) and High Resolution Camera (HRC) at below $1''$ angular precision, the European Space Agency (ESA) run *XMM-Newton* observatory prioritizes sensitivity using its Reflection Grating Spectrometer (RSG) and European Photon Imaging Camera (EPIC) instruments. Recently launched ongoing missions include the *Spectrum Röntgen Gamma* (*SRG*) satellite which carries the eROSITA instrument for compiling catalogs and conducting wide field surveys of cosmic filaments in the soft X-ray band, as well as the *X-Ray Imaging and Spectroscopy Mission* (*XRISM*) headed by the Japan Aerospace Exploration Agency (JAXA) that uses a cooled microcalorimeter array and wide field CCD imager to reach unmatched spectral resolutions for the study of particle and enrichment dynamics in cluster cores (Predehl et al. 2021, Tashiro et al. 2024, Zhang et al. 2026). Perhaps the most capable upcoming mission is *NewAthena* which will feature a massive cryogenic microcalorimeter array with its X-Ray Integral Field Unit (XIFU) to achieve extremely resolved spectra over large patches of clusters and filaments, accompanied by the Wide Field Imager (WFI) camera for deep surveys (Cruise et al. 2025). An exciting prospect of these newer highly sensitive multicalorimeters is the isolation of soft X-rays from Warm-Hot Intergalactic Medium (WHIM) filaments that bridge the otherwise empty gaps in space between galaxy clusters sitting at the nodes of the cosmic web and are predicted to contain a significant fraction of the overall baryon population. Better precision brings with it the possibility of resolving line shapes in order to understand the likely responsible broadening mechanisms (Böhringer & Werner 2010).

4 Spectral Characteristics

The appearance of X-ray spectra measured as the emission from galaxy clusters is governed by the physics at their origin in a high temperature, diffuse plasma. When working with data that have not been fully cleaned, such as those shown in Figure 5 from the *Chandra* observatory, we are additionally including instrument noise that has to be disentangled from real physical effects before the analysis. In our case, peaks that look like emission lines are in fact fluorescence artifacts from cosmic ray particles hitting the spacecraft materials. A telltale sign for their contaminating nature is the perfect alignment in frequency where we would expect displacement caused by the different z values of sources at varying cosmic redshifts. Practically, this means that the spiking relative count rates at 2.0 keV and 7.4 keV as well as near 9.7 keV should be subtracted from the continuum. Furthermore, the flattening or even slight increase of the smooth component beyond about 6.6 keV is likewise a result of the gentle power law slope of the cosmic ray spectrum overshooting the signal (Böhringer & Werner 2010, Weisskopf et al. 2000).




Figure 5: Binned raw *Chandra* spectra for cluster selection with uncorrected systematics (Weisskopf et al. 2000).

4.1 Continuum

For galaxy clusters, the baseline continuum is virtually fully dominated by ICM thermal bremsstrahlung. This prediction is supported by Figure 5 and allows the reconstruction of an average temperature from its slope by assuming a roughly isothermal system. Some observations also hint at possible excesses, both in the soft and hard X-ray regimes. Provided these signals are not some kind of calibration error, possible causes are the coexistence of multiphase plasma components, a type of phenomenon that seems especially likely in the case of stratified CC systems, weak emission from WHIM bremsstrahlung, or upscattering of background photons in shocks such as those in NCC clusters, all of which could be introducing nonthermal signatures in a single temperature model (Böhringer & Werner 2010).

4.2 Line Emission

Superimposed on the continuous bremsstrahlung spectrum are narrow emission lines from highly ionized ICM elements. The only prominent physical feature in Figure 5 is the Fe K complex spread between 6 and 7 keV depending on redshift. In it, the most important lines stem from the hydrogen like Fe XXVI and helium like Fe XXV ions with 6.70 keV and 6.97 keV restframe energies, respectively. Since the potential of 8.83 keV required to remove the last electron from the Fe nucleus is extremely high, there are very few electrons in the Maxwell distribution tail with velocities energetic enough at typical CC central temperatures, whereas massive NCC clusters routinely exceed the required thermal energy. Due to this fact, the line ratio of both dominant Fe K lines is an excellent thermometer since it can be well fitted under CIE assumptions. Analogously, we can use the relative overall abundances of the high temperature Fe K to low energy Fe L line forests to infer ICM thermal properties.

Another kind of analysis is enabled by taking an accounting of the O and Ne abundances through their H and He like lines in the coldest regions of cluster cores. Their ratio to the iron peak elements like Fe and Ni traces the occurrence rate between thermonuclear SNIa and core collapse SNII supernovae throughout all evolutionary history of the system.

Where the first type produces only negligible traces of O and Ne but massive amounts of Fe and Ni elements, the second kind does release large quantities of these lighter alpha process elements, which therefore represent a unique fingerprint. Correlating the relationship between supernova types with redshift, we can reveal the Initial Mass Function (IMF) of stellar populations throughout cosmic time. Studies that make use of this approach usually identify little to no deviation from the Salpeter IMF

$$\xi(M_*) = dN/dM_* \propto M_*^{-\alpha}$$

with $\alpha = 2.3$ which holds for the local universe. Lastly, by instead analyzing ratios such as Ni to Fe within the iron peak elements themselves, it is possible to distinguish between different SNIa models. From this, slow deflagration models assuming a subsonic flame can be ruled out, leaving delayed detonations with a supersonic shock burning through the near Chandrasekhar mass limit white dwarf as the probable main drivers for pollution of the ICM with iron peak elements (Böhringer & Werner 2010).

5 Plasma Diagnostics

As a fundamental theoretical baseline, galaxy clusters are modeled as systems governed purely by gravitation. Due to the scale invariance of gravity, they should then only differ by predictable mathematical scalings that depend solely on mass or temperature. In reality, Supermassive Black Hole (SMBH) accretion through AGN feedback and cooling in the cluster core disturb this idealized view. To understand these physics, we compare the deviations of several key ICM properties from our naive expectation about their behavior.

Most generally, the procedure of how to extract properties from a measured X-ray spectrum involves a computational CIE plasma model fitting the best estimators for temperature, density, abundances and redshift to the given. Perhaps the most famous ones are the legacy MEKAL and modern APEC codes, which are both implemented in the XSPEC software. Historical observations could often not resolve the extended ICM and used simpler isothermal models to extract data. As discussed previously, this is at best a rough approximation and at worst completely inaccurate. Thanks to advances in current X-ray telescopes, we have access to resolved ICM halo datasets from which we can extract radial distributions, although there are still large uncertainties about nonthermal support from turbulent eddies or bulk movement indicated by hydrodynamical simulations plaguing the field that bias our results towards lower total masses. Due to the revolution of microcalorimeter spectrographs on board newer missions, it is however expected that these unknowns will be solved with higher precision measurements (Zhang et al. 2026).

To translate our two dimensional images to radial profiles, we need to correct for the projection from three dimensions along the line of sight to the detector. The basic algorithm do this decomposes our cluster into multiple annuli around the central core. Starting from the outermost annulus, we fit our spectrum and since there is little pollution from foreground or background layers, we assume that it represents a uniform shell around the whole system, taken as radially symmetric.




Figure 6: Theoretical galaxy cluster radial profiles assuming universal selfsimilar scaling relations. *Top Left:* Electron number density (Vikhlinin et al. 2006). *Top Right:* Gas pressure (Arnaud et al. 2010). *Bottom Left:* Gas temperature (Vikhlinin et al. 2006). *Bottom Right:* Gas entropy (Cavagnolo et al. 2009). Compare to Figure 7 for real observations.

Using this, we compute the corrections which have to be subtracted from the adjacent inner annulus and fit our plasma model to its now cleaned spectrum. By iteratively peeling layer by layer from the cluster, we deproject our image onto the radius of the system. For CC clusters this is enough, but for highly disrupted NCC cases more sophisticated methods are needed (Böhringer & Werner 2010).

From the employed plasma model, we know the best fitting density n and temperature T parameters. With these, we can simply apply the ideal gas law

$$P = nkT$$

to calculate the pressure. Similarly, we get

$$K = kTn^{-2/3}$$

for the astrophysical entropy. This quantity is related to its thermodynamic counterpart as $S \propto \ln(K)$ and tracks parcels under adiabatic evolution. To prove this statement, we can take the relation $P \propto V^{-5/3} \propto n^{5/3}$ for a blob expanding without the exchange of heat to its environment and find the proportionality $kT \propto n^{2/3}$ by plugging in the ideal gas law, meaning that K indeed remains constant. This makes it an incredibly useful indicator of the dynamical state of a cluster and allows us to draw conclusions on its recent past.

While the purely gravitational baseline assumption produces smoothly falling entropy profiles towards the core, lower values are typically a result of radiative cooling in a relaxed configuration, whereas a flat or elevated profile indicates an injection of energy from central engines like SMBH driven feedback and galactic winds from star formation, or merger events with other galaxy clusters (Böhringer & Werner 2010, Sarazin 1986, Rosati et al. 2002).

We can reproduce qualitatively and even to a certain extent quantitatively the shapes of radial profiles using fairly simple equations. For a typical NCC cluster, number density is sufficiently well described by a single beta model

$$n(r) = n_0 \left(1 + r^2/r_c^2\right)^{-3\beta/2}$$

in the entire system. For the CC case, we superimpose the brightness $B \propto n^2$ from an inner cool and dense core with the extended sparse halo. We use a double beta model

$$n(r) = \sqrt{n_1^2 + n_2^2}$$

where n_1 and n_2 correspond to their respective substructure. When modeling the temperature profile, a broken power law is a decent fit for NCC systems. To incorporate the decline towards the CC central regions, we include a cooling factor.




Figure 7: Sample of galaxy cluster radial profiles from *Chandra* observations (Weisskopf et al. 2000, Cavagnolo et al. 2009). *Top Left:* Electron number density. *Top Right:* Gas pressure. *Bottom Left:* Gas temperature. *Bottom Right:* Gas entropy. Compare to Figure 6 for a visualization of theoretical and empirical fits with realistic parameter choices.

We then write the combined formula

$$T(r) = T_0 \frac{x_{\text{cool}} + T_{\text{min}}/T_0}{x_{\text{cool}} + 1} \frac{(r/r_t)^{-a}}{(1 + (r/r_t)^b)^{c/b}}$$

with $x_{\text{cool}} = (r/r_{\text{cool}})^{a_{\text{cool}}}$ and where the cooling term is only significant in CC clusters that have $T_{\text{min}} < T_0$ since for NCC we set $T_{\text{min}} \approx T_0$ so that it is unity (Vikhlinin et al. 2006). As defined before, pressure P and entropy K can be derived directly from the n and T profiles, or we can introduce their own scaling relations. First, with an averaged overdensity concentration C_ℓ and $x_\ell = r/R_\ell$ we use

$$P(r) = P_0 ((C_\ell x_\ell)^c (1 + (C_\ell x_\ell)^a))^{-(b-c)/a}$$

as a universal pressure profile (Arnaud et al. 2010). Second, we get a simple power law with constant floor

$$K(r) = K_0 + K_\ell x_\ell^\gamma$$

to describe entropy (Cavagnolo et al. 2009). This function agrees with both empirical and theoretical models. Selecting parameter values motivated by realistic ranges and readable graphical appearances, Figure 6 depicts radial profiles for the introduced quantities. Accounting for their simplicity, they match the observed distributions in Figures 7 and 8 remarkably well, with most of the deviation stemming from the inconsistent normalizations and radial distance ranges.

6 Conclusion

The spectroscopic study of X-ray emitting plasma in galaxy clusters yields valuable insights into the evolution of cosmic structures. Classifying systems into CC and NCC categories enables us to effectively reconstruct distinct thermodynamic histories. Furthermore, analyzing thermal bremsstrahlung and discrete line emission serves as a reliable diagnostic tool for determining ICM properties, including local temperature and electron density, as well as entropy and gas pressure.

At the same time, the purely gravitational baseline model is faced with some clear observational contradictions. The predicted catastrophic cooling flows for relaxed clusters are completely absent, requiring heating mechanisms like AGN feedback and galactic winds to stabilize cores. Additionally, deriving radial profiles in three dimensions from projected two dimensional spectral maps is complicated even further by lingering uncertainties regarding nonthermal support.

Modern X-ray missions, including XRISM and New Athena, are deploying highly sensitive cryogenic microcalorimeters with excellent spectral resolution expected to resolve such uncertainties. These instruments will rigorously map core enrichment dynamics and potentially detect the faint WHIM X-rays along the bridges of cosmic filaments.

References

- Aghanim, N. et al. *Astronomy and Astrophysics* (2020)
 Arnaud, M. et al. *Astronomy and Astrophysics* (2005)
 Arnaud, M. et al. *Astronomy and Astrophysics* (2010)
 Birkinshaw, M. *Physics Reports* (1999)
 Böhringer, H. & Werner, N. *The Astronomy and Astrophysics Review* (2010)
 Cavagnolo, K. W. et al. *The Astrophysical Journal Supplement* (2009)
 Cruise, M. et al. *Nature Astronomy* (2025)
 Eckert, D. et al. *Astronomische Nachrichten* (2017)
 Jansen, F. et al. *Astronomy and Astrophysics* (2001)
 Kaastra, J. S. et al. *Space Science Reviews* (2008)
 Lasker, B. M. *Bulletin of the American Astronomical Society* (1994)
 Pratt, G. W. et al. *Astronomy and Astrophysics* (2010)
 Pratt, G. W. et al. *Space Science Reviews* (2019)
 Predehl, P. et al. *Astronomy and Astrophysics* (2021)
 Rosati, P. et al. *Annual Review of Astronomy and Astrophysics* (2002)
 Rybicki, G. B. & Lightman, A. P. *Radiative Processes in Astrophysics* (1979)
 Sarazin, C. L. *Reviews of Modern Physics* (1986)
 Sazonov, S. Y. et al. *Monthly Notices of the Royal Astronomical Society* (2002)
 Tashiro, M. et al. *Proceedings of the Society of Photo Optical Instrumentation Engineers* (2024)
 Truemper, J. *Advances in Space Research* (1982)
 Vikhlinin, A. et al. *The Astrophysical Journal* (2006)
 Voit, M. G. *Reviews of Modern Physics* (2005)
 Weisskopf, M. C. et al. *Proceedings of the Society of Photo Optical Instrumentation Engineers* (2000)
 Zhang, C. et al. *Astronomy and Astrophysics* (2026)

Appendix

A Cluster Profiles

Many more quantities and modifications could be added to the previously defined radial profiles. Noting an interesting consequence of the self similarity between galaxy clusters, different characteristic overdensity averages are determined by R_ℓ and M_ℓ alone. Accordingly, combining values listed in Table 1 allows us to estimate averages for T_ℓ as well as P_ℓ and K_ℓ where in addition we must only account for the simplified cosmological evolution parameter

$$h(z) = \frac{H(z)}{H_0} = \sqrt{\Omega_M(1+z)^3 + \Omega_\Lambda}$$

under the condition $\Omega_M + \Omega_\Lambda = 1$ for a flat universe. Exact equations based on the REXCESS sample among others are given in the literature (Arnaud et al. 2005, Arnaud et al. 2010, Pratt et al. 2010).

B Reference Images

To supplement the composites in Figure 2 with intermediate pipeline data, Figure 9 displays the raw count rates after an initial smoothing.

C Methodology

All figures included in this text are generated using a semi automated Python pipeline that matches and pulls from the NASA High Energy Astrophysics Science Archive Research Center (HEASARC) or SkyView databases, and extracts locally saved archival data files saved manually directly from the different collaborations cited in this work. The relevant components are then cleaned and processed to be ingested by the plotting script itself. To reproduce and modify the implementation, consult this repository.




Figure 8: Plots of extracted radial profile projections for available *X-COP* galaxy clusters. *Upper Left:* Electron number density. *Upper Middle:* Gas pressure. *Upper Right:* Gas temperature. *Lower Right:* Gas entropy. *Lower Middle:* Gas mass fraction. *Lower Left:* Metal abundance. Consult the text for definitions of characteristic quantities and scaling. Compare as well to reconstructed distributions in Figure 7 which focus on the inner regions instead of the cluster outskirts.




Figure 9: Depicted are *RASS* counts used for X-ray contours shown in Figure 2 after applying a stretch and Gaussian blur (Truemper 1982). The sampled clusters are selected based on brightness and morphology, as well as data accessibility in the *Archive of Chandra Cluster Entropy Profile Tables (ACCEPT)* and *XMM Cluster Outskirts Project (X-COP)* catalogs (Cavagnolo et al. 2009, Eckert et al. 2017).