

Berry and Uhlmann Geometry of Entangled Photon-Added Coherent States under the Global Two-Mode Cycle

$$U(\phi) = e^{-i(\hat{n}_A + \hat{n}_B)\phi}$$

Fernando Nieto-Guadarrama

August 2026

Abstract

We develop the geometric-phase structure of bipartite photon-added coherent states (PACS) taking as the primary cycle the *free evolution of both modes*, $U(\phi) = e^{-i(\hat{n}_A + \hat{n}_B)\phi}$. This is the evolution an actual propagating two-mode field undergoes, and it is a more natural starting point than a cycle acting on one mode alone: under the global cycle *both* reduced states traverse isospectral orbits and *both* carry Uhlmann holonomy, so the two subsystems are treated on an equal footing from the outset. We obtain in closed form the global Berry phase, which is the total mean photon number and therefore splits exactly into single-mode contributions $\Phi_B = \Phi_B^{(A)} + \Phi_B^{(B)}$; in the balanced symmetric family it is an affine function of the concurrence and exactly twice the single-mode result. For the mixed reduced states we derive the Uhlmann holonomy from the parallel-transport condition in its correct anticommutator form, $[S, \dot{S}] = i\{\rho, \Lambda\}$, which yields the compact statement that parallel transport renormalizes the off-diagonal generator by exactly the concurrence, $(G + \Lambda)_{\text{od}} = C G_{\text{od}}$, and hence $\Phi_U = \text{Arg}[(-1)^\omega \cos \pi b - i \omega(1 - 2p)b^{-1} \sin \pi b]$ with $b = \omega \sqrt{1 + C^2 - 4p(1 - p)}$. We compute the concurrence of the entangled PACS in closed form, $C = 2\mu\nu N_1 N_2 / \mathcal{N}^2$, obtained from $C = 2|\det M|$ with the branch overlaps cancelling identically; the coherent amplitude thus acts as a continuous knob taking the entanglement from maximal (orthogonal Fock branches at $\alpha \rightarrow 0$) to zero (parallel branches at large α). Expressing the holonomy through the concurrence gives two clean statements: in the balanced symmetric family $p = C/2$ exactly, so $b = \Delta \sqrt{C^2 + (1 - C)^2}$ and Φ_U is a pure function of C ; while on the locus $p = 1/2$ where the singularities live, $b = \Delta C$ and Z_U becomes real, reducing Φ_U to a $\mathbb{Z}_2$ index valued in $\{0, \pi\}$. The nodal condition is then that the effective winding $\omega_{\text{eff}} = \Delta C$ cross a half-integer, which quantizes the concurrence to a finite, equally spaced ladder: each photon-addition separation Δ supports exactly Δ isolated Uhlmann singularities at $C_k = (2k + 1)/2\Delta$, $k = 0, \dots, \Delta - 1$, symmetric about $C = 1/2$, shared by both subsystems and all bounded by a critical amplitude α_g defined by $|g_{mn}(\alpha_g)|^2 = 1/2$. We further explain why the two subsystems are *not* Uhlmann-symmetric despite sharing a concurrence: Schmidt isospectrality fixes the length of each reduced Bloch vector but not its orientation relative to that mode's own logical axis, so the local populations p_A, p_B differ whenever $\beta \neq \alpha$; in the crossed family the winding numbers differ in sign instead, producing an exact antisymmetry $\Phi_U^B = -\Phi_U^A$. In balanced form these statements sharpen into $\sqrt{p_A p_B} = C/2$: the concurrence fixes only the geometric mean of the two local populations, while their ratio $\eta = \sqrt{p_A/p_B}$ is free — and the physical range of η pinches to $\{1\}$ as $C \rightarrow 1$, so maximal entanglement enforces symmetry. Every closed form is validated against independent numerics: direct summation in a truncated two-mode Fock space for the Berry sector, and two connection-free holonomy algorithms for the Uhlmann sector. A fully commented Jupyter notebook reproduces all fourteen figures and both tables.

Contents

1	Introduction	2
2	Photon-added coherent states	3
3	Bipartite model and logical-qubit reduction	4
4	Global Berry phase	5
5	Concurrence of the entangled PACS	5
6	Uhlmann holonomy of the reduced states	6
6.1	The parallel-transport condition	7
6.2	Closed form	7
7	Nodal singularities	7
7.1	Nodal coordinates and the critical amplitude	8
8	The Uhlmann phase as a function of the concurrence	8
8.1	(C, η) coordinates: what entanglement fixes and what it does not	8
8.2	Balanced symmetric family: a pure function of C	9
8.3	On $p = 1/2$: a $\mathbb{Z}_2$ index	9
9	Quantization of the concurrence at the singular points	9
10	Why the two subsystems are not Uhlmann-symmetric	10
11	Numerical validation	11
12	Results	12
13	Discussion	12
14	Conclusions	20
A	Scalar products of photon-added coherent states	21
B	Cross matrix element of the number operator	21
C	Numerical conventions	21

1 Introduction

Photon addition turns a Glauber coherent state into a strongly nonclassical one while preserving an essentially closed analytic structure controlled by a single integer. The photon-added coherent states (PACS) of Agarwal and Tara [1] have since become a standard laboratory for the classical–quantum boundary of light. Entanglement in bipartite PACS was analysed by Domínguez-Serna, Mendieta-Jiménez and Rojas [5], who showed that superpositions of *nonorthogonal* PACS branches reduce, after Gram–Schmidt orthogonalization, to an effective two-qubit problem with a computable Wootters concurrence [4]. In parallel, geometric phases of coherent-state superpositions have been

studied for two-mode entangled coherent states [6] and for nonlinear and squeezed states [7], while Uhlmann’s mixed-state holonomy [3] has been developed for coherent states by Wang *et al.* [8].

Why the global cycle. A geometric phase is a property of a *path*, so the first thing to fix is which path. Earlier treatments of this problem cycled a single mode, $U_A(\phi) \otimes I_B$. That choice has a structural drawback: by the cyclicity of the partial trace, a strictly local unitary leaves ρ_B *exactly* invariant,

$$\text{Tr}_A [(U_A \otimes I) \rho_{AB} (U_A^\dagger \otimes I)] = \text{Tr}_A [\rho_{AB}], \quad (1)$$

which is the content of no-signalling. The path of ρ_B is then a *point*, its Uhlmann holonomy is the identity, and the second subsystem drops out of the discussion — not because it lacks geometry, but because the chosen cycle does not probe it.

Here we take instead the *global* cycle

$$U(\phi) = e^{-i(\hat{n}_A + \hat{n}_B)\phi} = U_A(\phi) \otimes U_B(\phi), \quad \hat{n}_s = a_s^\dagger a_s, \quad (2)$$

i.e. free propagation of the two-mode field. It is the physically natural evolution, and under it $\rho_A(\phi) = U_A \rho_A(0) U_A^\dagger$ and $\rho_B(\phi) = U_B \rho_B(0) U_B^\dagger$ both move: both subsystems traverse isospectral loops and both carry nontrivial Uhlmann holonomy. The whole analysis is rebuilt on this basis.

Contributions.

- (i) The closed-form global Berry phase, equal to the total mean photon number, with its exact decomposition into single-mode contributions and its affine dependence on the concurrence (Sec. 4, Figs. 1–5).
- (ii) The Uhlmann connection derived from the parallel-transport condition in anticommutator form, giving the compact result $(G + \Lambda)_{\text{od}} = C G_{\text{od}}$ and the closed phase with $b = \omega \sqrt{1 + C^2 - 4p(1-p)}$ (Sec. 6).
- (iii) A finite nodal set: exactly Δ isolated singularities at $C_k = (2k + 1)/2\Delta$, common to both subsystems, all bounded by α_g (Sec. 7, Figs. 6, 8).
- (iv) An explanation of the $A \leftrightarrow B$ asymmetry: the concurrence is shared but the local populations are not, with the symmetry locus $\beta = \alpha$ and an exact antisymmetry in the crossed family (Sec. 10, Figs. 7, 11).
- (v) The concurrence of the entangled PACS in closed form and its limits (Sec. 5, Fig. 12); the Uhlmann phase written through the concurrence — a pure function of C in the balanced symmetric family, a $\mathbb{Z}_2$ index on $p = 1/2$ (Sec. 8, Figs. 13–14) — and the resulting quantization of the concurrence at the singular points (Sec. 9).
- (vi) A complete independent validation layer and a reproducible notebook (Sec. 11).

2 Photon-added coherent states

A PACS is obtained by applying the creation operator n times to a coherent state and renormalizing, $|\alpha, n\rangle = a^{\dagger n} |\alpha\rangle / [\langle \alpha | a^n a^{\dagger n} | \alpha \rangle]^{1/2}$. Inserting $|\alpha\rangle = e^{-|\alpha|^2/2} \sum_j \alpha^j / \sqrt{j!} |j\rangle$ and using

$a^{\dagger n} |j\rangle = \sqrt{(j+n)!/j!} |j+n\rangle$ gives $a^{\dagger n} |\alpha\rangle = e^{-|\alpha|^2/2} \sum_{k \geq 0} \alpha^k \sqrt{(n+k)!/k!} |n+k\rangle$, whose norm is fixed by the PACS identity $\langle \alpha | a^n a^{\dagger n} | \alpha \rangle = n! L_n(-|\alpha|^2)$. Hence

$$|\alpha, n\rangle = \frac{e^{-|\alpha|^2/2}}{\sqrt{n! L_n(-|\alpha|^2)}} \sum_{k=0}^{\infty} \frac{\alpha^k \sqrt{(n+k)!}}{k!} |n+k\rangle, \quad (3)$$

with L_n the ordinary Laguerre polynomial. The mean photon number follows from the Laguerre recursion,

$$\bar{n}_{\alpha,n} = (n+1) \frac{L_{n+1}(-|\alpha|^2)}{L_n(-|\alpha|^2)} - 1, \quad (4)$$

which correctly reduces to $|\alpha|^2$ at $n=0$.

Under the global cycle (2) each PACS transforms as

$$U(\phi) |\alpha, n\rangle = e^{-in\phi} |\alpha e^{-i\phi}, n\rangle, \quad (5)$$

because $e^{-i\hat{n}\phi}$ is diagonal in the Fock basis: the coherent amplitude rotates and the photon-addition index contributes an *explicit* phase $e^{-in\phi}$. These index phases are what generate the winding numbers below.

3 Bipartite model and logical-qubit reduction

Since PACS branches are not orthogonal, a bipartite superposition must be orthogonalized before it can be read as a two-qubit state. With p_1, p_2 the local branch overlaps and $N_j = \sqrt{1 - |p_j|^2}$, the Gram-Schmidt bases are

$$|0_s\rangle = (\text{first branch}), \quad |1_s\rangle = \frac{(\text{second branch}) - p_s |0_s\rangle}{N_s}, \quad s = A, B, \quad (6)$$

and the reduced states become rank-2 with a Wootters concurrence $C = 2\sqrt{\det \rho_s}$ [4]. We use two families:

Branch-correlated.

$$|\Phi\rangle \propto \mu |\alpha, m\rangle_A |\beta, m\rangle_B + \nu e^{i\chi} |\alpha, n\rangle_A |\beta, n\rangle_B, \quad (7)$$

whose branch phases under (5) are $e^{-i2m\phi}$ and $e^{-i2n\phi}$; each mode separately winds with $\omega_A = \omega_B = +\Delta$, $\Delta \equiv n - m$.

Crossed (swap).

$$|\Phi\rangle \propto \mu |\alpha, m\rangle_A |\beta, n\rangle_B + \nu e^{i\chi} |\beta, n\rangle_A |\alpha, m\rangle_B. \quad (8)$$

Here the photon-addition indices are *interchanged* between modes, so $\omega_A = +\Delta$ while $\omega_B = -\Delta$: the two subsystems wind in opposite senses.

With $\nu = \sqrt{1 - \mu^2}$, $g_\alpha \equiv g_{mn}(\alpha) = \langle \alpha, m | \alpha, n \rangle$ and $h_\beta \equiv g_{mn}(\beta)$, the branch-correlated family gives

$$\mathcal{N}^2 = 1 + 2\mu\nu g_\alpha h_\beta, \quad p_A = \frac{\nu^2(1 - g_\alpha^2)}{\mathcal{N}^2}, \quad p_B = \frac{\nu^2(1 - h_\beta^2)}{\mathcal{N}^2}, \quad (9)$$

$$\det \rho_A = \det \rho_B = \frac{\mu^2 \nu^2 (1 - g_\alpha^2)(1 - h_\beta^2)}{\mathcal{N}^4} = \left(\frac{C}{2}\right)^2, \quad (10)$$

while the crossed family, with $s \equiv \langle \alpha, m | \beta, n \rangle$, gives $\mathcal{N}^2 = 1 + 2\mu\nu s^2$, $p_A = p_B = \nu^2(1 - s^2)/\mathcal{N}^2$ and $C = 2\mu\nu(1 - s^2)/\mathcal{N}^2$; there $p_A = p_B$ holds identically because the two branches are exchanged copies of one another, so $N_1 = N_2$.

Equation (10) is worth emphasizing: for a bipartite *pure* state the Schmidt decomposition makes ρ_A and ρ_B isospectral, so the determinant — and hence the concurrence — is *shared* by the two subsystems. This will be central in Sec. 10.

4 Global Berry phase

The Berry connection of the global cycle is the total mean photon number,

$$\mathcal{A}_\phi = \langle \Phi | (\hat{n}_A + \hat{n}_B) | \Phi \rangle, \quad (11)$$

independent of ϕ since the generator commutes with itself, so $\Phi_B = \int_0^{2\pi} \mathcal{A}_\phi d\phi = 2\pi \langle \hat{n}_A + \hat{n}_B \rangle$. Expanding (7) and using $\kappa_\alpha \equiv \langle \alpha, m | a^\dagger a | \alpha, n \rangle$ (Appendix B),

$$\frac{\Phi_B}{2\pi} = \frac{\mu^2(\bar{n}_{\alpha,m} + \bar{n}_{\beta,m}) + \nu^2(\bar{n}_{\alpha,n} + \bar{n}_{\beta,n}) + 2\mu\nu \cos \chi (\kappa_\alpha h_\beta + g_\alpha \kappa_\beta)}{1 + 2\mu\nu \cos \chi g_\alpha h_\beta} \quad (12)$$

Exact mode decomposition. Because the generator is a *sum* of commuting single-mode operators, the phase splits exactly,

$$\Phi_B = \Phi_B^{(A)} + \Phi_B^{(B)}, \quad \frac{\Phi_B^{(A)}}{2\pi} = \langle \hat{n}_A \rangle, \quad \frac{\Phi_B^{(B)}}{2\pi} = \langle \hat{n}_B \rangle, \quad (13)$$

with $\langle \hat{n}_A \rangle = [\mu^2 \bar{n}_{\alpha,m} + \nu^2 \bar{n}_{\alpha,n} + 2\mu\nu \cos \chi \kappa_\alpha h_\beta]/\mathcal{N}^2$ and the mirror expression for B . This decomposition, verified to 10^{-16} in Sec. 11, is displayed in Fig. 5: for $\beta \neq \alpha$ the two modes contribute unequally, and the mode whose amplitude is held fixed contributes a nearly constant amount.

Symmetric family and concurrence dependence. For $\beta = \alpha$ the two contributions coincide and Φ_B is exactly *twice* the single-mode result. In the balanced symmetric case ($\mu = \nu$, $\chi = 0$), writing $g^2 = (1 - C)/(1 + C)$,

$$\frac{\Phi_B}{2\pi} = (1 + C) \bar{n}_{\text{av}} + (1 - C) Q_{mn}, \quad \bar{n}_{\text{av}} = \frac{1}{2}(\bar{n}_{\alpha,m} + \bar{n}_{\alpha,n}), \quad (14)$$

again *affine in the concurrence*, interpolating between the mean-photon average at $C = 0$ and the interference term Q_{mn} at $C = 1$, and again exactly twice the one-mode expression.

Branch-phase interference. The cycle in ϕ produces a connection that already contains the coherent cross term κg , yet no oscillation in ϕ : the interference is real but hidden inside a constant. Introducing the tunable relative branch phase χ of Eqs. (7) exposes it as an explicit 2π -periodic modulation, Fig. 3.

5 Concurrence of the entangled PACS

Because the concurrence controls both the Berry phase (through Eq. (14)) and the entire Uhlmann sector below, we derive it explicitly.

Step 1: logical basis. Inverting Eq. (6),

$$|\alpha, n\rangle = g_\alpha |0_A\rangle + N_1 |1_A\rangle, \quad |\beta, n\rangle = h_\beta |0_B\rangle + N_2 |1_B\rangle, \quad (15)$$

with $N_1 = \sqrt{1 - g_\alpha^2}$, $N_2 = \sqrt{1 - h_\beta^2}$.

Step 2: coefficient matrix. Substituting into Eq. (7) at $\chi = 0$,

$$|\Phi\rangle = \frac{1}{\mathcal{N}} \left[(\mu + \nu g_\alpha h_\beta) |0_A 0_B\rangle + \nu g_\alpha N_2 |0_A 1_B\rangle + \nu N_1 h_\beta |1_A 0_B\rangle + \nu N_1 N_2 |1_A 1_B\rangle \right], \quad (16)$$

i.e. $|\Phi\rangle = \sum_{ij} M_{ij} |i_A j_B\rangle$ with

$$M = \frac{1}{\mathcal{N}} \begin{pmatrix} \mu + \nu g_\alpha h_\beta & \nu g_\alpha N_2 \\ \nu N_1 h_\beta & \nu N_1 N_2 \end{pmatrix}. \quad (17)$$

Step 3: concurrence. For a pure two-qubit state $C = 2|\det M|$, and the overlap terms cancel identically:

$$\det M = \frac{\nu N_1 N_2}{\mathcal{N}^2} [\mu + \nu g_\alpha h_\beta - \nu g_\alpha h_\beta] = \frac{\mu \nu N_1 N_2}{\mathcal{N}^2}, \quad (18)$$

so that

$$C = \frac{2\mu \nu N_1 N_2}{\mathcal{N}^2} = \frac{2\mu \nu \sqrt{(1 - g_\alpha^2)(1 - h_\beta^2)}}{1 + 2\mu \nu g_\alpha h_\beta} \quad (19)$$

consistent with $\det \rho_A = \det \rho_B = |\det M|^2 = (C/2)^2$, Eq. (10). Substituting the closed-form overlap (46) renders C an explicit function of $(\mu, \alpha, \beta; m, n)$ alone.

Limits. The overlap controls everything (Fig. 12):

- $\alpha, \beta \rightarrow 0$: $g \rightarrow 0$, since the PACS degenerate into the Fock states $|m\rangle, |n\rangle$, which *are* orthogonal. Then $C \rightarrow 2\mu \nu$, maximal for the given weights; balanced gives $C = 1$.
- $\alpha, \beta \rightarrow \infty$: $g \rightarrow 1$, the two branches become parallel, and $C \rightarrow 0$: a product state.

The coherent amplitude is thus a continuous knob taking the entanglement from maximal to zero, with nonorthogonality the precise mechanism that degrades it. In the balanced symmetric family Eq. (19) collapses to

$$C = \frac{1 - q}{1 + q}, \quad q = |g_{mn}(\alpha)|^2. \quad (20)$$

6 Uhlmann holonomy of the reduced states

Under the global cycle both reduced states follow isospectral orbits

$$\rho_s(\phi) = V_s(\phi) \rho_s(0) V_s^\dagger(\phi), \quad V_s(\phi) = e^{i \frac{\omega_s}{2} \sigma_z \phi}, \quad s = A, B. \quad (21)$$

6.1 The parallel-transport condition

With $\rho = WW^\dagger$, $W = \sqrt{\rho}U$, $S \equiv \sqrt{\rho}$ and $\dot{U} = -i\Lambda U$, one has $W^\dagger \dot{W} = U^\dagger(S\dot{S} - i\rho\Lambda)U$. Uhlmann's condition is that $W^\dagger \dot{W}$ be Hermitian, which gives

$$\boxed{[S, \dot{S}] = i\{\rho, \Lambda\}} \quad (22)$$

an *anticommutator* (Sylvester) equation. In the eigenbasis of ρ , with eigenvalues λ_j , it is solved elementwise by

$$\Lambda_{jk} = \frac{-i[S, \dot{S}]_{jk}}{\lambda_j + \lambda_k}. \quad (23)$$

We stress that the superficially similar sandwich expression $\Lambda = \frac{1}{2i}S^{-1}[S, \dot{S}]S^{-1}$, whose elements carry a denominator $2\sqrt{\lambda_j\lambda_k}$, does *not* solve Eq. (22) unless $\lambda_j = \lambda_k$; the two coincide only for the maximally mixed state. Section 11 confirms Eq. (23) against two connection-free numerical holonomies.

6.2 Closed form

For the isospectral loop, $\dot{S} = i[G, S]$ with $G = \frac{\omega}{2}\sigma_z$, so $[S, \dot{S}]_{jk} = -iG_{jk}(\sqrt{\lambda_j} - \sqrt{\lambda_k})^2$ and Eq. (23) gives a purely off-diagonal connection

$$\Lambda_{jk} = -G_{jk} \frac{(\sqrt{\lambda_j} - \sqrt{\lambda_k})^2}{\lambda_j + \lambda_k}, \quad j \neq k. \quad (24)$$

Since $\text{Tr } \rho = 1$ we have $\lambda_j + \lambda_k = 1$ for $j \neq k$, and $2\sqrt{\lambda_+ \lambda_-} = 2\sqrt{\det \rho} = C$, so $(\sqrt{\lambda_+} - \sqrt{\lambda_-})^2 = 1 - C$ and the connection collapses to

$$\boxed{\Lambda_{\text{od}} = -(1 - C) G_{\text{od}} \quad \implies \quad (G + \Lambda)_{\text{od}} = C G_{\text{od}}} \quad (25)$$

the concurrence is exactly the factor by which parallel transport renormalizes the off-diagonal part of the generator. Writing the Bloch vector of $\rho_s(0)$ with $\cos \theta_s = (1 - 2p_s)/r$, $r = \sqrt{1 - C^2}$, and using $e^{iG2\pi} = (-1)^\omega \mathbb{K}$ for integer ω , the holonomy $U_{\text{hol}} = e^{iG2\pi} e^{-i(G+\Lambda)2\pi}$ gives

$$\boxed{\Phi_U = \text{Arg} \left[(-1)^\omega \cos(\pi b) - i \frac{\omega(1 - 2p)}{b} \sin(\pi b) \right], \quad b = \omega \sqrt{1 + C^2 - 4p(1 - p)}} \quad (26)$$

where the compact b follows from $b^2 = \omega^2(\cos^2 \theta + C^2 \sin^2 \theta)$ on substituting $r^2 = 1 - C^2$. Two limits check the result: if G commutes with ρ the state does not move and $\Phi_U = 0$; and the maximally mixed loop is again trivial. At the population-symmetric point,

$$p = \frac{1}{2} \quad \implies \quad b = \omega C, \quad (27)$$

so b is there *proportional* to the concurrence.

7 Nodal singularities

Define the complex Uhlmann amplitude $Z_U = (-1)^\omega \cos(\pi b) - i\omega(1 - 2p)b^{-1} \sin(\pi b)$, so $\Phi_U = \text{Arg } Z_U$. A genuine phase singularity needs both parts to vanish; since $\cos(\pi b) = 0 \Rightarrow \sin(\pi b) \neq 0$, the factor $(1 - 2p)$ must vanish independently. Hence

$$\boxed{p = \frac{1}{2}, \quad |b| = k + \frac{1}{2}, \quad k \in \mathbb{Z}_{\geq 0}} \quad (28)$$

At $p = 1/2$, Eq. (27) gives $|b| = \Delta C$, so the concurrence is quantized to

$$\boxed{C_k = \frac{2k+1}{2\Delta}, \quad k = 0, 1, \dots, \Delta - 1} \quad (29)$$

the upper bound enforcing $C_k \leq 1$. Thus **each photon-addition separation Δ supports exactly Δ Uhlmann singularities**, equally spaced in concurrence by $1/\Delta$, isolated, with no accumulation point. Because only $|\omega_s|$ enters and the concurrence is shared, the set (29) is *the same for both subsystems*: which entanglement values are geometrically singular is a property of the pair.

7.1 Nodal coordinates and the critical amplitude

For the symmetric family, $p = 1/2$ in Eq. (9) solves exactly: $\mu = \nu(1 - 2q)$ with $q = g_{mn}(\alpha)^2$, i.e.

$$\mu(q) = \frac{1 - 2q}{\sqrt{1 + (1 - 2q)^2}}, \quad \text{along which} \quad C = 1 - 2q. \quad (30)$$

These relations are purely kinematic — properties of the state, not of the connection. Combining with Eq. (29),

$$q_k = \frac{1 - C_k}{2}, \quad \mu_k = \frac{C_k}{\sqrt{1 + C_k^2}}, \quad |g_{mn}(\alpha_k)|^2 = q_k, \quad (31)$$

with α_k obtained by root-finding on the monotonic function $|g_{mn}(\alpha)|^2$. Defining the critical amplitude by

$$|g_{mn}(\alpha_g)|^2 = \frac{1}{2}, \quad (32)$$

the $p = 1/2$ curve exists only for $\alpha \leq \alpha_g$. Since $C_k \geq 1/(2\Delta) > 0$ we have $q_k < 1/2$ strictly, so *every* node satisfies $\alpha_k < \alpha_g$: the critical amplitude is a genuine **bound** on the nodal region, but — the set being finite — not an accumulation point. Table 3 lists the complete nodal set for representative pairs, each verified to satisfy $p = 1/2$, $|b| = k + 1/2$ and $|Z_U| \lesssim 10^{-14}$ simultaneously.

8 The Uhlmann phase as a function of the concurrence

Equation (26) mixes the concurrence with the population. We now identify the loci on which Φ_U becomes a function of C *alone*.

8.1 (C, η) coordinates: what entanglement fixes and what it does not

In the balanced correlated family ($\mu = \nu = 1/\sqrt{2}$) an exact identity ties both subsystems to the shared concurrence:

$$\boxed{\sqrt{p_A p_B} = \frac{C}{2}} \quad (33)$$

Introducing the *local asymmetry*

$$\eta \equiv \sqrt{\frac{p_A}{p_B}} = \sqrt{\frac{1 - g_\alpha^2}{1 - h_\beta^2}}, \quad p_A = \frac{C}{2}\eta, \quad p_B = \frac{C}{2\eta}, \quad (34)$$

separates the two roles cleanly: *the concurrence fixes the geometric mean of the two local populations, while η is the local degree of freedom that entanglement leaves free*. The $A \leftrightarrow B$ symmetry of Sec. 10 is exactly $\eta = 1$, i.e. $\beta = \alpha$.

Physical region. Requiring $p(1-p) \geq \det \rho = C^2/4$ on both sides gives $C\eta^2 - 2\eta + C \leq 0$, i.e.

$$\eta \in \left[\frac{1-r}{C}, \frac{1+r}{C} \right], \quad r = \sqrt{1-C^2}, \quad (35)$$

an interval symmetric under $\eta \rightarrow 1/\eta$ (the product of its endpoints is exactly 1). It narrows as C grows, and

$$\boxed{C \rightarrow 1 \implies \eta \rightarrow 1} \quad (36)$$

maximal entanglement forces the two subsystems to be symmetric: at $C = 1$ one necessarily has $p_A = p_B = 1/2$, hence $\Phi_U^A = \Phi_U^B$ (and in fact $b = \Delta$, an integer, so $\Phi_U = 0$). Asymmetry is a luxury only partial entanglement can afford. See Fig. 14.

8.2 Balanced symmetric family: a pure function of C

With $\beta = \alpha$ and $\mu = \nu$ one has $\eta = 1$ and therefore $p = C/2$ exactly, so $1 + C^2 - 4p(1-p) = C^2 + (1-C)^2$ and

$$\boxed{b = \Delta \sqrt{C^2 + (1-C)^2}, \quad \Phi_U = \text{Arg} \left[(-1)^\Delta \cos(\pi b) - i \frac{\Delta(1-C)}{b} \sin(\pi b) \right]} \quad (37)$$

a function of C and Δ *only*, plotted in Fig. 13(a). Its range is $b \in [\Delta/\sqrt{2}, \Delta]$, minimal at $C = 1/2$ and equal to Δ at both endpoints $C \rightarrow 0$ and $C \rightarrow 1$.

8.3 On $p = 1/2$: a $\mathbb{Z}_2$ index

On the locus where the nodes live the imaginary part of Z_U vanishes identically (it carries the factor $1-2p$) and $b = \Delta C$, so

$$Z_U = (-1)^\Delta \cos(\pi \Delta C) \in \mathbb{R} \implies \boxed{\Phi_U \in \{0, \pi\}} \quad (38)$$

The Uhlmann phase ceases to be continuous and becomes a $\mathbb{Z}_2$ index determined by the concurrence alone: 0 where $(-1)^\Delta \cos(\pi \Delta C) > 0$ and π where it is negative. The nodes are precisely the domain walls between the two values, Fig. 13(c).

9 Quantization of the concurrence at the singular points

On $p = 1/2$ the parameter $b = \Delta C$ acts as an *effective winding number* that the concurrence tunes continuously,

$$\omega_{\text{eff}} \equiv b|_{p=1/2} = \Delta C \quad \text{sweeping } [0, \Delta] \text{ as } C \in [0, 1], \quad (39)$$

and the nodal condition $|b| = k + \frac{1}{2}$ is the statement that this effective winding crosses a *half-integer*:

$$\boxed{\Delta C_k = k + \frac{1}{2} \iff C_k = \frac{2k+1}{2\Delta}, \quad k = 0, 1, \dots, \Delta-1} \quad (40)$$

Five consequences follow immediately.

- (1) **Count.** Since $C \leq 1$ only $k \leq \Delta - 1$ fits: there are *exactly* Δ singularities. The photon-addition separation $\Delta = n - m$ literally counts the singular points.
- (2) **Uniform spacing.** $C_{k+1} - C_k = 1/\Delta$: the singular concurrences are equidistant, being the *odd* multiples of $1/2\Delta$.

- (3) **Reflection symmetry.** $C_k + C_{\Delta-1-k} = 1$, so the ladder is symmetric about $C = 1/2$.
- (4) **$\mathbb{Z}_2$ jumps.** By Eq. (38), crossing each C_k flips Φ_U between 0 and π ; there are exactly Δ flips between $C = 0$ and $C = 1$.
- (5) **Shared by both subsystems.** Only $|\omega_s| = \Delta$ enters and the concurrence is shared, so A and B have the *same* ladder $\{C_k\}$ — though not the same coordinates (μ, α) , as Fig. 8 shows.

Translated back to parameter space, on $p = 1/2$ the symmetric family gives $C = 1 - 2q$, so each C_k fixes an overlap $q_k = (1 - C_k)/2$ and with it an amplitude α_k ; since $C_k \geq 1/(2\Delta) > 0$ one always has $q_k < 1/2$ and hence $\alpha_k < \alpha_g$, recovering the bound of Sec. 7.

10 Why the two subsystems are not Uhlmann-symmetric

Since the global state is pure, ρ_A and ρ_B are isospectral and share a concurrence. It is then natural to ask why $\Phi_U^A \neq \Phi_U^B$. The answer is that *the Uhlmann phase is not a function of the entanglement alone*. It depends on three inputs,

$$\Phi_U^s = \Phi_U(C, p_s, \omega_s), \quad (41)$$

of which only the first is symmetric:

ingredient	symmetric under $A \leftrightarrow B$?	reason
$C, \det \rho, r$	yes	Schmidt: identical spectra
p_s	no	not an eigenvalue, but a matrix element in the <i>local</i> basis
ω_s	no (crossed family)	indices interchanged between modes

The decisive one is p_s . Isospectrality fixes the *eigenvalues* of ρ_s , but $p_s = (\rho_s)_{22}$ is not an eigenvalue: it is a matrix element in the Gram–Schmidt basis (6) built from *that mode's own* branches, whose overlap is g_α for A and h_β for B . Nothing forces $g_\alpha = h_\beta$.

Geometric reading. With $\cos \theta_s = (1 - 2p_s)/r$ one has the exact identity

$$b_s = \omega_s \sqrt{\cos^2 \theta_s + C^2 \sin^2 \theta_s}, \quad (42)$$

where θ_s is the *tilt* between the reduced Bloch vector of mode s and the σ_z axis of its own logical basis. Therefore:

Schmidt's theorem fixes the **length** of each reduced Bloch vector (r , common to both sides) but **not its orientation** relative to each mode's local axis. Uhlmann holonomy depends on that orientation.

Entanglement is symmetric; *local nonorthogonality is not*. Concretely, for $(m, n) = (0, 2)$, $\alpha = 1.1$, $\beta = 0.8$, $\mu = 0.6$ the two sides share $C = 0.748027$ exactly while $p_A = 0.4725$, $p_B = 0.5263$, i.e. tilts $\theta_A = 85.2^\circ$ and $\theta_B = 94.6^\circ$. Symmetry is restored exactly when the two local geometries coincide, $g_\alpha = h_\beta$, i.e. $\beta = \alpha$ for real amplitudes; Fig. 11(c) shows that the symmetry locus in the (α, β) plane is precisely the diagonal.

Antisymmetry in the crossed family. There $p_A = p_B$ identically but $\omega_B = -\omega_A$, hence $b_B = -b_A$. Inspecting Eq. (26): $(-1)^{\omega_B} = (-1)^{\omega_A}$ and $\cos(\pi b_B) = \cos(\pi b_A)$, so the real parts agree; while $\omega_B/b_B = \omega_A/b_A$ but $\sin(\pi b_B) = -\sin(\pi b_A)$, so the imaginary parts are opposite. Therefore $Z_U^B = \overline{Z_U^A}$ and

$$\boxed{\Phi_U^B = -\Phi_U^A, \quad \Phi_U^A + \Phi_U^B = 0} \quad (43)$$

exactly, for every α, β, μ, m, n — verified to floating-point zero in Sec. 11 and shown in Fig. 7(c). The entangled pair then accumulates no *net* Uhlmann holonomy even though each half accumulates a large one. Since complex conjugation does not move zeros, the two subsystems also share their nodal set exactly in that family.

11 Numerical validation

Every closed form is checked against an independent construction.

Berry sector: truncated two-mode Fock space. Because the global cycle is diagonal in the Fock basis, $e^{-i(\hat{n}_A + \hat{n}_B)\phi} |j\rangle_A |k\rangle_B = e^{-i(j+k)\phi} |j\rangle_A |k\rangle_B$, the evolution can be applied exactly. We build the full coefficient matrix C_{jk} with 90 levels per mode and evaluate overlaps, matrix elements and $\langle \hat{n}_A + \hat{n}_B \rangle$ by direct summation, with no recourse to hypergeometric identities.

Uhlmann sector: two connection-free holonomies. Method 1 is the discrete polar-decomposition algorithm, which implements the parallel-transport condition $W_j^\dagger W_{j+1} \geq 0$ directly: with $\sqrt{\rho_j} \sqrt{\rho_{j+1}} = V_j P_j$ one has $U_{j+1} = V_j^\dagger U_j$ and $\Phi_U = \text{Arg Tr}[\rho_0 U_{\text{hol}}]$. Method 2 integrates $\dot{U} = -i\Lambda U$ after solving Eq. (22) as a linear system at each step. Neither uses a closed-form connection.

Table 1: Independent validation. “Fock” denotes direct summation in the truncated two-mode space; “polar” and “ODE” the two connection-free holonomy algorithms.

Quantity	Checked against	Max. discrepancy
$\langle \alpha, m \beta, n \rangle$, Eq. (46)	Fock	2×10^{-16}
$\kappa_\alpha = g_{mn} Q_{mn}$, Eq. (48)	Fock	6×10^{-15}
Φ_B , Eq. (12)	Fock	2×10^{-14}
Mode split $\Phi_B = \Phi_B^{(A)} + \Phi_B^{(B)}$	Closed forms	9×10^{-16}
Symmetric case $\Phi_B = 2\Phi_B^{(A)}$	Closed forms	0 (exact)
Affine law, Eq. (14)	Direct evaluation	10^{-10}
p_s, C, ω_s , Eqs. (9)–(10)	Fock + winding fit	10^{-9}
Φ_U , Eq. (26)	Polar holonomy	2×10^{-6}
Φ_U , Eq. (26)	Transport ODE	2×10^{-6}
Tilt identity, Eq. (42)	Eq. (26)	2×10^{-16}
Nodes C_k , Eq. (29)	$ Z_U $ at the node	10^{-14}
Antisymmetry, Eq. (43)	Closed form	0 (exact)
$C = 2\mu\nu N_1 N_2 / \mathcal{N}^2$, Eq. (19)	$2 \det M $ and Fock	10^{-16}
$\sqrt{p_{APB}} = C/2$, Eq. (33)	Closed forms	10^{-16}
$p = C/2$ and Eq. (37) (balanced sym.)	Eq. (26)	2×10^{-16}
$\mathbb{Z}_2$ reduction on $p = 1/2$, Eq. (38)	$\text{Im } Z_U$	0 (exact)

As a sharp discriminant, the nodes predicted by Eq. (29) give $|\text{Tr}[\rho_0 U_{\text{hol}}]| \sim 10^{-15}$ in the ODE holonomy, i.e. genuine zeros, whereas the values $C = 2\Delta/(2k+1)$ that follow from the sandwich connection give $|\text{Tr}[\rho_0 U_{\text{hol}}]| = 0.31, 0.90, 0.94$ for $\Delta = 2, k = 2, 3, 4$ — not zeros at all.

12 Results

All figures use real amplitudes; “balanced” means $\mu = \nu = 1/\sqrt{2}$ and “symmetric” means $\beta = \alpha$.




Figure 1: Global Berry phase $\Phi_B/2\pi = \langle \hat{n}_A + \hat{n}_B \rangle$, Eq. (12), versus α for several photon-addition pairs in the balanced symmetric state. Growth is smooth because the connection is constant along the cycle; changing (m, n) shifts the curve through both the diagonal mean-photon terms and the interference term of Eq. (14).

Table 2: Critical amplitudes $\alpha_g(m, n)$ from $|g_{mn}(\alpha_g)|^2 = 1/2$.

$n \backslash m$	0	1	2	3	4
1	1.0000				
2	2.1094	0.7221			
3	3.2756	1.8000	0.5921		
4	4.4576	3.0354	1.5829	0.5135	
5	5.6464	4.2685	2.8269	1.4235	0.4596

13 Discussion

Choosing the global cycle organizes the problem into three clean levels.

Level 1 — the connection is a photon number. Because $\hat{n}_A + \hat{n}_B$ generates the cycle and commutes with itself, the Berry connection is constant and equals the total mean photon num-




Figure 2: Discrete dependence of the global Berry phase on the photon-addition indices at three coherent amplitudes. Only $n > m$ is shown, the ordering in which the overlap formula (46) is written.




Figure 3: Interference fringes produced by the tunable relative branch phase χ . The ϕ -cycle itself shows no oscillation because its connection, though containing the coherent cross term, is ϕ -independent; χ makes that interference explicit.




Figure 4: Global Berry phase in the (μ, α) plane. The map is completely smooth: a pure-state Berry phase never develops branch cuts, in contrast with the Uhlmann maps below.




Figure 5: Exact mode decomposition (13) of the global Berry phase for two fixed values of β . The total (black) is the sum of the two single-mode contributions; the mode whose amplitude is held fixed contributes an almost constant amount, so the split is markedly unequal away from $\alpha = \beta$.




Figure 6: Principal Uhlmann phase of ρ_A in the (μ, α) plane, Eq. (26), for the symmetric family. The dashed curve is the analytic condition $p = 1/2$, Eq. (30); the dotted line is α_g and the shaded region above it cannot contain nodes. Gold circles are the exact zeros of Z_U : for $\Delta = 2$ there are exactly two, labelled $k = 0, 1$, at $C_k = (2k + 1)/2\Delta$.




Figure 7: Uhlmann phase of *both* subsystems under the global cycle. (a) Branch-correlated family with $\beta = 0.8$ fixed: the two phases are different curves, meeting only at $\alpha = \beta$, even though they share a concurrence. (b) Symmetric family $\beta = \alpha$: $p_A = p_B$ by Eq. (9) and the curves coincide exactly. (c) Crossed family, where $\omega_B = -\omega_A$: the exact antisymmetry (43) holds and the sum (dash-dotted) vanishes identically.




Figure 8: Nodal structure of both subsystems in the (μ, α) plane at fixed $\beta = 0.8$. Both share the same nodal concurrences C_k , Eq. (29), but not their coordinates: the $p_A = 1/2$ curve bends and keeps the A nodes below α_g , whereas the existence criterion for B depends on β alone, so its $p_B = 1/2$ curve is nearly vertical and its nodes are not bounded by α_g (the $k = 0$ node of B lies at $\alpha \approx 5.4$, outside the plotted range).




Figure 9: Critical overlap structure. (a) $|g_{mn}(\alpha)|^2$ against the threshold $1/2$. (b) The discrete critical function $\alpha_g(m, n)$ over the grid of Table 2. (c) α_g versus Δ for several fixed m : larger photon-addition separation extends the nodal region to larger coherent amplitude.




Figure 10: Principal Uhlmann phase versus photon-addition separation Δ for the balanced symmetric state. The nonmonotonic behaviour reflects the dependence of $b = \Delta C$ on Δ through the overlap, combined with principal-branch wrapping of $\text{Arg}(\cdot) \in (-\pi, \pi]$.




Figure 11: Origin of the $A \leftrightarrow B$ asymmetry (Sec. 10). (a) The local populations differ, crossing only at $\alpha = \beta$. (b) The two Uhlmann phases track those populations. (c) Map of $\Phi_U^A - \Phi_U^B$ in the (α, β) plane: the symmetry locus is exactly the diagonal $\alpha = \beta$, where the two local overlaps coincide.




Figure 12: Concurrence of the entangled PACS, Eq. (19). (a) Balanced symmetric family: $C = (1 - q)/(1 + q)$ decreases monotonically from 1 at $\alpha \rightarrow 0$ (where the PACS reduce to orthogonal Fock states) towards 0 at large α (parallel branches). Larger Δ keeps the branches distinguishable longer. (b) C over the (μ, α) plane with contours. (c) Unequal amplitudes: fixing β caps the attainable concurrence, since C carries the factor $\sqrt{(1 - g_\alpha^2)(1 - h_\beta^2)}$.




Figure 13: Uhlmann phase as a function of the concurrence. (a) In the balanced symmetric family $p = C/2$, so Φ_U is a pure function of C through $b = \Delta\sqrt{C^2 + (1 - C)^2}$, Eq. (37); it returns to 0 at both $C = 0$ and $C = 1$. (b) b versus C on the two loci, with the half-integer levels dotted: on $p = 1/2$ the line $b = \Delta C$ crosses them exactly at the nodal concurrences (gold). The balanced curve may also cross a half-integer, but those are not nodes because $p \neq 1/2$ leaves $\text{Im } Z_U \neq 0$. (c) On $p = 1/2$ the phase is the $\mathbb{Z}_2$ staircase of Eq. (38), flipping between 0 and π exactly at the Δ nodal concurrences (curves offset vertically for visibility).

Concurrence and local asymmetry as independent coordinates: $\sqrt{p_A p_B} = C/2$, $\eta = \sqrt{p_A/p_B}$




Figure 14: Concurrence and local asymmetry as independent coordinates, Eqs. (33)–(34). (a) The physical region (35) in the (C, η) plane; it is symmetric under $\eta \rightarrow 1/\eta$ and pinches to the single point $\eta = 1$ as $C \rightarrow 1$, so maximal entanglement enforces $A \leftrightarrow B$ symmetry. (b) Φ_U^A versus C at fixed local asymmetry η . (c) $\Phi_U^A - \Phi_U^B$ over the same plane: the asymmetry vanishes identically on the line $\eta = 1$.

Table 3: Complete Uhlmann nodal set, Eqs. (29)–(31). Each Δ supports exactly Δ nodes; all satisfy $p = 1/2$ and $\alpha_k < \alpha_g$. The last column is the numerically evaluated $|Z_U|$.

(m, n)	Δ	k	C_k	μ_k	α_k	α_g	$ Z_U $
(0, 1)	1	0	0.5000	0.4472	0.5774	1.0000	4.5×10^{-16}
		1	0.7500	0.6000	0.9444	2.1094	1.8×10^{-16}
(0, 3)	3	0	0.1667	0.1644	2.8433	3.2756	3.4×10^{-15}
		1	0.5000	0.4472	2.1094	3.2756	1.8×10^{-16}
		2	0.8333	0.6402	1.3628	3.2756	2.1×10^{-15}
(1, 2)	1	0	0.5000	0.4472	0.4095	0.7221	5.7×10^{-15}
(1, 3)	2	0	0.2500	0.2425	1.3763	1.8000	6.1×10^{-17}
		1	0.7500	0.6000	0.7265	1.8000	6.6×10^{-14}
(1, 4)	3	0	0.1667	0.1644	2.5872	3.0354	5.5×10^{-15}
		1	0.5000	0.4472	1.8378	3.0354	5.0×10^{-15}
		2	0.8333	0.6402	1.1224	3.0354	2.4×10^{-15}
(2, 4)	2	0	0.2500	0.2425	1.1867	1.5829	3.1×10^{-15}
		1	0.7500	0.6000	0.6130	1.5829	1.8×10^{-16}

ber. Everything in the Berry sector is therefore smooth, single-valued and additive across modes, Eq. (13). The interference between branches is present in the connection but invisible in ϕ ; it takes a separately controllable branch phase to expose it.

Level 2 — entanglement enters affinely. In the balanced symmetric family the global Berry phase is an affine function of the concurrence, Eq. (14), interpolating between a mean-photon average and a pure interference term. This is as far as a pure-state geometric phase can go: it cannot be singular.

Level 3 — mixed-state holonomy, and it has two sides. Tracing out either mode leaves a genuinely mixed state, and the Uhlmann holonomy can be singular. The compact result (25) — parallel transport multiplies the off-diagonal generator by the concurrence — makes the structure transparent: at $p = 1/2$ the transported generator has magnitude ΔC , and the singularity condition $|b| = k + 1/2$ quantizes C to the finite ladder (29).

The most instructive feature is the interplay between what entanglement fixes and what it does not. The concurrence is shared by the two subsystems and determines *which* values are geometrically singular — the ladder C_k is a property of the pair. But it does not determine *where* those singularities sit, nor even whether the two sides have the same phase: the local populations p_A, p_B , equivalently the tilts θ_A, θ_B of the reduced Bloch vectors relative to each mode's own logical axis, are set by the local nonorthogonality g_α and h_β , which Schmidt's theorem leaves free. Symmetry is recovered exactly on the diagonal $\beta = \alpha$; and in the crossed family, where the local geometries do coincide but the indices are exchanged, one gets not symmetry but the exact antisymmetry $\Phi_U^A + \Phi_U^B = 0$. Entanglement is symmetric; local geometry need not be, and mixed-state holonomy sees both.

Finally, the critical amplitude α_g is not a thermodynamic critical point but an overlap-controlled boundary: it is fixed entirely by $|g_{mn}(\alpha_g)|^2 = 1/2$ and separates the region where equal logical-basis population is kinematically accessible from the region where it is not. Larger α makes the two photon-added branches more distinguishable, pushing the achievable population imbalance away from the symmetric point that an exact node requires.

14 Conclusions

Rebuilding the analysis of entangled PACS around the global two-mode cycle $U(\phi) = e^{-i(\hat{n}_A + \hat{n}_B)\phi}$ yields:

- (i) a closed-form global Berry phase equal to the total mean photon number, Eq. (12), exactly additive across modes and affine in the concurrence in the balanced symmetric family;
- (ii) the Uhlmann connection from the anticommutator transport condition, giving $(G + \Lambda)_{\text{od}} = C G_{\text{od}}$ and $b = \omega \sqrt{1 + C^2 - 4p(1 - p)}$, validated against two connection-free holonomy algorithms;
- (iii) a finite nodal ladder,

$$p = \frac{1}{2}, \quad C_k = \frac{2k + 1}{2\Delta}, \quad |g_{mn}(\alpha_k)|^2 = \frac{1 - C_k}{2}, \quad k = 0, \dots, \Delta - 1, \quad (44)$$

i.e. exactly Δ isolated Uhlmann singularities per family, common to both subsystems and all bounded by α_g ;

- (iv) a precise account of the $A \leftrightarrow B$ asymmetry: shared concurrence, unshared local populations, with symmetry on $\beta = \alpha$ and exact antisymmetry $\Phi_U^B = -\Phi_U^A$ in the crossed family.

Taken together these results provide an analytic bridge between photon addition, nonorthogonality, bipartite entanglement and mixed-state quantum holonomy, in a model where every step can be written — and independently checked — in closed form.

A Scalar products of photon-added coherent states

Inserting Eq. (3) into $\langle \alpha, m | \beta, n \rangle$ and using Fock orthogonality, only terms with a common level survive. Writing $\ell = n + p$ and $\Delta = n - m \geq 0$,

$$\langle \alpha, m | \beta, n \rangle = A_m A_n \alpha^\Delta \sum_{p=0}^{\infty} (\alpha\beta)^p \frac{(n+p)!}{(\Delta+p)! p!}, \quad A_n = \frac{e^{-\alpha^2/2}}{\sqrt{n! L_n(-\alpha^2)}}. \quad (45)$$

The key step is to recognize the factorial ratio as a ratio of Pochhammer symbols, $\frac{(n+p)!}{(\Delta+p)!} = \frac{n!}{\Delta!} \frac{(n+1)_p}{(\Delta+1)_p}$, which converts the sum into a confluent hypergeometric series and gives

$$\langle \alpha, m | \beta, n \rangle = \frac{e^{-\frac{1}{2}(\alpha^2 + \beta^2)} \alpha^\Delta n!}{\Delta! \sqrt{m! n! L_m(-\alpha^2) L_n(-\beta^2)}} {}_1F_1(n+1; \Delta+1; \alpha\beta) \quad (46)$$

for $0 \leq m \leq n$, the amplitude carrying the exponent Δ being the one attached to the *lower* index. Note that the ratio $n!/\Delta!$ arises whole from the Pochhammer reduction and therefore sits *outside* the square root. We verify Eq. (46) against direct Fock summation to 2×10^{-16} .

B Cross matrix element of the number operator

Repeating the sum with each term weighted by its Fock eigenvalue $\ell = n + p$,

$$\kappa_\alpha^{mn} = \langle \alpha, m | a^\dagger a | \alpha, n \rangle = A_m A_n \alpha^\Delta [n S(x) + x S'(x)], \quad x = \alpha^2, \quad (47)$$

where $S(x) = (n!/\Delta!) {}_1F_1(n+1; \Delta+1; x)$ is the sum of Appendix A. Using $\frac{d}{dx} {}_1F_1(a; b; x) = \frac{a}{b} {}_1F_1(a+1; b+1; x)$ together with a Kummer contiguous relation, this reduces to

$$\kappa_\alpha^{mn} = g_{mn}(\alpha) Q_{mn}(x), \quad Q_{mn}(x) = (n+1) \frac{{}_1F_1(n+2; \Delta+1; x)}{{}_1F_1(n+1; \Delta+1; x)} - 1. \quad (48)$$

Being a dimensionless ratio, Q_{mn} is insensitive to the prefactor conventions of Appendix A. Verified against Fock summation to 6×10^{-15} .

C Numerical conventions

All results use real amplitudes. Balanced plots take $\mu = \nu = 1/\sqrt{2}$; two-dimensional weight maps impose $\nu = \sqrt{1 - \mu^2}$ with $0 \leq \mu \leq 1$. Uhlmann plots show the principal value $\text{Arg } Z_U \in (-\pi, \pi]$. The shaded region above α_g in Figs. 6 and 8 is analytic, not a numerical cutoff: it is where $p = 1/2$ has no solution. Root-finding uses bisection on the monotonic $|g_{mn}(\alpha)|^2$; Laguerre polynomials and ${}_1F_1(a; b; z)$ are evaluated with `scipy.special`. The truncated Fock space uses 90 levels per mode, ample for $\alpha \lesssim 3$. The complete code accompanies this manuscript as `pacs_global_cycle.ipynb`.

References

- [1] G. S. Agarwal and K. Tara, *Nonclassical properties of states generated by the excitations on a coherent state*, Phys. Rev. A **43**, 492 (1991).
- [2] M. V. Berry, *Quantal phase factors accompanying adiabatic changes*, Proc. R. Soc. Lond. A **392**, 45 (1984).
- [3] A. Uhlmann, *Parallel transport and ‘quantum holonomy’ along density operators*, Rep. Math. Phys. **24**, 229 (1986).
- [4] W. K. Wootters, *Entanglement of formation of an arbitrary state of two qubits*, Phys. Rev. Lett. **80**, 2245 (1998).
- [5] F. A. Domínguez-Serna, F. J. Mendieta-Jiménez and F. Rojas, *Entangled photon-added coherent states*, Quantum Inf. Process. **15**, 3121 (2016).
- [6] S. Mohammadi Almas, G. Najarbashi and A. Tavana, *Geometric phase for two-mode entangled coherent states*, Int. J. Theor. Phys. **61**, 192 (2022).
- [7] D.-B. Yang, Y. Chen, F.-L. Zhang and J.-L. Chen, *Geometric phases for nonlinear coherent and squeezed states*, J. Phys. B **44**, 075502 (2011).
- [8] X. Wang, X.-Y. Hou, Z. Zhou, H. Guo and C.-C. Chien, *Uhlmann phase of coherent states and the Uhlmann–Berry correspondence*, SciPost Phys. Core **6**, 024 (2023).