




Figure D.259: Velocity profiles at $x = 0$ for the 1×1 SRG at $\phi = 0.15$ and $\text{Re}_H = 100$, comparing the SSF, DFM, SSF-VCDFM, TIDFT-VCDFM, and ML-VCDFM. Profiles of $w(y)$, $v(y)$, and $u(y)$ are shown at $z = (\Gamma + 6)/2$ in (a–c) and $z = (\Gamma + 8)/2$ in (d–f). ML-VCDFM is evaluated outside the Reynolds-number range used for training.




Figure D.260: Detailed comparison between the geometry-resolved small-scale flow (SSF) and SSF-VCDFM using F_z^{SSF} for the 1×1 SRG at $\phi = 0.15$ and $Re_H = 100$. The upper panels compare streamwise velocity w and transverse velocity v in the central (z, y) -plane, restricted to the correction region. The lower panels compare the corresponding transverse (x, y) -plane distributions of w , v , and u , from left to right, at $z = (\Gamma + 6)/2$ in the first row and $z = (\Gamma + 8)/2$ in the second row.




Figure D.261: Detailed comparison between the geometry-resolved small-scale flow (SSF) and TIDFT-VCDFM using F_z^{TIDFT} for the 1×1 SRG at $\phi = 0.15$ and $\text{Re}_H = 100$. The upper panels compare streamwise velocity w and transverse velocity v in the central (z, y) -plane, restricted to the correction region. The lower panels compare the corresponding transverse (x, y) -plane distributions of w , v , and u , from left to right, at $z = (\Gamma + 6)/2$ in the first row and $z = (\Gamma + 8)/2$ in the second row.




Figure D.262: Detailed comparison between the geometry-resolved small-scale flow (SSF) and ML-VCDFM using F_z^{ML} for the 1×1 SRG at $\phi = 0.15$ and $Re_H = 100$. The upper panels compare streamwise velocity w and transverse velocity v in the central (z, y) -plane, restricted to the correction region. The lower panels compare the corresponding transverse (x, y) -plane distributions of w , v , and u , from left to right, at $z = (\Gamma + 6)/2$ in the first row and $z = (\Gamma + 8)/2$ in the second row. This case tests extrapolation beyond the Reynolds-number range used for training; deviations from the SSF appear in the velocity fields.




Figure D.263: Streamwise evolution of (a) $\|w\|_{L_2}$, (b) $\|p\|_{L_2}$, (c) $\|v\|_{L_2}$, and (d) centerline velocity $w(0,0,z)$ for the 1×1 SRG at $\phi = 0.2$ and $\text{Re}_H = 100$. The figure compares the SSF, DFM, SSF-VCDFM, TIDFT-VCDFM, and ML-VCDFM. Shading marks the grid and the characteristic entrance, exit, peak, and decay regions. Only the SSF-based $\mathcal{L}_{peak}$ and $\mathcal{L}_{decay}$ are marked on the plot; the corresponding TIDFT-VCDFM and ML-VCDFM values appear in the legend. The ML-VCDFM result tests extrapolation beyond the Reynolds-number range used for training and deviates from the reference SSF solution.




Figure D.264: Flow organization in the streamwise (z, y) -plane for the 1×1 SRG at $\phi = 0.2$, $\text{Re}_H = 100$, and $\alpha = 0^\circ$. The blocks show (a) pressure p with streamlines, (b) streamwise velocity w , and (c) transverse velocity v . Within each block, the rows show, from top to bottom, the SSF, DFM, SSF-VCDFM, TIDFT-VCDFM, and ML-VCDFM. The ML-VCDFM result tests extrapolation beyond the Reynolds-number range used for training and shows deviations from the reference SSF. Pink shading marks the grid walls in the SSF; olive shading marks the equivalent porous zone in the model solutions.




Figure D.265: Velocity profiles at $x = 0$ for the 1×1 SRG at $\phi = 0.2$ and $\text{Re}_H = 100$, comparing the SSF, DFM, SSF-VCDFM, TIDFT-VCDFM, and ML-VCDFM. Profiles of $w(y)$, $v(y)$, and $u(y)$ are shown at $z = (\Gamma + 6)/2$ in (a–c) and $z = (\Gamma + 8)/2$ in (d–f). ML-VCDFM is evaluated outside the Reynolds-number range used for training.




Figure D.266: Detailed comparison between the geometry-resolved small-scale flow (SSF) and SSF-VCDFM using F_z^{SSF} for the 1×1 SRG at $\phi = 0.2$ and $\text{Re}_H = 100$. The upper panels compare streamwise velocity w and transverse velocity v in the central (z, y) -plane, restricted to the correction region. The lower panels compare the corresponding transverse (x, y) -plane distributions of w , v , and u , from left to right, at $z = (\Gamma + 6)/2$ in the first row and $z = (\Gamma + 8)/2$ in the second row.




Figure D.267: Detailed comparison between the geometry-resolved small-scale flow (SSF) and TIDFT-VCDFM using F_z^{TIDFT} for the 1×1 SRG at $\phi = 0.2$ and $\text{Re}_H = 100$. The upper panels compare streamwise velocity w and transverse velocity v in the central (z, y) -plane, restricted to the correction region. The lower panels compare the corresponding transverse (x, y) -plane distributions of w , v , and u , from left to right, at $z = (\Gamma + 6)/2$ in the first row and $z = (\Gamma + 8)/2$ in the second row.




Figure D.268: Detailed comparison between the geometry-resolved small-scale flow (SSF) and ML-VCDFM using F_z^{ML} for the 1×1 SRG at $\phi = 0.2$ and $\text{Re}_H = 100$. The upper panels compare streamwise velocity w and transverse velocity v in the central (z, y) -plane, restricted to the correction region. The lower panels compare the corresponding transverse (x, y) -plane distributions of w , v , and u , from left to right, at $z = (\Gamma + 6)/2$ in the first row and $z = (\Gamma + 8)/2$ in the second row. This case tests extrapolation beyond the Reynolds-number range used for training; deviations from the SSF appear in the velocity fields.




Figure D.269: Streamwise evolution of (a) $\|w\|_{L_2}$, (b) $\|p\|_{L_2}$, (c) $\|v\|_{L_2}$, and (d) centerline velocity $w(0,0,z)$ for the 1×1 SRG at $\phi = 0.25$ and $\text{Re}_H = 100$. The figure compares the SSF, DFM, SSF-VCDFM, TIDFT-VCDFM, and ML-VCDFM. Shading marks the grid and the characteristic entrance, exit, peak, and decay regions. Only the SSF-based $\mathcal{L}_{peak}$ and $\mathcal{L}_{decay}$ are marked on the plot; the corresponding TIDFT-VCDFM and ML-VCDFM values appear in the legend. The ML-VCDFM result tests extrapolation beyond the Reynolds-number range used for training and deviates from the reference SSF solution.




Figure D.270: Flow organization in the streamwise (z, y) -plane for the 1×1 SRG at $\phi = 0.25$, $\text{Re}_H = 100$, and $\alpha = 0^\circ$. The blocks show (a) pressure p with streamlines, (b) streamwise velocity w , and (c) transverse velocity v . Within each block, the rows show, from top to bottom, the SSF, DFM, SSF-VCDFM, TIDFT-VCDFM, and ML-VCDFM. The ML-VCDFM result tests extrapolation beyond the Reynolds-number range used for training and shows deviations from the reference SSF. Pink shading marks the grid walls in the SSF; olive shading marks the equivalent porous zone in the model solutions.




Figure D.271: Velocity profiles at $x = 0$ for the 1×1 SRG at $\phi = 0.25$ and $\text{Re}_H = 100$, comparing the SSF, DFM, SSF-VCDFM, TIDFT-VCDFM, and ML-VCDFM. Profiles of $w(y)$, $v(y)$, and $u(y)$ are shown at $z = (\Gamma + 6)/2$ in (a–c) and $z = (\Gamma + 8)/2$ in (d–f). ML-VCDFM is evaluated outside the Reynolds-number range used for training.




Figure D.272: Detailed comparison between the geometry-resolved small-scale flow (SSF) and SSF-VCDFM using F_z^{SSF} for the 1 × 1 SRG at $\phi = 0.25$ and $\text{Re}_H = 100$. The upper panels compare streamwise velocity w and transverse velocity v in the central (z, y) -plane, restricted to the correction region. The lower panels compare the corresponding transverse (x, y) -plane distributions of w , v , and u , from left to right, at $z = (\Gamma + 6)/2$ in the first row and $z = (\Gamma + 8)/2$ in the second row.




Figure D.273: Detailed comparison between the geometry-resolved small-scale flow (SSF) and TIDFT-VCDFM using F_z^{TIDFT} for the 1×1 SRG at $\phi = 0.25$ and $Re_H = 100$. The upper panels compare streamwise velocity w and transverse velocity v in the central (z, y) -plane, restricted to the correction region. The lower panels compare the corresponding transverse (x, y) -plane distributions of w , v , and u , from left to right, at $z = (\Gamma + 6)/2$ in the first row and $z = (\Gamma + 8)/2$ in the second row.




Figure D.274: Detailed comparison between the geometry-resolved small-scale flow (SSF) and ML-VCDFM using F_z^{ML} for the 1×1 SRG at $\phi = 0.25$ and $\text{Re}_H = 100$. The upper panels compare streamwise velocity w and transverse velocity v in the central (z, y) -plane, restricted to the correction region. The lower panels compare the corresponding transverse (x, y) -plane distributions of w , v , and u , from left to right, at $z = (\Gamma + 6)/2$ in the first row and $z = (\Gamma + 8)/2$ in the second row. This case tests extrapolation beyond the Reynolds-number range used for training; deviations from the SSF appear in the velocity fields.

