

In this episode...

- Linear Independence

In this episode...

- ▶ Linear Independence
- ▶ Linear Transformation

In this episode...

- ▶ Linear Independence
- ▶ Linear Transformation
- ▶ Existence and One to One Mapping

In this episode...

In this episode...



In this episode...



In this episode...

In this episode...



In this episode...



In this episode...

In this episode...



In this episode...



In this episode...

In this episode...



In this episode...



Linear Independence

1.7 Linear Independence

Homogeneous Vector Equations

The homogeneous equations in Section 1.5 can be studied from a different perspective by writing them as vector equations. In this way, the focus shifts from the unknown solutions of $A\mathbf{x} = \mathbf{0}$ to the vectors that appear in the vector equations.

Homogeneous Vector Equations

The homogeneous equations in Section 1.5 can be studied from a different perspective by writing them as vector equations. In this way, the focus shifts from the unknown solutions of $A\mathbf{x} = \mathbf{0}$ to the vectors that appear in the vector equations.

For instance, consider

$$x_1 \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} + x_2 \begin{bmatrix} 4 \\ 5 \\ 6 \end{bmatrix} + x_3 \begin{bmatrix} 2 \\ 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}. \quad (1)$$

Homogeneous Vector Equations

The homogeneous equations in Section 1.5 can be studied from a different perspective by writing them as vector equations. In this way, the focus shifts from the unknown solutions of $A\mathbf{x} = \mathbf{0}$ to the vectors that appear in the vector equations.

For instance, consider

$$x_1 \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} + x_2 \begin{bmatrix} 4 \\ 5 \\ 6 \end{bmatrix} + x_3 \begin{bmatrix} 2 \\ 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}. \quad (1)$$

This equation has the trivial solution $x_1 = x_2 = x_3 = 0$. The main issue is whether the trivial solution is the only solution.

Linear Independence

Definition

An indexed set of vectors $\{\mathbf{v}_1, \dots, \mathbf{v}_p\}$ in $\mathbb{R}^n$ is **linearly independent** if

$$x_1\mathbf{v}_1 + x_2\mathbf{v}_2 + \cdots + x_p\mathbf{v}_p = \mathbf{0}$$

has only the trivial solution. The set is **linearly dependent** if there exist weights $c_1, \dots, c_p$, not all zero, such that

$$c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + \cdots + c_p\mathbf{v}_p = \mathbf{0}. \quad (2)$$

Linear Independence

Definition

An indexed set of vectors $\{\mathbf{v}_1, \dots, \mathbf{v}_p\}$ in $\mathbb{R}^n$ is **linearly independent** if

$$x_1\mathbf{v}_1 + x_2\mathbf{v}_2 + \cdots + x_p\mathbf{v}_p = \mathbf{0}$$

has only the trivial solution. The set is **linearly dependent** if there exist weights $c_1, \dots, c_p$, not all zero, such that

$$c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + \cdots + c_p\mathbf{v}_p = \mathbf{0}. \quad (2)$$

Equation (2) is called a **linear dependence relation** among the vectors when the weights are not all zero.

Example 1

Let

$$\mathbf{v}_1 = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}, \quad \mathbf{v}_2 = \begin{bmatrix} 4 \\ 5 \\ 6 \end{bmatrix}, \quad \mathbf{v}_3 = \begin{bmatrix} 2 \\ 1 \\ 0 \end{bmatrix}.$$

Example 1

Let

$$\mathbf{v}_1 = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}, \quad \mathbf{v}_2 = \begin{bmatrix} 4 \\ 5 \\ 6 \end{bmatrix}, \quad \mathbf{v}_3 = \begin{bmatrix} 2 \\ 1 \\ 0 \end{bmatrix}.$$

- Determine whether $\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$ is linearly independent.

Example 1

Let

$$\mathbf{v}_1 = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}, \quad \mathbf{v}_2 = \begin{bmatrix} 4 \\ 5 \\ 6 \end{bmatrix}, \quad \mathbf{v}_3 = \begin{bmatrix} 2 \\ 1 \\ 0 \end{bmatrix}.$$

- a. Determine whether $\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$ is linearly independent.
- b. If possible, find a linear dependence relation among the three vectors.

Example 1

Let

$$\mathbf{v}_1 = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}, \quad \mathbf{v}_2 = \begin{bmatrix} 4 \\ 5 \\ 6 \end{bmatrix}, \quad \mathbf{v}_3 = \begin{bmatrix} 2 \\ 1 \\ 0 \end{bmatrix}.$$

- a. Determine whether $\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$ is linearly independent.
- b. If possible, find a linear dependence relation among the three vectors.

Solution idea

We must determine if there is a nontrivial solution of the equation on the previous slide.

Example 1: Row Reduction

Row operations on the associated augmented matrix show that

Example 1: Row Reduction

Row operations on the associated augmented matrix show that

$$\left[\begin{array}{ccc|c} 1 & 4 & 2 & 0 \\ 2 & 5 & 1 & 0 \\ 3 & 6 & 0 & 0 \end{array} \right] \sim \left[\begin{array}{ccc|c} 1 & 4 & 2 & 0 \\ 0 & -3 & -3 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right].$$

Example 1: Row Reduction

Row operations on the associated augmented matrix show that

$$\left[\begin{array}{ccc|c} 1 & 4 & 2 & 0 \\ 2 & 5 & 1 & 0 \\ 3 & 6 & 0 & 0 \end{array} \right] \sim \left[\begin{array}{ccc|c} 1 & 4 & 2 & 0 \\ 0 & -3 & -3 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right].$$

- x_1 and x_2 are basic variables, and x_3 is free.

Example 1: Row Reduction

Row operations on the associated augmented matrix show that

$$\left[\begin{array}{ccc|c} 1 & 4 & 2 & 0 \\ 2 & 5 & 1 & 0 \\ 3 & 6 & 0 & 0 \end{array} \right] \sim \left[\begin{array}{ccc|c} 1 & 4 & 2 & 0 \\ 0 & -3 & -3 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right].$$

- ▶ x_1 and x_2 are basic variables, and x_3 is free.
- ▶ Each nonzero value of x_3 determines a nontrivial solution of (1).

Example 1: Row Reduction

Row operations on the associated augmented matrix show that

$$\left[\begin{array}{ccc|c} 1 & 4 & 2 & 0 \\ 2 & 5 & 1 & 0 \\ 3 & 6 & 0 & 0 \end{array} \right] \sim \left[\begin{array}{ccc|c} 1 & 4 & 2 & 0 \\ 0 & -3 & -3 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right].$$

- ▶ x_1 and x_2 are basic variables, and x_3 is free.
- ▶ Each nonzero value of x_3 determines a nontrivial solution of (1).
- ▶ Hence $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3$ are linearly dependent.

Example 1: A Dependence Relation

To find a linear dependence relation among $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3$, completely row reducing the augmented matrix gives

Example 1: A Dependence Relation

To find a linear dependence relation among $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3$, completely row reducing the augmented matrix gives

$$\left[\begin{array}{ccc|c} 1 & 0 & -2 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right], \quad \begin{aligned} x_1 - 2x_3 &= 0, \\ x_2 + x_3 &= 0. \\ 0 &= 0. \end{aligned}$$

Example 1: A Dependence Relation

To find a linear dependence relation among $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3$, completely row reducing the augmented matrix gives

$$\left[\begin{array}{ccc|c} 1 & 0 & -2 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right], \quad \begin{array}{l} x_1 - 2x_3 = 0, \\ x_2 + x_3 = 0. \\ 0 = 0. \end{array}$$

► Thus $x_1 = 2x_3$, $x_2 = -x_3$, and x_3 is free.

Example 1: A Dependence Relation

To find a linear dependence relation among $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3$, completely row reducing the augmented matrix gives

$$\left[\begin{array}{ccc|c} 1 & 0 & -2 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right], \quad \begin{array}{l} x_1 - 2x_3 = 0, \\ x_2 + x_3 = 0. \\ 0 = 0. \end{array}$$

- ▶ Thus $x_1 = 2x_3$, $x_2 = -x_3$, and x_3 is free.
- ▶ Choose, for example, $x_3 = 5$.

Example 1: A Dependence Relation

To find a linear dependence relation among $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3$, completely row reducing the augmented matrix gives

$$\left[\begin{array}{ccc|c} 1 & 0 & -2 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right], \quad \begin{aligned} x_1 - 2x_3 &= 0, \\ x_2 + x_3 &= 0. \\ 0 &= 0. \end{aligned}$$

- ▶ Thus $x_1 = 2x_3$, $x_2 = -x_3$, and x_3 is free.
- ▶ Choose, for example, $x_3 = 5$.
- ▶ Then $x_1 = 10$ and $x_2 = -5$.

Example 1: Conclusion

- ▶ Substituting the values into the vector equation gives

Example 1: Conclusion

- Substituting the values into the vector equation gives

$$10\mathbf{v}_1 - 5\mathbf{v}_2 + 5\mathbf{v}_3 = \mathbf{0}.$$

Example 1: Conclusion

- ▶ Substituting the values into the vector equation gives

$$10\mathbf{v}_1 - 5\mathbf{v}_2 + 5\mathbf{v}_3 = \mathbf{0}.$$

- ▶ This is one of infinitely many possible linear dependence relations among $\mathbf{v}_1$, $\mathbf{v}_2$, and $\mathbf{v}_3$.

Columns of a Matrix

Suppose we begin with a matrix $A = [\mathbf{a}_1 \ \cdots \ \mathbf{a}_n]$ instead of a set of vectors.

Columns of a Matrix

Suppose we begin with a matrix $A = [\mathbf{a}_1 \cdots \mathbf{a}_n]$ instead of a set of vectors.
Then the matrix equation

$$A\mathbf{x} = \mathbf{0}$$

Columns of a Matrix

Suppose we begin with a matrix $A = [\mathbf{a}_1 \cdots \mathbf{a}_n]$ instead of a set of vectors.
Then the matrix equation

$$A\mathbf{x} = \mathbf{0}$$

can be written as

$$x_1\mathbf{a}_1 + x_2\mathbf{a}_2 + \cdots + x_n\mathbf{a}_n = \mathbf{0}.$$

Columns of a Matrix

Suppose we begin with a matrix $A = [\mathbf{a}_1 \cdots \mathbf{a}_n]$ instead of a set of vectors.
Then the matrix equation

$$A\mathbf{x} = \mathbf{0}$$

can be written as

$$x_1\mathbf{a}_1 + x_2\mathbf{a}_2 + \cdots + x_n\mathbf{a}_n = \mathbf{0}.$$

- *Each linear dependence relation among the columns of A corresponds to a nontrivial solution of $A\mathbf{x} = \mathbf{0}$.*

Columns of a Matrix

Suppose we begin with a matrix $A = [\mathbf{a}_1 \cdots \mathbf{a}_n]$ instead of a set of vectors.
Then the matrix equation

$$A\mathbf{x} = \mathbf{0}$$

can be written as

$$x_1\mathbf{a}_1 + x_2\mathbf{a}_2 + \cdots + x_n\mathbf{a}_n = \mathbf{0}.$$

- ▶ *Each linear dependence relation among the columns of A corresponds to a nontrivial solution of $A\mathbf{x} = \mathbf{0}$.*
- ▶ The columns of A are linearly independent if and only if $A\mathbf{x} = \mathbf{0}$ has *only the trivial solution*.

Sets of One Vector

- A set containing only one vector, say $\mathbf{v}$, is linearly independent if and only if $\mathbf{v}$ is not the zero vector $\mathbf{0}$.

Sets of One Vector

- ▶ A set containing only one vector, say $\mathbf{v}$, is linearly independent if and only if $\mathbf{v}$ is not the zero vector $\mathbf{0}$.
- ▶ This is because the vector equation $x_1 \mathbf{v} = \mathbf{0}$ has only the trivial solution when $\mathbf{v} \neq \mathbf{0}$.

Sets of One Vector

- ▶ A set containing only one vector, say $\mathbf{v}$, is linearly independent if and only if $\mathbf{v}$ is not the zero vector $\mathbf{0}$.
- ▶ This is because the vector equation $x_1\mathbf{v} = \mathbf{0}$ has only the trivial solution when $\mathbf{v} \neq \mathbf{0}$.
- ▶ The zero vector is linearly dependent because $x_1\mathbf{0} = \mathbf{0}$ has many nontrivial solutions.

Sets of Two Vectors

- A set $\{\mathbf{v}_1, \mathbf{v}_2\}$ is linearly independent if and only if neither vector is a multiple of the other.

Sets of Two Vectors

- ▶ A set $\{\mathbf{v}_1, \mathbf{v}_2\}$ is linearly independent if and only if neither vector is a multiple of the other.
- ▶ The set is linearly dependent if at least one vector is a scalar multiple of the other.

Sets of Two or More Vectors

THEOREM 7

Characterization of Linearly Dependent Sets. An indexed set $S = \{\mathbf{v}_1, \dots, \mathbf{v}_p\}$ of two or more vectors is linearly dependent if and only if at least one vector in S is a linear combination of the others. In fact, if S is linearly dependent and $\mathbf{v}_1 \neq \mathbf{0}$, then some $\mathbf{v}_j$ with $j > 1$ is a linear combination of the preceding vectors $\mathbf{v}_1, \dots, \mathbf{v}_{j-1}$.

Using Theorem 7

- Theorem 7 *does not* say that *every* vector in a linearly dependent set is a linear combination of the preceding vectors.

Using Theorem 7

- ▶ Theorem 7 *does not* say that *every* vector in a linearly dependent set is a linear combination of the preceding vectors.
- ▶ A vector in a linearly dependent set may fail to be a linear combination of the other vectors.

Using Theorem 7

- ▶ Theorem 7 *does not* say that *every* vector in a linearly dependent set is a linear combination of the preceding vectors.
- ▶ A vector in a linearly dependent set may fail to be a linear combination of the other vectors.

Example 4. Let

$$\mathbf{u} = \begin{bmatrix} 3 \\ 1 \\ 0 \end{bmatrix}, \quad \mathbf{v} = \begin{bmatrix} 1 \\ 6 \\ 0 \end{bmatrix}.$$

Using Theorem 7

- ▶ Theorem 7 *does not* say that *every* vector in a linearly dependent set is a linear combination of the preceding vectors.
- ▶ A vector in a linearly dependent set may fail to be a linear combination of the other vectors.

Example 4. Let

$$\mathbf{u} = \begin{bmatrix} 3 \\ 1 \\ 0 \end{bmatrix}, \quad \mathbf{v} = \begin{bmatrix} 1 \\ 6 \\ 0 \end{bmatrix}.$$

Describe $\text{Span}\{\mathbf{u}, \mathbf{v}\}$, and explain why $\mathbf{w}$ is in this span if and only if $\{\mathbf{u}, \mathbf{v}, \mathbf{w}\}$ is linearly dependent.

Example 4: Solution

- **Solution:** $\mathbf{u}$ and $\mathbf{v}$ are linearly independent because neither is a multiple of the other; consequently, they span a plane in

Example 4: Solution

- ▶ **Solution:** $\mathbf{u}$ and $\mathbf{v}$ are linearly independent because neither is a multiple of the other; consequently, they span a plane in $\mathbb{R}^3$.
- ▶ $\text{Span}\{\mathbf{u}, \mathbf{v}\}$ is the x_1x_2 -plane, where $x_3 = 0$.


Example 4: Solution

- ▶ **Solution:** $\mathbf{u}$ and $\mathbf{v}$ are linearly independent because neither is a multiple of the other; consequently, they span a plane in $\mathbb{R}^3$.
- ▶ $\text{Span}\{\mathbf{u}, \mathbf{v}\}$ is the x_1x_2 -plane, where $x_3 = 0$.
- ▶ If $\mathbf{w}$ is a linear combination of $\mathbf{u}$ and $\mathbf{v}$, then $\{\mathbf{u}, \mathbf{v}, \mathbf{w}\}$ is linearly dependent by Theorem 7.
- ▶ Conversely, if $\{\mathbf{u}, \mathbf{v}, \mathbf{w}\}$ is dependent

Example 4: Solution


- ▶ **Solution:** $\mathbf{u}$ and $\mathbf{v}$ are linearly independent because neither is a multiple of the other; consequently, they span a plane in $\mathbb{R}^3$.
- ▶ $\text{Span}\{\mathbf{u}, \mathbf{v}\}$ is the x_1x_2 -plane, where $x_3 = 0$.
- ▶ If $\mathbf{w}$ is a linear combination of $\mathbf{u}$ and $\mathbf{v}$, then $\{\mathbf{u}, \mathbf{v}, \mathbf{w}\}$ is linearly dependent by Theorem 7.
- ▶ Conversely, if $\{\mathbf{u}, \mathbf{v}, \mathbf{w}\}$ is dependent
- ▶ By Theorem 7, some vector in $\{\mathbf{u}, \mathbf{v}, \mathbf{w}\}$ is a linear combination of the preceding vectors. Since $\mathbf{u} \neq 0$ and $\mathbf{u}, \mathbf{v}$ are independent, that vector must be $\mathbf{w}$.

Example 4: Geometric View




Linearly dependent: $\mathbf{w} \in \text{Span}\{\mathbf{u}, \mathbf{v}\}$.

Example 4: Geometric View




Linearly dependent: $\mathbf{w} \in \text{Span}\{\mathbf{u}, \mathbf{v}\}$.




Linearly independent: $\mathbf{w} \notin \text{Span}\{\mathbf{u}, \mathbf{v}\}$.

Example 4: Geometric View




Linearly dependent: $\mathbf{w} \in \text{Span}\{\mathbf{u}, \mathbf{v}\}$.




Linearly independent: $\mathbf{w} \notin \text{Span}\{\mathbf{u}, \mathbf{v}\}$.

- The example 4 generalizes to any set $\{\mathbf{u}, \mathbf{v}, \mathbf{w}\}$ in $\mathbb{R}^3$ with $\mathbf{u}$ and $\mathbf{v}$ linearly independent.

Example 4: Geometric View



Linearly dependent: $\mathbf{w} \in \text{Span}\{\mathbf{u}, \mathbf{v}\}$.



Linearly independent: $\mathbf{w} \notin \text{Span}\{\mathbf{u}, \mathbf{v}\}$.

- The example 4 generalizes to any set $\{\mathbf{u}, \mathbf{v}, \mathbf{w}\}$ in $\mathbb{R}^3$ with $\mathbf{u}$ and $\mathbf{v}$ linearly independent.
- The set $\{\mathbf{u}, \mathbf{v}, \mathbf{w}\}$ will be linearly dependent if and only if $\mathbf{w}$ is in the plane spanned by $\mathbf{u}, \mathbf{v}$.

A Set Containing the Zero Vector

THEOREM 9

If a set $S = \{\mathbf{v}_1, \dots, \mathbf{v}_p\}$ in $\mathbb{R}^n$ contains the zero vector, then the set is linearly dependent.

A Set Containing the Zero Vector

THEOREM 9

If a set $S = \{\mathbf{v}_1, \dots, \mathbf{v}_p\}$ in $\mathbb{R}^n$ contains the zero vector, then the set is linearly dependent.

Proof. Renumber the vectors so that $\mathbf{v}_1 = \mathbf{0}$. Then

$$1\mathbf{v}_1 + 0\mathbf{v}_2 + \dots + 0\mathbf{v}_p = \mathbf{0}$$

is a linear dependence relation because the weights are not all zero.


► LINEAR TRANSFORMATIONS

(Section 1.8)

Transformations

A **transformation** (function or mapping) T from $\mathbb{R}^n \rightarrow \mathbb{R}^m$ is a rule that assigns to each vector $\mathbf{x} \in \mathbb{R}^n$ and a vector $T(\mathbf{x}) \in \mathbb{R}^m$.

- ▶ The set $\mathbb{R}^n$ is the **domain** of T , and the set $\mathbb{R}^m$ is the **codomain** of T .
- ▶ The notation $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ indicates that the domain of T is $\mathbb{R}^n$ and the codomain is $\mathbb{R}^m$.
- ▶ For $\mathbf{x} \in \mathbb{R}^n$, the vector $T(\mathbf{x}) \in \mathbb{R}^m$ is called the **image of $\mathbf{x}$ (under the action of T)**.



Matrix Transformations

- For each $\mathbf{x} \in \mathbb{R}^n$, $T(\mathbf{x})$ is computed as $A\mathbf{x}$, where A is an $m \times n$ matrix.

Matrix Transformations

- ▶ For each $\mathbf{x} \in \mathbb{R}^n$, $T(\mathbf{x})$ is computed as $A\mathbf{x}$, where A is an $m \times n$ matrix.
- ▶ For simplicity, we *denote such a matrix transformation by* $x \mapsto Ax$

Matrix Transformations

- ▶ For each $\mathbf{x} \in \mathbb{R}^n$, $T(\mathbf{x})$ is computed as $A\mathbf{x}$, where A is an $m \times n$ matrix.
- ▶ For simplicity, we *denote such a matrix transformation by* $x \mapsto Ax$
- ▶ Observe that the domain of T is $\mathbb{R}^n$ when A has n columns and the codomain of T is $\mathbb{R}^m$ when each column of A has m entries.

Matrix Transformations

- ▶ For each $\mathbf{x} \in \mathbb{R}^n$, $T(\mathbf{x})$ is computed as $A\mathbf{x}$, where A is an $m \times n$ matrix.
- ▶ For simplicity, we denote such a *matrix transformation* by $x \mapsto Ax$
- ▶ Observe that the domain of T is $\mathbb{R}^n$ when A has n columns and the codomain of T is $\mathbb{R}^m$ when each column of A has m entries.
- ▶ The range of T is the set of all linear combinations of the columns of A , because each image $T(\mathbf{x})$ is of the form $A\mathbf{x}$.

Transforming Vectors by Matrix Multiplication

For instance, the equations

Transforming Vectors by Matrix Multiplication

For instance, the equations

$$\underbrace{\begin{bmatrix} 4 & -3 & 1 & 3 \\ 2 & 0 & 5 & 1 \end{bmatrix}}_A \underbrace{\begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}}_{\mathbf{x}} = \underbrace{\begin{bmatrix} 5 \\ 8 \end{bmatrix}}_{\mathbf{b}}, \quad A \underbrace{\begin{bmatrix} 4 \\ -1 \\ 3 \\ -3 \end{bmatrix}}_{\mathbf{u}} = \underbrace{\begin{bmatrix} 0 \\ 0 \end{bmatrix}}_{\mathbf{0}}.$$

Transforming Vectors by Matrix Multiplication

For instance, the equations

$$\underbrace{\begin{bmatrix} 4 & -3 & 1 & 3 \\ 2 & 0 & 5 & 1 \end{bmatrix}}_A \underbrace{\begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}}_{\mathbf{x}} = \underbrace{\begin{bmatrix} 5 \\ 8 \end{bmatrix}}_{\mathbf{b}}, \quad A \underbrace{\begin{bmatrix} 4 \\ -1 \\ 3 \\ -3 \end{bmatrix}}_{\mathbf{u}} = \underbrace{\begin{bmatrix} 0 \\ 0 \end{bmatrix}}_{\mathbf{0}}.$$


say that multiplication by A transforms $\mathbf{x}$ into $\mathbf{b}$ and transforms $\mathbf{u}$ into the zero vector. See Figure below.

Transforming Vectors by Matrix Multiplication

For instance, the equations

$$\underbrace{\begin{bmatrix} 4 & -3 & 1 & 3 \\ 2 & 0 & 5 & 1 \end{bmatrix}}_A \underbrace{\begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}}_{\mathbf{x}} = \underbrace{\begin{bmatrix} 5 \\ 8 \end{bmatrix}}_{\mathbf{b}}, \quad A \underbrace{\begin{bmatrix} 4 \\ -1 \\ 3 \\ -3 \end{bmatrix}}_{\mathbf{u}} = \underbrace{\begin{bmatrix} 0 \\ 0 \end{bmatrix}}_{\mathbf{0}}.$$

say that multiplication by A transforms $\mathbf{x}$ into $\mathbf{b}$ and transforms $\mathbf{u}$ into the zero vector. See Figure below.



Matrix Transformations: Example 1

Let

$$A = \begin{bmatrix} 1 & -3 \\ 3 & 5 \\ -1 & 7 \end{bmatrix}, \quad \mathbf{u} = \begin{bmatrix} 2 \\ -1 \end{bmatrix}, \quad \mathbf{b} = \begin{bmatrix} 3 \\ 2 \\ -5 \end{bmatrix}, \quad \mathbf{c} = \begin{bmatrix} 3 \\ 2 \\ 5 \end{bmatrix}.$$

and define a transformation $T : \mathbb{R}^2 \rightarrow \mathbb{R}^3$ by $T(\mathbf{x}) = A\mathbf{x}$. Thus

$$T(\mathbf{x}) = \begin{bmatrix} 1 & -3 \\ 3 & 5 \\ -1 & 7 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} x_1 - 3x_2 \\ 3x_1 + 5x_2 \\ -x_1 + 7x_2 \end{bmatrix}.$$

Matrix Transformations: Questions

- a. Find $T(\mathbf{u})$, the image of $\mathbf{u}$ under the transformation T .
- b. Find an $\mathbf{x}$ in $\mathbb{R}^2$ whose image under T is $\mathbf{b}$.
- c. Is there more than one $\mathbf{x}$ whose image under T is $\mathbf{b}$?
- d. Determine whether $\mathbf{c}$ is in the range of the transformation T .

Matrix Transformations: Solution

Solution:

Matrix Transformations: Solution

Solution:

- a. Compute

$$T(\mathbf{u}) = A\mathbf{u} = \begin{bmatrix} 1 & -3 \\ 3 & 5 \\ -1 & 7 \end{bmatrix} \begin{bmatrix} 2 \\ -1 \end{bmatrix} = \begin{bmatrix} 5 \\ 1 \\ -9 \end{bmatrix}.$$

Matrix Transformations: Solution

Solution:

- a. Compute

$$T(\mathbf{u}) = A\mathbf{u} = \begin{bmatrix} 1 & -3 \\ 3 & 5 \\ -1 & 7 \end{bmatrix} \begin{bmatrix} 2 \\ -1 \end{bmatrix} = \begin{bmatrix} 5 \\ 1 \\ -9 \end{bmatrix}.$$

- b. Solve $T(\mathbf{x}) = \mathbf{b}$ for $\mathbf{x}$. That is, solve $Ax = b$, or

$$\begin{bmatrix} 1 & -3 \\ 3 & 5 \\ -1 & 7 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 3 \\ 2 \\ -5 \end{bmatrix}. \quad (1)$$

Matrix Transformations: Row Reduction

Row reduce the augmented matrix:

$$\left[\begin{array}{cc|c} 1 & -3 & 3 \\ 3 & 5 & 2 \\ -1 & 7 & -5 \end{array} \right] \sim \left[\begin{array}{cc|c} 1 & -3 & 3 \\ 0 & 14 & -7 \\ 0 & 4 & -2 \end{array} \right] \sim \left[\begin{array}{cc|c} 1 & -3 & 3 \\ 0 & 1 & -.5 \\ 0 & 0 & 0 \end{array} \right] \sim \left[\begin{array}{cc|c} 1 & 0 & 1.5 \\ 0 & 1 & -.5 \\ 0 & 0 & 0 \end{array} \right] \quad (2)$$

Matrix Transformations: Row Reduction

Row reduce the augmented matrix:

$$\left[\begin{array}{cc|c} 1 & -3 & 3 \\ 3 & 5 & 2 \\ -1 & 7 & -5 \end{array} \right] \sim \left[\begin{array}{cc|c} 1 & -3 & 3 \\ 0 & 14 & -7 \\ 0 & 4 & -2 \end{array} \right] \sim \left[\begin{array}{cc|c} 1 & -3 & 3 \\ 0 & 1 & -.5 \\ 0 & 0 & 0 \end{array} \right] \sim \left[\begin{array}{cc|c} 1 & 0 & 1.5 \\ 0 & 1 & -.5 \\ 0 & 0 & 0 \end{array} \right] \quad (2)$$

Hence $x_1 = 1.5$, $x_2 = -.5$, and

$$\mathbf{x} = \begin{bmatrix} 1.5 \\ -.5 \end{bmatrix}.$$

The image of this $\mathbf{x}$ under T is the given vector $\mathbf{b}$.

Matrix Transformations: Uniqueness and Range

- Any $\mathbf{x}$ whose image under T is $\mathbf{b}$ must satisfy equation (1).

Matrix Transformations: Uniqueness and Range

- Any $\mathbf{x}$ whose image under T is $\mathbf{b}$ must satisfy equation (1).
 - From (2), it is clear that equation (1) has a unique solution.

Matrix Transformations: Uniqueness and Range

- Any $\mathbf{x}$ whose image under T is $\mathbf{b}$ must satisfy equation (1).
 - ▶ From (2), it is clear that equation (1) has a unique solution.
 - ▶ So there is exactly one $\mathbf{x}$ whose image is $\mathbf{b}$.

Matrix Transformations: Uniqueness and Range

- c. Any $\mathbf{x}$ whose image under T is $\mathbf{b}$ must satisfy equation (1).
 - ▶ From (2), it is clear that equation (1) has a unique solution.
 - ▶ So there is exactly one $\mathbf{x}$ whose image is $\mathbf{b}$.
- d. The vector $\mathbf{c}$ is in the range of T if $\mathbf{c}$ is the image of some $\mathbf{x}$ in $\mathbb{R}^2$, that is, if $\mathbf{c} = T(\mathbf{x})$ for some $\mathbf{x} \in \mathbb{R}^2$.

Matrix Transformations: Uniqueness and Range

- c. Any $\mathbf{x}$ whose image under T is $\mathbf{b}$ must satisfy equation (1).
 - ▶ From (2), it is clear that equation (1) has a unique solution.
 - ▶ So there is exactly one $\mathbf{x}$ whose image is $\mathbf{b}$.
- d. The vector $\mathbf{c}$ is in the range of T if $\mathbf{c}$ is the image of some $\mathbf{x}$ in $\mathbb{R}^2$, that is, if $\mathbf{c} = T(\mathbf{x})$ for some $\mathbf{x} \in \mathbb{R}^2$.

This is another way of asking if the system

$$A\mathbf{x} = \mathbf{c}$$

is consistent.

Matrix Transformations: Is $\mathbf{c}$ in the Range?

To find the answer, row reduce the augmented matrix:

$$\left[\begin{array}{cc|c} 1 & -3 & 3 \\ 3 & 5 & 2 \\ -1 & 7 & 5 \end{array} \right] \sim \left[\begin{array}{cc|c} 1 & -3 & 3 \\ 0 & 14 & -7 \\ 0 & 4 & 8 \end{array} \right] \sim \left[\begin{array}{cc|c} 1 & -3 & 3 \\ 0 & 1 & 2 \\ 0 & 14 & -7 \end{array} \right] \sim \left[\begin{array}{cc|c} 1 & -3 & 3 \\ 0 & 1 & 2 \\ 0 & 0 & -35 \end{array} \right].$$

The third equation, $0 = -35$, shows that the system is inconsistent.

So $\mathbf{c}$ is not in the range of T .

Four Basic Questions About a Transformation


For $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$:

- ❶ Question 1: for any $\mathbf{u} \in \mathbb{R}^n$, what is the image of $T(\mathbf{u})$? (function value)
- ❷ Question 2: find an $\mathbf{x} \in \mathbb{R}^n$ whose image under T is $\mathbf{b}$. (independent value)
- ❸ Question 3: is there more than one $\mathbf{x}$ whose image under T is $\mathbf{b}$?
- ❹ Determine if $\mathbf{c}$ is in the range of the transformation T .

Shear Transformation

Example 3 Let $A = \begin{bmatrix} 1 & 3 \\ 0 & 1 \end{bmatrix}$. The transformation $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ defined by $T(\mathbf{x}) = A\mathbf{x}$ is called a **shear transformation**.

- It can be shown that if T acts on each point in the 2×2 square shown below, then the set of images forms the shaded parallelogram.



Shear Transformation

The key idea is to show that T maps line segments onto line segments, and then to check that the corners of the square map onto the vertices of the parallelogram.

For instance, the image of the point $u = \begin{bmatrix} 0 \\ 2 \end{bmatrix}$

$$T(u) = \begin{bmatrix} 1 & 3 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 0 \\ 2 \end{bmatrix} = \begin{bmatrix} 6 \\ 2 \end{bmatrix},$$

$$\begin{bmatrix} 2 \\ 2 \end{bmatrix} = \begin{bmatrix} 1 & 3 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 2 \\ 2 \end{bmatrix} = \begin{bmatrix} 8 \\ 2 \end{bmatrix}.$$

T deforms the square as if its top were pushed right while its base stayed fixed.

Linear Transformations

Definition

A transformation (or mapping) T is **linear** if

- ① $T(\mathbf{u} + \mathbf{v}) = T(\mathbf{u}) + T(\mathbf{v})$ for all $\mathbf{u}, \mathbf{v}$ in its domain;
- ② $T(c\mathbf{u}) = cT(\mathbf{u})$ for all scalars c and all $\mathbf{u}$ in its domain.

Every matrix transformation is linear. Important examples of linear transformations that are not matrix transformations will be discussed in Chapters 4 and 5.

Linear Transformations Preserve Operations

Linear transformations *preserve the operations of vector addition and scalar multiplication*.

- ▶ $T(\mathbf{u} + \mathbf{v})$ can be found either by adding first $\mathbf{u}$ and $\mathbf{v}$ and then applying T to $\mathbf{u} + \mathbf{v}$, or by applying T first towards $T(\mathbf{u})$ and $T(\mathbf{v})$, and adding the images to $T(\mathbf{u} + \mathbf{v})$.
- ▶ These two properties lead to the following useful facts

$$T(\mathbf{0}) = \mathbf{0}, \quad (3)$$

and, for all scalars c, d ,

$$T(c\mathbf{u} + d\mathbf{v}) = cT(\mathbf{u}) + dT(\mathbf{v}). \quad (4)$$

Why Properties (3) and (4) Hold

Property (3) follows from condition (2) in the definition because:

$$T(\mathbf{0}) = T(0\mathbf{u}) = 0T(\mathbf{u}) = \mathbf{0}.$$

Property (4) requires both defining properties:

$$T(c\mathbf{u} + d\mathbf{v}) = T(c\mathbf{u}) + T(d\mathbf{v}) = cT(\mathbf{u}) + dT(\mathbf{v}).$$

A useful converse

If a transformation satisfies (4) for all $\mathbf{u}, \mathbf{v}$ and c, d , it must be linear.

(Set $c = d = 1$ for preservation of addition; set $d = 0$ for preservation of scalar multiplications.)

The Superposition Principle

Repeated application of (4) gives

$$T(c_1\mathbf{v}_1 + \cdots + c_p\mathbf{v}_p) = c_1T(\mathbf{v}_1) + \cdots + c_pT(\mathbf{v}_p). \quad (5)$$

In engineering and physics, (5) is called the *superposition principle*.

Contractions and Dilations

Given a scalar r , define $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ by

$$T(\mathbf{x}) = r\mathbf{x}.$$

Contraction

$0 \leq r \leq 1$: vectors keep their direction and shrink toward the origin.


Dilation

$r > 1$: vectors keep their direction and stretch away from the origin.

From a Matrix to a Geometric Description

Let $A = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$. Given a geometric description of the transformation $\mathbf{x} \mapsto A\mathbf{x}$.

Plot some random points (vectors) on graph paper to see what happens. A point such as $(4,1)$ maps into $(4,-1)$. The transformation $\mathbf{x} \mapsto A\mathbf{x}$ reflects points through the x -axis, or x_1 -axis.



A question

Given a geometric description of a linear transformation,

how do we describe the transformation?

(How to give an explicit function relationship between $\mathbf{x}$ and $\mathbf{y}$)

A “formula” for $T(\mathbf{x})$.

Transformation and Matrix

Every linear transformation from $\mathbb{R}^n$ to $\mathbb{R}^m$



a matrix transformation $\mathbf{x} \mapsto A\mathbf{x}$



the matrix A

The Matrix of a Linear Transformation (Section 1.9)

Theorem 10. Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation. Then there exists a unique matrix A such that

$$T(\mathbf{x}) = A\mathbf{x} \quad \text{for all } \mathbf{x} \in \mathbb{R}^n. \quad (3)$$

In fact, A is the $m \times n$ matrix whose j th column is the vector $T(\mathbf{e}_j)$, where $\mathbf{e}_j$ is the j th column of the identity matrix in $\mathbb{R}^n$:

The Matrix of a Linear Transformation: Proof

Proof: Write

$$\mathbf{x} = I_n \mathbf{x} = [\mathbf{e}_1 \ \cdots \ \mathbf{e}_n] \mathbf{x} = x_1 \mathbf{e}_1 + \cdots + x_n \mathbf{e}_n.$$

and use the linearity of T to compute

$$T(x) = T(x_1 e_1 + \cdots + x_n e_n) = x_1 T(e_1) + \cdots + x_n T(e_n)$$

The Standard Matrix

The matrix A in (3) is called the **standard matrix for the linear transformation T** .

We know now that every linear transformation from $\mathbb{R}^n$ to $\mathbb{R}^m$ can be viewed as a matrix transformation, and vice versa.

- ▶ The term *Linear transformation* focuses on a property of a mapping.
- ▶ The term *Matrix transformation* describes how such a mapping is implemented, as the example on the next slide illustrates.

Example 2: Standard Matrix of a Dilation

Find the standard matrix A for the dilation transformation $T(\mathbf{x}) = 3\mathbf{x}$ in $\mathbb{R}^2$.

Solution: Write

$$T(\mathbf{e}_1) = 3\mathbf{e}_1 = \begin{bmatrix} 3 \\ 0 \end{bmatrix}, \quad T(\mathbf{e}_2) = 3\mathbf{e}_2 = \begin{bmatrix} 0 \\ 3 \end{bmatrix}.$$


Therefore

$$A = [T(\mathbf{e}_1) \ T(\mathbf{e}_2)] = \begin{bmatrix} 3 & 0 \\ 0 & 3 \end{bmatrix}.$$

Geometric Linear Transformations of $\mathbb{R}^2$


Tables 1–4 illustrate common geometric linear transformations of the plane.

TABLE 1 Reflections

Transformation	Image of the Unit Square	Standard Matrix
Reflection through the x_1 -axis		$\begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$


Reflections: Table 1 Continued

Reflection through
the x_2 -axis



$$\begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix}$$


Reflection through
the line $x_2 = x_1$



$$\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$$


Reflections: Table 1 Continued

Reflection through
the line $x_2 = -x_1$



$$\begin{bmatrix} 0 & -1 \\ -1 & 0 \end{bmatrix}$$

Reflection through
the origin



$$\begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix}$$

Table 2: Contractions and Expansions



Table 2: Contractions and Expansions



Table 2: Contractions and Expansions

TABLE 2 Contractions and Expansions





Transformation	Image of the Unit Square		Standard Matrix
Horizontal contraction and expansion	 $0 < k < 1$	 $k > 1$	$\begin{bmatrix} k & 0 \\ 0 & 1 \end{bmatrix}$
Vertical contraction and expansion	 $0 < k < 1$	 $k > 1$	$\begin{bmatrix} 1 & 0 \\ 0 & k \end{bmatrix}$

Table 3: Shears



Table 3: Shears

TABLE 3 Shears





Transformation	Image of the Unit Square	Standard Matrix
Horizontal shear	 $k < 0$  $k > 0$	$\begin{bmatrix} 1 & k \\ 0 & 1 \end{bmatrix}$
Vertical shear	 $k < 0$  $k > 0$	$\begin{bmatrix} 1 & 0 \\ k & 1 \end{bmatrix}$

Table 4: Projections






Table 4: Projections

TABLE 4 Projections

Transformation	Image of the Unit Square	Standard Matrix
Projection onto the x_1 -axis		$\begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$
Projection onto the x_2 -axis		$\begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}$



Existence and Uniqueness Questions

Definition

A mapping $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is said to be **onto** $\mathbb{R}^m$ if each $\mathbf{b} \in \mathbb{R}^m$ is the image of at least one $\mathbf{x} \in \mathbb{R}^n$.

- ▶ Equivalently, T is onto $\mathbb{R}^m$ when the range of T is all of the codomain $\mathbb{R}^m$.
- ▶ That is, T maps $\mathbb{R}^n$ onto $\mathbb{R}^m$ if, for each $\mathbf{b}$ in the codomain $\mathbb{R}^m$, there exists at least one solution of $T(\mathbf{x})=\mathbf{b}$.
- ▶ "Does T map $\mathbb{R}^n$ onto $\mathbb{R}^m$?" is an existence question.
- ▶ The mapping T is not onto $\mathbb{R}^m$ when there is some $\mathbf{b}$ in $\mathbb{R}^m$ for which the equation $T(\mathbf{x})=\mathbf{b}$ has no solution. See the figure on the next slide.

Onto and One-to-One



Definition of One-to-one

A mapping $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is said to be **one-to-one** if every $\mathbf{b} \in \mathbb{R}^m$ is the image of *at most one* $\mathbf{x} \in \mathbb{R}^n$.

Example 4: Onto and One-to-One

Let T be the linear transformation whose standard matrix is

$$A = \begin{bmatrix} 1 & -4 & 8 & 1 \\ 0 & 2 & -1 & 3 \\ 0 & 0 & 0 & 5 \end{bmatrix}.$$

Does T map $\mathbb{R}^4$ onto $\mathbb{R}^3$? Is T a one-to-one mapping?

Example 4: Solution

Solution:

- ▶ Since A is in echelon form, we can see at once that A has a pivot position in each row.
- ▶ By Theorem 4 in Section 1.4, for each $\mathbf{b}$ in $\mathbb{R}^3$, the equation $A\mathbf{x}=\mathbf{b}$ is consistent.
- ▶ In other words, the linear transformation T maps $\mathbb{R}^4$ (its domain) onto $\mathbb{R}^3$.

However, since $A\mathbf{x} = \mathbf{b}$ has a **free variable** because there are four variables and only three basic variables, each $\mathbf{b}$ is the image of more than one $\mathbf{x}$. Thus T is not one-to-one.

Example 4: recap

- ▶ Check the echelon form of linear transformation:
- ▶ If consistent: maps domain to codomain.
- ▶ If and only if no free variable: one-to-one mapping

Kernel and Range of a Linear Transformation

Subspaces of vector spaces other than $\mathbb{R}^n$ are often described in terms of a linear transformations instead of matrices.

Definition

A **linear transformation** T from a vector space V into a vector space W is a rule that assigns to each $\mathbf{x} \in V$ a unique vector $T(\mathbf{x}) \in W$, such that

- ❶ $T(\mathbf{u} + \mathbf{v}) = T(\mathbf{u}) + T(\mathbf{v})$ for all $\mathbf{u}, \mathbf{v} \in V$;
- ❷ $T(c\mathbf{u}) = cT(\mathbf{u})$ for every $\mathbf{u} \in V$ and scalar c .

The **kernel** is $\{\mathbf{x} \in V : T(\mathbf{x}) = \mathbf{0}\}$; the **range** is $\{T(\mathbf{x}) : \mathbf{x} \in V\}$.