Midterm 1, Math 33b, Winter 2013 Instructor: Tonći Antunović

| Printed name and student ID: | |
|-------------------------------|----|
| | |
| Signed name: | |
| Section number time and TA na | mo |

Instructions:

- Read problems very carefully. Please raise your hand if you have questions at any time.
- The correct final answer alone is not sufficient for full credit try to explain your answers as much as you can. This way you are minimizing the chance of losing points for not explaining details, and maximizing the chance of getting partial credit if you fail to solve the problem completely. However, try to save enough time to seriously attempt to solve all the problems.
- If it's obvious that you final answers can be simplified, please simplify them. Otherwise, your final answers need to be simplified only if this is required in the statement of the problem.
- You are allowed to use only the items necessary for writing. Notes, books, sheets, calculators, phones are not allowed, please put them away. If you need more paper please raise your hand (you can write on the back pages).

| Question | Points | Score |
|----------|--------|-------|
| 1 | 10 | |
| 2 | 10 | |
| 3 | 10 | |
| 4 | 10 | |
| 5 | 10 | |
| Total: | 50 | |

1. (a) (2 points) Find all values of constant r such that the function $y(t) = (t+r)e^t$ is a solution of the equation

$$2ty' - e^{-t}y^2 = (t^2 + 2t - 4)e^t.$$

(b) (2 points) General solution of the equation $x^2y' = y^2 + 2xy$ is given by $y = \frac{x^2}{C-x}$. Find the solution of the initial value problem

$$x^2y' = y^2 + 2xy, \quad y(1) = 2.$$

(c) (2 points) Is the following differential equation exact

$$(e^x y + y^2) dx + (e^x + x^2) dy = 0?$$

- (d) (2 points) The graph of the function y passes through the point (1,2). Moreover, for x>0 the tangent line at the point (x,y(x)) hits the y axis at the point (0,y(x)/2). Write down an initial value problem whose solution is y.
- (e) (2 points) Does the following differential equation have any solutions $e^{y'} + x^2 = 0$?

 $2.\ (10\ \mathrm{points})\ \mathrm{Find}$ the solution of the initial value problem

$$y' - e^{-y} = \frac{e^{-y}}{x+1}, \quad y(0) = 1.$$

3. (10 points) Find the general solution of the equation

$$t^2y' - y = e^{-1/t}.$$

| 4. | (10 points) A large tank contains only 10 gallons of water with 1 pound of salt. A drain is opened which allows the mixture to leave the tank at the rate 0.5 gallons per second. At the same time pure water |
|----|---|
| | starts pouring in the tank at the rate 1 gallon per second raising the water level. Find the concentration of salt in the tank after time t . |
| | |
| | |
| | |
| | |

5. (10 points) Solve the initial value problem

$$(e^x + 2xy) dx + (x^2 + 1) dy = 0, \quad y(0) = 3.$$