1 Explain carefully how the classical "divergence theorem"

$$\iint\limits_S \vec{V}\cdot\vec{n}d(\text{area}) = \iiint\limits_V \operatorname{div}\vec{V}d(\text{volume})$$

(V a bounded volume in $\mathbb{R}^3, S =$ boundary of V) follows from Stokes' Theorem for differential forms.

2 Without using deRham's Theorem, prove:

2.1 every closed 1-form on S^2 is exact.

2.2 a two-form Ω is exact on S^2 if and only if

$$\int_{S^2}\Omega = 0.$$

3 Show that the set of all lines in $\mathbb{R}^2$ has a natural structure as a differentiable manifold. What (already familiar) manifold is it?

4 Show that S^3 is the union of two solid tori ($S^1\times$ 2-disc) with an embedded torus ($S^1\times S^1$) as their common boundary. (Hint: Express $\mathbb{R}^3$ with a solid torus removed as a union of circles and a single straight line and then add a point at infinity.)

5 Suppose M is a compact manifold (with empty boundary).

5.1 Prove that, if $f : M \to \mathbb{R}$ is a C^∞ function, then f has at least two critical points.

5.2 A C^∞ function on $S^1\times S^1$ cannot have only two critical points. Prove this (e.g.) by deforming a homotopically nontrivial S' along the gradient flow of $f : M \to \mathbb{R}$.

6.1 Prove carefully that a group of homeomorphisms of S^{2n}, each of which has no fixed points (unless it is the identity map) contains at most two elements.

6.2 Give a counterexample for some S^{2n+1}, $n \geq 1$.

7 Find the homology groups with $\mathbb{Z}$ coefficients, of $\mathbb{RP}^n$, $n = 2, 3, 4, \ldots$ by some systematic rigorous method.

8 Find the homology and the fundamental group of $S^1\times S^1$ with two points removed.

9 Prove that if a compact (empty boundary) manifold X has S^{2n+1}, $n \geq 1$, as a covering space, then X is orientable.

10 Suppose M is a compact manifold (empty boundary). Prove that

$$H_n(M, \mathbb{Z}) \simeq \mathbb{Z}.$$

(You may assume M is triangulated.)