

V123

Template

Fritz Ali Agildere
fritz.agildere@udo.edu

Jan Lucca Viola
janlucca.viola@udo.edu

Experiment: October 14, 2024
Submission: October 21, 2024

TU Dortmund – Department of Physics

Contents

1	Objective	1
2	Theory	1
2.1	General	1
2.2	Specific	1
3	Procedure	1
4	Results	2
5	Discussion	3
	References	3
	Appendix	4

1 Objective

This experiment serves as a template for a report in the advanced lab course.

2 Theory

The Pythagorean theorem describes aspect ratios.

2.1 General

We can always write

$$a^2 + b^2 = c^2 \quad (1a)$$

$$a^2 = c^2 - b^2 \quad (1b)$$

for any right triangle.

2.2 Specific

With $a = b$ follows from the general case (1a) that

$$2a^2 = c^2 \quad (2)$$

must hold.

3 Procedure

A *Geodreieck* is used.




Figure 1: Schematic depiction of a *Geodreieck*.

4 Results

Table 1: General measurements.

a / mm	b / mm	c / mm
1	2	2.25
2	3	3.60
3	4	5.00
4	5	6.40

Table 2: Specific measurements.

a / mm	c / mm
1	1.4
2	2.8
3	4.2
4	5.7
5	7.1




Figure 2: Measurements and theory prediction.

For $c^2 = ma^2 + n$ and using `numpy.polyfit` [1] we find

$$m = 2.03 \pm 0.01 \qquad n = (-0.24 \pm 0.21) \text{ mm}^2$$

as parameters.

5 Discussion

The measurements are in good agreement with predictions derived from theory.

References

- [1] Charles R. Harris et al. “Array programming with NumPy”. In: *Nature* 585.7825 (Sept. 2020), pp. 357–362. DOI: 10.1038/s41586-020-2649-2. URL: <https://doi.org/10.1038/s41586-020-2649-2>.

Appendix

