V47

Molar Heat of Copper

Fritz Agildere Amelie Strathmann fritz.agildere@udo.edu amelie.strathmann@udo.edu

Experiment: May 13, 2024 Submission: May 13, 2024

TU Dortmund – Department of Physics

Contents

1	Objective	2
2	Theory 2.1 Heat Capacity	2
3	Setup	3
4	Procedure	3
5	Results	3
6	Discussion	4
Re	ferences	4
Αp	pendix	5

1 Objective

By investigating the molar heat capacity of copper, we test some of the most prominent models for their validity and determine select characteristic quantities of solids.

2 Theory [1]

In the description of condensed matter, tools from classical thermodynamics and statistical physics as well as quantum mechanical considerations need to be applied. Throughout the following paragraphs, we explore some of the more well known models in the subfield of thermal properties, specifically for the explanation of the heat capacity in solids.

2.1 Heat Capacity
Molar Heat
Isobaric
Isochoric
2.2 Models
2.2.1 Classical Physics
2.2.2 Quantum Mechanics
Phonons
Einstein
Debye
Electrons

3 Setup




Figure 1: [2]

4 Procedure

5 Results

T[°C]	-200	-190	-180	-170	-160	-150	-140	-130	-120
R[Ω]	18,44	22,71	27,03	31,28	35,48	39,65	43,80	47,93	52,04
T[°C]	-110	-100	-90	-80	-70	-60	-50	-40	-30
$R[\Omega]$	56,13	60,20	64,25	68,28	72,29	76,28	80,25	84,21	88,17
T[°C]	-20	-10	0	+10	+20	+30	+40		
$R[\Omega]$	92,13	96,07	100	103,90	107,79	111,67	115,54		

Figure 2: [2]

т [к]	70	80	90	100	110	120	130	140
α [10 ⁻⁶ grd ⁻¹]	7,00	8,50	9,75	10,70	11,50	12,10	12,65	13,15
т [к]	150	160	170	180	190	200	210	220
α [10 ⁻⁶ grd ⁻¹]	13,60	13,90	14,25	14,50	14,75	14,95	15,20	15,40
т [к]	230	240	250	260	270	280	290	300
α [10 ⁻⁶ grd ⁻¹]	15,60	15,75	15,90	16,10	16,25	16,35	16,50	16,65

Figure 3: [2]

θ _D /T	0	1	2	3	4	5	6	7	8	9
0	24,9430	24,9310	24,8930	24,8310	24,7450	24,6340	24,5000	24,3430	24,1630	23,9610
1	23,7390	23,4970	23,2360	22,9560	22,6600	22,3480	22,0210	21,6800	21,3270	20,9630
2	20,5880	20,2050	19,8140	19,4160	19,0120	18,6040	18,1920	17,7780	17,3630	16,9470
3	16,5310	16,1170	15,7040	15,2940	14,8870	14,4840	14,0860	13,6930	13,3050	12,9230
4	12,5480	12,1790	11,8170	11,4620	11,1150	10,7750	10,4440	10,1190	9,8030	9,4950
5	9,1950	8,9030	8,6190	8,3420	8,0740	7,8140	7,5610	7,3160	7,0780	6,8480
6	6,6250	6,4090	6,2000	5,9980	5,8030	5,6140	5,4310	5,2550	5,0840	4,9195
7	4,7606	4,6071	4,4590	4,3160	4,1781	4,0450	3,9166	3,7927	3,6732	3,5580
8	3,4468	3,3396	3,2362	3,1365	3,0403	2,9476	2,8581	2,7718	2,6886	2,6083
9	2,5309	2,4562	2,3841	2,3146	2,2475	2,1828	2,1203	2,0599	2,0017	1,9455
10	1,8912	1,8388	1,7882	1,7393	1,6920	1,6464	1,6022	1,5596	1,5184	1,4785
11	1,4400	1,4027	1,3667	1,3318	1,2980	1,2654	1,2337	1,2031	1,1735	1,1448
12	1,1170	1,0900	1,0639	1,0386	1,0141	0,9903	0,9672	0,9449	0,9232	0,9021
13	0,8817	0,8618	0,8426	0,8239	0,8058	0,7881	0,7710	0,7544	0,7382	0,7225
14	0,7072	0,6923	0,6779	0,6638	0,6502	0,6368	0,6239	0,6113	0,5990	0,5871
15	0,5755	0,5641	0,5531	0,5424	0,5319	0,5210	0,5117	0,5020	0,4926	0,4834

Figure 4: [2]

6 Discussion

References

- [1] Rudolf Gross and Achim Marx. Festkörperphysik. Berlin, Boston: De Gruyter Oldenbourg, 2022. ISBN: 9783110782394. DOI: doi:10.1515/9783110782394. URL: https://doi.org/10.1515/9783110782394.
- [2] Manual for Experiment 47, Molar Heat of Copper. TU Dortmund, Department of Physics. 2024.

Appendix