

MATH 121: COMMUTATIVE ALGEBRA

PRISHITA DHARAMPAL

1. RINGS

Important Example: Let X be a “geometric set”, and consider the ring

$$\text{Hom}(X, \mathbb{R}) = \{X \xrightarrow{f} \mathbb{R}\}$$

with pointwise addition and multiplication

$$(f + g)(x) = f(x) + g(x), \quad (fg)(x) = f(x)g(x)$$

$r \in \text{Hom}(X, \mathbb{R})$ is a unit if and only if there exists $g \in \text{Hom}(X, \mathbb{R})$ such that $fg = 1$ if and only if $\forall x \in X, (fg)(x) = f(x)g(x) = 1$

$$\text{Hom}(X, \mathbb{R})^\times = \text{Hom}(X, \mathbb{R}^\times)$$

Definition 1.1. Let $a \in R, a \neq 0$

- a is a unit if and only if $\exists b \in R$ such that $ab = 1$,
- a is a zero divisor if and only if $\exists b \in R \setminus \{0\}$ such that $ab = 0$,

Lemma 1.2. The set of units $R^\times = \{a \in R \mid a \text{ is a unit}\}$ is a group under multiplication.

Proof: Worksheet (03/31).

Lemma 1.3 (Useful Lemma). If $\phi : R \rightarrow S$ is a ring homomorphism, then it follows from the definition that

- $\phi(0_R) = 0_S$
- $\phi(-a) = -\phi(a)$
- $\phi(a^{-1}) = \phi(a)^{-1}$

Remark 1.4. $\mathbb{Z}$ is initial in the category of rings. There is a unique homomorphism from the initial object to every other object. For any ring R ,

$$\text{Hom}(\mathbb{Z}, R) = \{1 \rightarrow 1_R\}$$

We don't know anything about homomorphisms to the initial object.

2. IDEALS

Lemma 2.1. If $\phi : R \rightarrow S$ is any ring homomorphism and $J \subset S$ is an ideal $\phi^{-1}(J)$ is also an ideal.

Proof: Worksheet Q4.1.

Theorem 2.2. If $I \subset R$ is an ideal then,

- (1) There exists a ring R/I and a surjective ring homomorphism $\pi : R \rightarrow R/I$ such that $\ker(\pi) = I$. Moreover R/I and π are unique upto isomorphism.
(2) There is an inclusion preserving bijection

$$\{\text{Ideals in } R/I\} \leftrightarrow \{\text{Ideals in } R \text{ containing } I\}$$

$$\bar{J} \leftrightarrow \pi^{-1}(\bar{J})$$

Proof.

- (1) Construct R/I explicitly.
(2) I gives an equivalence relation $\sim$ on R such that

$$a \sim b \iff a - b \in I \iff a = b + I$$

Show that equivalence classes $R/\sim$ is a ring.

$$R \rightarrow R/I = R/\sim$$

$$a \mapsto \bar{a} = a + I$$

Let $\bar{J}$ be an ideal in R/I , and J be an ideal in R and define the maps

$$\phi : \bar{J} \rightarrow \pi^{-1}(\bar{J}), \quad \psi : J \rightarrow \pi(J)$$

Now we want to show that $\phi \circ \psi = \psi \circ \phi = id$. Consider $\phi \circ \psi(J)$ and $\psi \circ \phi(\bar{J})$. Note that

$$\psi \circ \phi(\bar{J}) = \pi(\pi^{-1}(\bar{J})) = \bar{J}$$

since π is surjective. Thus $\psi \circ \phi = id$.

Now, note that $J \subseteq \phi \circ \psi(J) = \pi^{-1}(\pi(J))$, and since $\bar{0} \in \pi(J)$, $I \subseteq \pi^{-1}(\pi(J))$. An element $x \in \pi^{-1}(\bar{J})$ looks like $x + I$, and $\pi^{-1}(\bar{J}) = J + I$. Moreover since $\bar{0} \in \bar{J}$, we have $\pi^{-1}(\bar{0}) \subseteq \pi^{-1}(\bar{J})$ which implies $I \subseteq J$. Thus, $\psi \circ \phi(J) = J$ and $\psi \circ \phi = id$.

□

Exercise: Describe the ideals in

- (1) $\mathbb{Z}/\langle 24 \rangle = \{\langle 0 \rangle, \langle 1 \rangle, \langle 2 \rangle, \langle 3 \rangle, \langle 4 \rangle, \langle 6 \rangle, \langle 8 \rangle, \langle 12 \rangle\}$
(2) $\mathbb{Z}/\langle n \rangle = \{\langle d \rangle : d|n\}$
(3) $\mathbb{Z}/\langle 3 \rangle = \{\langle 0 \rangle, \langle 1 \rangle\}$
(4) $\mathbb{Z}/\langle 5 \rangle = \{\langle 0 \rangle, \langle 1 \rangle\}$
(5) $\mathbb{C}[x, y]/\langle y - x^2 \rangle \cong \mathbb{C}[x] = \{\text{Ideals in } \mathbb{C}[x]\}$
(6) $\mathbb{C}[x]/\langle x^n \rangle = \{\langle x^i \rangle : 0 \leq i < n\}$

Lemma 2.3. If $\phi : R \rightarrow S$ is a ring homomorphism with $\ker(\phi) \supseteq I$ then there exists a unique morphism $\tilde{\phi} : R/I \rightarrow S$

$$\begin{array}{ccc} R & \xrightarrow{\pi} & R/I \\ \phi \downarrow & \swarrow \tilde{\phi} & \\ S & & \end{array}$$

Universal Property of Quotients: Upto isomorphism R/I is the ring such that

$$\text{Hom}(R/I, S) = \{\phi : R \rightarrow S : I \subset \ker(\phi)\}$$

We want to define $\tilde{\phi}(\bar{a}) = \tilde{\phi}(a + I) = \phi(a)$ and $\tilde{\phi}(\bar{0}) = 0$.

Lemmact 2.4. Maximal Ideal $\implies$ Prime Ideals.

Definition 2.5. The spectrum of a ring R is the set

$$\text{Spec}(R) = \{\text{Prime Ideals in } R\}$$

the maximal spectrum of R is the set

$$m\text{Spec}(R) = \{\text{Maximal Ideals in } R\}$$

$\text{Spec}(R)$ is where the 'geometry lives'. Note that $m\text{Spec}(R) \subset \text{Spec}(R)$.

Example: $R = \mathbb{Z}[\sqrt{-5}]$. 2 is irreducible but not prime. $R/\langle 2 \rangle = \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$.

Example: $R = \mathbb{C}[x, y]/\langle x^3 - y^2 \rangle$. x is irreducible, but $\langle x \rangle$ is not prime. $R/\langle x \rangle = \mathbb{C}[y]/\langle y \rangle$ and y is a zero divisor.

Theorem 2.6. Let R be a non-zero ring.

- (1) R has a maximal ideal.
- (2) If $I \subsetneq R$ is a proper ideal there exists a prime ideal $P \subseteq R$ such that $I \subset P \subsetneq R$.

Proof. Zorn's Lemma: Let $(P, \leq)$ be a po-set.

If $P \neq \emptyset$ and if $C = C_1 \leq C_2 \leq C_3 \leq \dots$ is a chain in $(P, \leq)$ then there exists $B \in P$ such that $C_i \leq B$ for all i , then there exists $M \in P$ such that if $S \leq M$ then $M = S$.

Let $P = \{\text{Proper ideals in } R\}$ and $\leq$ be inclusion. We claim that $(P, \leq)$ is a poset.

- (1)
 - $P \neq \emptyset$ since $\langle 0 \rangle \in P$.
 - Given a chain in P

$$I_1 \subseteq I_2 \subseteq \dots$$

Then, define $B = \bigcup I_j$. Note that B is an ideals since union of nested ideals is an ideal. And B is proper since $1 \notin I_j$ for all j , and hence $1 \notin B$. Then there exists some ideal $M \subset R$ such that M is maximal.

- (2) Follows from (1). Let $R = R/I$. Pre-image of prime ideals is prime.

Thus, $m\text{Spec}(R) \neq \emptyset$ and $\text{Spec}(R) \neq \emptyset$. □

3. ZARISKI TOPOLOGY

Definition 3.1. If $S \subseteq R$ is any subset of R ,

$$\mathbb{V}(S) = \{[P] \in \text{Spec}(R) | S \subseteq P\} \subseteq \text{Spec}(R)$$

Definition 3.2. An algebraic subset of $\mathbb{K}^n$ of some field K is a set of the form

$$\mathbb{V}(S) = \{P \in \mathbb{K}^n | f(p) = 0, \forall f \in S\}$$

for a subset $S \subset \mathbb{K}[x_1, \dots, x_n]$

Examples:

- $\mathbb{V}(x^2 + y^2 - 1) \subseteq \mathbb{R}^2$

$$\{(x, y) \in \mathbb{R}^2 \mid x^2 + y^2 - 1 = 0\}$$

- $\mathbb{V}(x^2 + y^2 - 1) \subseteq \mathbb{R}^3$

$$\{(x, y, z) \in \mathbb{R}^3 \mid x^2 + y^2 - 1 = 0\}$$

- $\mathbb{V}(x^2 - y) \subseteq \mathbb{R}[x, y]$

$$\{(x, y) \in \mathbb{R}[x, y] \mid x^2 = y\}$$

- $\mathbb{V}(x^2 - xy) \subseteq \mathbb{R}[x, y]$

$$\{(x, y) \in \mathbb{R}[x, y] \mid x = 0, x - y = 0\}$$

- $\mathbb{V}(x^2 + 1) \subseteq \mathbb{R}[x]$

$$\mathbb{V}(x^2 + 1) = \emptyset$$

- $\mathbb{V}(x^2 + 1) \subseteq \mathbb{C}[x]$

$$\mathbb{V}(x^2 + 1) = \{i, -i\}$$

- $\mathbb{V}(x^2 + 4) \subseteq \mathbb{F}_3[x]$

$$\mathbb{V}(x^2 + 4) = \emptyset$$

- $\mathbb{V}(x^2 + 4) \subseteq \mathbb{F}_p[x]$

TODO:

- $\mathbb{V}(x^6 - 3x^5y + 3x^4y^2 - x^3y^3) \subseteq \mathbb{R}[x, y]$

$$\mathbb{V}(x^6 - 3x^5y + 3x^4y^2 - x^3y^3) = \mathbb{V}(x^3(x - y)^3) = \{(x, y) \in \mathbb{R}[x, y] \mid x = 0, x - y = 0\}$$

- $\mathbb{V}(n(x^2 + y^2 + z^2), n \in \mathbb{Z}) \subseteq \mathbb{R}^3$

TODO:

- $\mathbb{V}(\text{a collection of any linear polynomials in } \mathbb{K}[x_1, \dots, x_n])$

TODO:

Definition 3.3. Let $V \subseteq \mathbb{K}^n$ be an algebraic set. The defining ideal of V is

$$\mathbb{I}(V) = \{f \in \mathbb{K}[x_1, \dots, x_n] \mid f(p) = 0, \forall p \in V\}$$

Definition 3.4. If $\mathfrak{M} \subseteq R$ is a maximial ideal, the residue field of $\mathfrak{M}$ is $R / \mathfrak{M}$.

Definition 3.5. Let $\mathbb{K}$ be a field. A $\mathbb{K}$ algebra is the data of a ring R together with a fixed ring homomorphism

$$\alpha : \mathbb{K} \rightarrow R$$

Definition 3.6. If $(R, \mathbb{K} \xrightarrow{\alpha} R)$ and $(S, \mathbb{K} \xrightarrow{\beta} S)$ are $\mathbb{K}$ -algebras, a ring map $\phi : R \rightarrow S$ is a morphism of $\mathbb{K}$ -algebras iff

$$\begin{array}{ccc}
R & \xrightarrow{\phi} & S \\
\alpha \uparrow & \nearrow \beta & \\
\mathbb{K} & &
\end{array}$$

Lemmact 3.7. If R is k -algebra, R/I has a natural k -algebra structure compatible with the quotient map.

$$\begin{array}{ccc}
R & \xrightarrow{\pi} & R/I \\
\alpha \uparrow & \nearrow \exists! & \\
k & &
\end{array}$$

Definition 3.8. A valuation ring of a field $\mathbb{K}$ is a subring $R \subset \mathbb{K}$ with the property that for all $x \in \mathbb{K}^\times$ either $x \in R$ or $x^{-1} \in R$.

4. GRÖBNER BASIS

Remark 4.1. Ideal Membership Problem: Suppose $I \subseteq R[x_1, \dots, x_n]$ an ideal. How can we algorithmically determine if $f \in I$?

Definition 4.2. A ring R is Noetherian if and only if every ascending chain of ideals stabilizes (Ascending Chain Condition), i.e.

$$I_1 \subseteq I_2 \subseteq \dots$$

there exists $N \in \mathbb{Z}$ such that $I_k = I_{k+1}$ for all $k \geq N$.

Lemma 4.3. R is Noetherian if and only if every ideal is finitely generated.

Lemma 4.4 (Dickson's Lemma). Every monomial ideal in $\mathbb{K}[x_1, \dots, x_n]$ is finitely generated.

Theorem 4.5 (Hilbert's Basis Theorem). If R is a Noetherian ring, then $R[x]$ is also Noetherian.

5. NULLSTELLENSATZ

Theorem 5.1 (Hilbert's Nullstellensatz I). If $\mathbb{K}$ is algebraically closed, then every maximal ideal of $\mathbb{K}[x_1, \dots, x_n]$ is of the form $\langle x_1 - a_1, \dots, x_n - a_n \rangle$ for some point $(a_1, \dots, a_n) \in \mathbb{K}^n$.

Theorem 5.2 (Hilbert's Nullstellensatz II). If $\mathbb{K}$ is algebraically closed and $V = \mathbb{V}(\mathcal{S}) \subset \mathbb{K}^n$ is an algebraic set, for some subset $\mathcal{S} \subset \mathbb{K}[x_1, \dots, x_n]$, then $\mathbb{I}(V) = \sqrt{\langle \mathcal{S} \rangle}$.

Theorem 5.3 (Hilbert's Weak Nullstellensatz). If $\mathbb{K}$ is algebraically closed and $I \subset \mathbb{K}[x_1, \dots, x_n]$ is an ideal then $\mathbb{V}(I) = \emptyset$ if and only if $I = \langle 1 \rangle$.

Theorem 5.4 (Hilbert's Nullstellensatz III). Let $\mathbb{K}$ be an algebraically closed field. If $V \subset \mathbb{K}^n$ is an algebraic subset then the maps $\mathbb{I}_V$ and $\mathbb{V}_V$ induce inverse, inclusion-reversing bijections:

$$\begin{array}{ccc}
\left\{ \begin{array}{c} \text{algebraic subsets} \\ \text{in } V \end{array} \right\} & \begin{array}{c} \xrightarrow{\mathbb{I}_V} \\ \xleftarrow{\mathbb{V}_V} \end{array} & \left\{ \begin{array}{c} \text{radical ideals} \\ \text{in } \mathbb{K}[V] \end{array} \right\} \\
& 5 &
\end{array}$$

Remark 5.5. $\mathbb{I}$ and $\mathbb{V}$ properties:

- (1) If $\mathcal{S} \subset \mathcal{T}$ then $\mathbb{V}(\mathcal{T}) \subset \mathbb{V}(\mathcal{S})$
- (2)

$$\mathbb{V}(\mathcal{S}) = \mathbb{V}(\langle \mathcal{S} \rangle) = \mathbb{V}(\sqrt{\langle \mathcal{S} \rangle})$$

- (3) $\mathbb{I}(V)$ is radical.
- (4) $\langle \mathcal{S} \rangle \subset \sqrt{\langle \mathcal{S} \rangle} \subset \mathbb{I}(\mathbb{V}(\mathcal{S}))$
- (5) $\mathbb{V}(\mathcal{S}) = \mathbb{V}(\mathbb{I}(\mathbb{V}(\mathcal{S})))$
- (6) If $fg \in \mathbb{I}(V)$ then $V \subseteq \mathbb{V}(g) \cap \mathbb{V}(f)$.

6. LOCALIZATION

Definition 6.1. A subset $U \subset R$ is **multiplicatively closed** if and only if

- (1) $1_R \in U$
- (2) and if $s, u \in U$ then $su \in U$.

Example 6.1.

- (1) $R = \mathbb{Z}$, $U = \{1, 3, 9, \dots\}$ is multiplicatively closed.
- (2) $R = \mathbb{Z}$, $U = \{1, 3, 9\}$ is not multiplicatively closed.
- (3) $R = \mathbb{R}[x, y]$, $U = \{f \in R \mid f(1, 0) \neq 0\}$ is multiplicatively closed.
- (4) $R = \mathbb{R}[x, y]$, $U = \{f \in R \mid f(1, 1) \neq 0, \text{ or } f(0, 0) \neq 0\}$ is not multiplicatively closed.

Definition 6.2. Let R be a ring and $U \subseteq R$ be a multiplicative closed set. The **localization** of R at U is a ring $U^{-1}R$ together with a ring map $\eta : R \rightarrow U^{-1}R$ that is universal in the following sense:

- (1) Given a ring map $\psi : R \rightarrow T$ such that $\psi(U) \subseteq T^\times$ there exists a unique ring map $\tilde{\psi} : U^{-1}R \rightarrow T$ such that

$$\begin{array}{ccc} R & \xrightarrow{\eta} & U^{-1}R \\ & \searrow \psi & \downarrow \tilde{\psi} \\ & & T \end{array}$$

Construction of $U^{-1}R$:

Consider $R \times U$ as a set (r_1, u_1) where $r_1 \in R, u_1 \in U$ “ $=$ ” $\frac{r_1}{u_1}$. Define an equivalence relation

$$(r_1, u_1) \sim (r_2, u_2) \iff t(u_2r_1 - u_1r_2) = 0, \quad t \in U$$

Note if R is a domain $t = 1_R$.

Define $\frac{r}{u}$ to be the equivalence class of (r, u) in $R \times U / \sim$.

Set $U^{-1}R = R \times U / \sim = \left\{ \frac{r}{u} \mid r \in R, u \in U \right\} / \sim$

Make $U^{-1}R$ a ring with

$$\left(\frac{r_1}{u_1}\right)\left(\frac{r_2}{u_2}\right) = \frac{r_1 r_2}{u_1 u_2}, \quad \frac{r_1}{u_1} + \frac{r_2}{u_2} = \frac{r_1 u_2 + r_2 u_1}{u_1 u_2}$$

Additive Identity: $0/1$, **Multiplicative Identity:** $1/1$

$$\eta : R \rightarrow U^{-1}R$$

Then

$$\begin{aligned} \ker(\eta) &= \{r \in R \mid \eta(r) = 0\} \\ &= \{r \in R \mid \frac{r}{1} = \frac{0}{1}\} \\ &= \{r \in R \mid t(r \cdot 1 - 1 \cdot 0) = 0, t \in U\} \\ &= \{r \in R \mid tr = 0, t \in U\} \end{aligned}$$

If R is a domain, $\ker(\eta) = \langle 0 \rangle \implies \eta$ is injective, unless $0 \in U$.

Example 6.2. TODO: Let R be any ring and $U = \{0, 1\}$. Then

$$U^{-1}R = \left\{\frac{r}{0} \mid r \in R\right\} \cup \left\{\frac{r}{1} \mid r \in R\right\}$$

Let $\frac{r}{1} \stackrel{?}{=} \frac{0}{1} \iff t(r - 0) = 0 \iff tr = 0$ for some $t \in U$. We can take $t = 0$ and make this always true (same argument $\frac{r}{0} = \frac{0}{1}$). Hence $U^{-1}R$ is the zero ring.

Lemma 6.3. Let $U \subseteq R$ be multiplicatively closed. If U contains a nilpotent then $U^{-1}R$ is the zero ring.

Definition 6.4. If $P \subseteq R$ is a prime ideal then $U = R \setminus P$ is multiplicatively closed. Here we write $U^{-1}R = R_P$ and say R **localized at** P .

Lemma 6.5.

$$PR_P = \left\{\frac{r}{u} \mid r \in P, u \in R \setminus P\right\} \subseteq R_P$$

is a maximal ideal.

Proof. Let $PR_P \subsetneq I \subseteq R_P$. Then for $r \in I \setminus PR_P$ with $r \notin P \implies \frac{r}{u}$ with $r, u \in R \setminus P \implies \frac{u}{r} \in R_P \implies \frac{r}{u}$ is a unit in $R_P \implies I = R_P$. **TODO:** □

Proposition 6.6. R_P is a local ring (i.e., PR_P is the unique maximal ideal in R_P).

Proof. If $I \subseteq R_P$ is a maximal ideal, then $I \cap R \setminus P \neq \emptyset$, that is I contains a unit, and hence $I = R_P$. □

Example 6.3. Let $f \in R$, and $U = \{1, f, f^2, \dots\}$, then

$$U^{-1}R = R\left[\frac{1}{f}\right] = \left\{\frac{r}{u} \mid r \in R, u \in U\right\} = \left\{\frac{r}{f^k} \mid r \in R, k \in \mathbb{Z}_{\geq 0}\right\}$$

that is $U^{-1}R$ is the ring of polynomials in $1/f$ with coefficients in R .

Lemma 6.7.

$$R \left[\frac{1}{f} \right] \cong \frac{R[x]}{\langle xf - 1 \rangle}$$

Proof. $R[x] \rightarrow R \left[\frac{1}{f} \right], x \mapsto \frac{1}{f}$ and $1 \mapsto \frac{1}{1}$. This is surjective because for every $rx^k \in U^{-1}R$ we have $rx^k \mapsto \frac{r}{f^k}$. To show injectivity we must show that $\ker(\varphi) \supseteq \langle xf - 1 \rangle$ is trivial

TODO: . □

Theorem 6.8. *There is an inclusion preserving bijection*

$$\begin{aligned} \left\{ \text{prime ideals of } U^{-1}R \right\} &\xleftrightarrow{1:1} \left\{ \text{prime ideals } P \subseteq R \text{ such that } P \cap U = \emptyset \right\} \\ P &\mapsto \eta(P) \end{aligned}$$

Let $Q \subseteq R$ be a prime ideal such that $Q \cap U \neq \emptyset$. Consider $\eta(Q) \cdot U^{-1}R = \langle \eta(Q) \rangle$. Let $r \in Q \cap U, \eta(r) \in \eta(Q) \subseteq \eta(Q) \cdot U^{-1}R$. Since $r \in U, \eta(r)$ is a unit in $U^{-1}R$, which implies

$$\eta(Q) \cdot U^{-1}R = U^{-1}R$$

i.e. it's not prime. **TODO:**

Theorem 6.9.

$$\left\{ \text{Spec}(U^{-1}R) \right\} \xleftrightarrow{1:1} \left\{ P \in \text{Spec}(R) \mid P \cap U = \emptyset \right\} \subseteq \text{Spec}(R)$$

$$P \mapsto \eta^{-1}(P)$$

Definition 6.10. Let R be a ring, then a basic open subset of $\text{Spec}(R)$ is of the form $D(f) = \{P \in \text{Spec}(R) \mid f \notin P\}$ for $f \in R$.

And,

$$D(f) = \text{Spec}(R) \setminus V(I) = \text{Spec}(R) \setminus V(\langle f \rangle)$$

The intersection of basic open sets $D(f) \cap D(g)$ is

$$D(f) \cap D(g) = \{P \in \text{Spec}(R) \mid f \notin P, g \notin P\} = \{P \in \text{Spec}(R) \mid fg \notin P\} = D(fg)$$

Remark 6.11. We know that there is a bijection,

$$\text{Spec} \left(R \left[\frac{1}{f} \right] \right) \leftrightarrow \{P \in \text{Spec}(R) \mid \{1, f, f^2, \dots\} \cap P \neq \emptyset\} = \{P \in \text{Spec}(R) \mid f \notin P\} = D(f)$$

Hence there is a bijection,

$$\text{Spec} \left(R \left[\frac{1}{f} \right] \right) \leftrightarrow D(f)$$

Theorem 6.12. Let $f \in R$ and $\eta : R \rightarrow R \left[\frac{1}{f} \right]$ be a localization map. The induced map

$$\text{Spec} \left(R \left[\frac{1}{f} \right] \right) \xrightarrow{\eta^\#} \text{Spec}(R)$$

is a homeomorphism onto $D(f)$.

Definition 6.13. The nilradical of R is the set of nilpotents

$$\mathfrak{N}(R) = \{f \in R \mid f^k = 0, \text{ for some } k\} = \sqrt{\langle 0 \rangle}$$

Proposition 6.14. $\text{Spec}(R) \cong \text{Spec}(R/\mathfrak{N}(R))$ induced by the quotient map

$$R \xrightarrow{\pi} R/\mathfrak{N}(R)$$

such that

$$\text{Spec}(R/\mathfrak{N}(R)) \xrightarrow{\pi^\#} \text{Spec}(R)$$

Proposition 6.15. $\mathfrak{N}(R) = \bigcap_{P \in \text{Spec}(R)} P$

Proof. ($\implies$)

Let $f \in \mathfrak{N}(R)$ so $f^n = 0$. Pick any prime ideal $P \subseteq R$, $f^n = 0 \in P \implies f \in P$ since P is prime.

($\impliedby$)

We show given $f \notin \mathfrak{N}(R)$ there exists a prime $P \subseteq R$ with $f \notin P$ (i.e. $D(f) \neq \emptyset$). $\square$

7. NONZERODIVISORS

Remark 7.1. If M is a R -module we say an element $r \in R$ is a nonzerodivisor on $M \iff M \xrightarrow{\times r} M$ is injective $\iff r \notin \text{Ann}(m), \forall m \in M \iff (0 :_M r) = \{m \in M \mid rm = 0\} = 0$

Definition 7.2. Let M be a R -module. A sequence of elements $x_1, \dots, x_t \in R$ is a **regular sequence** on M if and only if

- (1) x_i is a nonzerodivisor on $M/\langle x_1, \dots, x_{i-1} \rangle M$
- (2) $M \neq \langle x_1 \dots x_t \rangle M$ (non-degeneracy condition)

Lemma 7.3. If $(R, \mathfrak{m})$ is a local ring, M is a finitely generated non-zero R -module, and $x_1, \dots, x_t \in \mathfrak{m}$ then the conditions (1) and (2) above are equivalent to just (1).

Proof. Nakayama $\square$

Theorem 7.4 (Krull's Intersection Theorem). Let R be a Noetherian ring, $I \subseteq R$ an ideal, and M a finitely-generated R -module. There exists an element $r \in I$ such that

$$(1-r) \bigcap_{k=1}^{\infty} I^k M = 0$$

Corollary 7.4.1. If $(R, \mathfrak{m})$ is a local Noetherian ring, $I \subsetneq R$ then

$$\bigcap_{k=1}^{\infty} I^k = 0$$

Theorem 7.5. Let $(R, \mathfrak{m})$ be a local ring, R is Noetherian, and M is a non-zero finitely generated R -module. If $x_1, \dots, x_t \in \mathfrak{m}$ then $x_1, \dots, x_t$ is a regular sequence on M if and only if every permutation is a regular sequence on M .

Proof. Let x, y be a regular sequence on M . We want to show that y, x is a regular sequence on M .

Assume $ym = 0$ for some $m \in M$. Then we have $y\bar{m} = 0 \in M/xM \implies \bar{m} = 0 \implies m = xm_1$ for some m_1 .

That is,

$$0 = ym = yxm_1 \implies ym_1 = 0 \in M \implies y\bar{m}_1 = 0 \in M/xM$$

Then by inducting on the m_i s, we get $m \in \bigcap_{k=1}^{\infty} \langle x \rangle^k M$. By KIT, this implies $m = 0$, a contradiction, and thus y is a NZD on M .

Now, assume $x\bar{m} = 0 \in M/yM$. Then, $xm = ym_1 \in M \implies y\bar{m}_1 = 0 \in M/xM \implies \bar{m}_1 = 0 \implies m_1 = xm_2 \in M$. So in M ,

$$xm = ym_1 = xym_2 \implies m = ym_2 \implies \bar{m} = 0 \in M/yM$$

Thus x is a nonzerodivisor on M/yM , and y, x is a regular sequence. $\square$

Definition 7.6. Given an ideal $I \subseteq R$ the height of I is

$$\begin{aligned} ht(I) &= \sup\{d \in \mathbb{Z}_{\geq 0} \mid \text{there exists a chain of prime ideals } P_0 \subsetneq P_1 \subsetneq \cdots \subsetneq P_d \subset I\} \\ &= \dim R - \dim \mathbb{V}(I) \\ &= \dim R - \dim R/I \end{aligned}$$

Theorem 7.7 (Krull's Principal Ideal Theorem). Let R be a Noetherian ring, $x \in R$. If $P \subseteq R$ is a prime ideal minimal over $\langle x \rangle$ then $ht(P) \leq 1$.

Corollary 7.7.1. Let R be a Noetherian ring $I = \langle r_1, \dots, r_n \rangle \subseteq R$. If $P \subseteq R$ is a prime ideal minimal over I then $ht(P) \leq n$.

Lemma 7.8. Let x be a nonzerodivisor on a Noetherian ring R .

$$\dim R/x = \dim R - 1$$

Corollary 7.8.1. If $x_1, \dots, x_n$ is a regular sequence on R then

$$\dim (R/\langle x_1, \dots, x_n \rangle) = \dim R - n$$

Definition 7.9. Let R be a ring, $I \subseteq R$ an ideal, M a module, The depth of M with respect to I is

$$\text{depth}_I(M) = \{\text{maximal length of a regular sequence on } M \text{ contained in } I\}$$

If $(R, \mathfrak{m})$ is local,

$$\text{depth}(M) = \text{depth}_{\mathfrak{m}}(M)$$

Example 7.1. Depth is not always finite. For instance, if $R = M = \mathbb{K}[x_1, x_2, \dots]$ where $x_1, \dots, x_n$ is a regular sequence for all n .

Proposition 7.10. Let M be a R -module, and $x \in R$, then TFAE,

(1) x is a regular sequence on M .

(2) $k(x, M) = 0 \rightarrow M \xrightarrow{\times x} M \rightarrow 0$

Where $H_1(k(x, M)) = 0$ and $H_0(k(x, M)) \neq 0$. In general,

$$H_1(k(x, M)) = \ker(M \xrightarrow{\times x} M) = (0 :_M x)$$

and,

$$H_0(k(x, M)) = M/xM$$

If $x, y \in R$.

$$k(x, y, M) = 0 \rightarrow M \xrightarrow{\begin{bmatrix} y \\ -x \end{bmatrix}} M \oplus M \xrightarrow{\begin{bmatrix} x & y \end{bmatrix}} M \rightarrow 0$$

In general we need,

$$H_0(k(x, y, m)) = M/(xf + yg \mid f, g \in M) = M/\langle x, y \rangle M \neq 0$$

$$H_2(k(x, y, m)) = \ker \left(\begin{bmatrix} y \\ -x \end{bmatrix} \right) = \{m \in M \mid m(-x) = 0, my = 0\}$$