

Math 121: Commutative Algebra

Prishita Dharampal

Throughout this course, “ring” means *commutative* ring with unity. In the previous worksheets we introduced modules, homomorphisms, quotient modules, free modules, Hom, tensor products, and localization. The goal of this worksheet is to organize these constructions around one central idea: *exactness*. Exactness is a way to say that one map accounts for exactly the elements killed by the next map. It packages kernels, images, quotient modules, presentations, and many theorems into one flexible language.

Let R be a ring. A *sequence* of R -modules is $\dots, M_0, M_1, M_2, \dots$ together with R -module homomorphisms $\phi_i : M_i \rightarrow M_{i-1}$. We often arrange a sequence into a diagram like

$$\dots \xrightarrow{\phi_{i+1}} M_i \xrightarrow{\phi_i} M_{i-1} \xrightarrow{\phi_{i-1}} M_{i-2} \longrightarrow \dots$$

We are most interested in sequences when the maps are related in some nice way that allows us to compare the image of one map with the kernel of the next. A *chain complex*, or simply a *complex*, of R -modules is a sequence in which $\phi_{i-1} \circ \phi_i = 0$ for every i . Equivalently, $\text{img}(\phi_i) \subset \ker(\phi_{i-1})$ for all i . We often write $(M_\bullet, \phi_\bullet)$ for a complex and call the maps ϕ_i *differentials*. A complex is *exact at M_i* if $\text{img}(\phi_{i+1}) = \ker(\phi_i)$. One way to measure the failure, or success, of a complex to be exact at M_i is by looking at the *i -th homology group* of $(M_\bullet, \phi_\bullet)$, which is defined to be the quotient R -module:

$$H_i(M_\bullet, \phi_\bullet) := \frac{\ker(\phi_i)}{\text{img}(\phi_{i+1})}.$$

The definition of a complex guarantees this quotient makes sense, since $\text{img}(\phi_{i+1}) \subset \ker(\phi_i)$. Equivalently, a complex is exact at M_i if and only if $H_i(M_\bullet, \phi_\bullet) = 0$. (Here 0 denotes the zero R -module.) A complex is *exact* if it is exact at every module where this condition makes sense. Thus every exact sequence is a complex, but a complex need not be exact.

Complexes appear throughout commutative algebra and other areas of mathematics. Perhaps the most important special case of a complex is a short exact sequence. A *short sequence* is a complex of the form

$$0 \longrightarrow A \xrightarrow{i} B \xrightarrow{p} C \longrightarrow 0.$$

It is called a *short exact sequence* if it is exact at A , B , and C . Strictly speaking, a complex extends infinitely in both directions, so the diagram above is shorthand for

$$\cdots \longrightarrow 0 \longrightarrow A \xrightarrow{i} B \xrightarrow{p} C \longrightarrow 0 \longrightarrow \cdots,$$

where all modules outside the displayed range are zero. We will follow the standard convention of suppressing these trailing and leading zero modules. Saying that this sequence is exact is the same as saying that i is injective, p is surjective, and $\text{img}(i) = \ker(p)$. Thus a short exact sequence says that A is being identified with a submodule of B , and that the quotient of B by this submodule is isomorphic to C .

Problem 1. First Encounters with Exactness.

Let A, B, C be R -modules.

1. Prove that a sequence

$$0 \longrightarrow A \xrightarrow{i} B$$

is exact at A if and only if i is injective.

2. Prove that a sequence

$$B \xrightarrow{p} C \longrightarrow 0$$

is exact at C if and only if p is surjective.

3. Prove carefully that

$$0 \longrightarrow A \xrightarrow{i} B \xrightarrow{p} C \longrightarrow 0$$

is exact if and only if i is injective, p is surjective, and $\text{img}(i) = \ker(p)$.

4. If

$$A \xrightarrow{f} B \xrightarrow{g} C$$

is exact at B , prove that $g \circ f = 0$.

5. Give an example of maps of R -modules

$$A \xrightarrow{f} B \xrightarrow{g} C$$

such that $g \circ f = 0$ but the sequence is not exact at B .

6. Consider the complex below. Compute the homology at B in each of the following cases:

$$A \xrightarrow{f} B \xrightarrow{g} C$$

- (a) $f = 0$ and $g = 0$.
- (b) f is injective and $g = 0$.
- (c) g is injective.

Solution.

1. ($\implies$)

Assume the sequence is exact at A . Then image of the 0 map is equal to $\ker(i)$. But the image of 0 under any map is 0. So we have $\ker(i) = \{0\} \implies i$ is injective.

($\Leftarrow$)

Assume i is injective. Then $\ker(i) = 0 = \text{im}(0)$, thus making the sequence exact at A .

2. ($\implies$)

Assume the sequence is exact at C . Then $\text{im}(p) = C$, thus p is surjective.

($\Leftarrow$)

Assume p is surjective. Then $\text{im}(p) = C$ which is the kernel of the map from $C \rightarrow 0$ thus making the sequence exact at C .

3. ($\implies$)

Assume the sequence is exact. Then it is exact at A and C . From parts (1) and (2) this implies that i is injective, p is surjective. Since it is also exact at B we must also have $\ker(p) = \text{im}(i)$.

($\Leftarrow$)

Assume i is injective, p is surjective, and $\text{im}(i) = \ker(p)$. Then by parts (1), (2), the sequence is exact at A and C . And since $\text{im}(i) = \ker(p)$, the sequence is exact at B by definition. Thus the sequence is exact.

4. Let $a \in A$. Then $f(a) \in \text{im}(f)$, and since the sequence is exact we have $\text{im}(f) = \ker(g)$. So, $f(a) \in \ker(g) \implies g(f(a)) = 0$. Since a was arbitrary this implies that $g \circ f = 0$.

5. Consider the $\mathbb{Z}$ -modules $A = C = \mathbb{Z}$, $B = \mathbb{Z}^2$ and the morphisms $f = 0$ and $g = \pi$. Then, we have $g \circ f = 0$ but $\ker(g) \subset \text{im}(f)$, so the sequence is not exact.

6. (a) Note that $\text{im}(f) = \{0\}$, $\ker(g) = B$. Then,

$$H = \ker(g) / \text{im}(f) = B / \{0\} = B$$

(b) Note that $\text{im}(f) \cong A$, $\ker(g) = B$. Then,

$$H = \ker(g) / \text{im}(f) \cong B / A$$

(c) Note that $\ker(g) = \{0\}$. Then,

$$H = \ker(g) / \text{im}(f) = \{0\} / \text{im}(f) = \{0\}$$

Problem 2. Examples of Short Exact Sequences.

Let R be a ring. In this problem we will see that a number of constructions we are quite familiar with can be framed in terms of short exact sequences.

1. Let $I \subset R$ be an ideal. Construct a short exact sequence

$$0 \longrightarrow I \xrightarrow{\iota} R \xrightarrow{\pi} R/I \longrightarrow 0$$

where π is the natural quotient map and ι is the inclusion I into R .

2. Let $N \subset M$ be a submodule. Prove that the sequence

$$0 \longrightarrow N \xrightarrow{\iota} M \xrightarrow{\pi} M/N \longrightarrow 0$$

is exact, where ι is the inclusion of N into M .

3. Let $I \subset J \subset R$ be ideals. Construct a short exact sequence

$$0 \longrightarrow J/I \longrightarrow R/I \longrightarrow R/J \longrightarrow 0.$$

Hint: Define $J/I \rightarrow R/I$ by $j + I \mapsto j + I$ and define $R/I \rightarrow R/J$ by $r + I \mapsto r + J$. Check well-definedness before checking exactness.

4. For a positive integer n , prove that multiplication by n gives

$$0 \longrightarrow \mathbb{Z} \xrightarrow{\times n} \mathbb{Z} \longrightarrow \mathbb{Z}/n\mathbb{Z} \longrightarrow 0$$

which is a short exact sequence of $\mathbb{Z}$ -modules.

Solution.

1. **TODO:**
2. Note that $\text{im}(\iota) \cong N$ and $\ker(\pi) = N$. That is, $\text{im}(\iota) = \ker(\pi)$. Moreover since ι is injective and π is surjective the sequence is exact (using Problem 1.3).
3. We define the maps ι and π as follows:
 - $\iota : J/I \rightarrow R/I$ where $\iota(j + I) = j + I$.
 - $\pi : R/I \rightarrow R/J$ where $\pi(r + I) = \bar{r} + J$.

TODO: check well-definedness

$$0 \longrightarrow J/I \xrightarrow{\iota} R/I \xrightarrow{\pi} R/J \longrightarrow 0.$$

The sequence is exact because ι is injective, π is surjective, and $\text{img}(\iota) \cong J/I = \ker(\pi)$ by the first and third isomorphism theorems.

4. The map $\times n$ is injective because $\ker(\times n) = \{0\}$, $\mathbb{Z} \rightarrow \mathbb{Z}/n\mathbb{Z}$ is surjective because it is the natural quotient map, and $\text{im}(\times n) = \langle n \rangle = \ker(\mathbb{Z} \rightarrow \mathbb{Z}/n\mathbb{Z})$, thus the sequence is exact.

Problem 3. Kernels, Cokernels, and Images.

Let $\phi : A \rightarrow B$ be a homomorphism of R -modules. The *cokernel* of ϕ is the R -module $\text{coker}(\phi) := B / \text{img}(\phi)$.

1. Prove that the sequence

$$0 \longrightarrow \ker(\phi) \xrightarrow{\iota} A \xrightarrow{\phi} B$$

where ι is the inclusion of $\ker(\phi)$ into A is exact.

2. Suppose

$$0 \longrightarrow A \xrightarrow{f} B \xrightarrow{g} C$$

is an exact sequence of R -modules. Prove that f induces an isomorphism $A \cong \ker(g)$.

3. Prove that the sequence

$$A \xrightarrow{\phi} B \xrightarrow{\pi} \text{coker}(\phi) \longrightarrow 0$$

where π is the natural quotient map is exact.

4. Suppose

$$A \xrightarrow{f} B \xrightarrow{g} C \longrightarrow 0$$

is an exact sequence of R -modules. Prove that g induces an isomorphism $\text{coker}(f) \cong C$.

5. Deduce there is an exact sequence of R -modules

$$0 \longrightarrow \ker(\phi) \xrightarrow{\iota} A \xrightarrow{\phi} B \xrightarrow{\pi} \text{coker}(\phi) \longrightarrow 0 .$$

6. Prove that ϕ induces an isomorphism $A / \ker(\phi) \cong \text{img}(\phi)$, and deduce that the exact sequence in part (5) factors through $\text{img}(\phi)$ as two short exact sequences

$$0 \longrightarrow \ker(\phi) \xrightarrow{\iota} A \xrightarrow{\bar{\phi}} \text{img}(\phi) \longrightarrow 0$$

and

$$0 \longrightarrow \text{img}(\phi) \xrightarrow{j} B \xrightarrow{\pi} \text{coker}(\phi) \longrightarrow 0$$

where $\bar{\phi}$ is ϕ with codomain restricted to $\text{img}(\phi)$ and j is the inclusion.

Solution.

1. • $\text{im}(0 \rightarrow \ker(\phi)) = 0 = \ker(\iota)$ (inclusion map)
 • $\text{im}(\iota) = \ker(\phi)$

Thus, the sequence is exact.

2. If the sequence is exact, then we have $\text{im}(0 \rightarrow A) = \{0\} = \ker(f)$ and $\text{im}(f) = \ker(g)$. Now we define $\psi : A \rightarrow \ker(g)$ where $\psi(a) = f(a)$. This is well-defined since $\text{im}(f) = \ker(g)$.

- ψ is injective since $\psi(a) = 0 \iff f(a) = 0$ that is $\ker(\psi) = \ker(f)$ but $\ker(f) = \{0\}$.
- ψ is surjective; since $\text{im}(f) = \ker(g)$ we have for any $b \in \ker(g)$ we have $b = f(a) = \psi(a)$ for some $a \in A$.

Thus we have a bijection $\psi : A \rightarrow \ker(g)$ giving us $A \cong \ker(g)$.

3. • $\text{im}(\phi) = \ker(\pi)$ (by definition)
 • $\text{im}(\pi) = \text{coker}(\phi) = \ker(\text{coker}(\phi) \rightarrow 0)$

Thus, the sequence is exact.

4. If the sequence is exact, then we have $\text{im}(f) = \ker(g)$ and $\text{im}(g) = \ker(C \rightarrow 0) = C$. By definition $\text{coker}(f) = B / \text{im}(f)$ Now we define a map $\psi : B / \text{im}(f) \rightarrow C$ by $\psi(\bar{b}) = g(b)$. Then,

- ψ is well-defined since if $\bar{b} = \bar{b}'$ then $b - b' \in \text{im}(f) = \ker(g)$, so $g(b) = g(b')$.
- ψ is injective since if $\psi(\bar{b}) = g(b) = 0 \implies b \in \text{im}(f) \implies \bar{b} = 0$.
- ψ is surjective since $\text{im}(g) = C$.

Thus we have a bijection $\psi : B / \text{im}(f) \rightarrow C$ giving us $\text{coker}(f) \cong C$.

5. From parts (1) and (3) we can conclude that the sequence is exact.

6. We define $\psi : A / \ker(\phi) \rightarrow \text{img}(\phi)$ by $\psi(\bar{a}) = \phi(a)$. Then,

- ψ is well-defined since if $\bar{a} = \bar{a}'$ then $a - a' \in \ker(\phi) \implies \psi(a - a') = 0 \implies \psi(a) = \psi(a')$.
- ψ is injective since if $\psi(\bar{a}) = \phi(a) = 0 \implies a \in \ker(\phi) \implies \bar{a} = 0$.
- ψ is surjective since $\text{im}(\psi) = \text{im}(\phi)$.

Thus we have a bijection $\psi : A / \ker(\phi) \rightarrow \text{im}(\phi)$ giving us $A / \ker(\phi) \cong \text{im}(\phi)$.

Then we have note that the following sequences are exact:

$$0 \longrightarrow \ker(\phi) \xrightarrow{\iota} A \xrightarrow{\bar{\phi}} \text{img}(\phi) \longrightarrow 0$$

- At $\ker(\phi)$: since the inclusion map is injective we have $\text{im}(0 \rightarrow \ker(\phi)) = 0 = \ker(\iota)$.
- At A : $\text{im}(\iota) \cong \ker(\phi) = \ker(\bar{\phi})$
- At $\text{img}(\phi)$: $\bar{\phi}$ is surjective, and $\ker(\text{im}(\phi) \rightarrow 0) = \text{im}(\phi) = \text{im}(\bar{\phi})$.

And

$$0 \longrightarrow \text{img}(\phi) \xrightarrow{j} B \xrightarrow{\pi} \text{coker}(\phi) \longrightarrow 0$$

- At $\text{img}(\phi)$: since the inclusion map j is injective we have $\text{im}(0 \rightarrow \text{im}(\phi)) = 0 = \ker(j)$.
- At B : $\text{im}(j) \cong \text{im}(\phi) = \ker(\pi)$ since $\text{coker}(\phi) = B / \text{im}(\phi)$
- At $\text{coker}(\phi)$: π is surjective by definition, and $\ker(\text{coker}(\phi) \rightarrow 0) = \text{coker}(\phi) = \text{im}(\pi)$.

Thus, the exact sequence in part (5) factors through $\text{img}(\phi)$.

Problem 4. Presentations of Modules.

Let R be a ring. A *presentation* of an R -module M is an exact sequence of the form

$$F_1 \xrightarrow{\varphi} F_0 \longrightarrow M \longrightarrow 0$$

where F_0 and F_1 are free modules.

1. Explain why a presentation of M as above says that M is obtained from the generators of F_0 by imposing the relations coming from $\varphi(F_1)$. Hint: Use exactness to identify M with $F_0 / \text{img}(\varphi)$.
2. Let $R = \mathbb{K}[x]$ and $M = R / \langle x^3 \rangle$. Write a presentation of M using finite free R -modules.
3. Let $R = \mathbb{K}[x, y]$ and $M = R / \langle x^2, xy, y^2 \rangle$. Write a presentation of M using one free generator and three relations.
4. Let A be an $m \times n$ matrix with entries in R , viewed as a map $R^n \rightarrow R^m$. Interpret the sequence

$$R^n \xrightarrow{A} R^m \longrightarrow \text{coker}(A) \longrightarrow 0$$

as a presentation of $\text{coker}(A)$.

Solution.

1. From problem 3.4 we know that $\text{coker}(\varphi) = F_0 / \text{im}(\varphi) \cong M$. And $F_0 / \text{im}(\varphi)$ is determined by what quotienting by $\text{im}(\varphi)$ does to F_0 's distinguished elements. Since the sequence is exact at F_0 , we have $\ker(F_0 \rightarrow M) = \text{im}(\varphi) = \varphi(F_1)$, so these are precisely the relations between the generators. Thus, $M = F_0 / \text{im}(\varphi) = F_0 / \varphi(F_1)$.

2.

$$\mathbb{K}[x] \xrightarrow{\varphi} \mathbb{K}[x] \longrightarrow \mathbb{K}[x] / \langle x^3 \rangle \longrightarrow 0$$

where $\varphi(e_1) = x^3$.

3.

$$(\mathbb{K}[x])^3 \xrightarrow{\varphi} \mathbb{K}[x] \longrightarrow \mathbb{K}[x] / \langle x^2, xy, y^2 \rangle \longrightarrow 0$$

where $\varphi(e_1) = x^2$, $\varphi(e_2) = xy$, $\varphi(e_3) = y^2$.

4. We have $\text{coker}(A) = R^m / \text{im}(A)$, which is a module with m -generators and n relations (the n columns of A give n relations).

We just saw that a short exact sequence $0 \rightarrow A \xrightarrow{i} B \xrightarrow{p} C \rightarrow 0$ of R -modules says that B has a submodule isomorphic to A (namely $\text{img } i$) whose quotient is isomorphic to C (namely $B/\text{img } i \cong C$). One might hope that knowing A and C determines B up to isomorphism, and in the simplest cases it does: $B \cong A \oplus C$. When this happens – and when the isomorphism is compatible with the maps i and p – the sequence is called *split*. More precisely, a short exact sequence is *split* (also called *split exact*) if there exists an isomorphism of R -modules $h : B \rightarrow A \oplus C$ such that the following diagram commutes

$$\begin{array}{ccccccccc} 0 & \longrightarrow & A & \xrightarrow{i} & B & \xrightarrow{p} & C & \longrightarrow & 0 \\ & & \parallel & & \downarrow h & & \parallel & & \\ 0 & \longrightarrow & A & \xrightarrow{\iota} & A \oplus C & \xrightarrow{\pi} & C & \longrightarrow & 0 \end{array}$$

where $\iota(a) = (a, 0)$ is the natural inclusion and $\pi(a, c) = c$ is the natural projection. Saying that the diagram commutes means that the two squares commute. This is the generic situation for vector spaces: every subspace has a complement, so every short exact sequence of vector spaces splits. Over more general rings, however, splitting is the exception rather than the rule.

We can characterize when a short exact sequence splits in terms of the existence of either a section of p or a retraction of i . A *section* of $p : B \rightarrow C$ is an R -module homomorphism $s : C \rightarrow B$ such that $p \circ s = \text{Id}_C$. A *retraction* of $i : A \rightarrow B$ is an R -module homomorphism $r : B \rightarrow A$ such that $r \circ i = \text{Id}_A$. Note that in general s is not an inverse to p , and r is not an inverse to i ; the compositions in the other order need not be the identity.

Lemma 0.1 (Splitting Lemma). *Let R be a ring, and consider a short exact sequence of R -modules*

$$0 \longrightarrow A \xrightarrow{i} B \xrightarrow{p} C \longrightarrow 0.$$

The following are equivalent:

- (i) *The sequence splits, i.e. $B \cong A \oplus C$ in a way compatible with i and p .*
- (ii) *The map p admits a section $s : C \rightarrow B$.*
- (iii) *The map i admits a retraction $r : B \rightarrow A$.*

Condition (ii) is often called *right split*, while condition (iii) is called *left split*. Thus, for R -modules, “right split,” “left split,” and “split” are all equivalent. Warning: This equivalence is special to the setting of R -modules (more generally, abelian categories). For example, in the category of non-abelian groups, a sequence may admit a section without being a direct product: such extensions correspond to semidirect products. Fortunately, none of these issues arise for R -modules.

Problem 5. Split Short Exact Sequences.

Let

$$0 \longrightarrow A \xrightarrow{i} B \xrightarrow{p} C \longrightarrow 0$$

be a short exact sequence of R -modules.

1. Suppose there exists a homomorphism $s : C \rightarrow B$ such that $p \circ s = \text{Id}_C$. Prove that the map defined below is an isomorphism of R -modules:

$$\Phi : A \oplus C \longrightarrow B, \quad (a, c) \longmapsto i(a) + s(c).$$

2. Conversely, suppose there is an isomorphism $B \cong A \oplus C$ under which i becomes the standard inclusion $A \rightarrow A \oplus C$ and p becomes the standard projection $A \oplus C \rightarrow C$. Construct a homomorphism $s : C \rightarrow B$ with $p \circ s = \text{Id}_C$.
3. Suppose there exists a homomorphism $r : B \rightarrow A$ such that $r \circ i = \text{Id}_A$.

- (a) Prove that the map

$$B \longrightarrow A \oplus C, \quad b \longmapsto (r(b), p(b))$$

is injective.

- (b) Prove that it is surjective. Hint: Given $(a, c) \in A \oplus C$, choose $b_0 \in B$ with $p(b_0) = c$ and modify b_0 by an element of $i(A)$.
 - (c) Conclude that $B \cong A \oplus C$.
4. Conversely, suppose there is an isomorphism $B \cong A \oplus C$ under which i becomes the standard inclusion $a \mapsto (a, 0)$ and p becomes the standard projection $(a, c) \mapsto c$. Construct a homomorphism $r : B \rightarrow A$ with $r \circ i = \text{Id}_A$.

Solution.

1. Φ is an R -module homomorphism:

- $\Phi((a + b, c + d)) = i(a + b) + s(c + d) = i(a) + i(b) + s(c) + s(d) = \Phi(a, c) + \Phi(b, d)$.
- $\Phi(r(a, c)) = i(ra) + s(rc) = ri(a) + rs(c) = r(i(a) + s(c)) = r(\Phi(a, c))$.

Φ is injective:

Suppose $\Phi((a, c)) = 0$, then $i(a) + s(c) = 0$. We apply p to both sides. $p(\Phi(a, c)) = p(i(a) + s(c)) = p(i(a)) + c$. But $i(a) \in \ker(p)$, so we have $p(i(a)) + c = 0 + c = 0$. Hence, $c = 0$. Substituting back, $i(a) + s(0) = 0 \implies i(a) = 0 \implies a = 0$ since i is injective. Thus, if $\Phi((a, c)) = 0$ then $(a, c) = (0, 0)$.

Φ is surjective:

Let $b \in B$. Then we have, $p(b) = c$ for some $c \in C$. Now consider $p(b - s(c))$,

$$p(b - s(c)) = p(b) - p(s(c)) = c - c = 0$$

That is $b - s(c) \in \ker(p) = \text{im}(A)$, i.e. there exists a $a \in A$ such that $i(a) = b - s(c)$. Upon re-arranging we get,

$$b = i(a) + s(c) = \Phi((a, c))$$

Thus the map Φ is an isomorphism, and $A \oplus C \cong B$.

2. Let $\phi : B \rightarrow A \oplus C$ be the isomorphism, and let $j : C \rightarrow A \oplus C$ be the standard inclusion. Then we define $s : C \rightarrow B$ as $s(c) = \phi^{-1}(j(c)) = \phi^{-1}((0, c))$ for all $c \in C$. Then

$$p \circ s = p \circ \phi^{-1} \circ j = \pi \circ j = \text{id}_C$$

3. (a) Let ϕ denote the map from $B \rightarrow A \oplus C$. Then, if $\phi(b) = (0, 0) \implies (r(b), p(b)) = (0, 0)$.
- $p(b) = 0 \implies b \in \ker(p) = \text{im}(i)$. So $b = i(a)$ for some a .
 - $r(b) = 0 \implies r(i(a)) = 0$ but $r \circ i = \text{id}_A$ so $a = 0$.

Thus the map ϕ is injective.

- (b) Let $(a, c) \in A \oplus C$ be an arbitrary element. Since p is surjective we have $c = p(b)$ for some $b \in B$. Let $b' = b - i(r(b)) + i(a)$. Then $\phi(b') = (r(b'), p(b'))$. Where,
- $r(b') = r(b) - r(i(r(b))) + r(i(a)) = r(b) - r(b) + a = a$
 - $p(b') = p(b) - p(i(r(b))) + p(i(a)) = c + 0 + 0 + c$ since $\text{im}(i) = \ker(p)$.

Thus the map ϕ is surjective.

- (c) Therefore we conclude $B \cong A \oplus C$.

4. Let $\phi : B \rightarrow A \oplus C$ be the isomorphism. Then we define $r = \pi_A \circ \phi$, where $\pi_A : A \oplus C \rightarrow A$ be the standard projection to A . Then,

$$r \circ i = \pi_A \circ \phi \circ i = \pi_A \circ \iota = \text{id}_A$$

Problem 6. Examples of Split & Non-Split Exact Sequences.

1. Prove that every short exact sequence of vector spaces over a field $\mathbb{K}$ splits.
Hint: Choose a basis for the first term, extend it to a basis for the middle term, and identify the remaining basis vectors with a basis for the third term.
2. Explain why the Rank-Nullity theorem for finite dimensional vector spaces over $\mathbb{K}$ is a consequence of the splitting theorem.
3. Prove that the sequence

$$0 \longrightarrow \mathbb{Z} \xrightarrow{\cdot 2} \mathbb{Z} \longrightarrow \mathbb{Z}/2\mathbb{Z} \longrightarrow 0$$

does not split as a sequence of $\mathbb{Z}$ -modules.

4. Let p be a prime number. Prove that the sequence of $\mathbb{Z}$ -modules:

$$0 \longrightarrow \mathbb{Z}/p\mathbb{Z} \longrightarrow \mathbb{Z}/p^2\mathbb{Z} \longrightarrow \mathbb{Z}/p\mathbb{Z} \longrightarrow 0,$$

where the first map sends $\bar{1} \mapsto \bar{p}$ and the second is reduction modulo p , does not split. Hint: A section would send $\bar{1} \in \mathbb{Z}/p\mathbb{Z}$ to an element of $\mathbb{Z}/p^2\mathbb{Z}$ whose reduction modulo p is $\bar{1}$. What is the order of such an element?

5. Let $R = \mathbb{K}[x]$. Prove that the sequence below is exact, but does not split as a sequence of R -modules:

$$0 \longrightarrow \langle x \rangle \longrightarrow \mathbb{K}[x] \longrightarrow \mathbb{K} \longrightarrow 0.$$

Hint: If a section $s : \mathbb{K} \rightarrow \mathbb{K}[x]$ existed as a $\mathbb{K}[x]$ -module map, what could $s(1)$ be? Use the fact that $x \cdot 1 = 0$ in $\mathbb{K} \cong \mathbb{K}[x]/\langle x \rangle$.

6. Let R be a ring and let $e \in R$ be an idempotent, meaning $e^2 = e$. Prove that

$$0 \longrightarrow eR \hookrightarrow R \twoheadrightarrow R/eR \longrightarrow 0$$

splits. Then prove that the map $R \rightarrow (1-e)R$ given by $r \mapsto (1-e)r$ induces an isomorphism $R/eR \cong (1-e)R$.

Solution.

1.

We now introduce just enough categorical language to talk cleanly about Hom, tensor product, and localization. A *category* is a collection of objects together with, for each pair of objects, a set of morphisms between them, equipped with an associative composition law and an identity morphism for each object. The quintessential category, both in this class and arguably beyond, is the category of R -modules, denoted $R - \mathbf{Mod}$, whose objects are R -modules and whose morphisms are R -module homomorphisms.

A *functor* between categories $\mathbf{C}$ and $\mathbf{D}$ is a mapping that assigns to each object of $\mathbf{C}$ an object of $\mathbf{D}$ and to each morphism in $\mathbf{C}$ a morphism in $\mathbf{D}$, preserving identity morphisms and composition. A functor is contravariant if it reverses the order of composition and covariant if it preserves the order of composition. More precisely, a *covariant functor* $F : \mathbf{C} \rightarrow \mathbf{D}$ is the data of

1. for every object X of $\mathbf{C}$, $F(X)$ is an object of $\mathbf{D}$,
2. for every morphism $f : X \rightarrow Y$ in $\mathbf{C}$, $F(f) : F(X) \rightarrow F(Y)$ is a morphism in $\mathbf{D}$,

which satisfies the following conditions:

- (i) $F(\text{Id}_X) = \text{Id}_{F(X)}$ for every object X of $\mathbf{C}$,
- (ii) $F(g \circ f) = F(g) \circ F(f)$ for all composable morphisms $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ in $\mathbf{C}$.

Similarly a *contravariant functor* $F : \mathbf{C} \rightarrow \mathbf{D}$ is the data of:

1. for every object X of $\mathbf{C}$, $F(X)$ is an object of $\mathbf{D}$,
2. for every morphism $f : X \rightarrow Y$ in $\mathbf{C}$, $F(f) : F(Y) \rightarrow F(X)$ is a morphism in $\mathbf{D}$,

which satisfies the following conditions:

- (i) $F(\text{Id}_X) = \text{Id}_{F(X)}$ for every object X of $\mathbf{C}$,
- (ii) $F(g \circ f) = F(f) \circ F(g)$ for all composable morphisms $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ in $\mathbf{C}$.

We have already constructed several operations that take modules as input and produce new modules. For example, after fixing an R -module T , the assignments

$$M \longmapsto \text{Hom}_R(T, M), \quad M \longmapsto \text{Hom}_R(M, T), \quad M \longmapsto M \otimes_R T$$

each send an object of $R - \mathbf{Mod}$ to another object of $R - \mathbf{Mod}$. Similarly, if $U \subset R$ is a multiplicatively closed set, localization gives an assignment

$$M \longmapsto U^{-1}M,$$

which sends an R -module to a module over the localized ring $U^{-1}R$. We would like to view each of these constructions as defining functors:

$$R\text{-}\mathbf{Mod} \xrightarrow{\mathrm{Hom}_R(T,-)} R\text{-}\mathbf{Mod} \qquad R\text{-}\mathbf{Mod} \xrightarrow{\mathrm{Hom}_R(-,T)} R\text{-}\mathbf{Mod}$$

$$R\text{-}\mathbf{Mod} \xrightarrow{-\otimes_R T} R\text{-}\mathbf{Mod} \qquad R\text{-}\mathbf{Mod} \xrightarrow{U^{-1}-} U^{-1}R\text{-}\mathbf{Mod}$$

However, a functor is not merely a rule on objects. To view these constructions functorially, we must also understand what they do to morphisms. Thus, given an R -module homomorphism $f : M \rightarrow N$, we must define induced maps between the corresponding Hom modules, tensor products, and localizations, and then check that these induced maps respect identities and composition. The following exercise asks you to make these constructions explicit and verify that they really do define functors.

Problem 7. Hom, Tensor, and Localization as Functors.

Throughout let R be a ring and fix an R -module T .

1. Let T be a fixed R -module. If $f : M \rightarrow N$ is a homomorphism, define

$$f_* : \text{Hom}_R(T, M) \rightarrow \text{Hom}_R(T, N), \quad \phi \mapsto f \circ \phi.$$

Prove that $\text{Hom}_R(T, -)$ is a covariant functor from $R - \mathbf{Mod}$ to $R - \mathbf{Mod}$.

2. If $f : M \rightarrow N$ is a homomorphism, define

$$f^* : \text{Hom}_R(N, T) \rightarrow \text{Hom}_R(M, T), \quad \psi \mapsto \psi \circ f.$$

Prove that $\text{Hom}_R(-, T)$ is a contravariant functor from $R - \mathbf{Mod}$ to $R - \mathbf{Mod}$.

3. If $f : M \rightarrow N$ is a homomorphism, use the universal property of tensor products to construct

$$f \otimes \text{Id}_T : M \otimes_R T \rightarrow N \otimes_R T, \quad m \otimes t \mapsto f(m) \otimes t.$$

Prove that $- \otimes_R T$ is a covariant functor from $R - \mathbf{Mod}$ to $R - \mathbf{Mod}$.

4. Let $U \subset R$ be a multiplicatively closed set. If $f : M \rightarrow N$ is a homomorphism, define

$$U^{-1}f : U^{-1}M \rightarrow U^{-1}N, \quad \frac{m}{u} \mapsto \frac{f(m)}{u}.$$

Prove that this is well-defined and that $U^{-1}(-)$ is a covariant functor from $R - \mathbf{Mod}$ to $U^{-1}R - \mathbf{Mod}$.

Solution.

1. • We know that $\text{Hom}_R(T, -)$ is a $R - \mathbf{Mod}$. That is, for every R -module M we have $\text{Hom}_R(T, M)$ another R -module.
• Let $f : M \rightarrow N, g : N \rightarrow O$ be R -module homomorphisms. Then, we need to show that f_* and g_* are R -module homomorphisms.

$$- r \cdot f_*(\phi) = r \cdot f \circ \phi = f \circ (r\phi) = f_*(r\phi)$$

$$- f_*(\phi + \psi) = f \circ (\phi + \psi) = f \circ \phi + f \circ \psi = f_*(\phi) + f_*(\psi)$$

A similar proof works to show g_* is a R -module homomorphism.

- $(g \circ f)_*(\phi) = g \circ f \circ \phi = g_*(f \circ \phi) = g_*(f_*(\phi)) = (g_* \circ f_*)(\phi)$
• Let $h : M \rightarrow M$ be a R -module homomorphism, where $h = \text{id}_M$. Then

$$h_* = h \circ \phi = \text{id}_M \circ \phi = \text{id}_{\text{Hom}_R(T, M)}$$

Thus, $\text{Hom}_R(T, -)$ is a covariant functor from $R - \mathbf{Mod}$ to $R - \mathbf{Mod}$.

2. • We know that $\text{Hom}_R(-, T)$ is a $R - \mathbf{Mod}$. That is, for every R -module M we have $\text{Hom}_R(M, T)$ another R -module.
- Let $f : M \rightarrow N, g : N \rightarrow O$ be R -module homomorphisms. Then, we need to show that f^* and g^* are R -module homomorphisms.

$$- r \cdot f^*(\phi) = r \cdot \phi \circ f = (r\phi) \circ f = f^*(r\phi)$$

$$- f^*(\phi + \psi) = (\phi + \psi) \circ f = \phi \circ f + \psi \circ f = f^*(\phi) + f^*(\psi)$$

A similar proof works to show g^* is a R -module homomorphism.

- $(g \circ f)^*(\phi) = \phi \circ g \circ f = f^*(\phi \circ g) = f^*(g^*(\phi)) = (f^* \circ g^*)(\phi)$
- Let $h : M \rightarrow M$ be a R -module homomorphism, where $h = \text{id}_M$. Then

$$h_* = \phi \circ h = \phi \circ \text{id}_M = \text{id}_{\text{Hom}_R(M, T)}$$

Thus, $\text{Hom}_R(T, -)$ is a contravariant functor from $R - \mathbf{Mod}$ to $R - \mathbf{Mod}$.

3. • We know that for any R -module $M, M \otimes_R T$ is a R -module.
- Let $f : M \rightarrow N, g : N \rightarrow O$ be R -module homomorphisms. Then, we need to show that $f \otimes \text{Id}_T$ and $g \otimes \text{Id}_T$ are R -module homomorphisms.

$$- r \cdot f \otimes \text{Id}_T(m \otimes t) = r \cdot f(m) \otimes t = f(rm) \otimes t = f \otimes \text{Id}_T(rm \otimes t)$$

-

$$\begin{aligned} f \otimes \text{Id}_T((m + n) \otimes t) &= f(m + n) \otimes t \\ &= f(m) \otimes t + f(n) \otimes t \\ &= f \otimes \text{Id}_T(m \otimes t) + f \otimes \text{Id}_T(n \otimes t) \end{aligned}$$

A similar proof works to show $g \otimes \text{Id}_T$ is a R -module homomorphism.

•

$$\begin{aligned} (g \circ f) \otimes \text{Id}_T(m \otimes t) &= (g \circ f)(m) \otimes t = g(f(m)) \otimes t \\ &= g \otimes \text{Id}_T(f(m) \otimes t) \\ &= (g \otimes \text{Id}_T) \circ (f \otimes \text{Id}_T)(m \otimes t) \end{aligned}$$

- Let $h : M \rightarrow M$ be a R -module homomorphism, where $h = \text{Id}_M$. Then

$$h \otimes \text{Id}_T(m \otimes t) = h(m) \otimes t = m \otimes t = (\text{Id}_M \otimes \text{Id}_T)(m \otimes t)$$

Thus, $- \otimes_R T$ is a covariant functor from $R - \mathbf{Mod}$ to $R - \mathbf{Mod}$.

4. • We know that $U^{-1}(-)$ is a $R - \mathbf{Mod}$. That is, for every R -module M we have $U^{-1}M$ another R -module.

- Let $f : M \rightarrow N, g : N \rightarrow O$ be R -module homomorphisms. Then, we need to show that $U^{-1}f$ and $U^{-1}g$ are R -module homomorphisms.

$$\begin{aligned}
- r \cdot U^{-1}f\left(\frac{m}{u}\right) &= \frac{rf(m)}{u} = \frac{f(rm)}{u} = U^{-1}f\left(\frac{rm}{u}\right) \\
- U^{-1}f\left(\frac{m}{u}\right) + U^{-1}f\left(\frac{n}{s}\right) &= \frac{f(m)}{u} + \frac{f(n)}{s} = \frac{f(ms)+f(nu)}{us} = U^{-1}f\left(\frac{ms+nu}{us}\right) = \\
&U^{-1}f\left(\frac{m}{u} + \frac{n}{s}\right)
\end{aligned}$$

A similar proof works to show g^* is a R -module homomorphism.

- $(U^{-1}g \circ U^{-1}f)\left(\frac{m}{u}\right) = U^{-1}g\left(U^{-1}f\left(\frac{m}{u}\right)\right) = \frac{g(f(m))}{u} = U^{-1}(f \circ g)\left(\frac{m}{u}\right)$
- Let $h : M \rightarrow M$ be a R -module homomorphism, where $h = \text{id}_M$. Then

$$U^{-1}h\left(\frac{m}{u}\right) = \frac{h(m)}{u} = \frac{m}{u}$$

Thus, $U^{-1}(-)$ is a covariant functor from $R - \mathbf{Mod}$ to $R - \mathbf{Mod}$.

The functors constructed above let us ask a new version of the exactness question. Instead of asking whether a given sequence of modules is exact, we can ask whether a construction preserves exactness. Roughly speaking, a functor is exact if applying it to an exact sequence does not introduce any new homology: the kernel of one map is still the image of the previous. This is a very strong condition, and many of the most important functors in algebra preserve only part of an exact sequence.

For the functors in this worksheet, applying the functor to a sequence again produces a sequence: these functors are additive, so they send zero maps to zero maps, and they respect composition. Let $F : R\text{-}\mathbf{Mod} \rightarrow S\text{-}\mathbf{Mod}$ be one of the covariant functors under consideration, where S is some ring. We say that F is *exact* if, whenever

$$M' \xrightarrow{f} M \xrightarrow{g} M''$$

is exact at M , the transformed sequence

$$F(M') \xrightarrow{F(f)} F(M) \xrightarrow{F(g)} F(M'')$$

is exact at $F(M)$. Equivalently, F sends exact sequences to exact sequences. In particular, an exact covariant functor sends every short exact sequence

$$0 \longrightarrow A \xrightarrow{i} B \xrightarrow{p} C \longrightarrow 0$$

to a short exact sequence

$$0 \longrightarrow F(A) \xrightarrow{F(i)} F(B) \xrightarrow{F(p)} F(C) \longrightarrow 0.$$

Thus exactness means that $F(i)$ remains injective, $F(p)$ remains surjective, and $\text{img}(F(i)) = \ker(F(p))$. A covariant functor F is called *left exact* if every exact sequence

$$0 \longrightarrow A \xrightarrow{i} B \xrightarrow{p} C$$

is sent to a sequence

$$0 \longrightarrow F(A) \xrightarrow{F(i)} F(B) \xrightarrow{F(p)} F(C)$$

which is exact at $F(A)$ and $F(B)$. So a left exact functor preserves the injectivity at the left end and preserves the kernel condition in the middle, but it may fail to make $F(p)$ surjective. Dually, F is called *right exact* if every exact sequence

$$A \xrightarrow{i} B \xrightarrow{p} C \longrightarrow 0$$

is sent to a sequence

$$F(A) \xrightarrow{F(i)} F(B) \xrightarrow{F(p)} F(C) \longrightarrow 0$$

which is exact at $F(B)$ and $F(C)$. Thus a right exact functor preserves the quotient at the right end, but it may fail to preserve injectivity at the left end. A functor that is both left exact and right exact is exact. One can also define exactness, left exactness, and right exactness for contravariant functors; in that case, the arrows are reversed.

The next exercises focus on three fundamental examples. The Hom functors are naturally left exact. For $\text{Hom}_R(T, -)$, the reason is that a map $T \rightarrow B$ whose composite with $B \rightarrow C$ is zero is exactly a map whose image lands in the submodule $A \subseteq B$. For the contravariant functor $\text{Hom}_R(-, T)$, the same idea appears in quotient form: a map $B \rightarrow T$ that kills A factors uniquely through $C = B/A$. In both cases, surjectivity can fail; a map into or out of one end of a short exact sequence need not lift or extend across the middle term.

Tensor product behaves in the opposite direction. The functor $- \otimes_R T$ is always right exact: it preserves cokernels and therefore preserves quotient presentations. This is why tensor product is so useful for computing modules from generators and relations. But tensor product need not be left exact; after tensoring, an injective map can acquire a kernel. The modules T for which tensoring with T is exact are precisely the flat modules, which will be introduced below.

Localization is the especially well-behaved example in commutative algebra. Since $U^{-1}M \cong U^{-1}R \otimes_R M$, localization is right exact for the same reason tensor product is right exact. What makes localization better than an arbitrary tensor product is that it is also left exact: if a fraction becomes zero after applying a localized map, then after clearing denominators its numerator already came from the previous module. Thus localization preserves all exact sequences, and this exactness is one of the main reasons localization is such a powerful tool.

Problem 8. Left Exactness of $\text{Hom}_R(T, -)$.

Let T be an R -module and let

$$0 \longrightarrow A \xrightarrow{i} B \xrightarrow{p} C \longrightarrow 0$$

be a short exact sequence.

1. Define the maps i_* and p_* by postcomposition, and prove that $p_* \circ i_* = 0$.
2. Prove that $i_* : \text{Hom}_R(T, A) \rightarrow \text{Hom}_R(T, B)$ is injective.
3. Suppose $\phi : T \rightarrow B$ satisfies $p \circ \phi = 0$. Prove that ϕ factors uniquely through $i : A \rightarrow B$. In other words, prove that there is a unique homomorphism $\tilde{\phi} : T \rightarrow A$ such that $\phi = i \circ \tilde{\phi}$.
4. Conclude that the sequence below is exact

$$0 \longrightarrow \text{Hom}_R(T, A) \xrightarrow{i_*} \text{Hom}_R(T, B) \xrightarrow{p_*} \text{Hom}_R(T, C).$$

5. Suppose the original short exact sequence splits. Prove that $p_* : \text{Hom}_R(T, B) \rightarrow \text{Hom}_R(T, C)$ is surjective. Thus $\text{Hom}_R(T, -)$ sends split short exact sequences to short exact sequences.
6. Let $R = \mathbb{Z}$ and apply $\text{Hom}_{\mathbb{Z}}(\mathbb{Z}/2\mathbb{Z}, -)$ to

$$0 \longrightarrow \mathbb{Z} \xrightarrow{\cdot 2} \mathbb{Z} \longrightarrow \mathbb{Z}/2\mathbb{Z} \longrightarrow 0$$

Compute the resulting sequence and explain why $\text{Hom}_{\mathbb{Z}}(\mathbb{Z}/2\mathbb{Z}, -)$ is not right exact.

Solution.

1. We define $i_*(\phi) = \phi \circ i$ and $p_*(\psi) = \psi \circ p$ for $\phi \in \text{Hom}_R(T, A)$ and $\psi \in \text{Hom}_R(T, B)$. Let $i(a) = b$. Then,

$$p_* \circ i_*(\phi) = \psi \circ p \circ \phi \circ i$$

1. **Left Exactness of $\text{Hom}_R(-, T)$.** Let T be an R -module and consider the short exact sequence below:

$$0 \longrightarrow A \xrightarrow{i} B \xrightarrow{p} C \longrightarrow 0.$$

- (a) Using precomposition define the maps as below. Prove that $i^* \circ p^* = 0$:

$$\text{Hom}_R(C, T) \xrightarrow{p^*} \text{Hom}_R(B, T) \xrightarrow{i^*} \text{Hom}_R(A, T) :$$

- (b) Prove that p^* is injective.
- (c) Suppose $\psi : B \rightarrow T$ satisfies $\psi \circ i = 0$. Prove that ψ factors uniquely through $p : B \rightarrow C$. In other words, prove that there is a unique homomorphism $\tilde{\psi} : C \rightarrow T$ such that $\psi = \tilde{\psi} \circ p$.
- (d) Conclude that the sequence below is exact:

$$0 \longrightarrow \text{Hom}_R(C, T) \xleftarrow{p^*} \text{Hom}_R(B, T) \xrightarrow{i^*} \text{Hom}_R(A, T).$$

- (e) Suppose the original short exact sequence splits. Prove that $i^* : \text{Hom}_R(B, T) \rightarrow \text{Hom}_R(A, T)$ is surjective. Thus $\text{Hom}_R(-, T)$ sends split short exact sequences to short exact sequences.
- (f) Let $R = \mathbb{Z}$ and apply $\text{Hom}_{\mathbb{Z}}(-, \mathbb{Z})$ to

$$0 \longrightarrow \mathbb{Z} \xrightarrow{\cdot 2} \mathbb{Z} \longrightarrow \mathbb{Z}/2\mathbb{Z} \longrightarrow 0$$

Compute the resulting sequence and explain why $\text{Hom}_{\mathbb{Z}}(-, \mathbb{Z})$ is not exact.

2. Hom Computations from Exactness.

- (a) Let M be an R -module and let $I \subseteq R$ be an ideal. Use the short exact sequence

$$0 \longrightarrow I \longrightarrow R \longrightarrow R/I \longrightarrow 0$$

together with $\text{Hom}_R(-, M)$ to recover the description

$$\text{Hom}_R(R/I, M) \cong \{m \in M \mid Im = 0\}.$$

Hint: Identify $\text{Hom}_R(R, M)$ with M by evaluating a homomorphism at $1 \in R$. Under this identification, what does the condition that a map $R \rightarrow M$ vanish on I become?

- (b) Let N be an abelian group and n a positive integer. Apply $\text{Hom}_{\mathbb{Z}}(-, N)$ to

$$0 \longrightarrow \mathbb{Z} \xrightarrow{\cdot n} \mathbb{Z} \longrightarrow \mathbb{Z}/n\mathbb{Z} \longrightarrow 0$$

Show that there is an exact sequence

$$0 \longrightarrow \text{Hom}_{\mathbb{Z}}(\mathbb{Z}/n\mathbb{Z}, N) \longrightarrow N \xrightarrow{\cdot n} N$$

Deduce that $\text{Hom}_{\mathbb{Z}}(\mathbb{Z}/n\mathbb{Z}, N)$ is naturally isomorphic to the subgroup of n -torsion elements of N .

- (c) For positive integers m, n , use the previous part to compute $\text{Hom}_{\mathbb{Z}}(\mathbb{Z}/m\mathbb{Z}, \mathbb{Z}/n\mathbb{Z})$.

- (d) Let $R = \mathbb{K}[x]$ and $M = R/\langle x^b \rangle$. Compute $\text{Hom}_R(R/\langle x^a \rangle, M)$ for positive integers a and b .
- (e) Let A be an $m \times n$ matrix with entries in R , let $M = \text{coker}(A)$, and let N be an R -module. Use the presentation

$$R^n \xrightarrow{A} R^m \twoheadrightarrow M \longrightarrow 0$$

to describe $\text{Hom}_R(M, N)$ as the kernel of an explicit map $N^m \rightarrow N^n$. Hint: Identify $\text{Hom}_R(R^m, N)$ with N^m and $\text{Hom}_R(R^n, N)$ with N^n . The induced map is precomposition with A ; in coordinates it is given by the transpose matrix A^t acting on column vectors, with the usual convention.

Problem 11. Right Exactness of Tensor Product.

Let T be an R -module and let the below be an exact sequence:

$$A \xrightarrow{f} B \xrightarrow{g} C \longrightarrow 0$$

1. Prove that $g \otimes \text{Id}_T : B \otimes_R T \rightarrow C \otimes_R T$ is surjective.
2. Construct a bilinear map $C \times T \rightarrow (B \otimes_R T) / \langle f(a) \otimes t \mid a \in A, t \in T \rangle$ by choosing a representative $b \in B$ of $c \in C$ and sending (c, t) to the class of $b \otimes t$. Check that this is well-defined.
3. Since $C \cong B / \text{img}(f)$, use the universal property of quotients and tensor products to show that

$$C \otimes_R T \cong (B \otimes_R T) / \langle f(a) \otimes t \mid a \in A, t \in T \rangle.$$

4. Prove that $\ker(g \otimes \text{Id}_T) = \text{img}(f \otimes \text{Id}_T)$.
5. Conclude that the sequence below is exact:

$$A \otimes_R T \longrightarrow B \otimes_R T \longrightarrow C \otimes_R T \longrightarrow 0.$$

6. Suppose the original short exact sequence splits. Prove that $g \otimes \text{Id}_T$ is injective. Thus $- \otimes_R T$ sends split short exact sequences to short exact sequences.

Solution.

1. We know that for all $c \in C$, there exists $b \in B$ such that $g(b) = c$. Now, let $c \otimes_R t \in C \otimes_R T$. Then we have,

$$g \otimes \text{Id}_T(b \otimes_R t) = g(b) \otimes_R t = c \otimes_R t$$

Hence the map $g \otimes_R \text{Id}_T$ is surjective.

2. Let $\phi : C \times T \rightarrow (B \otimes_R T) / \langle f(a) \otimes t \rangle$, where $(c, t) \mapsto \overline{b \otimes t}$ where $g(b) = c$. To check well-definedness let $g(b) = g(b')$. Then we have $g(b - b') = 0$, i.e. $b - b' \in \ker(g) = \text{im}(f)$. So $b - b' = f(a)$ for some $a \in A$.

$$(b - b') \otimes t = f(a) \otimes t \in \langle f(a) \otimes t \rangle \implies b \otimes t = b' \otimes t$$

3.

Problem 12. Tensor Product Need Not Be Left Exact.

The next example shows that tensor product can turn a nonzero injective map into the zero map.

1. Tensor the short exact sequence

$$0 \longrightarrow \mathbb{Z} \xrightarrow{\cdot 2} \mathbb{Z} \longrightarrow \mathbb{Z}/2\mathbb{Z} \longrightarrow 0$$

with $\mathbb{Z}/2\mathbb{Z}$. Compute the resulting sequence explicitly.

2. Explain why the map $\mathbb{Z} \otimes_{\mathbb{Z}} \mathbb{Z}/2\mathbb{Z} \rightarrow \mathbb{Z} \otimes_{\mathbb{Z}} \mathbb{Z}/2\mathbb{Z}$ induced by multiplication by 2 is the zero map.
3. Conclude that $- \otimes_{\mathbb{Z}} \mathbb{Z}/2\mathbb{Z}$ is not left exact.
4. Let n be a positive integer. Compute $(\mathbb{Z}/n\mathbb{Z}) \otimes_{\mathbb{Z}} (\mathbb{Z}/2\mathbb{Z})$ and use this to predict what happens when one tensors the exact sequence below with $\mathbb{Z}/2\mathbb{Z}$:

$$0 \longrightarrow \mathbb{Z} \xrightarrow{\cdot n} \mathbb{Z} \longrightarrow \mathbb{Z}/n\mathbb{Z} \longrightarrow 0.$$

Solution.

1. We get,

$$0 \longrightarrow \mathbb{Z}/2\mathbb{Z} \xrightarrow{2 \otimes \text{id}_{\mathbb{Z}/2\mathbb{Z}}} \mathbb{Z}/2\mathbb{Z} \longrightarrow \mathbb{Z}/2\mathbb{Z} \longrightarrow 0$$

Since $\mathbb{Z} \otimes_{\mathbb{Z}} \mathbb{Z}/2\mathbb{Z} \cong \mathbb{Z}/2\mathbb{Z}$ and $\mathbb{Z}/\mathbb{Z} \otimes_{\mathbb{Z}} \mathbb{Z}/2\mathbb{Z} \cong \mathbb{Z}/2\mathbb{Z}$.

2. Let $a \otimes_{\mathbb{Z}} b \in \mathbb{Z} \otimes_{\mathbb{Z}} \mathbb{Z}/2\mathbb{Z}$. Then,

$$\cdot 2 \otimes_{\mathbb{Z}} \text{id}_{\mathbb{Z}/2\mathbb{Z}}(a \otimes_{\mathbb{Z}} b) = 2a \otimes_{\mathbb{Z}} b = a \otimes_{\mathbb{Z}} 2b = a \otimes_{\mathbb{Z}} 0 = 0(a \otimes_{\mathbb{Z}} b) = 0$$

Hence the map induced by multiplication by 2 is the zero map.

3. Since the map induced by multiplication by 2 is the zero map, it is not injective and hence $-\otimes_{\mathbb{Z}} \mathbb{Z}/2\mathbb{Z}$ is not left exact.
4. $(\mathbb{Z}/n\mathbb{Z}) \otimes_{\mathbb{Z}} (\mathbb{Z}/2\mathbb{Z}) \cong \mathbb{Z}/(n\mathbb{Z} + 2\mathbb{Z}) = \mathbb{Z}/(\gcd(n, 2)\mathbb{Z})$. That is if n is even, we have $(\mathbb{Z}/n\mathbb{Z}) \otimes_{\mathbb{Z}} (\mathbb{Z}/2\mathbb{Z}) \cong \mathbb{Z}/2\mathbb{Z}$ and the map induced by multiplication by n is the zero map and the sequence is not exact at the first term.

$$\mathbb{Z}/2\mathbb{Z} \xrightarrow{0} \mathbb{Z}/2\mathbb{Z} \longrightarrow \mathbb{Z}/2\mathbb{Z} \longrightarrow 0$$

If n is odd, we have $(\mathbb{Z}/n\mathbb{Z}) \otimes_{\mathbb{Z}} (\mathbb{Z}/2\mathbb{Z}) \cong 0$, the map induced by multiplication by n remains injective, and the sequence is exact.

$$\mathbb{Z}/2\mathbb{Z} \xrightarrow{\cdot n} \mathbb{Z}/2\mathbb{Z} \longrightarrow 0 \longrightarrow 0$$

Problem 13. Flat Modules.

An R -module T is called *flat* if the functor $- \otimes_R T$ is exact, or equivalently if tensoring with T preserves injective maps.

1. Explain why right exactness of tensor product implies that T is flat if and only if, for every injection $A \hookrightarrow B$, the induced map $A \otimes_R T \rightarrow B \otimes_R T$ is injective.
2. Prove that R is flat as an R -module.
3. Prove that a finite free module R^n is flat.
4. Prove that every free R -module is flat. Hint: Tensor product commutes with direct sums.

Solution.

1. By right exactness of tensor products we have two of the three sufficient conditions for exactness. The third condition requires injective maps to map to injective maps. Thus, T is flat iff for every injection $A \hookrightarrow B$, the induced map $A \otimes_R T \rightarrow B \otimes_R T$ is injective.
2. We need to show that tensoring with R preserves exactness, but since $A \otimes_R R \cong A$ gives us back the original sequence. Hence the functor $\otimes_R R$ is exact and R is flat.
3. For any exact sequence $0 \rightarrow A \rightarrow B$, we need to show $0 \rightarrow A \otimes_R R^n \rightarrow B \otimes_R R^n$ is exact. But $A \otimes_R R^n \cong A^n$, this becomes $0 \rightarrow A^n \rightarrow B^n$, and the map $A^n \rightarrow B^n$ is just $i^{\oplus n}$ (applying i componentwise), which is injective since i is injective. Hence a finite free module R^n is flat.
4. We know $A \otimes R^{(I)} \cong \bigoplus_{i \in I} A$, then the map $\bigoplus_{i \in I} A \rightarrow \bigoplus_{i \in I} B$ is injective because $A \rightarrow B$ is injective componentwise.

Localization is the most important exact functor in commutative algebra. Let $U \subset R$ be a multiplicatively closed set. The localization construction sends an R -module M to a $U^{-1}R$ -module $U^{-1}M$, and sends a homomorphism $f : M \rightarrow N$ to the homomorphism

$$U^{-1}M \xrightarrow{U^{-1}f} U^{-1}N \quad \frac{m}{u} \mapsto \frac{f(m)}{u}.$$

We already saw that $U^{-1}M \cong U^{-1}R \otimes_R M$. Since tensor product is right exact, localization is automatically right exact. The special feature of localization is that it is also left exact. The reason is the denominator-clearing argument: if a fraction maps to zero after localization, then some element of U already kills the relevant numerator.

Problem 14. Exactness of Localization.

Let $U \subset R$ be a multiplicatively closed set and let the sequence below be exact at M :

$$M' \xrightarrow{f} M \xrightarrow{g} M''.$$

1. Prove that

$$U^{-1}M' \xrightarrow{U^{-1}f} U^{-1}M \xrightarrow{U^{-1}g} U^{-1}M''$$

is a sequence of $U^{-1}R$ -modules whose consecutive composition is zero.

2. Prove that $\text{img}(U^{-1}f) \subseteq \ker(U^{-1}g)$.
3. Let $m/u \in U^{-1}M$ satisfy $(U^{-1}g)(m/u) = 0$. Unpack the definition of equality to zero in $U^{-1}M''$ to show that there exists $t \in U$ such that $g(tm) = 0$.
4. Use exactness of the original sequence at M to find $m' \in M'$ with $f(m') = tm$.
5. Prove that $\frac{m}{u} = \frac{f(m')}{tu}$ in $U^{-1}M$, and conclude that $m/u \in \text{img}(U^{-1}f)$.
6. Deduce that localization preserves exactness of arbitrary exact sequences.

Solution.

1. Let $\frac{m'}{u} \in U^{-1}M'$. Then the composition $U^{-1}g \circ U^{-1}f$ gives us,

$$U^{-1}g \circ U^{-1}f \left(\frac{m'}{u} \right) = \frac{g(f(m'))}{u} = \frac{0}{u}$$

since $g(f(m)) = 0$. Thus the consecutive composition is zero.

2. Let $\frac{m}{u} \in \text{img}(U^{-1}f)$. Then we know that $m \in \text{img}(f) = \ker(g)$. That is $g(m) = 0$ and thus $\frac{m}{u} \in \ker(U^{-1}g)$.

3. If $U^{-1}g \left(\frac{m}{u} \right) = \frac{g(m)}{u} = 0$. Then there exists $t \in U$ such that

$$t(g(m) \cdot 1 - 0 \cdot u) = t \cdot g(m) = g(tm) = 0$$

4. Since the original sequence is exact at M we have $\text{im}(f) = \ker(g)$, since $tm \in \ker(g)$ we must have some $m' \in M'$ such that $f(m') = tm$.

5. Then, $\frac{f(m')}{tu} = \frac{tm}{tu} = \frac{m}{u} \in U^{-1}M$, since $tu \in U$, we have $\frac{m}{u} \in \text{img}(U^{-1}f)$.

6. Since $\frac{m}{u}$ is an arbitrary element in $\ker(U^{-1}g)$, we have $\ker(U^{-1}g) \subseteq \text{img}(U^{-1}f)$. And part (2) gives us $\text{img}(U^{-1}f) \subseteq \ker(U^{-1}g)$, so $\ker(U^{-1}g) = \text{img}(U^{-1}f)$, and the sequence is exact at M . Thus localization preserves exactness of arbitrary exact sequences.

Problem 15. Localization Preserves Short Exact Sequences.

Let

$$0 \longrightarrow A \xrightarrow{i} B \xrightarrow{p} C \longrightarrow 0$$

be a short exact sequence and let $U \subset R$ be a multiplicatively closed set.

1. Prove directly that $U^{-1}i : U^{-1}A \rightarrow U^{-1}B$ is injective.
2. Prove directly that $U^{-1}p : U^{-1}B \rightarrow U^{-1}C$ is surjective.
3. Use the previous exercise to conclude that

$$0 \longrightarrow U^{-1}A \longrightarrow U^{-1}B \longrightarrow U^{-1}C \longrightarrow 0$$

is a short exact sequence of $U^{-1}R$ -modules.

4. Combine this with the isomorphism $U^{-1}M \cong U^{-1}R \otimes_R M$ to prove that $U^{-1}R$ is a flat R -module.

Solution.

1. Let $U^{-1}i\left(\frac{a}{u}\right) = \frac{i(a)}{u} = \left(\frac{0}{1}\right)$. Then there exists a $t \in U$ such that

$$t(i(a) \cdot 1 - u \cdot 0) = ti(a) = i(ta) = 0$$

Since i is injective, this implies $ta = 0$. But then, $t(a \cdot 1 - u \cdot 0) = ta = 0$, so $\frac{a}{u} \sim \frac{0}{1} \in U^{-1}A$. Hence $U^{-1}i$ is injective.

2. Since p is surjective, for every $c \in C$ there exists some $b \in B$ such that $p(b) = c$. Then, for some $\frac{c}{u} \in U^{-1}C$ we have, $U^{-1}p\left(\frac{b}{u}\right) = \frac{p(b)}{u} = \frac{c}{u}$. Thus $U^{-1}p$ is surjective.
3. Since $U^{-1}i$ is injective, $U^{-1}p$ is surjective, and $\ker(U^{-1}p) = \text{img}(U^{-1}i)$ by Problem 14, the sequence is exact at every term and hence is a short exact sequence of $U^{-1}R$ modules.
4. Since $U^{-1}M \cong U^{-1}R \otimes_R M$ we have the resulting exact sequence,

$$0 \longrightarrow U^{-1}R \otimes_R A \longrightarrow U^{-1}R \otimes_R B \longrightarrow U^{-1}R \otimes_R C \longrightarrow 0$$

Thus $U^{-1}R \otimes_R -$ preserves exact sequences and hence $U^{-1}R$ is a flat R -module.

Problem 16. Computing Localized Exact Sequences.

1. Let $I \subseteq R$ be an ideal. Use exactness of localization to prove that $U^{-1}(R/I) \cong U^{-1}R/IU^{-1}R$.
2. Localize the short exact sequence at $U = \{1, 2, 2^2, 2^3, \dots\}$. Identify each term explicitly

$$0 \longrightarrow \mathbb{Z} \xrightarrow{\cdot 6} \mathbb{Z} \longrightarrow \mathbb{Z}/6\mathbb{Z} \longrightarrow 0.$$

3. Localize the same sequence at $V = \mathbb{Z} \setminus \langle 2 \rangle$. Identify $V^{-1}(\mathbb{Z}/6\mathbb{Z})$.
4. Let $R = \mathbb{K}[x]$ and $M = R/\langle x^2(x-1) \rangle$. Compute $M[1/x]$ using the localized presentation

$$R[1/x] \xrightarrow{\cdot x^2(x-1)} R[1/x] \longrightarrow M[1/x] \longrightarrow 0$$

5. Explain why localizing $R/\langle x^2(x-1) \rangle$ at the prime ideal $\langle x \rangle$ “keeps” the $x = 0$ component and “removes” the $x = 1$ component.
6. With R and M as in the previous part, compute $M_{\langle x \rangle}$. Hint: What happens to $x - 1$ in $R_{\langle x \rangle}$?
7. Let $R = \mathbb{K}[x, y]$ and $M = R/\langle xy \rangle$. Compute $M[1/x]$ and $M[1/y]$.

Solution.

1. Using previous exercises, we get the exact sequence,

$$0 \longrightarrow U^{-1}I \xrightarrow{U^{-1}\iota} U^{-1}R \xrightarrow{U^{-1}\pi} U^{-1}(R/I) \longrightarrow 0$$

Then by the first isomorphism theorem we have,

$$U^{-1}(R/I) \cong U^{-1}R / \ker(U^{-1}\pi) = U^{-1}R / \text{img}(U^{-1}\iota) \cong U^{-1}R / U^{-1}I = U^{-1}R / IU^{-1}R$$

2. Localizing at U gives us,

$$0 \longrightarrow \mathbb{Z} \begin{bmatrix} 1 \\ 2 \end{bmatrix} \xrightarrow{\cdot 6} \mathbb{Z} \begin{bmatrix} 1 \\ 2 \end{bmatrix} \longrightarrow \mathbb{Z}/3\mathbb{Z} \longrightarrow 0.$$

Since $U^{-1}\mathbb{Z}$ gives us $\mathbb{Z} \left[\frac{1}{2} \right]$ and since $\bar{2}$ is a unit in $\mathbb{Z}/3\mathbb{Z}$,

$$\begin{aligned} U^{-1}(\mathbb{Z}/6\mathbb{Z}) &\cong U^{-1}\mathbb{Z}/(6U^{-1}\mathbb{Z}) \\ &= \mathbb{Z} \left[\frac{1}{2} \right] / \left(6\mathbb{Z} \left[\frac{1}{2} \right] \right) \\ &= \mathbb{Z} \left[\frac{1}{2} \right] / \left(3\mathbb{Z} \left[\frac{1}{2} \right] \right) \\ &\cong U^{-1}(\mathbb{Z}/3\mathbb{Z}) = \mathbb{Z}/3\mathbb{Z} \end{aligned}$$

3. Since $\frac{3}{1} \in U^{-1}(\mathbb{Z}/6\mathbb{Z})$ is a unit, we have $\frac{2}{1} = \frac{3 \cdot 2}{3} = \frac{0}{1} \in U^{-1}(\mathbb{Z}/6\mathbb{Z})$. Hence, $U^{-1}(\mathbb{Z}/6\mathbb{Z}) \cong \mathbb{Z}/2\mathbb{Z}$.
4. Note that in $R \left[\frac{1}{x} \right] = \mathbb{K}[x, x^{-1}]$, x is a unit. Then

$$\text{img}(\cdot x^2(x-1)) = \langle x^2(x-1) \rangle = \langle x-1 \rangle$$

So, $M \left[\frac{1}{x} \right] = R \left[\frac{1}{x} \right] / \langle x-1 \rangle = \mathbb{K}$.

5. Localizing at $\langle x \rangle$ means inverting $U = \mathbb{R} \setminus \langle x \rangle$, i.e. all polynomials that do not vanish at $x = 0$. Since $x-1 \neq 0$ when $x = 0$ it gets inverted killing the $x = 1$ component. But since $x \notin U$, we preserve the $x = 0$ component.
6. Since $\langle x-1 \rangle$ is now a unit we get $M_{\langle x \rangle} \cong R / \langle x^2 \rangle$.
7. We have $R \left[\frac{1}{x} \right] = \mathbb{K}[x, y, x^{-1}]$. In $M \left[\frac{1}{x} \right]$, x is a unit, so $\langle xy \rangle = \langle y \rangle$. Then $M \left[\frac{1}{x} \right] = \mathbb{K}[x, y, x^{-1}] / \langle y \rangle \cong \mathbb{K}[x, x^{-1}]$. Similarly, in $M \left[\frac{1}{y} \right]$, y is a unit, so $\langle xy \rangle = \langle x \rangle$, and $M \left[\frac{1}{y} \right] = \mathbb{K}[x, y, y^{-1}] / \langle x \rangle \cong \mathbb{K}[y, y^{-1}]$.