

Math 121: Commutative Algebra

Prishita Dharampal

Worksheet 4.2: Localization

Throughout this course, “ring” means *commutative* ring with unity. In the previous worksheet we built the basic language of modules, quotient modules, free modules, Hom, and tensor products. The goal of this worksheet is to introduce one of the central operations in commutative algebra: *localization*. Informally, localization is the process of forcing a chosen collection of elements of a ring to become invertible. This construction simultaneously generalizes the passage from $\mathbb{Z}$ to $\mathbb{Q}$, the localization $\mathbb{Z}_{\langle p \rangle}$ at a prime number p , and the algebraic operation of restricting attention to a principal open subset $D(f)$ of $\text{Spec}(R)$.

Let R be a ring. A subset $U \subset R$ is called *multiplicatively closed* if $1 \in U$ and $u, t \in U$ implies $ut \in U$. Given a multiplicative set $U \subset R$, the localization of R at U is a ring $U^{-1}R$ together with a ring homomorphism $\eta : R \rightarrow U^{-1}R$ satisfying the following universal property: For all ring homomorphisms $\phi : R \rightarrow T$ such that every element of $\phi(U)$ is a unit in T there exists a unique homomorphism $\tilde{\phi} : U^{-1}R \rightarrow T$ such that the following diagram commutes:

$$\begin{array}{ccc} R & \xrightarrow{\eta} & U^{-1}R \\ \phi \downarrow & \nearrow \tilde{\phi} & \\ T & & \end{array} .$$

Roughly speaking we should think of $U^{-1}R$ as being the “smallest” ring where every element of U is a unit. As always the universal property guarantees that $U^{-1}R$ and the map η are unique up to unique isomorphism (assuming that they exist, which we will now discuss).

Informally, we think of elements of $U^{-1}R$ as obtained by formally inverting every element of U and imposing the necessary equivalence conditions to make it into a well-behaved ring. Elements of $U^{-1}R$ are formal fractions r/u where $r \in R$ and $u \in U$ subject to some equivalence relation. If R is an integral domain and $0 \notin U$, equality of fractions is the familiar cross-multiplication relation, i.e. $r_1/u_1 = r_2/u_2$ as elements in $U^{-1}R$ if and only if $r_1u_2 = r_2u_1$ as elements in R . In general, because zero divisors may be present, one must allow extra clearing denominators:

$$\frac{r_1}{u_1} = \frac{r_2}{u_2} \text{ in } U^{-1}R \iff t(r_1u_2 - r_2u_1) = 0 \text{ in } R \text{ for some } t \in U$$

Multiplication and addition in $U^{-1}R$ are then defined in the usual way for fractions:

$$\left(\frac{r_1}{u_1}\right)\left(\frac{r_2}{u_2}\right) = \frac{r_1 r_2}{u_1 u_2} \quad \text{and} \quad \frac{r_1}{u_1} + \frac{r_2}{u_2} = \frac{r_1 u_2 + r_2 u_1}{u_1 u_2}$$

The additive identity is then the equivalence class of $0/1$ and the multiplicative identity is $1/1$. The function $\eta : R \rightarrow U^{-1}R$ is defined by $r \mapsto \frac{r}{1}$, which formally makes sense because $1 \in U$. Although this construction is relatively easy to define, there is work involved in checking that the operations and η are well-defined, that $U^{-1}R$ is a ring, and that it satisfies the correct universal property. You will carefully go through this construction in the exercises.

A few warnings: when R is not a domain localization can behave somewhat strangely. For example, from our experience with fractions since childhood and its unassuming definition one might expect η to be injective; this is not the case. The element $\eta(r) = r/1$ can be equal to zero in $U^{-1}R$. By definition this occurs when $r/1 = 0/1$, which translates to $tr = 0$ in R for some element $t \in U$. Even worse, if $0 \in U$ then the ring $U^{-1}R$ becomes the zero ring.

Problem 1. Multiplicative Sets.

1. Verify using the definition directly which of the following subsets are multiplicatively closed:

(a) $R = \mathbb{Z}$ and $U = \{1, 3, 9, 27, \dots\}$.	(f) $R = \mathbb{R}[x, y]$ and $U = \{f \in R \mid f(1, 0) \neq 0\}$.
(b) $R = \mathbb{Z}$ and $U = \{1, 2, 3\}$.	(g) $R = \mathbb{R}[x, y] / \langle y^2 - x^3 + x \rangle$ and $U = R \setminus \langle \bar{x}, \bar{y} \rangle$.
(c) $R = \mathbb{Z}$ and $U = \mathbb{Z} \setminus \langle 2 \rangle$.	(h) $R = \mathbb{R}[x, y]$ and $U = \{f \in R \mid f(0, 0) \neq 0 \text{ and } f(1, 1) \neq 0\}$.
(d) $R = \mathbb{Z}$ and $U = \{n \geq 1\}$.	
(e) $R = \mathbb{R}[x, y]$ and $U = R \setminus \langle x, y \rangle$.	
2. Let $I \subset R$ be an ideal. Prove that $U = R \setminus I$ is multiplicatively closed if and only if I is prime.
3. Let $U \subset R$ be a multiplicatively closed set. If $\phi : R \rightarrow S$ is a ring homomorphism, is $\phi(U) \subset S$ a multiplicatively closed subset?

Solution.

1. (a) U is multiplicatively closed.
 (b) U is not multiplicatively closed; $2, 3 \in U$ but $2 \cdot 3 \notin U$.

- (c) U is multiplicatively closed.
 - (d) U is multiplicatively closed.
 - (e) $U = \mathbb{R}$ and hence is multiplicatively closed.
 - (f) U is multiplicatively closed.
 - (g) U is multiplicatively closed.
 - (h) U is multiplicatively closed.
2. Let $I \subset R$ ideal.
- ($\Rightarrow$)
- Assume $U = R \setminus I$ is multiplicatively closed. Assume $a, b \notin I$. Then $a, b \in U$. Since U is multiplicatively closed, $ab \in U$, hence $ab \notin I$. That is if $a, b \notin I$ then $ab \notin I$. Thus, I is prime.
- ($\Leftarrow$)
- Assume I is prime. Then, we know that if $a, b \notin I$, $ab \notin I$. Thus, $a, b, ab \in U$, and U is multiplicatively closed.
3. Let $U \subset R$ be multiplicatively closed. Let $\phi : R \rightarrow S$ be a ring homomorphism. Then for any $r, s \in U$ we have,

$$\phi(r) \cdot \phi(s) = \phi(rs) \in \phi(U)$$

Hence, $\phi(U) \subset S$ is multiplicatively closed.

Problem 2. First Examples.

Despite not having formally verified that our construction is well-defined, discuss ways to explicitly understand $U^{-1}R$ in each of the following cases.

1. $R = \mathbb{Z}$ and $U = \mathbb{Z} \setminus \{0\}$.
2. $R = \mathbb{Z}$ and $U = \{2^n \mid n \in \mathbb{Z}_{\geq 0}\}$.
3. $R = \mathbb{Z}$ and $U = \mathbb{Z} \setminus \langle 2 \rangle$.
4. $R = \mathbb{K}[x]$ and $U = \mathbb{K}[x] \setminus \{0\}$.
5. $R = \mathbb{K}[x]$ and $U = \{1, x, x^2, x^3, \dots\}$.
6. $R = \mathbb{Z}/12\mathbb{Z}$ and $U = \{\bar{1}, \bar{2}, \bar{4}, \bar{8}\}$.
7. $R = \mathbb{Z}/12\mathbb{Z}$ and $U = \{\bar{1}, \bar{3}, \bar{9}\}$.
8. $R = \mathbb{Z}[x]/\langle nx - 1 \rangle$ and $U = \{n^k \mid k \in \mathbb{Z}_{\geq 0}\}$ for a fixed non-zero integer n .
9. R an arbitrary ring and U a multiplicative closed set containing a nilpotent.

Solution.

1. $U^{-1}R = \mathbb{Q}$
2. $U^{-1}R = \{m/2^n \mid m \in \mathbb{Z}\}$. These are the dyadic integers.
3. $U^{-1}R = \{m/n \mid m \in \mathbb{Z}, n \text{ odd}\}$
4. $U^{-1}R = \mathbb{K}(x)$
5. $U^{-1}R = \mathbb{K}[x, x^{-1}]$
6. $U^{-1}R = \mathbb{Z}/3\mathbb{Z}$
7. $U^{-1}R = \mathbb{Z}/4\mathbb{Z}$
8. $U^{-1}R = \mathbb{Z} \left[\frac{1}{n} \right] = R$
9. If $u \in U$ is nilpotent, then for some $k \geq 1$ $u^k = 0 \in U$. Then for any two arbitrary fractions $a/b, c/d \in U$, pick $w = 0$. Then we have

$$w(ad - cd) = 0(ad - cd) = 0$$

that is, all fractions are equivalent and $U^{-1}R = \{0\}$.

Problem 3. Constructing $U^{-1}R$.

Let R be a ring and $U \subset R$ a multiplicatively closed set. Define a relation on $R \times U$ by saying $(r_1, u_1) \sim (r_2, u_2)$ if and only if $t(r_1u_2 - r_2u_1) = 0$ in R for some $t \in U$.

1. Prove that $\sim$ is an equivalence relation on $R \times U$.
2. Let $\frac{r}{u}$ denote the equivalence class of a pair (r, u) under the equivalence relation $\sim$ defined in the previous part. Prove that the operations:

$$\left(\frac{r_1}{u_1}\right) \left(\frac{r_2}{u_2}\right) = \frac{r_1r_2}{u_1u_2} \quad \text{and} \quad \frac{r_1}{u_1} + \frac{r_2}{u_2} = \frac{r_1u_2 + r_2u_1}{u_1u_2}$$

are well-defined. Hint: You must show that changing the representative of either input fraction does not change the resulting equivalence class.

3. Define $U^{-1}R$ to be the set of equivalence classes of $\sim$. Show that the operations above make $U^{-1}R$ into a ring where $\frac{0}{1}$ is the zero element and $\frac{1}{1}$ is the multiplicative identity.
4. Show that the function $\eta : R \rightarrow U^{-1}R$ given by $r \mapsto \frac{r}{1}$ is a well-defined ring homomorphism.
5. Prove that if $u \in U$ then $\eta(u)$ is a unit in $U^{-1}R$. Explicitly what is its inverse?
6. Where in this construction did you use that U was multiplicatively closed and contains 1?

Solution.

1. Let $(r_1, u_1) \sim (r_2, u_2) \sim (r_3, u_3) \in R \times U$.

(a) **Reflexivity:** $(r_1, u_1) \sim (r_1, u_1)$ since $1(r_1u_1 - r_1u_1) = 0$.

(b) **Symmetry:** If $(r_1, u_1) \sim (r_2, u_2)$ then there exists a $t \in U$ such that

$$t(r_1u_2 - r_2u_1) = 0 \implies tr_1u_2 = tr_2u_1 \implies t(r_2u_1 - r_1u_2) = 0$$

Hence, $(r_2, u_2) \sim (r_1, u_1)$.

(c) **Transitivity:** If $(r_1, u_1) \sim (r_2, u_2)$ and $(r_2, u_2) \sim (r_3, u_3)$ then there exist s, t such that

$$s(r_1u_2 - r_2u_1) = 0, \quad t(r_2u_3 - r_3u_2) = 0$$

Note that

$$stu_3(r_1u_2 - r_2u_1) = 0, \quad stu_1(r_2u_3 - r_3u_2) = 0$$

Adding,

$$0 = st(r_1u_2u_3 - r_2u_1u_3 + r_2u_3u_1 - r_3u_2u_1) = stu_2(r_1u_3 - r_3u_1)$$

Since $stu_2 \in U$ we have $(r_1, u_1) \sim (r_3, u_3)$.

Thus, $\sim$ is an equivalence relation on $R \times U$.

2. Let $(r, u) \sim (r', u')$ and $(s, t) \sim (s', t')$. Then there exist $p, q \in U$ such that

$$p(ru' - r'u) = 0, \quad q(st' - s't) = 0$$

Multiplying,

$$pqst'(ru' - r'u) = 0, \quad pqr'u(st' - s't) = 0$$

Adding,

$$pq(st'ru' - st'r'u + r'ust' - r'us't) = pq(rsu't' - r's'ut) = 0$$

That is,

$$\left(\frac{r}{u}\right) \left(\frac{s}{t}\right) = \frac{rs}{ut} \sim \frac{r's'}{u't'} = \left(\frac{r'}{u'}\right) \left(\frac{s'}{t'}\right)$$

TODO:

3. Note that associativity, commutativity, and distributivity are inherited from R .

(a) **Additive Identity:**

$$\frac{r}{u} + \frac{0}{1} = \frac{r+0}{u} = \frac{r}{u}$$

(b) **Additive Inverses:**

$$\frac{r}{u} + \frac{-r}{u} = \frac{ru - ru}{u} = \frac{0}{1}$$

(c) **Multiplicative Identity:**

$$\frac{r}{u} \cdot \frac{1}{1} = \frac{r \cdot 1}{u \cdot 1} = \frac{r}{u}$$

Hence, $U^{-1}R$ is a ring.

4. Let $\eta : R \rightarrow U^{-1}R$, where $r \mapsto \frac{r}{1}$.

- $\eta(r) + \eta(s) = \frac{r}{1} + \frac{s}{1} = \frac{r+s}{1} = \eta(r+s)$
- $\eta(r) \cdot \eta(s) = \frac{r}{1} \cdot \frac{s}{1} = \frac{rs}{1} = \eta(rs)$

Hence, η is a ring homomorphism.

5. For any $u \in U$ we have,

$$\eta(u) \cdot \frac{1}{u} = \frac{u}{1} \cdot \frac{1}{u} = \frac{u}{u} \sim \frac{1}{1}$$

6. In proving transitivity and verifying that the operations are well-defined.

Problem 4. First Properties of Localization.

Let R be a ring and $U \subset R$ a multiplicatively closed set.

1. Prove that if $0 \in U$ then $U^{-1}R$ is the zero ring.
2. Prove that if R is an integral domain and $0 \notin U$ then $\frac{r_1}{u_1} = \frac{r_2}{u_2}$ in $U^{-1}R$ if and only if $r_1u_2 = r_2u_1$.
3. Let R be an integral domain and $U = R \setminus \{0\}$. Prove that $U^{-1}R$ is a field.
4. Find an example of a non-zero ring R for which $R \setminus \{0\}$ is not multiplicatively closed. What goes wrong with trying to form $(R \setminus \{0\})^{-1}R$?

Solution.

1. If $0 \in U$. Then for any two arbitrary fractions $a/b, c/d \in U$, pick $w = 0$. And we have

$$w(ad - cd) = 0(ad - cd) = 0$$

that is, all fractions are equivalent and $U^{-1}R = \{0\}$.

2. Assume R be an integral domain, and $0 \notin U$.

($\implies$)

Assume $\frac{r_1}{u_1} = \frac{r_2}{u_2}$ in $U^{-1}R$. Then there exists a $t \neq 0 \in U$ such that

$$t(r_1u_2 - u_1r_2) = 0$$

Since R is an integral domain this implies that

$$r_1u_2 - u_1r_2 = 0 \implies r_1u_2 = u_1r_2$$

($\impliedby$)

Assume $r_1u_2 = r_2u_1$. Then with $w = 1 \in U$,

$$r_1u_2 - r_2u_1 = 0 \implies \frac{r_1}{u_1} = \frac{r_2}{u_2}$$

Hence proved.

3. Let $\frac{r}{u} \approx \frac{0}{1} \in U^{-1}R$. From (2) we know that $r \neq 0$. Then $\frac{u}{r} \in U^{-1}R$ and hence, $U^{-1}R$ is a field.
4. Let $R = \mathbb{Z}/6\mathbb{Z}$. Then, we have $\frac{1}{2}, \frac{1}{3} \approx \frac{0}{1} \in (R \setminus \{0\})^{-1}R$. But $\frac{1}{2} \cdot \frac{1}{3} = \frac{1}{6} \sim \frac{1}{0}$ which is not defined.

Problem 5. The Kernel of the Localization Map.

Let R be a ring, $U \subset R$ be a multiplicatively closed set, and $\eta : R \rightarrow U^{-1}R$ the localization map as described in Problem 3.

1. Prove that η is injective if and only if every $u \in U$ is a nonzerodivisor on R , i.e. $ur = 0$ implies $r = 0$ for all $r \in R$.
2. Let $I = \{r \in R \mid ur = 0 \text{ for some } u \in U\}$. Prove that I is an ideal.
3. Show that there exists an injective ring homomorphism $R/I \rightarrow U^{-1}R$ such that the following diagram commutes:

$$\begin{array}{ccc} R & \xrightarrow{\pi} & R/I \\ & \searrow \eta & \downarrow \\ & & U^{-1}R \end{array} .$$

4. Prove that $\ker(\eta) = I$.
5. Revisit the examples in Problem 2. For which of them is the localization map $R \rightarrow U^{-1}R$ injective?

Solution.

1. ($\implies$)

Assume η is injective. Suppose $ur = 0$ for some $u \in U$. Then,

$$\eta(r) = \frac{r}{1} = \frac{ur}{u} = \frac{0}{u} \sim \frac{0}{1}$$

Since $\ker(\eta) = \{0\}$, this implies that either $r = 0$.

(We cannot have $u = 0$ since that would make $U^{-1}R$ into the zero ring, and η is injective.) Hence, U contains no zero-divisors on R .

($\impliedby$)

Assume U has no zero-divisors in R . An element r is in $\ker(\eta)$ iff $\eta(r) \sim \frac{0}{1}$. That is there exists $u \in U$ such that,

$$u(r - 0) = 0$$

But by assumption, u is not a zero-divisor and hence r must be 0. Since r was an arbitrary element, this implies that $\ker(\eta) = \{0\}$ and η is injective.

2. Since $I \subseteq R$ we only need to show that I is closed under addition and multiplication (from R).

(a) **Addition:** Let $r, s \in I$. Then there exist u, v such that $ru = 0, sv = 0$. Since U is multiplicatively closed, $uv \in U$. Then we have,

$$(r + s)uv = (ru)v + (sv)u = 0$$

Hence, $r + s \in I$.

(b) Let $r \in R$ and $s \in I$. Then there exist some $u \in U$ such that $su = 0$. Consider rs ,

$$(rs)u = r(su) = 0$$

Hence, $rs \in I$.

Thus I is an ideal.

3. Define $\phi : R/I \rightarrow U^{-1}R$, where $\bar{r} \mapsto \frac{r}{1}$. Now we check for the following:

(a) **Well-Definedness:** Let $\bar{r} = \bar{s}$. Then, $\bar{r} - \bar{s} \in I \implies \exists u \in U$ such that

$$u(r - s) = 0 \implies u(r \cdot 1 - s \cdot 1) \implies \frac{r}{1} \sim \frac{s}{1}$$

(b) **Ring Homomorphism:**

- $\phi(\bar{r} + \bar{s}) = \frac{r+s}{1} = \frac{r}{1} + \frac{s}{1} = \phi(\bar{r}) + \phi(\bar{s})$
- $\phi(\bar{r} \cdot \bar{s}) = \frac{r \cdot s}{1} = \frac{r}{1} \cdot \frac{s}{1} = \phi(\bar{r}) \cdot \phi(\bar{s})$

(c) **Injectivity:** Let $\bar{r} \in \ker(\phi)$. Then $\phi(\bar{r}) = \frac{r}{1} \sim \frac{0}{1}$. That is, there exists $u \in U$ such that $u(r) = 0$. Then $\bar{r} \in I$ and hence $\bar{r} = \bar{0} \in R/I$.

(d) **Commutates:** Let $r \in R$. Then we have,

$$\eta(r) = \frac{r}{1}, \quad \phi(\pi(r)) = \phi(\bar{r}) = \frac{r}{1}$$

Hence, there exists an injective ring homomorphism $R/I \rightarrow U^{-1}R$

4. $\ker(\eta) = \ker(\phi \circ \pi) = \pi^{-1}(\ker(\phi)) = \pi^{-1}(\langle 0 \rangle) = I$.

5. η is injective for subparts 1, 2, 3, 4, 5, 8.

Problem 6. Field of Fractions.

Let R be an integral domain. The *field of fractions* of R is the ring $U^{-1}R$ where $U = R \setminus \{0\}$. We denote the field of fractions of R by $\text{Frac}(R)$.

1. Show that $\text{Frac}(R)$ is a field containing R .
2. Describe $\text{Frac}(R)$ when $R = \mathbb{Z}$ and $R = \mathbb{K}[x]$.
3. If R is an integral domain and $V \subset R$ is any multiplicatively closed set not containing zero, explain how we can view $V^{-1}R$ as a subring of $\text{Frac}(R)$.

Solution.

1. From Question 4.3 we know that $\text{Frac}(R)$ is a field, and contains R since $\eta(r) = \frac{r}{1} \in \text{Frac}(R)$ for all $r \in R$.
2. $\text{Frac}(\mathbb{Z}) = \mathbb{Q}$, and $\text{Frac}(\mathbb{K}[x]) = \mathbb{K}(x)$.
3. Note that $V \subseteq U$. Then every element $a = \frac{r}{v} \in V^{-1}R$, is also an element of $\text{Frac}(R)$. Then we can define an embedding $\iota : V^{-1}R \rightarrow U^{-1}R$ where $\iota\left(\frac{r}{v}\right) = \left(\frac{r}{v}\right)$, showing that $V^{-1}R$ is a subring of $U^{-1}R = \text{Frac}(R)$.

Problem 7. Universal Property of Localization.

In this exercise we shall show that $U^{-1}R$ and η as constructed in Problem 3 satisfy the universal property of localization. Let R be a ring and $U \subset R$ a multiplicatively closed subset. Let $\phi : R \rightarrow T$ be a ring homomorphism such that $\phi(U) \subset T^\times$.

1. Define a function $\tilde{\phi} : U^{-1}R \rightarrow T$ by $\frac{r}{u} \mapsto \phi(r)\phi(u)^{-1}$. Explain why $\tilde{\phi}$ is well-defined and prove it is a ring homomorphism.
2. Show that $\tilde{\phi}$ makes the following diagram commute:

$$\begin{array}{ccc} R & \xrightarrow{\eta} & U^{-1}R \\ \phi \downarrow & \swarrow \tilde{\phi} & \\ T & & \end{array} .$$

3. Prove that $\tilde{\phi}$ is the unique ring homomorphism $U^{-1}R \rightarrow T$ making the diagram commute. Hint: If $\psi : U^{-1}R \rightarrow T$ is any such map, first determine $\psi\left(\frac{1}{u}\right)$ for $u \in U$, and then use the equality $\frac{r}{u} = \frac{r}{1} \frac{1}{u}$.

Solution.

1. Let $\frac{r}{u} \sim \frac{s}{v} \in U^{-1}R$, we want to show that $\phi(r)\phi(u)^{-1} = \phi(s)\phi(v)^{-1}$. Then there exists some $t \in U$ such that $t(rv - su) = 0$. Applying ϕ ,

$$\phi(t)(\phi(r)\phi(v) - \phi(s)\phi(u)) = \phi(0) = 0$$

Since $\phi(t) \in \phi(U) \subseteq T^\times$ we can multiply both sides by $\phi(t)^{-1}$,

$$\phi(r)\phi(v) = \phi(s)\phi(u) \implies \phi(r)\phi(u)^{-1} = \phi(s)\phi(v)^{-1}$$

Hence $\tilde{\phi}$ is well-defined.

Ring homomorphism:

• **Addition:**

$$\begin{aligned} \tilde{\phi}\left(\frac{r}{u} + \frac{s}{v}\right) &= \tilde{\phi}\left(\frac{rv + su}{uv}\right) \\ &= \phi(rv + su)\phi(uv)^{-1} \\ &= \phi(r)\phi(v)\phi(u)^{-1}\phi(v)^{-1} + \phi(s)\phi(u)\phi(u)^{-1}\phi(v)^{-1} \\ &= \phi(r)\phi(u)^{-1} + \phi(s)\phi(v)^{-1} \\ &= \tilde{\phi}\left(\frac{r}{u}\right) + \tilde{\phi}\left(\frac{s}{v}\right) \end{aligned}$$

• **Multiplication:**

$$\begin{aligned}
\tilde{\phi}\left(\frac{r}{u} \cdot \frac{s}{v}\right) &= \tilde{\phi}\left(\frac{rs}{uv}\right) \\
&= \phi(rs)\phi(uv)^{-1} \\
&= \phi(r)\phi(s)\phi(u)^{-1}\phi(v)^{-1} \\
&= \phi(r)\phi(u)^{-1} \cdot \phi(s)\phi(v)^{-1} \\
&= \tilde{\phi}\left(\frac{r}{u}\right) \cdot \tilde{\phi}\left(\frac{s}{v}\right)
\end{aligned}$$

Hence $\tilde{\phi}$ is a well-defined ring homomorphism.

2. Let $r \in R$. Then we have,

$$\tilde{\phi}(\eta(r)) = \tilde{\phi}\left(\frac{r}{1}\right) = \phi(r)\phi(1)^{-1} = \phi(r)$$

3. Let $\psi : U^{-1}R \rightarrow T$ be a ring homomorphism that commutes with the given diagram. Then since $\phi(u) = \psi(\eta(u)) = \psi\left(\frac{u}{1}\right)$, and $\psi(u)\psi(u^{-1}) = 1$ we must have $\psi\left(\frac{u}{1}\right) \cdot \psi\left(\frac{1}{u}\right) = \phi(u) \cdot \phi(u)^{-1}$.

That is for an arbitrary element $\frac{r}{u} \in U^{-1}R$ we have,

$$\psi\left(\frac{r}{u}\right) = \psi\left(\frac{r}{1} \cdot \frac{1}{u}\right) = \psi\left(\frac{r}{1}\right) \cdot \psi\left(\frac{1}{u}\right) = \phi(r)\phi(u)^{-1} = \tilde{\phi}\left(\frac{r}{u}\right)$$

Problem 8. Applications of the Universal Property of Localization.

Let R be a ring and $U \subset R$ a multiplicatively closed subset.

1. Use the universal property of localization to show that $U^{-1}R$ and η are unique up to unique isomorphism.
2. Suppose every element of U is already a unit in R . Prove that the localization map $\eta : R \rightarrow U^{-1}R$ is an isomorphism.
3. Suppose $V \subset R$ is another multiplicatively closed set and $U \subset V$. Use the universal property to construct a natural homomorphism $U^{-1}R \rightarrow V^{-1}R$.
4. Let $\phi : R \rightarrow S$ be a ring homomorphism such that $\phi(U) \subset S^\times$. Let $\tilde{\phi} : U^{-1}R \rightarrow S$ be the induced map. Prove that if ϕ is surjective, then $\tilde{\phi}$ is surjective. Prove that if ϕ is injective, then $\tilde{\phi}$ is injective.

Solution.

1. **TODO:**

2. If every element of U is a unit in R then there exist natural maps $\eta : R \rightarrow U^{-1}R$ and $\tilde{\phi} : U^{-1}R \rightarrow R$. We then have the following diagram,

$$\begin{array}{ccc} R & \xrightarrow{\eta} & U^{-1}R \\ \phi \downarrow & \swarrow \tilde{\phi} & \\ R & & \end{array} .$$

This forces $\phi = \text{id}_R$ hence showing that the composition $\tilde{\phi} \circ \eta = \text{id}_R$. Now consider the maps $\eta \circ \tilde{\phi}, \text{id}_{U^{-1}R} : U^{-1}R \rightarrow U^{-1}R$. They agree on composition with η , and hence by uniqueness of the map from $U^{-1}R \rightarrow T$, they must be the same map.

$$(\eta \circ \tilde{\phi}) \circ \eta = \eta \circ (\tilde{\phi} \circ \eta) = \eta \circ \text{id}_R = \eta = \text{id}_{U^{-1}R} \circ \eta$$

Hence $R \cong U^{-1}R$.

3. If $U \subset V$, then every element in U is a unit in $V^{-1}R$, then by the universal property, there exists a map $\tilde{\phi} : U^{-1}R \rightarrow V^{-1}R$.
4. Assume ϕ is surjective. Then for all $s \in S$ there exists some $r \in R$ such that $\phi(r) = s$. Then,

$$\tilde{\phi}(\eta(r)) = \tilde{\phi}\left(\frac{r}{1}\right) = \phi(r) = s$$

Thus $\tilde{\phi}$ must be surjective.

Assume ϕ is injective. Suppose $\tilde{\phi}(\frac{r}{1}) = 0$, then $\tilde{\phi}(\frac{r}{1}) = \phi(r)\phi(1)^{-1} = \phi(r) = 0$. But since ϕ is injective, this implies that $r = 0$. Thus, $\tilde{\phi}$ is injective.

The two most important examples of localization are: i) when U is equal to the complement of a prime ideal and ii) when $U = \{1, f, f^2, f^3, \dots\}$ for some non-zero element $f \in R$. Given a prime ideal $\mathfrak{p} \subset R$ you showed in Question 2 that $U = R \setminus \mathfrak{p}$ is a multiplicatively closed set. The ring $(R \setminus \mathfrak{p})^{-1}R$ occurs so frequently that we give it its own notation and name. The *localization of R at $\mathfrak{p}$* is the ring:

$$R_{\mathfrak{p}} := (R \setminus \mathfrak{p})^{-1}R.$$

The guiding feature of $R_{\mathfrak{p}}$ is that all elements outside of $\mathfrak{p}$ become invertible, while the elements in $\mathfrak{p}$ remain non-units. As we shall see shortly, this implies that $R_{\mathfrak{p}}$ has a unique maximal ideal, namely,

$$\mathfrak{p} R_{\mathfrak{p}} := \left\{ \frac{r}{u} \mid r \in \mathfrak{p}, u \in R \setminus \mathfrak{p} \right\}.$$

A ring with a unique maximal ideal is called a *local ring*. We often denote a local ring by $(R, \mathfrak{m})$, where $\mathfrak{m}$ is its unique maximal ideal. Local rings are algebraic analogues of rings of germs near a point.

Turning towards the second important example, fix a non-zero element $f \in R$ and consider the set $U = \{1, f, f^2, f^3, \dots\}$. Note we should assume f is not a nilpotent element, since otherwise as we saw in Question 2, the localization $U^{-1}R$ will be the zero ring. The localization $U^{-1}R$ consists of fractions of the form $\frac{r}{f^n}$ with $r \in R$ and $n \geq 0$. Every such fraction can be rewritten as $r \cdot \frac{1}{f^n} = r \cdot \left(\frac{1}{f}\right)^n$, so every element of $U^{-1}R$ is a polynomial expression in $\frac{1}{f}$ with coefficients in R . In other words, $U^{-1}R$ is generated, as a ring, by the image of R and the single new inverse $\frac{1}{f}$. We therefore write:

$$R\left[\frac{1}{f}\right] := \left\{1, f, f^2, \dots\right\}^{-1}R = \left\{\frac{r}{f^n} \mid r \in R, n \geq 0\right\}.$$

We will see shortly that we can make the idea that $R[\frac{1}{f}]$ is “polynomials in $\frac{1}{f}$ with coefficients in R ” precise by showing $R[\frac{1}{f}]$ is isomorphic to $R[x]/\langle xf - 1 \rangle$. Intuitively this makes sense since quotienting by $\langle xf - 1 \rangle$ means setting $xf = 1$, i.e., let the formal variable x equal the inverse of f . (Note we saw something similar to this when we used the Rabinowitsch trick on the Nullstellensatz worksheet.)

Problem 9. Localization at a Prime Ideal.

Let R be a ring and $\mathfrak{p} \subset R$ be a prime ideal.

1. Prove that an element $\frac{r}{u} \in R_{\mathfrak{p}}$ is a unit if and only if $r \notin \mathfrak{p}$.
2. Deduce that the non-units of $R_{\mathfrak{p}}$ are precisely the elements

$$\mathfrak{p} R_{\mathfrak{p}} := \left\{ \frac{r}{u} \mid r \in \mathfrak{p}, u \in R \setminus \mathfrak{p} \right\}.$$

3. Prove that $\mathfrak{p} R_{\mathfrak{p}}$ is an ideal in $R_{\mathfrak{p}}$.
4. Show that $\mathfrak{p} R_{\mathfrak{p}}$ is a maximal ideal in $R_{\mathfrak{p}}$. Hint: Show that any ideal strictly containing $\mathfrak{p} R_{\mathfrak{p}}$ must contain a unit.
5. Show that $R_{\mathfrak{p}}$ does not have any other maximal ideal, i.e. $(R_{\mathfrak{p}}, \mathfrak{p} R_{\mathfrak{p}})$ is a local ring.
6. More generally, show that a ring R is a local ring if and only if the set of non-units in R is an ideal.
7. Let $p \in \mathbb{Z}$ be a prime number. Describe the units and unique maximal ideal of $\mathbb{Z}_{\langle p \rangle}$.
8. Let $R = \mathcal{C}[x]$ and $\mathfrak{m} = \langle x - a \rangle$ for $a \in \mathcal{C}$. Which rational functions in $\mathcal{C}(x)$ lie in $R_{\mathfrak{m}}$? Which of them are units?

Solution.

1. ($\implies$)

Assume $\frac{r}{u} \in R_{\mathfrak{p}}$ is a unit. Assume for contradiction that $r \in \mathfrak{p}$, and let $\frac{s}{v}$ be the inverse of $\frac{r}{u}$. Then $\exists t \in U$ such that

$$t(rs - vu) = 0 \implies trs = tvu \implies tvu \in \mathfrak{p}$$

But $t, v, u \notin \mathfrak{p}$ by assumption and since $\mathfrak{p}$ is prime $tvu \notin \mathfrak{p}$, thus a contradiction.

Hence $r \notin \mathfrak{p}$.

($\impliedby$)

Assume $r \notin \mathfrak{p}$. Then $r \in (R \setminus \mathfrak{p})$. That is there exists $\frac{u}{r} \in R_{\mathfrak{p}}$ and thus $\frac{r}{u}$ is a unit.

2. Then we can conclude that the non-units of $R_{\mathfrak{p}}$ are the precisely elements $r \in \mathfrak{p}$, which is the definition of $\mathfrak{p} R_{\mathfrak{p}}$.

3. To show that $\mathfrak{p} R_{\mathfrak{p}}$ is an ideal we need to show that it is closed under addition and closed under multiplication from $R_{\mathfrak{p}}$. Let $\frac{r}{u}, \frac{s}{u} \in \mathfrak{p} R_{\mathfrak{p}}$, and $\frac{a}{v} \in R_{\mathfrak{p}}$, with $u, v \in R \setminus \mathfrak{p}$.

- $\frac{r}{u} + \frac{s}{u} = \frac{r+s}{u} \in \mathfrak{p} R_{\mathfrak{p}}$ since $\mathfrak{p}$ is closed under addition.
- $\frac{r}{u} \cdot \frac{a}{v} = \frac{ra}{uv} \in \mathfrak{p} R_{\mathfrak{p}}$ since both $R \setminus \mathfrak{p}$ and $\mathfrak{p}$ are closed under multiplication.

4. Assume for contradiction that there exists a maximal ideal M such that $\mathfrak{p} R_{\mathfrak{p}} \subset M \subset R_{\mathfrak{p}}$. Then there exists some element $\frac{a}{u} \in M$ that is not in $\mathfrak{p} R_{\mathfrak{p}}$. But from part (2) we know that if $\frac{a}{u} \in R_{\mathfrak{p}}$ but not in $\mathfrak{p} R_{\mathfrak{p}}$ it is a unit. So there exists some element $\frac{b}{v} \in R_{\mathfrak{p}}$ such that $\frac{a}{u} \cdot \frac{b}{v} = 1 \in M$ contradicting our assumption that M is a proper ideal.

Hence, $\mathfrak{p} R_{\mathfrak{p}}$ is maximal.

5. From part (2) we know that all elements not in $\mathfrak{p} R_{\mathfrak{p}}$ are units. Then by the argument in (4) is any ideal $I \subset R_{\mathfrak{p}}$ such that $I \neq \mathfrak{p} R_{\mathfrak{p}}$ contains 1 and hence is the whole ring. Hence, $(R_{\mathfrak{p}}, \mathfrak{p} R_{\mathfrak{p}})$ is a local ring.

6. So far we have shown that if the set of non-units is an ideal then R is a local ring, so we only need to show the opposite direction.

Assume R is a local ring with maximal ideal M . Let J be the set of non-units in R . The containment $M \subseteq J$ is obvious (because if not, we have $1 \in M$). To show the reverse containment, let $j \in J$. Consider the ideal $\langle j \rangle$. Then there must exist some maximal ideal in R containing $\langle j \rangle$. Since R is a local ring we must have $\langle j \rangle \subseteq M$. Since j was an arbitrary element in J , this implies that $J \subseteq M$. Thus, the set of non-units in R is an ideal.

7. The maximal ideal (and the set of units) of $\mathbb{Z}_{\langle p \rangle}$ is $\mathfrak{p} \mathbb{Z}_{\langle p \rangle} := \left\{ \frac{q}{u} \mid q \in \langle p \rangle, u \in \mathbb{Z} \setminus \langle p \rangle \right\}$.

8. All rational functions in $\mathbb{C}(x)$ that don't have f in the denominator lie in $R_{\mathfrak{m}}$ where $f \in \langle x - a \rangle$. The ones that don't have f in the numerator are units.

Problem 10. Inverting an Element.

Let R be a ring and let $f \in R$ be a nonzero, non-nilpotent element. Recall the ring $R\left[\frac{1}{f}\right]$ is defined to be the localization of R at $U = \{1, f, f^2, f^3, \dots\}$.

1. Show that the R -algebra homomorphism $\phi : R[x] \rightarrow R\left[\frac{1}{f}\right]$ determined by $\phi(r) = \frac{r}{1}$ for $r \in R$ and $\phi(x) = \frac{1}{f}$ is well-defined.
2. Prove that ϕ is surjective by explicitly finding an element that maps to $\frac{r}{f^k}$ for $r \in R$ and $k \in \mathbb{Z}_{\geq 0}$.
3. Show that $\ker(\phi) = \langle fx - 1 \rangle$.
4. Conclude that $R\left[\frac{1}{f}\right]$ is isomorphic to $R[x] / \langle fx - 1 \rangle$.
5. Give a different argument proving $R\left[\frac{1}{f}\right]$ and $R[x] / \langle fx - 1 \rangle$ are isomorphic using the universal property of localizations.
6. What changes if f is nilpotent? Show that both $R\left[\frac{1}{f}\right]$ and $R[x] / \langle fx - 1 \rangle$ are the zero ring.
7. Let $f, g \in R$. Use the universal property to prove that

$$\left(R\left[\frac{1}{f}\right]\right)\left[\frac{1}{g}\right] \cong R\left[\frac{1}{fg}\right].$$

Solution.

1. **TODO:**

2. For any element $\frac{r}{f^k} \in R\left[\frac{1}{f}\right]$ we have

$$\phi(rx^k) = \phi(r) \cdot \phi(x^k) = \frac{r}{1} \cdot \frac{1}{f^k} = \frac{r}{f^k}$$

Thus ϕ is surjective.

3. $\ker(\phi) \subseteq \langle fx - 1 \rangle$

Assume $a(x) \in \ker(\phi)$ non-zero polynomial. Then $a(x) = q(x)(fx - 1) + r(x)$ for

some $q(x), r(x) \in R[x]$. We know that $\phi(a(x)) = 0 = \phi(q(x)(fx - 1) + r(x))$. Then,

$$\begin{aligned}\phi(q(x)(fx - 1) + r(x)) &= 0 \\ \phi(q(x))(\phi(fx) - \phi(1)) + \phi(r(x)) &= 0 \\ \phi(q(x))\left(\frac{f}{1} \cdot \frac{1}{f} - 1\right) + \phi(r(x)) &= 0 \\ 0 + \phi(r(x)) &= 0\end{aligned}$$

Thus $\phi(r(x)) = 0$. But since we were dividing by a degree one polynomial, $\deg(r(x)) = 0$, i.e. r is a constant, that forces $r = 0$. Hence, $a(x) \in \langle fx - 1 \rangle$.

$$\langle fx - 1 \rangle \subseteq \ker(\phi)$$

Let $a \in \langle fx - 1 \rangle$. Then $a = g(fx - 1)$ for some $g \in R$. Consider $\phi(a)$.

$$\phi(a) = \phi(g(fx - 1)) = \phi(gfx) - \phi(g) = \frac{gf}{f} - \frac{g}{1} = 0$$

Since a was an arbitrary element, we have $\langle fx - 1 \rangle \subseteq \ker(\phi)$.

4. Then by the first isomorphism theorem we have,

$$R \left[\frac{1}{f} \right] \cong R[x] / \ker(\phi) = R[x] / \langle fx - 1 \rangle$$

5. We have $U = \{1, f, f^2, \dots\}$, and $\tilde{\phi}$ exists because $\phi(U) \subset (R[x] / \langle xf - 1 \rangle)^\times$ since f maps to a unit in that ring ($x \mapsto \frac{1}{f}$).

$$\begin{array}{ccc} R & \xrightarrow{\eta} & U^{-1}R[x] = R \left[\frac{1}{f} \right] \\ \phi \downarrow & \nwarrow \tilde{\phi} & \\ R[x] / \langle xf - 1 \rangle & & \end{array} .$$

Now, since $\langle xf - 1 \rangle = \ker(\eta)$ by the universal property of quotients we have the map $\tilde{\eta}$,

$$\begin{array}{ccc} R & \xrightarrow{\pi} & R[x] / \langle fx - 1 \rangle \\ \eta \downarrow & \nwarrow \tilde{\eta} & \\ R \left[\frac{1}{f} \right] & & \end{array}$$

Consider the following compositions:

- $\tilde{\eta} \circ \tilde{\phi} \left(\frac{r}{f^k} \right) = \tilde{\eta}(rx^k) = \frac{r}{f^k}$
- $\tilde{\phi} \circ \tilde{\eta}(rx^k) = \tilde{\phi} \left(\frac{r}{f^k} \right) = rx^k.$

Thus $R \left[\frac{1}{f} \right] \cong R[x] / \langle fx - 1 \rangle.$

6. If f is nilpotent, from Question 4(1) we know that $R \left[\frac{1}{f} \right] = \{0\}.$ By part (4), this implies that $R[x] / \langle fx - 1 \rangle = \{0\}.$

7. First consider the composition $R \rightarrow R[1/f] \rightarrow (R[1/f])[1/g].$

In $(R[1/f])[1/g],$ both f and g are units, so fg is a unit. Then by the universal property of localization (on $R[1/fg],$), there exists a unique map $\tilde{\phi} : R[1/fg] \rightarrow (R[1/f])[1/g].$

Now, note that in $R[1/fg], fg$ is a unit, so both f and g are units

$$f \cdot \frac{g}{fg} = 1, \quad g \cdot \frac{f}{fg} = 1$$

Then by the universal property of localization (on $(R[1/f])[1/g],$), there exists a unique map $\tilde{\psi} : (R[1/f])[1/g] \rightarrow R[1/fg].$ Both compositions $\tilde{\phi} \circ \tilde{\psi}$ and $\tilde{\psi} \circ \tilde{\phi}$ are the identity by uniqueness in the universal property, hence:

$$\left(R \left[\frac{1}{f} \right] \right) \left[\frac{1}{g} \right] \cong R \left[\frac{1}{fg} \right]$$

Given any ring homomorphism $\phi : R \rightarrow S$, there is a natural interplay between ideals of R and ideals of S . If $I \subset R$ is an ideal, its *extension* is the ideal of S generated by the image of I under ϕ :

$$\phi(I) \cdot S := \langle \phi(I) \rangle = \left\{ \sum_{i=1}^n s_i \phi(r_i) \mid r_i \in I, s_i \in S, n \geq 0 \right\}.$$

Note we must take the ideal generated by $\phi(I)$, not just the image of I itself, since as we previously saw $\phi(I)$ may not be an ideal of S .

Conversely, if $J \subseteq S$ is an ideal, its *contraction* is the preimage

$$\phi^{-1}(J) = \{r \in R \mid \phi(r) \in J\},$$

which is always an ideal of R . Note there are numerous different notations for extension and contraction of ideals: some (namely Atiyah–MacDonald) write I^e and J^c for the extension and contraction respectively, others write IS and $J \cap R$ (as a nod to what occurs when ϕ is an injection). We generally avoid these notations to emphasize that contraction and extension are dependent on the homomorphism ϕ .

Problem 11. Extension and Contraction of Ideals.

Let $\phi : R \rightarrow S$ be a homomorphism of rings. Let $I \subset R$ and $J \subset S$ be ideals.

1. Prove that $\phi^{-1}(J)$ is an ideal of R .
2. Give an explicit example showing that $\phi(I) \subset S$ need not be an ideal.
3. Prove that $I \subset \phi^{-1}(\phi(I)S)$.
4. Let $\phi : \mathbb{Z} \rightarrow \mathbb{Z}/\langle 6 \rangle$ be the natural quotient map and let $I = \langle 0 \rangle \subset \mathbb{Z}$. Explicitly describe $J = \phi(I)(\mathbb{Z}/\langle 6 \rangle)$ and $\phi^{-1}(J)$. Show that $I \subsetneq \phi^{-1}(J)$. Thus, the inclusion in the previous part can be strict.
5. Prove that $\phi(\phi^{-1}(J))S \subset J$.
6. Let $\phi : \mathbb{Z} \rightarrow \mathbb{Z}[x]$ be the inclusion map and let $J = \langle x \rangle \subset \mathbb{Z}[x]$. Explicitly describe $I = \phi^{-1}(J)$ and $\phi(I)\mathbb{Z}[x]$. Show that $\phi(I)\mathbb{Z}[x] \subsetneq J$. Thus, the inclusion in the previous part can be strict.
7. Prove that if $J \subset S$ is prime then the contraction $\phi^{-1}(J) \subset R$ is prime.
8. Give an example showing that if $J \subset S$ is maximal the contraction $\phi^{-1}(J) \subset R$ need not be maximal.
9. Give an example showing that if $I \subset R$ is prime then the extension $\phi(I)S \subset S$ need not be prime.
10. Let $\phi : R \hookrightarrow S$ be an injective ring homomorphism. Prove that

$$\phi^{-1}(J) = J \cap R \quad \text{and} \quad \phi(I)S = IS.$$

Solution.

1. We showed this in Worksheet 1.1 Question 4.1.
2. Let $R = \mathbb{Z}/4\mathbb{Z}$, $S = \mathbb{Z}$, $I = \langle 2 \rangle$ and ϕ be the inclusion map. Then we can see that $\phi(I) = \{0, 2\} \in S$ is not an ideal.
3. Let $a \in I$. Note that $\phi(a) \in \phi(I) \subseteq \phi(I)S$. Then we have,

$$\phi^{-1}(\phi(a)) = a \in \phi^{-1}(\phi(I)) \subseteq \phi^{-1}(\phi(I)S)$$

Since a was an arbitrary element, this implies that $I \subseteq \phi^{-1}(\phi(I)S)$.

4. We have $J = \phi(I)(\mathbb{Z}/\langle 6 \rangle) = \{\bar{0}\}$, and $\phi^{-1}(J) = \langle 6 \rangle$. Thus, there is a strict inclusion $I = \langle 0 \rangle \subsetneq \langle 6 \rangle = \phi^{-1}(J)$.

5. Let $I = \phi^{-1}(J)$, and let $a \in \phi(I)S$. Note that $s = \sum_i \phi(r_i)s_i$ where $r_i \in I$ and $s_i \in S$. Then $\phi(r_i) \in J$ and by definition of ideals, we have $s \in J$.
6. We have $I = \phi^{-1}J = \{r \in R \mid \phi(r) \in \langle x \rangle\} = \{0\}$, and $\phi(I)\mathbb{Z}[x] = \{0\}\mathbb{Z}[x] = \{0\}$. Thus, there is a strict inclusion $\phi(I)\mathbb{Z}[x] \subsetneq J$.
7. Let $ab \in \phi^{-1}(J)$. Then J is prime either $\phi(a) \in J$ or $\phi(b) \in J$. That is, either $a \in \phi^{-1}(J)$ or $b \in \phi^{-1}(J)$. Thus, $\phi^{-1}(J)$ is prime.
8. Let $R = \mathbb{Z}$, $S = \mathbb{Q}$, $J = \langle 0 \rangle$, and ϕ be the inclusion map. Then $\phi^{-1}(J) = \langle 0 \rangle$ which is not maximal in $\mathbb{Z}$.
9. Let $R = \mathbb{Z}$, $S = \mathbb{Z}/4\mathbb{Z}$, $I = \langle 0 \rangle$, and ϕ be the quotient map. Then $\phi(I)S = \langle 0 \rangle$ is not prime in $\mathbb{Z}/4\mathbb{Z}$ since $2 \times 2 = 0$.
10. Since ϕ is injective, R is isomorphic to a subring of S . Then $\phi^{-1}(J) = \{r \in R \mid r \in J\} = J \cap R$. Again since ϕ is injective $\phi(I) = I$ as a subset of S . So, $\phi(I)S = IS$.

As we saw in the previous exercise, in general, extension and contraction are far from being inverse operations: $\phi^{-1}(\phi(I)S)$ may strictly contain I , and $\phi(\phi^{-1}(J))S$ may be strictly smaller than J . In the case of localization, extension and contraction along the canonical localization map $\eta : R \rightarrow U^{-1}R$ sending $r \mapsto r/1$ have very nice forms. For an ideal $I \subset R$ the extension along η is

$$\eta(I) (U^{-1}R) = U^{-1}I := \left\{ \frac{r}{u} \mid r \in I, u \in U \right\} \subset U^{-1}R.$$

For an ideal $J \subset U^{-1}R$ the contraction along η is:

$$\eta^{-1}(J) = \left\{ r \in R \mid \frac{r}{1} \in J \right\}.$$

Although the extension–contraction correspondence remains lossy for arbitrary ideals, it becomes perfectly behaved for prime ideals. If $\mathfrak{q} \subset U^{-1}R$ is a proper ideal, then no element of U can lie in its contraction. (This is because $\frac{u}{1}$ is a unit in $U^{-1}R$ for every $u \in U$.) Thus every prime ideal of $U^{-1}R$ contracts to a prime ideal of R which is disjoint from U . Conversely, if $\mathfrak{p} \subset R$ is a prime ideal disjoint from U , then its extension

$$U^{-1}\mathfrak{p} = \left\{ \frac{r}{u} \mid r \in \mathfrak{p}, u \in U \right\}$$

is a prime ideal of $U^{-1}R$. In fact, extension and contraction give an order-preserving bijection

$$\left\{ \text{prime ideals of } U^{-1}R \right\} \longleftrightarrow \left\{ \mathfrak{p} \in \text{Spec}(R) \mid \mathfrak{p} \cap U = \emptyset \right\}.$$

Under this correspondence, a prime ideal $\mathfrak{q} \subset U^{-1}R$ is sent to its contraction $\eta^{-1}(\mathfrak{q})$, and a prime ideal $\mathfrak{p} \subset R$ disjoint from U is sent to its extension $U^{-1}\mathfrak{p}$.

This is especially important in the two basic examples above. If $U = R \setminus \mathfrak{p}$, then the prime ideals of $R_{\mathfrak{p}}$ correspond exactly to the prime ideals of R contained in $\mathfrak{p}$. If $U = \{1, f, f^2, \dots\}$, then the prime ideals of $R[\frac{1}{f}]$ correspond exactly to the prime ideals of R which do not contain f .

Problem 12. Ideals and Localization.

Let R be a ring, let $U \subset R$ be a multiplicatively closed set, and let $\eta : R \rightarrow U^{-1}R$ be the localization map.

1. Let $I \subset R$ be an ideal. Prove that $U^{-1}I := \{\frac{r}{u} \mid r \in I, u \in U\}$ is an ideal of $U^{-1}R$.
2. Prove that $U^{-1}I$ is the extension of I along the localization map $\eta : R \rightarrow U^{-1}R$.
3. Prove that $U^{-1}I = U^{-1}R$ if and only if $I \cap U \neq \emptyset$.
4. Prove that $\frac{r}{u} \in U^{-1}I$ if and only if $tr \in I$ for some $t \in U$.
5. Deduce that the contraction of $U^{-1}I$ back to R is

$$\eta^{-1}(U^{-1}I) = \{r \in R \mid tr \in I \text{ for some } t \in U\}.$$

6. Let $R = \mathbb{Z}$ and $U = \{1, 2, 4, 8, \dots\}$. If $I = \langle 6 \rangle$, compute $U^{-1}I \subset U^{-1}\mathbb{Z}$ and its contraction back to $\mathbb{Z}$. Show that $I \subsetneq \eta^{-1}(U^{-1}I)$.
7. Let $R = \mathbb{Z}$ and $U = \mathbb{Z} \setminus \langle 2 \rangle$. If $I = \langle 6 \rangle$, compute $U^{-1}I \subset \mathbb{Z}_{\langle 2 \rangle}$ and its contraction back to $\mathbb{Z}$.
8. Let $R = \mathbb{K}[x, y]$ and $U = \{1, x, x^2, \dots\}$. For each of the ideals $\langle x \rangle$, $\langle y \rangle$, and $\langle x, y \rangle$, describe its extension to $R[\frac{1}{x}]$. Which of these ideals become the whole ring after localization?

Solution.

1.

Problem 13. Prime Ideal Correspondence for Localization. Let R be a ring, let $U \subset R$ be a multiplicatively closed set, and let $\eta : R \rightarrow U^{-1}R$ be the localization map.

1. Let $\mathfrak{q} \subset U^{-1}R$ be a prime ideal. Prove that $\eta^{-1}(\mathfrak{q})$ is a prime ideal of R and that $\eta^{-1}(\mathfrak{q}) \cap U = \emptyset$.
2. Let $\mathfrak{p} \subset R$ be a prime ideal such that $\mathfrak{p} \cap U = \emptyset$. Prove that $U^{-1}\mathfrak{p}$ is a proper prime ideal of $U^{-1}R$. Hint: Use the criterion from the previous exercise for when a fraction belongs to $U^{-1}\mathfrak{p}$.
3. Let $\mathfrak{p} \subset R$ be a prime ideal such that $\mathfrak{p} \cap U = \emptyset$. Prove that the contraction of $U^{-1}\mathfrak{p}$ is $\mathfrak{p}$.

4. Let $\mathfrak{q} \subset U^{-1}R$ be a prime ideal. Prove that the extension of its contraction is again $\mathfrak{q}$, i.e.

$$U^{-1}\eta^{-1}(\mathfrak{q}) = \mathfrak{q}.$$

Hint: If $\frac{r}{u} \in U^{-1}R$, then $\frac{r}{u} \in \mathfrak{q}$ if and only if $\frac{r}{1} \in \mathfrak{q}$, since $\frac{u}{1}$ is a unit.

5. Conclude that extension and contraction give an order-preserving bijection

$$\text{Spec}(U^{-1}R) \longleftrightarrow \{\mathfrak{p} \in \text{Spec}(R) \mid \mathfrak{p} \cap U = \emptyset\}.$$

6. Specialize to the case $U = R \setminus \mathfrak{p}$. Prove that the prime ideals of $R_{\mathfrak{p}}$ correspond to the prime ideals of R contained in $\mathfrak{p}$.
7. Specialize to the case $U = \{1, f, f^2, \dots\}$. Prove that the prime ideals of $R[\frac{1}{f}]$ correspond to the prime ideals of R which do not contain f .
8. Describe the prime ideals of $\mathbb{Z}_{\langle p \rangle}$ for a prime number p .

Localization is not limited to rings: we can also localize modules in essentially the same way. Let R be a ring and let M be an R -module. If $U \subset R$ is a multiplicatively closed subset, we want to build a new module $U^{-1}M$ by formally dividing elements of M by the elements of U . The elements of $U^{-1}M$ are fractions $\frac{m}{u}$, where $m \in M$ and $u \in U$. So the numerators come from the module M , while the denominators come from the set U which lives in the ring. Of course, we must introduce a criterion of when two fractions $\frac{m_1}{u_1}$ and $\frac{m_2}{u_2}$ are equal. Just as when localizing a ring we say

$$\frac{m_1}{u_1} = \frac{m_2}{u_2} \quad \text{in } U^{-1}M \quad \Longleftrightarrow \quad t(u_2m_1 - u_1m_2) = 0 \quad \text{in } M \text{ for some } t \in U$$

The factor of $t \in U$ is needed in order to ensure this is a transitive relation. Define addition on $U^{-1}M$ by

$$\frac{m_1}{u_1} + \frac{m_2}{u_2} = \frac{m_1u_2 + m_2u_1}{u_1u_2}$$

which makes $U^{-1}M$ into an abelian group where the identity is $0/1$. The abelian group $U^{-1}M$ has the structure of both an R -module and a $U^{-1}R$ -module. The R -module structure and the $U^{-1}R$ -module structure are given by

$$r \left(\frac{m}{u} \right) = \frac{rm}{u} \quad \text{and} \quad \left(\frac{r}{u_1} \right) \left(\frac{m}{u_2} \right) = \frac{rm}{u_1u_2}$$

respectively. These module structures are compatible: the R -module structure on $U^{-1}M$ is the one obtained from the $U^{-1}R$ -module structure by restriction of scalars along the localization map $\eta : R \rightarrow U^{-1}R$.

In general, if $\phi : R \rightarrow S$ is any ring homomorphism and N is an S -module, then N becomes an R -module by defining $rn := \phi(r)n$ for $r \in R$ and $n \in N$. This construction is called *restriction of scalars*. One checks immediately that this gives N a well-defined R -module structure, since ϕ preserves addition, multiplication, and the identity. In our situation, the localization map $\eta : R \rightarrow U^{-1}R$ is defined by $\eta(r) = r/1$. Since $U^{-1}M$ is a $U^{-1}R$ -module, restriction of scalars along η gives $U^{-1}M$ an R -module structure via

$$r \cdot \frac{m}{u} = \eta(r) \cdot \frac{m}{u} = \frac{r}{1} \cdot \frac{m}{u} = \frac{rm}{u},$$

which is exactly the R -module structure we defined above. So there is really only one module structure at play: the $U^{-1}R$ -module structure is the fundamental one, as the R -module structure is inherited from it.

Just as the localization of a ring is characterized by a universal property, so is the localization of a module. There is a natural R -module homomorphism $\eta_M : M \rightarrow U^{-1}M$ given by $m \mapsto \frac{m}{1}$, which we call the *localization map*. Note that the target $U^{-1}M$ carries a $U^{-1}R$ -module structure, and in particular every element of U acts invertibly on $U^{-1}M$: for any $u \in U$, multiplication by $u/1 \in U^{-1}R$ is an isomorphism with inverse given by multiplication by $1/u$. The pair $U^{-1}M$ and η_M satisfies the following universal property: Let N be any $U^{-1}R$ -module, viewed as an R -module by restriction of scalars along $\eta : R \rightarrow U^{-1}R$. Then for every R -module homomorphism $\phi : M \rightarrow N$, there exists a unique $U^{-1}R$ -module homomorphism $\tilde{\phi} : U^{-1}M \rightarrow N$ such that the following diagram commutes:

$$\begin{array}{ccc} M & \xrightarrow{\eta_M} & U^{-1}M \\ \phi \downarrow & \swarrow \tilde{\phi} & \\ N & & \end{array} .$$

Problem 14. Constructing $U^{-1}M$. Let R be a ring, $U \subset R$ a multiplicatively closed set, and M be an R -module. Define a relation on $M \times U$ by $(m_1, u_1) \sim (m_2, u_2)$ if and only if $t(m_1u_2 - m_2u_1) = 0$ in M for some $t \in U$.

1. Prove that $\sim$ is an equivalence relation on $M \times U$.
2. Let $\frac{m}{u}$ denote the equivalence class of a pair (m, u) under the equivalence relation $\sim$ defined in the previous part. Prove that the operation:

$$\frac{m_1}{u_1} + \frac{m_2}{u_2} = \frac{m_1u_2 + m_2u_1}{u_1u_2}$$

is well-defined. Show that this makes $U^{-1}M$ an abelian group.

3. Show that the operations

$$r \left(\frac{m}{u} \right) = \frac{rm}{u} \quad \text{and} \quad \left(\frac{r}{u_1} \right) \left(\frac{m}{u_2} \right) = \frac{rm}{u_1 u_2}$$

are well-defined. Deduce that $U^{-1}M$ has the structure of both an R -module and a $U^{-1}R$ -module.

4. Let $\eta_M : M \rightarrow U^{-1}M$ be the function $m \mapsto \frac{m}{1}$. Show that η_M is an R -module homomorphism.
5. Prove that $\eta_M(m) = 0$ if and only if $um = 0$ for some $u \in U$. Deduce that η_M is injective if and only if no element of U acts as a zero-divisor on M .

Problem 15. Examples of Localizing Modules. Discuss ways to explicitly understand $U^{-1}M$ in each of the following cases. Describe the natural map $M \rightarrow U^{-1}M$ and determine whether it is injective.

1. $R = \mathbb{Z}, U = \mathbb{Z} \setminus \{0\}$, and $M = \mathbb{Z}$. $M = R/\langle x^3 \rangle$.
2. $R = \mathbb{Z}, U = \{1, 2, 4, 8, \dots\}$, and $M = \mathbb{Z}$. 6. $R = \mathbb{K}[x], U = R \setminus \langle x \rangle$, and $M = R/\langle x^3 \rangle$.
3. $R = \mathbb{Z}, U = \{1, 2, 4, 8, \dots\}$, and $M = \mathbb{Z}/\langle 6 \rangle$. 7. $R = \mathbb{K}[x, y], U = \{1, y, y^2, \dots\}$, and $M = R/\langle xy \rangle$.
4. $R = \mathbb{Z}, U = \mathbb{Z} \setminus \langle 2 \rangle$, and $M = \mathbb{Z}/\langle 6 \rangle$. 8. R is an arbitrary ring, M is an R -module, and U contains an element u such that $uM = 0$.
5. $R = \mathbb{K}[x], U = \{1, x, x^2, \dots\}$, and

Problem 16. Universal Property of $U^{-1}M$. Let R be a ring, let $U \subset R$ be a multiplicatively closed set, and let M be an R -module. Let N be a $U^{-1}R$ -module, which we view as an R -module via the localization map $\eta : R \rightarrow U^{-1}R$. Let $\psi : M \rightarrow N$ be an R -module homomorphism.

1. Define a function $\tilde{\psi} : U^{-1}M \rightarrow N$ by $\tilde{\psi}(\frac{m}{u}) = \frac{1}{u}\psi(m)$. Prove that $\tilde{\psi}$ is well-defined.
2. Prove that $\tilde{\psi}$ is a homomorphism of $U^{-1}R$ -modules.

3. Show that $\tilde{\psi}$ makes the following diagram commute:

$$\begin{array}{ccc} M & \xrightarrow{\eta_M} & U^{-1}M \\ \psi \downarrow & \swarrow \tilde{\psi} & \\ N & & \end{array}$$

4. Prove that $\tilde{\psi}$ is the unique $U^{-1}R$ -module homomorphism making the diagram commute.
5. Deduce that $U^{-1}M$ is unique up to unique isomorphism as a $U^{-1}R$ -module satisfying this universal property.

Problem 17. $U^{-1}M$ as a Tensor Product. Let R be a ring, $U \subset R$ be a multiplicatively closed set, and let M be an R -module. We view $U^{-1}R$ as an R -module via the localization map $\eta : R \rightarrow U^{-1}R$.

1. Show that the function given below is a well-defined bilinear map of R -modules:

$$\begin{aligned} U^{-1}R \times M &\xrightarrow{\bar{\alpha}} U^{-1}M \\ \left(\frac{r}{u}, m\right) &\longmapsto \frac{rm}{u} \end{aligned}$$

2. Use the universal property of tensor products to construct an R -module homomorphism $\alpha : U^{-1}R \otimes_R M \rightarrow U^{-1}M$. Find a formula for $\alpha\left(\frac{r}{u} \otimes m\right)$.
3. Show that the function β given below is a well-defined R -module homomorphism:

$$\begin{aligned} U^{-1}M &\xrightarrow{\beta} U^{-1}R \otimes_R M \\ \frac{m}{u} &\longmapsto \frac{1}{u} \otimes m \end{aligned}$$

4. Prove that α and β are mutual inverses. Conclude that $U^{-1}M \cong U^{-1}R \otimes_R M$ as R -modules.
5. Deduce that $U^{-1}(R^n) \cong (U^{-1}R)^n$ as $U^{-1}R$ -modules.
6. If $f : M \rightarrow N$ is an R -module homomorphism, define a natural $U^{-1}R$ -module homomorphism $U^{-1}f : U^{-1}M \rightarrow U^{-1}N$. What is its formula on fractions?