

Math 121: Commutative Algebra

Prishita Dharampal

Throughout this course, “ring” means a *commutative* ring with unity. Recall that, given an ideal $I \subset R$, we define the vanishing set of I to be

$$\mathbb{V}(I) := \{[P] \in \operatorname{Spec}(R) \mid I \subset P\}.$$

The *Zariski topology* on $\operatorname{Spec}(R)$ is then defined by declaring the closed sets to be the subsets of the form $\mathbb{V}(I)$ as I ranges over all ideals of R . In Worksheet 2.1 we verified that this is indeed a topology and studied it in detail. For example, we showed that any ring map $\phi : R \rightarrow S$ induces a continuous map of topological spaces $\phi^\# : \operatorname{Spec}(S) \rightarrow \operatorname{Spec}(R)$ given by $[P] \mapsto [\phi^{-1}(P)]$. (More precisely, Spec is a contravariant functor from the category of commutative rings to the category of topological spaces.) Further, we showed that the natural quotient map $\pi : R \rightarrow R/I$ induces a closed embedding $\pi^\# : \operatorname{Spec}(R/I) \rightarrow \operatorname{Spec}(R)$ whose image is $\mathbb{V}(I)$.

Now that we have localization in our toolbox, we return to studying the Zariski topology on $\operatorname{Spec}(R)$. The main slogan is that localizing a ring lets us study the part of $\operatorname{Spec}(R)$ where a given “function” is nonzero. We have not yet defined functions on $\operatorname{Spec}(R)$, but in many familiar examples elements of R really are functions—for example, when R is the coordinate ring of an algebraic set. Recall the following theorem from the last worksheet.

Theorem 0.1. *Let R be a ring and let $U \subset R$ be a multiplicatively closed set. The localization map $\eta : R \rightarrow U^{-1}R$ induces an inclusion-preserving bijection:*

$$\begin{array}{ccc} \left\{ \begin{array}{l} \text{prime ideals in} \\ U^{-1}R \end{array} \right\} & \xrightarrow{\sim} & \left\{ \begin{array}{l} \text{prime ideals in } R \\ \text{disjoint from } U \end{array} \right\} \\ P & \longmapsto & \eta^{-1}(P) \end{array}$$

We begin by using this theorem in the case that $U = \{1, f, f^2, f^3, \dots\}$ for an element $f \in R$ to study what are called the basic open sets of $\operatorname{Spec}(R)$. A *basic open subset* – also called a *distinguished open subset* – of $\operatorname{Spec}(R)$ is a set of the form

$$D(f) := \{[P] \in \operatorname{Spec}(R) \mid f \notin P\} \subset \operatorname{Spec}(R)$$

for some element $f \in R$. We will show that these subsets form a nice basis for the Zariski topology on $\operatorname{Spec}(R)$. A *basis* for a topological space X is a collection $\mathcal{B}$ of open subsets of

X such that for every open set $U \subset X$ and every point $x \in U$, there exists some $B \in \mathcal{B}$ with $x \in B \subset U$. Equivalently, every open subset of X can be written as a union of elements of $\mathcal{B}$.

Problem 1. Basic Open Sets.

Let R be a ring and $f, g \in R$.

1. Prove directly from the definition of the Zariski topology that $D(f)$ is an open set.
2. Compute $D(0)$ and $D(1)$.
3. Prove that $D(f) = D(f^n)$ for all $n \geq 1$.
4. Prove that $D(f) \cap D(g) = D(fg)$. Use induction to find a formula for the intersection of finitely many basic opens.
5. Prove that $D(f) \subset D(g)$ if and only if $f \in \sqrt{\langle g \rangle}$. Hint: Take complements to rewrite $D(f) \subset D(g)$ as $\mathbb{V}(g) \subset \mathbb{V}(f)$, and use the fact that $\mathbb{V}(I) \subset \mathbb{V}(J)$ if and only if $\sqrt{J} \subset \sqrt{I}$.
6. Prove that $D(f) = D(g)$ if and only if $\sqrt{\langle f \rangle} = \sqrt{\langle g \rangle}$.
7. Let $U \subset \text{Spec}(R)$ be an open subset and let $[\mathfrak{p}] \in U$. Show that there exists $f \in R$ such that $[\mathfrak{p}] \in D(f) \subset U$. Hint: Since U is open, write $\text{Spec}(R) \setminus U = \mathbb{V}(I)$ for some ideal I . Then use the fact that $[\mathfrak{p}] \notin \mathbb{V}(I)$ to choose $f \in I \setminus \mathfrak{p}$.
8. Deduce that the basic open sets form a basis for the Zariski topology on $\text{Spec}(R)$.

Solution.

1. $D(f)$ is the set of all prime ideals where $f \neq 0$, i.e. where f doesn't vanish. So, $\text{Spec}(R) \setminus D(f) = \mathbb{V}(f)$. Which by definition is a closed set, thus $D(f)$ is open.
2. $D(0) = \emptyset$, and $D(1) = \text{Spec}(R)$ since 0 is in every prime ideal, and all prime ideals are proper.
3. $D(f^n) \subseteq D(f)$ since P is prime and $f^n \notin P$ we must have $f \notin P$. And $D(f) \subseteq D(f^n)$ since if $f \notin P$ and P is prime, $f^n \notin P$. Thus $D(f) = D(f^n)$ for all $n \geq 1$.
4. $D(f) \cap D(g) = \{[P] \in \text{Spec}(R) \mid f, g \notin P\}$. Since P is prime, this implies that $fg \notin P$. Thus we have,

$$D(f) \cap D(g) = \{[P] \in \text{Spec}(R) \mid f, g \notin P\} = \{[P] \in \text{Spec}(R) \mid fg \notin P\} = D(fg)$$

5. ($\implies$)

Assume $D(f) \subset D(g)$.

Then we have $\mathbb{V}(g) \subset \mathbb{V}(f) \implies \sqrt{\langle f \rangle} \subset \sqrt{\langle g \rangle} \implies f \in \sqrt{\langle g \rangle}$.

($\Leftarrow$)

Assume $f \in \sqrt{\langle g \rangle}$. Then $f^n \in \langle g \rangle$ for some n . We first show that $\sqrt{\langle f \rangle} \subset \sqrt{\langle g \rangle}$. Let $h \in \sqrt{\langle f \rangle}$, then $h^m = rf \in \langle f \rangle$ for some $m \in \mathbb{N}$ and some $r \in R$. Then, we have $h^{nm} = r^n f^n \in \langle g \rangle$. Thus, $h \in \sqrt{\langle g \rangle}$ and $\sqrt{\langle f \rangle} \subset \sqrt{\langle g \rangle}$.

But that implies $\mathbb{V}(g) \subset \mathbb{V}(f) \implies D(f) \subset D(g)$.

6. ($\implies$)

Assume $D(f) = D(g)$. Then we have $\mathbb{V}(g) = \mathbb{V}(f) \implies \sqrt{\langle f \rangle} = \sqrt{\langle g \rangle}$.

($\Leftarrow$)

Assume $\sqrt{\langle f \rangle} = \sqrt{\langle g \rangle}$. Then we have $\mathbb{V}(g) = \mathbb{V}(f) \implies D(f) = D(g)$.

7. Let $U \in \text{Spec}(R)$ open set. Then its complement $\text{Spec}(R) \setminus U = \mathbb{V}(I)$ is a closed set for some $I \subset R$. Since $[\mathfrak{p}] \in U$ we know that $[\mathfrak{p}] \notin \mathbb{V}(I)$, then we have $I \not\subset \mathfrak{p}$. So there exists some $f \in I$ such that $f \notin \mathfrak{p}$ giving us $[\mathfrak{p}] \in D(f) \subset U$.
8. Since $[\mathfrak{p}]$ and U were arbitrary, by part (7) the basic open sets satisfy the basis criterion and hence form a basis.

Problem 2. Examples of Basic Opens.

Explicitly describe the basic open set $D(f) \subset \text{Spec}(R)$ in each of the following cases.

- | | |
|--|--|
| 1. $R = \mathbb{K}[x]$ and $f = x$. | 5. $R = \mathbb{K}[x, y] / \langle xy \rangle$ and $f = \bar{y}$. |
| 2. $R = \mathbb{K}[x]$ and $f = x - a$ for $a \in \mathbb{K}$. | 6. $R = \mathbb{Z} / \langle 12 \rangle$ and $f = \bar{2}$. |
| 3. $R = \mathbb{Z}$ and $f = p$ for a prime number p . | 7. $R = \mathbb{Z} / \langle 12 \rangle$ and $f = \bar{3}$. |
| 4. $R = \mathbb{K}[x, y] / \langle xy \rangle$ and $f = \bar{x}$. | 8. $R = \mathbb{Z} / \langle 12 \rangle$ and $f = \bar{6}$. |

Solution.

1. $D(x) = \text{Spec}(R) \setminus \langle x \rangle = \{ \langle 0 \rangle \cup \langle p(x) \rangle \mid p \neq x, p \text{ is irreducible} \}.$
2. $D(x - a) = \text{Spec}(R) \setminus \langle x - a \rangle = \{ \langle 0 \rangle \cup \langle p(x) \rangle \mid p \neq x - a, p \text{ is irreducible} \}.$
3. $D(p) = \text{Spec}(\mathbb{Z}) \setminus \langle p \rangle = \{ [\mathfrak{p}] \in \text{Spec}(\mathbb{Z}), \mathfrak{p} \neq \langle p \rangle \}.$
4. $D(\bar{x}) = \{ \langle \bar{y}, q(x) \rangle, q \text{ is irreducible or zero} \}.$
5. $D(\bar{y}) = \{ \langle \bar{x}, q(y) \rangle, q \text{ is irreducible or zero} \}.$
6. $D(\bar{2}) = \{ \langle \bar{3} \rangle \}.$
7. $D(\bar{3}) = \{ \langle \bar{2} \rangle \}.$
8. $D(\bar{6}) = \emptyset.$

Problem 3. Induced Map by Localization.

Let $U \subset R$ be a multiplicatively closed set and let $\eta : R \rightarrow U^{-1}R$ be the localization map. In this exercise we will study the induced map $\eta^\# : \text{Spec}(U^{-1}R) \rightarrow \text{Spec}(R)$.

1. Use the correspondence between prime ideals in $U^{-1}R$ and R given in Theorem 0.1 to prove that $\eta^\#$ is injective and the image of $\eta^\#$ is

$$X_U := \{[\mathfrak{p}] \in \text{Spec}(R) \mid \mathfrak{p} \cap U = \emptyset\} \subset \text{Spec}(R).$$

2. Let $R = \mathbb{Z}$ and $U = \mathbb{Z} \setminus \langle p \rangle$ for a prime $p \in \mathbb{Z}$. Show that $X_U = \{\langle 0 \rangle, \langle p \rangle\}$. Prove that this is not open in $\text{Spec}(\mathbb{Z})$.
3. Let $R = \mathbb{Z}$ and $U = \{2^i 3^j \mid i, j \in \mathbb{Z}_{\geq 0}\}$. Show that $X_U = \{\langle 0 \rangle\} \cup \{\langle p \rangle \mid p \text{ prime}, p \neq 2, 3\}$. Show that this is an open subset of $\text{Spec}(\mathbb{Z})$.
4. Describe X_U when $R = \mathbb{C}[x, y]$ and $U = \{(xy)^n \mid n \in \mathbb{Z}_{\geq 0}\}$.
5. Prove that $X_U = \bigcap_{f \in U} D(f)$.
6. Deduce that if U is a multiplicative set generated by finitely many elements then X_U is open.
7. Fix an element $\frac{r}{u} \in U^{-1}R$. Prove that $\eta^\#(D(\frac{r}{u})) = D(r) \cap X_U$. Hint: In $U^{-1}R$, the element u is a unit, so $D(\frac{r}{u}) = D(\frac{r}{1})$. Then use part (1) to identify the image of $\eta^\#$.
8. Prove that $\eta^\# : \text{Spec}(U^{-1}R) \rightarrow \text{Spec}(R)$ is a homeomorphism onto its image X_U (considered with the subspace topology).
9. Specializing to the case that $U = \{1, f, f^2, f^3, \dots\}$, prove that $\eta^\# : \text{Spec}(R[\frac{1}{f}]) \rightarrow \text{Spec}(R)$ is a homeomorphism onto the basic open subset $D(f)$.

Solution.

1. By Theorem 0.1 the map $\mathfrak{q} \rightarrow \eta^{-1}(\mathfrak{q})$ gives a bijection between prime ideals in $U^{-1}R$ to prime ideals disjoint from U in R . That is,

$$\text{im}(\eta^\#) = X_U := \{[\mathfrak{p}] \in \text{Spec}(R) \mid \mathfrak{p} \cap U = \emptyset\} \subset \text{Spec}(R).$$

and $\eta^\#$ is injective since the bijection is injective.

2. We have, by construction,

$$X_U = \{[\mathfrak{p}] \in \text{Spec}(R) \mid \mathfrak{p} \cap U = \emptyset\} = \{\langle p \rangle, \langle 0 \rangle\}$$

Assume X_U is open in $\text{Spec}(\mathbb{Z})$. Since distinguished open sets form a basis for the Zariski topology, there exists some $D(f) \subseteq X_U$ such that $\langle 0 \rangle \subseteq D(f)$ which means $f \neq 0$. So we must have $f = n$, for some $n \in \mathbb{Z}$. Since n is divisible by only a finite number of primes, we have $D(n) = \{[P] \in \text{Spec}(\mathbb{Z}) \mid P \nmid n\}$ which is an infinite set, and thus $D(f) \not\subseteq X_U$. Thus, a contradiction and X_U cannot be a closed set.

3. By definition,

$$\begin{aligned} X_U &= \{[p] \in \text{Spec}(\mathbb{Z}) \mid p \cap U = \emptyset\} \\ &= \{[p] \in \text{Spec}(\mathbb{Z}) \mid p \neq 2, 3\} \\ &= \{\langle 0 \rangle\} \cup \{\langle p \rangle \mid p \text{ prime}, p \neq 2, 3\} \end{aligned}$$

Since X_U is the complement of a closed set in $\text{Spec}(\mathbb{Z})$, namely, $V(\langle 6 \rangle) = \{\langle 2 \rangle, \langle 3 \rangle\}$, it is an open set of $\text{Spec}(\mathbb{Z})$.

4. By definition,

$$\begin{aligned} X_U &= \{[p] \in \text{Spec}(\mathbb{C}[x, y]) \mid p \cap U = \emptyset\} \\ &= \{[p] \in \text{Spec}(\mathbb{C}[x, y]) \mid p \neq x, y\} \\ &= \{\langle 0 \rangle\} \end{aligned}$$

5. $X_U \supseteq \cap_{f \in U} D(f)$ is clear by definition. To see the other containment, let $[p] \in \cap_{f \in U} D(f)$. Then $f \notin p$ for all f . Hence, $p \cap U = \emptyset$ and thus $[p] \in X_U$.

6. In this case, X_U is a finite intersection of open sets and thus is open.

7. We know that $D(\frac{r}{u}) = D(\frac{r}{1})$. For each prime $[p] \in D(\frac{r}{1})$,

$$\frac{r}{1} = \eta(r) \notin p \iff r \notin \eta^\#(p)$$

8. From part 1 we know that $\eta^\#$ is injective, thus it is a bijection onto its image. Let $D_u^r \subseteq \text{Spec}(U^{-1}R)$ be a basic open set. Then by part 7,

$$\eta^\#(D(\frac{r}{u})) = D(r) \cap X_U$$

which is open in the subspace topology on X_U .

9. If $U = \{1, f, f^2, \dots\}$, then

$$X_U = \cap_{f \in U} D(f) = D(f)$$

as $D(f)$ contains $D(f^n)$ for all n . Hence, $\eta(\text{Spec}(f[\frac{1}{f}])) \cong D(f)$.

Let X be a topological space. An *open cover* of X is a collection $\{U_i\}_{i \in \Lambda}$ of open subsets of X such that $X = \bigcup_{i \in \Lambda} U_i$. A *finite subcover* is a finite subcollection $\{U_{i_1}, \dots, U_{i_n}\}$ that still covers X , i.e. $X = U_{i_1} \cup \dots \cup U_{i_n}$.

With these terms in hand, recall that X is *quasi-compact* if every open cover admits a finite subcover. More generally, a subset $Y \subset X$ is *quasi-compact* if every cover of Y by open subsets of X admits a finite subcover; equivalently, Y is quasi-compact in the subspace topology. In the Hausdorff setting this is simply compactness; the prefix “quasi-” is standard in algebraic geometry because Zariski topologies are virtually never Hausdorff, and one wants to keep the two notions distinct.

Every affine spectrum $\text{Spec } R$ is quasi-compact. The proof is a clean illustration of how the basic opens $D(f)$ serve as a bridge between topology and commutative algebra. If $\{D(f_i)\}_{i \in \Lambda}$ covers $\text{Spec}(R)$, then no prime ideal contains every f_i , so the f_i generate the unit ideal. But any expression $1 = \sum a_i f_i$ involves only finitely many terms, so a finite subfamily already generates the unit ideal and the corresponding $D(f_i)$ ’s already cover. The argument generalizes immediately: any open subset that can be written as a finite union of basic opens is itself quasi-compact.

This kind of translation—reformulating a covering condition as an ideal-membership statement, then exploiting the finiteness built into ring equations—is one of the workhorses of scheme theory.

Problem 4. Quasi-compactness of the Zariski Topology.

Let R be a ring.

1. Let $\Lambda \subset R$ be a collection of elements of R . Prove that $\{D(f)\}_{f \in \Lambda}$ is an open cover of $\text{Spec}(R)$ if and only if $\langle f \mid f \in \Lambda \rangle$ is the unit ideal.
2. Let $\Lambda \subset R$ be a collection of elements of R . Explain why if $\langle f \mid f \in \Lambda \rangle$ is the unit ideal then there exist finitely many elements $f_1, \dots, f_n \in \Lambda$ such that $\langle f_1, \dots, f_n \rangle$ is also the unit ideal.
3. Deduce that every cover of $\text{Spec}(R)$ by basic open subsets has a finite subcover.
4. Use the fact that the $D(f)$ form a basis to prove that $\text{Spec}(R)$ is quasi-compact. Hint: Refine each open set in the cover by basic open neighborhoods using the basis property from Problem 1.
5. Prove that $\mathbb{V}(I)$ is quasi-compact for every ideal $I \subseteq R$. Hint: Use the homeomorphism $\text{Spec}(R/I) \cong \mathbb{V}(I)$ from Worksheet 2.1.
6. Let $I \subset R$ be an ideal and set $U := \text{Spec}(R) \setminus \mathbb{V}(I)$.
 - (a) Show that $U = \bigcup_{f \in I} D(f)$.
 - (b) Show that if U is quasi-compact, then $U = D(f_1) \cup \dots \cup D(f_n)$ for some $f_1, \dots, f_n \in I$.
 - (c) Deduce that U is quasi-compact if and only if $\sqrt{I} = \sqrt{\langle f_1, \dots, f_n \rangle}$ for some $f_1, \dots, f_n \in I$.

Equivalently, U is quasi-compact if and only if $\sqrt{I}$ is the radical of a finitely generated ideal.
7. Deduce that if R is Noetherian, every open subset of $\text{Spec}(R)$ is quasi-compact. Hint: First show that $\text{Spec}(R) \setminus \mathbb{V}(I) = \bigcup_{f \in I} D(f)$.

Solution.

1. ($\implies$)

Assume $\{D(f)\}_{f \in \Lambda}$ is an open cover of $\text{Spec}(R)$. Then $\langle f \mid f \in \Lambda \rangle \not\subseteq \mathfrak{p}$ for any prime ideal. In particular, $\langle f \mid f \in \Lambda \rangle$ is not contained in any maximal ideal, and hence must be the unit ideal.

($\impliedby$)

Assume $\langle f \mid f \in \Lambda \rangle$ is the unit ideal. Then it is not contained in any prime ideal. In particular, there exists at least one $f \in \Lambda$ such that $f \notin \mathfrak{p}$ for all $\mathfrak{p}$. Hence, $\{D(f)\}_{f \in \Lambda}$

is a open cover of $\text{Spec}(R)$.

2. If $\langle f \mid f \in \Lambda \rangle$ is the unit ideal, then $1 = \sum_i a_i f_i \in \langle f \mid f \in \Lambda \rangle$ is a finite sum. Then, these (finite) f_i s generate the unit ideal.
3. Thus, the corresponding finitely many $D(f_i)$ s generate a finite subcover for any given cover of $\text{Spec}(R)$ by basic open sets.
4. Every open set can be written as a union of the basic open subsets and hence every open cover is a cover of $\text{Spec}(R)$ by basic open subsets and hence has a finite subcover. Thus, $\text{Spec}(R)$ is quasi-compact.
5. Recall that, $\mathbb{V}(I) \cong \text{Spec}(R/I)$. Since R/I is a ring, and we just showed that the spectrum of a ring is quasi-compact, $\text{Spec}(R/I)$ is quasi-compact.
6. (a) $U = \text{Spec}(R) \setminus \mathbb{V}(I) = \text{Spec}(R) \setminus \text{Spec}(R/I)$. We know that ideals in $\text{Spec}(R/I)$ correspond to ideals in $\text{Spec}(R)$ that contain I . So, U is the set of ideals in $\text{Spec}(R)$ that don't contain I . Hence,

$$U = \text{Spec}(R) \setminus \text{Spec}(R/I) = \cup_{f \in I} D(f)$$

- (b) If U is quasi-compact, then any cover of U has a finite-subcover. Since $\cup_{f \in I} D(f)$ is a cover, it also admits a finite subcover,

$$U = D(f_1) \cup \dots \cup D(f_n)$$

- (c) ($\implies$)

Assume U is quasi-compact. Then,

$$\begin{aligned} U &= D(f_1) \cup \dots \cup D(f_n) \\ &= \text{Spec}(R) \setminus (\mathbb{V}(f_1) \cap \dots \cap \mathbb{V}(f_n)) \\ &= \text{Spec}(R) \setminus \mathbb{V}(\langle f_1, \dots, f_n \rangle) \\ &= \text{Spec}(R) \setminus \mathbb{V}(\sqrt{\langle f_1, \dots, f_n \rangle}) \\ &= \text{Spec}(R) \setminus \mathbb{V}(I) = \text{Spec}(R) \setminus \mathbb{V}(\sqrt{I}) \\ &\implies \sqrt{I} = \sqrt{\langle f_1, \dots, f_n \rangle} \end{aligned}$$

- ($\impliedby$)

Assume $\sqrt{I} = \sqrt{\langle f_1, \dots, f_n \rangle}$ for some $f_1, \dots, f_n \in I$. Then,

$$\begin{aligned} U &= \operatorname{Spec}(R) \setminus \mathbb{V}(I) = \operatorname{Spec}(R) \setminus \mathbb{V}(\sqrt{I}) \\ &= \operatorname{Spec}(R) \setminus \mathbb{V}(\sqrt{\langle f_1, \dots, f_n \rangle}) \\ &= \operatorname{Spec}(R) \setminus (\mathbb{V}(f_1) \cap \dots \cap \mathbb{V}(f_n)) \\ &= \operatorname{Spec}(R) \setminus \mathbb{V}(\langle f_1, \dots, f_n \rangle) \\ &= D(f_1) \cup \dots \cup D(f_n) \end{aligned}$$

7. If R is Noetherian, every ideal $\mathbb{V}(I)$ is finitely-generated. Then by part 6.b, every open subset of $\operatorname{Spec}(R)$ is quasi-compact.

Problem 5. Counterexamples to Quasi-compactness of Subsets.

In the previous problem we saw that $\text{Spec}(R)$ and every closed subset $V(I) \subset \text{Spec}(R)$ are quasi-compact in the Zariski topology. We also saw that when R is Noetherian every open subset of $\text{Spec}(R)$ is quasi-compact. In this exercise we explore some pathologies for open subsets when R is not Noetherian.

1. Let $R = \mathbb{K}[x_1, x_2, \dots]$ and $I = \langle x_1, x_2, \dots \rangle$. Consider the open subset $U := \text{Spec}(R) \setminus V(I)$. Prove that $\{D(x_i)\}_{i=1}^\infty$ is an open cover of U , which has no finite subcover. Hence an open subset of $\text{Spec}(R)$ need not be quasi-compact. Hint: If you try to cover U by only $D(x_1), \dots, D(x_n)$, consider the prime ideal $\langle x_1, \dots, x_n \rangle$.
2. Explain why each basic open set $D(x_i)$ is quasi-compact. Deduce that an arbitrary union of quasi-compact open subsets of $\text{Spec}(R)$ need not be quasi-compact.

Solution.

1. For the sake of contradiction, assume there exists a finite subcover of $\{D(x_i)\}_{i=1}^\infty$ that covers U , say $D(x_1) \cup \dots \cup D(x_n)$. Then, the ideal generated by $\langle x_1, \dots, x_n \rangle \subseteq I$ is in U . Which is a contradiction. Hence, no finite subcover exists.
2. We know that $D(x_i) \cong \text{Spec}\left(R\left[\frac{1}{x_i}\right]\right)$, since the spectrum of any ring is quasi-compact $D(x_i)$ is quasi compact. And in part (1) we see that arbitrary unions of quasi-compact open subsets of $\text{Spec}(R)$ are not quasi-compact, however finite unions are.

Problem 6. Checking Properties on a Finite Cover.

Let R be a ring. Suppose that $f_1, \dots, f_n \in R$ are such that $\{D(f_i)\}_{i=1}^n$ is a finite open cover of $\text{Spec}(R)$, i.e., $\text{Spec}(R) = D(f_1) \cup D(f_2) \cup \dots \cup D(f_n)$. We will see that certain ring-theoretic properties of R can be checked locally by looking at the rings $R[\frac{1}{f_i}]$.

1. Let $r \in R$. Prove that if $r/1 = 0$ in $R[\frac{1}{f_i}]$ for all i then $r = 0$ in R . Hint: In $R[\frac{1}{f_i}]$, the equality $r/1 = 0$ means that $f_i^{n_i}r = 0$ for some $n_i \geq 0$.
2. Deduce that if $R[\frac{1}{f_i}] = 0$ for all i then $R = 0$.
3. Prove that if $R[\frac{1}{f_i}]$ is reduced for all i , then R is reduced.

Solution.

1. Since $\{D(f_i)\}_{i=1}^n$ covers $\text{Spec}(R)$, we know $\langle f_1, \dots, f_n \rangle = R$. And for any prime $\mathfrak{p}$ we have $f_i \in \mathfrak{p} \iff f_i^{n_i} \in \mathfrak{p}$, so $D(f_i) = D(f_i^{n_i})$.

Hence $\{D(f_i^{n_i})\}_{i=1}^n$ also covers $\text{Spec}(R)$, and we have

$$1 = \sum_{i=1}^n b_i f_i^{n_i} \implies r = 1 \cdot r = \sum_{i=1}^n b_i f_i^{n_i} r = \sum_{i=1}^n b_i \cdot 0 = 0.$$

Thus, $r = 0$ in R .

2. If $R[\frac{1}{f_i}] = 0$ for all i then each f_i is nilpotent in R . But $D(f_i)_{1 \leq i \leq n}$ covers R . Since $f_i \in \sqrt{\langle 0 \rangle} \subseteq \mathfrak{p}$ for all primes, $D(f_i) = \emptyset$ for all i . Thus, $\text{Spec}(R) = \emptyset \implies R = 0$.
3. Let $r \in R$ such that $r^m = 0$. Then for all i , in $R[\frac{1}{f_i}]$, we have $\frac{r^m}{1} = 0$. Since this ring is reduced, this implies $\frac{r}{1} = 0 \implies r = 0$ (by part 1).

An element $r \in R$ is *nilpotent* if $r^n = 0$ for some integer $n \geq 1$. If $I \subset R$ is an ideal, the *radical* of I is

$$\sqrt{I} := \{r \in R \mid r^n \in I \text{ for some } n \geq 1\},$$

which we previously checked is an ideal of R containing I and $\sqrt{\sqrt{I}} = \sqrt{I}$. An ideal $I \subset R$ is *radical* if and only if $I = \sqrt{I}$. We saw that radical ideals play a crucial role in the study of algebraic sets as $\mathbb{I}(V)$ is always radical and Hilbert's Nullstellensatz gave an inclusion-reversing bijection between radical ideals in $\mathbb{K}[x_1, \dots, x_n]$ and algebraic subsets of $\mathbb{K}^n$ when $\mathbb{K}$ is an algebraically closed field. These facts highlight the following motto: *(classical) geometry does not see nilpotents*. (Here classical means anything one encounters pre-scheme theory; topology, classical algebraic geometry, complex geometry, etc.)

We shall see that the invisibility of nilpotents to geometry also applies to spectra of rings. The *nilradical* of R is the set of all nilpotent elements of R , equivalently the radical of the zero ideal:

$$\text{Nil}(R) := \sqrt{\langle 0 \rangle} = \{r \in R \mid r^n = 0 \text{ for some } n \geq 1\}.$$

If $f \in \text{Nil}(R)$ and $f^n = 0$, then $f^n \in I$ for every ideal $I \subset R$. In particular, if $P \subset R$ is a prime ideal, then $f^n = 0 \in P$, which implies that $f \in P$. Thus the nilradical of R is contained in every prime ideal of R . In the exercises below you will prove the reverse inclusion, giving the fundamental characterization

$$\text{Nil}(R) = \bigcap_{\mathfrak{p} \in \text{Spec}(R)} \mathfrak{p}.$$

Since prime ideals of $R/\text{Nil}(R)$ are in bijection with prime ideals of R containing $\text{Nil}(R)$, the equality above implies that prime ideals of R are in bijection with prime ideals of $R/\text{Nil}(R)$. Geometrically, this means that the quotient map $\pi : R \rightarrow R/\text{Nil}(R)$ induces a map of topological spaces

$$\text{Spec}(R/\text{Nil}(R)) \xrightarrow{\pi^\#} \text{Spec}(R)$$

that is bijective and continuous. We will see shortly that this map is actually a homeomorphism; hence, topologically $\text{Spec}(R)$ and $\text{Spec}(R/\text{Nil}(R))$ are indistinguishable. This is another instance of the motto: *geometry does not see nilpotents*. A ring R is called *reduced* if $\text{Nil}(R) = \langle 0 \rangle$, i.e., if the only nilpotent element is 0. We often call $R/\text{Nil}(R)$ the *reduced ring* associated to R and denote it $R_{\text{red}} := R/\text{Nil}(R)$.

More generally, the nilradical characterization extends to arbitrary ideals. One checks that $\text{Nil}(R/I) = \sqrt{I}/I$, since the elements of R/I whose powers lie in I are exactly the images of elements of $\sqrt{I}$. Applying the nilradical formula in the ring R/I and pulling back to R gives

$$\sqrt{I} = \bigcap_{\substack{\mathfrak{p} \in \text{Spec}(R) \\ \mathfrak{p} \supseteq I}} \mathfrak{p}.$$

This has immediate geometric consequences. For example, this gives another proof that $\mathbb{V}(I) = \mathbb{V}(\sqrt{I})$. Thus the Zariski-closed subsets of $\operatorname{Spec}(R)$ determined by I and $\sqrt{I}$ are identical. This interacts cleanly with quotient rings; in particular, an argument like the one above shows that $\operatorname{Spec}(R/I)$ and $\operatorname{Spec}(R/\sqrt{I})$ are homeomorphic.

Problem 7. Nilradicals and Reduced Rings.

Let R be a ring.

1. Explain why $\text{Nil}(R)$ is an ideal and $\text{Nil}(R) \subset \bigcap_{\mathfrak{p} \in \text{Spec}(R)} \mathfrak{p}$.
2. Prove that if $f \in R$ is not a nilpotent then $R[\frac{1}{f}] \neq 0$ and hence contains a prime ideal. Hint: If $R[\frac{1}{f}] = 0$, then $1 = 0$ in $R[\frac{1}{f}]$. What does that say about some power of f ?
3. Use the correspondence between prime ideals in $R[\frac{1}{f}]$ and R not containing f to show there is a prime ideal $\mathfrak{p} \subset R$ such that $f \notin \mathfrak{p}$.
4. Deduce the theorem

$$\text{Nil}(R) = \bigcap_{\mathfrak{p} \in \text{Spec}(R)} \mathfrak{p}.$$

5. Define R_{red} to be equal to $R/\text{Nil}(R)$. Prove that R_{red} is reduced.
6. Use the above characterization to prove that $D(f) = \emptyset$ if and only if f is a nilpotent element.
7. Show that $D(f) = \text{Spec}(R)$ if and only if f is a unit in R .
8. Let $\pi : R \rightarrow R_{\text{red}}$ be the natural quotient map. Prove that the induced map $\pi^\# : \text{Spec}(R_{\text{red}}) \rightarrow \text{Spec}(R)$ is a homeomorphism. (Hint: We already know it is continuous. Use part 4 to show it is a bijection. Finally, show that the map is closed by proving that for every ideal $I \supseteq \text{Nil}(R)$, the closed set $\mathbb{V}(I/\text{Nil}(R)) \subseteq \text{Spec}(R_{\text{red}})$ maps to $\mathbb{V}(I) \subseteq \text{Spec}(R)$.)
9. Compute R_{red} for each of the following rings:

(a) $\mathbb{K}[\epsilon]/\langle \epsilon^2 \rangle,$

(c) $\mathbb{K}[x, y]/\langle x^2, xy \rangle,$

(b) $\mathbb{K}[x]/\langle x^3 \rangle,$

(d) $\mathbb{Z}/12\mathbb{Z}.$

Solution.

1. We only need to show that $\text{Nil}(R)$ is closed under addition and multiplication from the ring.
 - (a) Let $a, b \in \text{Nil}(R)$ such that $a^n = b^m = 0$. Wlog assume $m > n$, then by the binomial theorem $(a + b)^{m+n} = 0$.
 - (b) Let $r \in R$, and $a \in \text{Nil}(R)$ such that $a^n = 0$. Then, $(ra)^n = r^n a^n = 0$.

Hence, $\text{Nil}(R)$ is an ideal. Let $a \in \text{Nil}(R)$ then $a^n = 0 \in \mathfrak{p}$ then by primeness of $\mathfrak{p}$ $a \in \mathfrak{p}$ for all $\mathfrak{p}$, hence $\text{Nil}(R) \subset \bigcap_{\mathfrak{p} \in \text{Spec}(R)} \mathfrak{p}$.

2. We will prove this by contrapositive. Assume $R[\frac{1}{f}] = 0$. Then $\frac{1}{1} = \frac{0}{1}$, and there exists f^n such that $f^n \cdot 1 = 0 \implies f^n = 0 \implies f$ is a nilpotent.
3. Prime ideals in $R[\frac{1}{f}]$ correspond to prime ideals in R not containing f . Since a non-zero ring has at least one maximal (and hence prime) ideal there is at least one $\mathfrak{p} \subset R$ such that $f \notin \mathfrak{p}$.
4. We only need to show the containment $\bigcap_{\mathfrak{p} \in \text{Spec}(R)} \mathfrak{p} \subseteq \text{Nil}(R)$. Let $f \in \bigcap_{\mathfrak{p} \in \text{Spec}(R)} \mathfrak{p}$ then $f \in \mathfrak{p}$ for all $\mathfrak{p}$. Hence $R[\frac{1}{f}] = 0$. By part (2), this implies that f is nilpotent. Hence,

$$\text{Nil}(R) = \bigcap_{\mathfrak{p} \in \text{Spec}(R)} \mathfrak{p}.$$

5. Suppose $\bar{a} \in R_{\text{red}} = R/\text{Nil}(R)$ is nilpotent, say $\bar{a}^k = 0$ in R_{red} . Then $a^k \in \text{Nil}(R)$, so $(a^k)^\ell = 0$ in R for some ℓ , i.e. $a^{k\ell} = 0$. Thus $a \in \text{Nil}(R)$, which means $\bar{a} = 0$ in R_{red} . Hence the only nilpotent of R_{red} is 0, so R_{red} is reduced.
6. By Problem 3.9, we know that $\eta^\# : \text{Spec}(R[\frac{1}{f}]) \rightarrow D(f)$ is a homeomorphism. If f is nilpotent, then $\text{Spec}(R[\frac{1}{f}]) = \emptyset$ and thus $D(f) = \emptyset$. Similarly, if $D(f) = \emptyset$ then $R[\frac{1}{f}] = 0$, then f is a nilpotent.
7. If f is unit, then $f \notin \mathfrak{p}$ for all $\mathfrak{p}$ hence $D(f) = \text{Spec}(R)$. And if $D(f) = \text{Spec}(R)$ then $f \notin \bigcup_{\mathfrak{p} \in \text{Spec}(R)} \mathfrak{p} \implies f$ is not in any maximal ideal and hence is a unit.
8. We know that $\pi^\#$ is continuous. It is a bijection because ideals in $\text{Spec}(R_{\text{red}})$ correspond to ideals in R containing $\text{Nil}(R)$, by part 4 this is every prime ideal. To see that this map is closed, let $\mathbb{V}(I_{\text{red}})$ be a closed set in R_{red} . Then $\pi^\#(\mathbb{V}(I_{\text{red}})) = \{\mathfrak{p} \mid \pi^{-1}(I_{\text{red}}) \subseteq \mathfrak{p}\} = \{\mathfrak{p} \mid I \subseteq \mathfrak{p}\} = \mathbb{V}(I)$, which is closed in $\text{Spec}(R)$.

9. (a) $\mathbb{K}$ (c) $\mathbb{K}[y]$
(b) $\mathbb{K}$ (d) $\mathbb{Z}/6\mathbb{Z}$

Given a ring homomorphism $\phi : R \rightarrow S$, we know that it induces a continuous map $\phi^\# : \text{Spec}(S) \rightarrow \text{Spec}(R)$ of topological spaces. A natural geometric question is: what is the fibre of this map over a point $[\mathfrak{p}] \in \text{Spec}(R)$? Recall that if $f : X \rightarrow Y$ is a map of topological spaces, then the fibre over a point $y \in Y$ is just the preimage $f^{-1}(\{y\}) = \{x \in X \mid f(x) = y\} \subset X$, considered with the subspace topology. In algebraic geometry, authors often write X_y for the fibre over y when the map f is clear. Returning to the induced map $\phi^\#$, since we know that $\phi^\#([\mathfrak{q}]) = [\phi^{-1}(\mathfrak{q})]$, we can directly describe the fibre as

$$(\phi^\#)^{-1}([\mathfrak{p}]) = \left\{ [\mathfrak{q}] \in \text{Spec}(S) \mid \phi^{-1}(\mathfrak{q}) = \mathfrak{p} \right\} \subset \text{Spec}(S).$$

However, we would like a better, more algebraic description of this set. Answering this involves two conceptual steps: localization and reduction to the residue field. It is worth pausing to try to get an intuitive understanding of each step. As a roadmap: to compute the fibre over $[\mathfrak{p}]$, we first localize at $\mathfrak{p}$, then quotient by $\mathfrak{p} R_{\mathfrak{p}}$; this is what leads naturally to the fibre ring $\kappa(\mathfrak{p}) \otimes_R S$.

Let $f : X \rightarrow Y$ be any continuous map of topological spaces. The fibre over a point $y \in Y$ is local on Y in the sense that it really only depends on the behavior of f at the point y , or perhaps in an open neighborhood of y . More precisely, if $U \subset Y$ contains y and $V = f^{-1}(U)$, then

$$f^{-1}(\{y\}) = (f|_V)^{-1}(\{y\}),$$

where $f|_V : V \rightarrow U$ is the restriction of f . If we wished to be fancy, we might draw a commutative diagram of topological spaces

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ \uparrow \iota & & \uparrow \\ V & \xrightarrow{f|_V} & U \end{array}$$

where ι is the inclusion of V into X . Since $V = f^{-1}(U)$ and $y \in U$, every point of the fibre over y already lies in V , so the displayed equality holds. In other words, restricting f to V does not change the fibre over y ; more generally, we may replace X by any subset that still contains the fibre over y .

Returning to the algebraic situation $\phi^\# : \text{Spec}(S) \rightarrow \text{Spec}(R)$, we happen to know from the previous exercise on localization a very nice subset of $\text{Spec}(R)$ containing the point $[\mathfrak{p}]$, namely

$$[\mathfrak{p}] \in X_U \subset \text{Spec}(R) \quad \text{where} \quad U = R \setminus \mathfrak{p}.$$

Even better, the localization map $\eta : R \rightarrow R_{\mathfrak{p}}$ induces a homeomorphism from $\text{Spec}(R_{\mathfrak{p}})$ onto X_U . We could thus hope to compute the fibre over $[\mathfrak{p}]$ by trying to factor $\phi^\#$ through

$\eta^\#$, i.e. to find a commutative diagram of topological spaces as shown on the left below for some subset $V \subset \text{Spec}(S)$.

$$\begin{array}{ccc}
 \text{Spec}(S) & \xrightarrow{\phi^\#} & \text{Spec}(R) \\
 \uparrow \text{---} & & \uparrow \eta^\# \\
 V & \text{-----} & \text{Spec}(R_{\mathfrak{p}})
 \end{array}
 \qquad
 \begin{array}{ccc}
 R & \xrightarrow{\phi} & S \\
 \downarrow \eta & & \downarrow \text{---} \\
 R_{\mathfrak{p}} & \text{-----} & ??
 \end{array}
 .$$

It turns out we can find such a subset using the universal property of localization. To see why, let us just hope that V can be realized as the spectrum of some ring and that the dashed maps arise from ring homomorphisms. This means we are trying to complete the diagram of rings above on the right. Recall that the image of a multiplicatively closed set is again multiplicatively closed, so we may localize S at

$$T := \phi(R \setminus \mathfrak{p}) \subset S.$$

Let $\eta_S : S \rightarrow T^{-1}S$ be the localization map. We now have a diagram in which the diagonal map is the composition of ϕ and η_S :

$$\begin{array}{ccc}
 R & \xrightarrow{\phi} & S \\
 \downarrow \eta & \searrow & \downarrow \eta_S \\
 R_{\mathfrak{p}} & \xrightarrow{\tilde{\phi}} & T^{-1}S
 \end{array}$$

where the diagonal map is just the composition of ϕ and η_S . Every element of $R \setminus \mathfrak{p}$ maps to a unit in $T^{-1}S$ under the diagonal map, so the universal property of localization gives a unique ring map $\tilde{\phi} : R_{\mathfrak{p}} \rightarrow T^{-1}S$ making the diagram commute. Passing to spectra gives a commutative diagram

$$\begin{array}{ccc}
 \text{Spec}(S) & \xrightarrow{\phi^\#} & \text{Spec}(R) \\
 \uparrow \eta_S^\# & & \uparrow \eta^\# \\
 \text{Spec}(T^{-1}S) & \xrightarrow{\tilde{\phi}^\#} & \text{Spec}(R_{\mathfrak{p}})
 \end{array}$$

and hence the fibre of $\phi^\#$ over $[\mathfrak{p}]$ is naturally identified with the fibre of $\tilde{\phi}^\#$ over the point $[\mathfrak{p} R_{\mathfrak{p}}]$. Since $\eta^\#$ and $\eta_S^\#$ are homeomorphisms onto their images, this reduces the computation of the fibre of $\phi^\#$ over $[\mathfrak{p}]$ to the computation of the fibre of $\tilde{\phi}^\#$ over $[\mathfrak{p} R_{\mathfrak{p}}]$.

Now we would like to understand this latter fibre algebraically. Since $R_{\mathfrak{p}}$ is a local ring with unique maximal ideal $\mathfrak{p} R_{\mathfrak{p}}$, a prime ideal $\mathfrak{q} \subset T^{-1}S$ lies over $\mathfrak{p} R_{\mathfrak{p}}$ if and only if it contains the extended ideal $\mathfrak{p}(T^{-1}S)$. Thus the fibre is exactly the closed subset

$$\mathbb{V}(\mathfrak{p}(T^{-1}S)) \subset \operatorname{Spec}(T^{-1}S),$$

so

$$(\phi^{\#})^{-1}([\mathfrak{p}]) \cong \operatorname{Spec}(T^{-1}S / \mathfrak{p}(T^{-1}S)).$$

At first glance this quotient ring looks a bit mysterious, but tensor products package it very nicely. In fact there is a natural isomorphism $R_{\mathfrak{p}} \otimes_R S \cong T^{-1}S$. The *residue field* at $\mathfrak{p}$ is $\kappa(\mathfrak{p}) := R_{\mathfrak{p}} / \mathfrak{p} R_{\mathfrak{p}}$, which one checks is isomorphic to $\operatorname{Frac}(R / \mathfrak{p})$. (Recall $R / \mathfrak{p}$ is a domain since $\mathfrak{p}$ is a prime ideal.) There is then an isomorphism

$$\kappa(\mathfrak{p}) \otimes_R S \cong (R_{\mathfrak{p}} \otimes_R S) / \mathfrak{p}(R_{\mathfrak{p}} \otimes_R S) \cong T^{-1}S / \mathfrak{p}(T^{-1}S).$$

Therefore the fibre of $\phi^{\#}$ over $[\mathfrak{p}]$ is naturally homeomorphic to $\operatorname{Spec}(\kappa(\mathfrak{p}) \otimes_R S)$. The ring $\kappa(\mathfrak{p}) \otimes_R S$ is called the *fibre ring* of ϕ at $[\mathfrak{p}]$. The next exercise asks you to prove this carefully.

Problem 8. Residue Field.

Let R be a ring. Given a prime ideal $\mathfrak{p} \subset R$, the residue field of R at $\mathfrak{p}$ is the ring $R_{\mathfrak{p}} / \mathfrak{p} R_{\mathfrak{p}}$, which we denote by $\kappa(\mathfrak{p})$.

1. Explain why $\mathfrak{p} R_{\mathfrak{p}}$ is the unique maximal ideal of $R_{\mathfrak{p}}$. Deduce that $\kappa(\mathfrak{p})$ is a field.
2. Show that the composite map

$$R \xrightarrow{\eta} R_{\mathfrak{p}} \xrightarrow{\pi} \kappa(\mathfrak{p})$$

has kernel $\mathfrak{p}$. Deduce that it induces an injective ring map $R / \mathfrak{p} \hookrightarrow \kappa(\mathfrak{p})$.

3. Prove that every nonzero element of $R / \mathfrak{p}$ becomes invertible in $\kappa(\mathfrak{p})$. Conclude that $\kappa(\mathfrak{p}) \cong \text{Frac}(R / \mathfrak{p})$.
4. Use part (3) to compute the residue field $\kappa(\mathfrak{p})$ in each of the following cases. In each case it may be helpful to first identify the quotient ring $R / \mathfrak{p}$:

(a) $R = \mathbb{Z}$ and $\mathfrak{p} = \langle p \rangle$ for a prime p .	(g) $R = \mathbb{K}[x, y]$ and $\mathfrak{p} = \langle x, y \rangle$.
(b) $R = \mathbb{Z}$ and $\mathfrak{p} = \langle 0 \rangle$.	(h) $R = \mathbb{Z}[x]$ and $\mathfrak{p} = \langle p, x \rangle$ for a prime p .
(c) $R = \mathbb{K}[x]$ and $\mathfrak{p} = \langle x - a \rangle$ for $a \in \mathbb{K}$.	(i) $R = \mathbb{Z}[i]$ and $\mathfrak{p} = \langle 1 + i \rangle$.
(d) $R = \mathbb{K}[x]$ and $\mathfrak{p} = \langle 0 \rangle$.	(j) $R = \mathbb{Z}[\sqrt{-5}]$ and $\mathfrak{p} = \langle 2, 1 + \sqrt{-5} \rangle$.
(e) $R = \mathbb{R}[x]$ and $\mathfrak{p} = \langle x^2 + 1 \rangle$.	
(f) $R = \mathbb{K}[x, y]$ and $\mathfrak{p} = \langle y - x^2 \rangle$.	

Solution.

1. Prime ideals of $R_{\mathfrak{p}}$ correspond to prime ideals $\mathfrak{q} \subseteq R$ such that $\mathfrak{q} \subseteq \mathfrak{p}$, since $\mathfrak{p}$ is the largest such ideal it is maximal. $\kappa(\mathfrak{p})$ is a field since quotienting by a maximal ideal gives us a field.
2. Consider $(\pi \circ \eta)^{-1}(0)$.

$$\eta^{-1}(\pi^{-1}(0)) = \eta^{-1}(\mathfrak{p} R_{\mathfrak{p}}) = \mathfrak{p}$$

by primeness. Hence, $\mathfrak{p} = \ker(\pi \circ \eta)$. Then by the first isomorphism theorem,

$$R / \mathfrak{p} \hookrightarrow \kappa(\mathfrak{p})$$

3. Let $0 \neq f \in R / \mathfrak{p}$. Then $\eta(\pi(f)) \neq 0$. Since $\kappa(\mathfrak{p})$ is a field, every element is a unit and hence $\eta(\pi(f))$ is invertible. Thus, $\text{Frac}(R / \mathfrak{p}) \subseteq \kappa(\mathfrak{p})$.

Every element of $\kappa(\mathfrak{p}) = R_{\mathfrak{p}}/\mathfrak{p}R_{\mathfrak{p}}$ is the class of some a/s with $a \in R$, $s \in R \setminus \mathfrak{p}$, and this class equals $\iota(\bar{a})\iota(\bar{s})^{-1}$ with $\bar{s} \neq 0$ (as $s \notin \mathfrak{p}$). That is exactly the image of $\bar{a}/\bar{s} \in \text{Frac}(R/\mathfrak{p})$. Hence there is an isomorphism, $\kappa(\mathfrak{p}) \cong \text{Frac}(R/\mathfrak{p})$.

4. (a) $R/\mathfrak{p} = \mathbb{Z}/\mathfrak{p}\mathbb{Z} = \mathbb{F}_{\mathfrak{p}} = \kappa(\mathfrak{p})$. (g) $R/\mathfrak{p} = \mathbb{K} = \kappa(\mathfrak{p})$.
- (b) $R/\mathfrak{p} = \mathbb{Z}$ and $\kappa(\mathfrak{p}) = \text{Frac}(\mathbb{Z}) = \mathbb{Q}$. (h) $R/\mathfrak{p} = \mathbb{Z}/\mathfrak{p}\mathbb{Z} = \mathbb{F}_{\mathfrak{p}} = \kappa(\mathfrak{p})$.
- (c) $R/\mathfrak{p} = \mathbb{K}[a] = \mathbb{K} = \kappa(\mathfrak{p})$.
- (d) $R/\mathfrak{p} = \mathbb{K}[x]$ and $\kappa(\mathfrak{p}) = \mathbb{K}(x)$. (i) $R/\mathfrak{p} = \mathbb{Z}/2\mathbb{Z} = \mathbb{F}_2 = \kappa(\mathfrak{p})$.
- (e) $R/\mathfrak{p} = \mathbb{R}[i] = \mathbb{C} = \kappa(\mathfrak{p})$. (j) $R/\mathfrak{p} = \mathbb{Z}/2\mathbb{Z}[\sqrt{-5}]/\langle 1 + \sqrt{5} \rangle = \mathbb{F}_2 = \kappa(\mathfrak{p})$.
- (f) $R/\mathfrak{p} = \mathbb{K}[x]$ and $\kappa(\mathfrak{p}) = \mathbb{K}(x)$.

Problem 9. Fibre Ring.

Let $\phi : R \rightarrow S$ be a ring map and fix a point $[\mathfrak{p}] \in \text{Spec}(R)$. Set $U := R \setminus \mathfrak{p}$ and $T := \phi(U) \subset S$. Let $\tilde{\phi} : R_{\mathfrak{p}} \rightarrow T^{-1}S$ be the map induced by the universal property of localization discussed above.

1. Prove that there is a natural isomorphism

$$R_{\mathfrak{p}} \otimes_R S \xrightarrow{\sim} T^{-1}S$$

$$\frac{r}{u} \otimes s \longmapsto \frac{\phi(r)s}{\phi(u)}.$$

2. Tensor the exact sequence

$$\mathfrak{p} R_{\mathfrak{p}} \rightarrow R_{\mathfrak{p}} \rightarrow \kappa(\mathfrak{p}) \rightarrow 0$$

with S over R to obtain an exact sequence

$$(\mathfrak{p} R_{\mathfrak{p}}) \otimes_R S \rightarrow R_{\mathfrak{p}} \otimes_R S \rightarrow \kappa(\mathfrak{p}) \otimes_R S \rightarrow 0.$$

Hint: View S as an R -module via ϕ , and use the right exactness of $- \otimes_R S$, which will be proved in Worksheet 5-1.

3. Show that the image of

$$(\mathfrak{p} R_{\mathfrak{p}}) \otimes_R S \rightarrow R_{\mathfrak{p}} \otimes_R S$$

is exactly the ideal $\mathfrak{p}(R_{\mathfrak{p}} \otimes_R S)$.

4. Deduce that there are natural isomorphisms

$$\kappa(\mathfrak{p}) \otimes_R S \cong (R_{\mathfrak{p}} \otimes_R S) / \mathfrak{p}(R_{\mathfrak{p}} \otimes_R S) \cong T^{-1}S / \mathfrak{p}(T^{-1}S).$$

Solution.

1. From the universal property of localization applied twice we have,

$$\begin{array}{ccc} R & \xrightarrow{\phi} & S \\ \eta \downarrow & \searrow f & \downarrow \\ R_{\mathfrak{p}} & \xrightarrow[\tilde{\phi}]{} & T^{-1}S \end{array}$$

Tensoring by $\otimes_R S$ gives us,

$$\begin{array}{ccc} R \otimes_R S & \xrightarrow{\phi \otimes \text{id}_S} & S \\ \eta \otimes \text{id}_S \downarrow & \searrow f \otimes \text{id}_S & \downarrow \\ R_{\mathfrak{p}} \otimes_R S & \xrightarrow[\tilde{\phi} \otimes \text{id}_S]{} & T^{-1}S \end{array}$$

Let $\tilde{\phi} \otimes \text{id}_S = \psi$.

ψ is injective: We show this for simple tensors. Let $\frac{\phi(r)s}{\phi(u)} = 0$. Then, there exists $t \in T$ such that $t\phi(r)s = 0$. Let $t = \phi(u)$ for some $u \in U$. Then, $\phi(vr)s = 0$. That is, in $R_{\mathfrak{p}} \otimes_R S$,

$$\frac{r}{u} \otimes s = \frac{vr}{vu} \otimes s = \frac{1}{vu} \otimes \phi(vr)s.$$

Since $\phi(vr)s = 0$,

$$\frac{1}{vu} \otimes \phi(vr)s = \frac{1}{vu} \otimes 0 = 0.$$

Therefore

$$\frac{r}{u} \otimes s = 0.$$

Thus $\ker(\psi) = 0$, and so ψ is injective.

ψ is surjective: An element in $T^{-1}S$ is of the form $\frac{s}{t}$, where $t = \phi(u)$ for some $u \in U$. Then,

$$\psi\left(\frac{1}{u} \otimes s\right) = \frac{1}{t} \otimes S = \frac{s}{t}$$

Hence ψ is surjective, and thus $R_{\mathfrak{p}} \otimes_R S \cong T^{-1}S$.

2. Since tensoring is right-exact, and the sequence

$$\mathfrak{p} R_{\mathfrak{p}} \longrightarrow R_{\mathfrak{p}} \longrightarrow \kappa(\mathfrak{p}) \longrightarrow 0$$

is exact, the obtained sequence is also exact,

$$(\mathfrak{p} R_{\mathfrak{p}}) \otimes_R S \xrightarrow{f} R_{\mathfrak{p}} \otimes_R S \xrightarrow{g} \kappa(\mathfrak{p}) \otimes_R S \longrightarrow 0.$$

3. Since the sequence is exact at $R_{\mathfrak{p}} \otimes_R S$, we have $\ker(g) = \text{im}(f)$. From question 8.2 we know that $\ker(g) = (\mathfrak{p} R_{\mathfrak{p}}) \otimes_R S$

Problem 10. Fibres of Spec Maps.

Let $\phi : R \rightarrow S$ be a ring map and fix a point $[\mathfrak{p}] \in \text{Spec}(R)$. Set $U := R \setminus \mathfrak{p}$ and

$T := \phi(U) \subset S$. Let $\tilde{\phi} : R_{\mathfrak{p}} \rightarrow T^{-1}S$ be the map induced by the universal property of localization discussed above.

1. Show that the fibre $(\phi^{\#})^{-1}([\mathfrak{p}])$ is naturally homeomorphic to the fibre of $\tilde{\phi}^{\#} : \text{Spec}(T^{-1}S) \rightarrow \text{Spec}(R_{\mathfrak{p}})$ over the point $[\mathfrak{p} R_{\mathfrak{p}}]$.
2. Let $\mathfrak{q} \in \text{Spec}(T^{-1}S)$. Prove that $\tilde{\phi}^{-1}(\mathfrak{q}) = \mathfrak{p} R_{\mathfrak{p}}$ if and only if $\mathfrak{p}(T^{-1}S) \subseteq \mathfrak{q}$. Hint: Use that $R_{\mathfrak{p}}$ is a local ring with unique maximal ideal $\mathfrak{p} R_{\mathfrak{p}}$.

3. Deduce that

$$(\phi^{\#})^{-1}([\mathfrak{p}]) \cong \mathbb{V}(\mathfrak{p}(T^{-1}S)) \subset \text{Spec}(T^{-1}S),$$

and hence that there is a homeomorphism of topological spaces $(\phi^{\#})^{-1}([\mathfrak{p}]) \cong \text{Spec}(T^{-1}S / \mathfrak{p}(T^{-1}S))$.

4. Use Exercise 9 to conclude that $(\phi^{\#})^{-1}([\mathfrak{p}]) \cong \text{Spec}(\kappa(\mathfrak{p}) \otimes_R S)$.
5. Deduce that the fibre over $[\mathfrak{p}]$ is empty if and only if $\kappa(\mathfrak{p}) \otimes_R S = 0$.

Solution.

1.

Problem 11. Computing Fibres of Spec Maps.

In this problem, we use the previous exercise to compute several examples:

1. Let $\phi : \mathbb{Z} \rightarrow \mathbb{Z}[x]$ be the inclusion. Compute the fibre of the induced map $\phi^{\#} : \text{Spec}(\mathbb{Z}[x]) \rightarrow \text{Spec}(\mathbb{Z})$ over the generic point $\langle 0 \rangle$ and over a closed point $\langle p \rangle$ for $p \in \mathbb{Z}$ prime.
2. Fix an integer $n \geq 2$, and consider the localization map $\eta : \mathbb{Z} \rightarrow \mathbb{Z}[\frac{1}{n}]$. For which points of $\text{Spec}(\mathbb{Z})$ is the fibre with respect to $\eta^{\#}$ empty? Compute the nonempty fibres.
3. Let $\mathbb{K}$ be a field and consider the map

$$\mathbb{K}[t] \rightarrow \mathbb{K}[x, y], \quad t \mapsto x.$$

Compute the fibre of the induced map over $\langle 0 \rangle \in \text{Spec}(\mathbb{K}[t])$ and the fibre over $\langle t - a \rangle$ for $a \in \mathbb{K}$.

4. Consider the inclusion of $\mathcal{C}$ -algebras $\mathcal{C}[t] \rightarrow \mathcal{C}[x, y, t] / \langle xy - t \rangle$ given by $t \mapsto t$.
 - (a) Compute the fibre over the generic point $\langle 0 \rangle$.

- (b) Compute the fibre over $\langle t - a \rangle$ for $a \neq 0$.
- (c) Compute the fibre over $\langle t \rangle$.
- (d) Draw the corresponding map on closed points and compare it with the fibres above.

Solution.

1.