

Math 121: Commutative Algebra

Prishita Dharampal

Worksheet 1.1: Basics of Rings

Problem 1. Let R be a ring and consider $R[[t]]$ the ring of formal power series with coefficients in R . The goal of this exercise is prove that:

$$(R[[t]])^\times = \{u + tf(t) \mid u \in R^\times, f(t) \in R[[t]]\}.$$

- (a) Elements of $R[[t]]$ can be written as $f = \sum_{k=0}^{\infty} f_k t^k$ for $f_k \in R$. We call f_0 the constant (or lead) term of f . If $f = \sum_{k=0}^{\infty} f_k t^k$ and $g = \sum_{k=0}^{\infty} g_k t^k$ then find a formula for the lead term of fg .
- (b) Using the formula from (a) prove that if f is a unit then $f_0 \in R^\times$.
- (c) Find a formula for the coefficient of t^k appearing in fg .
- (d) Assume that $f_0 \in R^\times$. Prove that f is a unit by explicitly constructing an element $g \in R[[t]]$ such that $fg = 1$. (Hint: the condition that $fg = 1$ places constraints on each coefficient of fg . Use the constraints together with part (c) to recursively solve for the coefficients of g .)

Solution.

- (a) Formula for the constant term of fg is $f_0 g_0$.
- (b) Assume f is a unit. Then there exists some $f' \in R[[t]]$ such that $ff' = 1$. By part (1), the constant term of ff' is $f_0 f'_0$. The constant term of 1 is 1, so $f_0 f'_0 = 1$. Hence f_0 is a unit, and $f_0 \in R^\times$.
- (c) The formula for the coefficient of t^k is $\sum_{i=0}^k f_i g_{k-i}$.
- (d) Assume $f_0 \in R^\times$. Let its inverse be g_0 . For f to be a unit, every non-leading coefficient of fg must be 0. That is,

$$fg = f_0 g_0 + \sum_{k=1}^{\infty} \left(\sum_{i=0}^k f_i g_{k-i} \right) t^k$$

where $\sum_{i=0}^k f_i g_{k-i} = 0$ for all k . Then solving for g_k gives us,

$$g_k = -\frac{1}{f_0} \sum_{i=0}^{k-1} f_i g_{k-i} = -g_0 \sum_{i=0}^{k-1} f_i g_{k-i}$$

Problem 2.

- (a) Prove that an element $u \in R$ is a unit if and only if the principal ideal $\langle u \rangle = R$.
- (b) Prove that if ab and b are units, then a is a unit.
- (c) Prove that if u is a unit and $a \in R$, then a and ua generate the same principal ideal.
- (d) In the ring $\mathbb{Z}/\langle 12 \rangle$, determine which elements are units.

Solution.

- (a) ($\implies$) Assume $u \in R$ is a unit.

Then there exists some $u' \in R$ such that $uu' = 1 \in \langle u \rangle$. If 1 is in the ideal generated by u then $\forall r \in R, r \in \langle u \rangle$. Hence $\langle u \rangle = R$.

($\impliedby$) Assume $\langle u \rangle = R$ for some $u \in R$.

Then $1 \in \langle u \rangle$. That is, by definition of an ideal, 1 is a multiple of u . Hence, there exists some $u' \in R$ such that $uu' = 1$. Hence, u is a unit.

- (b) Let $a, b \in R$, and ab, b be units. Then there exist $x, b^{-1} \in R$ such that

$$abx = 1, \quad bb^{-1} = 1$$

Then we have,

$$abx = bb^{-1} \implies \frac{abx}{bb^{-1}} = 1 \implies a \cdot \frac{x}{b^{-1}} = 1$$

Hence, a is a unit and $\frac{x}{b^{-1}}$ is its multiplicative inverse.

- (c) Let $u \in R^\times, a \in R$.

Note that the containment $\langle ua \rangle \subseteq \langle a \rangle$ holds by definition. So we only need to show that $\langle a \rangle \subseteq \langle ua \rangle$. Since u is a unit, there exists some $u^{-1} \in R$ such that $uu^{-1} = 1$. Then, note that $uu^{-1}a = a \in \langle ua \rangle$, i.e. $\langle a \rangle \subseteq \langle ua \rangle$.

Hence a and ua generate the same principal ideal.

- (d) $(\mathbb{Z}/\langle 12 \rangle)^\times = \{1, 5, 7, 11\}$

Problem 3. Given two rings R and S we let $\text{Hom}_{\text{Ring}}(R, S)$ be the set of ring homomorphism from R to S . When there is little room for confusion we will often drop the subscript Ring and simply write $\text{Hom}(R, S)$.

- (a) Prove or disprove: If R and S are rings, then $\text{Hom}(R, S)$ is a ring under point-wise multiplication and point-wise addition.
- (b) Give an explicit description of the hom-sets below:
 - (a) $\text{Hom}(\mathbb{Z}, \mathbb{Z})$
 - (b) $\text{Hom}(\mathbb{Z}/\langle 2 \rangle, \mathbb{Z})$
 - (c) $\text{Hom}(\mathbb{Z}, \mathbb{Z}/\langle 2 \rangle)$
 - (d) $\text{Hom}(\mathbb{Z}, \mathbb{C}[x])$
- (c) Based on part (b) state and prove a characterization of $\text{Hom}(\mathbb{Z}, R)$ for any ring R .

Solution.

- (a) Let $f, g \in \text{Hom}(R, S)$. Note that since ring homomorphisms are unital $f(1_R) = 1_S$ and $g(1_R) = 1_S$. Assume for the sake of contradiction that $\text{Hom}(R, S)$ is a ring under pointwise addition. Then,

$$1 = (f + g)(1) = f(1_R) + g(1_R) = 1 + 1 = 2$$

A contradiction! Hence $\text{Hom}(R, S)$ is not a ring under pointwise addition and pointwise multiplication.

- (b) (i) $\text{Hom}(\mathbb{Z}, \mathbb{Z}) = \{\text{id}\}$
- (ii) $\text{Hom}(\mathbb{Z}/\langle 2 \rangle, \mathbb{Z}) = \emptyset$
- (iii) $\text{Hom}(\mathbb{Z}, \mathbb{Z}/\langle 2 \rangle) = \{\mathbb{Z} \rightarrow \mathbb{Z}/\langle 2 \rangle\}$
- (iv) $\text{Hom}(\mathbb{Z}, \mathbb{C}[x]) = \{\mathbb{Z} \hookrightarrow \mathbb{C}[x]\}$
- (c) Note that since both $\mathbb{Z}$ and R are rings, any homomorphism $\phi : \mathbb{Z} \rightarrow R$ is unital, i.e. $\tau(1) = 1_R$. To be a homomorphism we must have $\forall a \in \mathbb{Z}$,

$$\tau(a) = \tau\left(\sum_{i=1}^a 1\right) = \sum_{i=1}^a \tau(1)$$

This completely determines the homomorphism, and thus there is a unique (unital) homomorphism from $\mathbb{Z}$ to any ring R .

Problem 4. Let $\phi : R \rightarrow S$ be a ring homomorphism.

- (a) Prove that if $J \subset S$ is an ideal then $\phi^{-1}(J) \subset R$ is an ideal.
- (b) Assuming ϕ is injective, give a nice description of $\phi^{-1}(J)$ for an ideal $J \subset S$.
- (c) Find a counterexample showing that if $I \subset R$ is an ideal $\phi(I)$ need not be an ideal.
- (d) Prove that if ϕ is surjective then $\phi(I)$ is an ideal of S .
- (e) Let $I \subset R$ be an ideal. Prove that if $\ker(\phi) \subset I$ then $\phi^{-1}(\phi(I)) = I$.
- (f) Is the assumption that $\ker(\phi) \subset I$ needed in part (5)?

Solution.

- (a) Let $a, b \in \phi^{-1}(J)$. Then $\phi(a), \phi(b) \in J$. Since ideals are closed under addition, $\phi(a) + \phi(b) = \phi(a + b) \in J$. Thus $\phi^{-1}(a + b) \in \phi^{-1}(J)$.
Similarly let $r \in R$, and $a \in \phi^{-1}(J)$. Then $\phi(r) \cdot \phi(a) = \phi(r \cdot a) \in J$. Thus, $\phi^{-1}(r \cdot a) \in \phi^{-1}(J)$.

- (b) If ϕ is injective then,

$$\phi^{-1}(J) = \{r \in R \mid \phi(r) \in J\}$$

$$\phi^{-1}(J) = J \cap \phi(R) = J \cap R$$

- (c) Let $R = \mathbb{Z}$, $S = \mathbb{Z}[x]$ and define the homomorphism to be $\phi : R \hookrightarrow S$. Then consider the image of the ideal generated by 3. It does not contain $3x$ and hence is not closed under multiplication.
- (d) Let $I \subset R$ be an ideal and assume that ϕ is surjective. Let $a, b \in I$. We want to show that $\phi(I)$ is a subgroup of the additive group of S and is closed under multiplication by elements of S .

Consider $\phi(a) - \phi(b)$. Note that since ϕ is a homomorphism the following holds,

$$\phi(a) - \phi(b) = \phi(a - b)$$

and since $a - b \in I$, $\phi(a - b) \in \phi(I)$. Moreover since $\phi(0) = 0 \in \phi(I)$, $\phi(I)$ is a subgroup of S under addition.

Now note that since ϕ is surjective, $\forall s \in S \exists r \in R$ such that $s = \phi(r)$. Then since $a \in I, r \in R$ and I is an ideal of R we have $ra \in I$. So,

$$s \cdot \phi(a) = \phi(r)\phi(a) = \phi(ra) \in \phi(I)$$

Hence, $\phi(I)$ is closed under multiplication by elements of S and thus is an ideal of S .

(e) Assume $\ker(\phi) \subset I$.

$$I \subset \phi^{-1}(\phi(I))$$

Since $\phi^{-1}(\phi(I))$ is the pre-image of $\phi(I)$, this containment is straightforward.

$$\phi^{-1}(\phi(I)) \subseteq I$$

Let $x \in \phi^{-1}(\phi(I))$. That is $\phi(x) = \phi(a)$ for some $a \in I$.

$$\phi(x - a) = 0 \implies x - a \in \ker(\phi) \subset I$$

Since I is closed under addition $x \in I$. Hence, $\phi^{-1}(\phi(I)) \subseteq I$

(f) Yes. As a counterexample, consider the quotient map $\phi : \mathbb{Z} \rightarrow \mathbb{Z}/\langle 3 \rangle$ and the ideal generated by 2, $I = \langle 2 \rangle$. Note that $0 \in \phi(I) \implies \ker(\phi) = 3\mathbb{Z} \subset \phi^{-1}(\phi(I))$. That is $3 \in \phi^{-1}(\phi(I))$ but $3 \notin I$.

Worksheet 1.2: Quotients, Primes, Maximal Ideals

Problem 1. Operations on Ideals.

Throughout this problem let I and J be ideals in a ring R .

- (a) Prove or disprove that $I \cap J$ is an ideal.
- (b) Prove or disprove that $I \cup J$ is an ideal.
- (c) Prove or disprove that the product of ideals, defined below, is an ideal:

$$IJ := \{a_1b_1 + \cdots + a_tb_t \mid a_i \in I, b_i \in J, \text{ for all } t\}.$$

- (d) Prove or disprove that the sum of ideals, defined below, is an ideal:

$$I + J := \{a + b \mid a \in I, b \in J\}.$$

- (e) Prove or disprove that the radical of an ideal, defined below, is an ideal:

$$\sqrt{I} := \{a \in R \mid a^k \in I \text{ for some } k \in \mathbb{Z}_{\geq 1}\}.$$

Solution.

- (a) $I \cap J$ is an ideal.

Let $a, b \in I \cap J, r \in R$.

- (i) $I \cap J$ is a subgroup because intersection of subgroups is a subgroup.
- (ii) Note that $a, b \in I, J$. Since I and J are closed under addition and inverses, $a - b \in I, J$. Hence $a - b \in I \cap J$.
- (iii) $r \cdot a \in I \cap J$ since $ra \in I$ and $ra \in J$.

Hence $I \cap J$ is an additive subgroup of R and is closed under multiplication from elements of R .

- (b) $I \cup J$ is not an ideal.

Consider the ideals $I = \langle 2 \rangle, J = \langle 3 \rangle$ in the ring $R = \mathbb{Z}$. Note that $5 \notin I \cup J$ even though $2, 3 \in I \cup J$. Hence, $I \cup J$ is not closed under addition and is not an ideal.

- (c) IJ is an ideal.

Let $s \in IJ, r \in R$.

- (i) By definition, IJ includes all finite sums of elements in I, J . So it contains 0 and is closed under addition and inverses. Hence it is an additive subgroup.
- (ii) Consider $r \cdot s \in IJ$.

$$r \cdot s = r(a_1b_1 + \cdots + a_tb_t) = r \cdot a_1b_1 + \cdots + r \cdot a_tb_t$$

Note that $r \cdot a_i \in I$ for all i and thus $rs \in IJ$ for all r .

Hence IJ is an additive subgroup of R and is closed under multiplication from elements of R .

- (d) $I + J$ is an ideal.

Let $p = a_1 + b_1, q = a_2 + b_2 \in I + J, r \in R$.

- (i) Note that $0 \in I + J$ since $0 \in I, J$.
- (ii) Consider $p - q$.

$$p - q = a_1 + b_1 - (a_2 + b_2) = (a_1 - a_2) + (b_1 - b_2)$$

Note that $(a_1 - a_2) = a$ for some $a \in I$ and $(b_1 - b_2) = b$ for some $b \in I$. Hence, $I + J$ contains zero and is closed under addition and inverses and thus is an additive subgroup.

- (iii) Consider $r \cdot p$.

$$r \cdot p = r \cdot (a_1 + b_1) = ra_1 + rb_1$$

Note that $ra_1 \in I$ and $rb_1 \in J$ by definition of ideals. Hence $rp \in I + J$.

Thus $I + J$ is an additive subgroup of R and is closed under multiplication from elements of R .

- (e) $\sqrt{I}$ is an ideal.

Let $a^n, b^m \in I$, and $a, b, r \in R$.

- (i) Note that $0 \in \sqrt{I}$ since $0^1 \in I$.
- (ii) Consider $(a - b)^{nm}$.

$$(a - b)^{nm} = \sum_{i=0}^{nm} \binom{nm}{i} (a)^i (-b)^{nm-i}$$

- If $i \geq n$, then $a^n \in I \implies \binom{nm}{i} (a)^i (-b)^{nm-i} \in I$.
- If $i < n$, then $nm - i > nm - n = n(m - 1) = nm - i \geq m$. And $b^m \in I \implies \binom{nm}{i} (a)^i (-b)^{nm-i} \in I$.

Since every term in the sum lies in I , the whole sum lies in I . So $a - b \in \sqrt{I}$.

- (iii) Note that if $a^n \in I$ then $(ra)^n \in I$. Hence $ra \in \sqrt{I}$.

Thus $\sqrt{I}$ is an additive subgroup of R and is closed under multiplication from elements of R .

Problem 2. Let $\mathbb{K}$ be a field. The rings $\mathbb{Z}$, $\mathbb{K}[x]$, and $\mathbb{K}[[t]]$ are all principal ideal domains (PIDs) meaning domains where every ideal is generated by a single element. If $I = \langle a \rangle$ and $J = \langle b \rangle$ give, with proof, a careful concrete description of generators for IJ , $I \cap J$, $I + J$, and $\sqrt{I}$ in each of these rings.

Solution.

Let $I = \langle a \rangle$, $J = \langle b \rangle$. Then, we have

(a) $\mathbb{Z}$

- (i) $IJ = \langle ab \rangle$
- (ii) $I \cap J = \langle \text{lcm}(a, b) \rangle$
- (iii) $I + J = \langle \text{gcd}(a, b) \rangle$
- (iv) Let $a = p_1^{e_1} \cdots p_k^{e_k}$, then $\sqrt{I} = \langle p_1 \cdots p_k \rangle$, where p_i is a prime element.

(b) $\mathbb{K}[x]$

- (i) $IJ = \langle ab \rangle$
- (ii) $I \cap J = \langle \text{lcm}(a, b) \rangle$
- (iii) $I + J = \langle \text{gcd}(a, b) \rangle$
- (iv) Let $a = p_1^{e_1} \cdots p_k^{e_k}$, then $\sqrt{I} = \langle p_1 \cdots p_k \rangle$, where p_i is an irreducible element.

(c) $\mathbb{K}[[t]]$ Note that elements of $\mathbb{K}[[t]]$ are of two forms: $a \in \mathbb{K}$ or $a \cdot t^n$, $a \in \mathbb{K}$. So all proper ideals are of the form $\langle 0 \rangle$ or $\langle t^n \rangle$. Let $I = \langle t^n \rangle$, $J = \langle t^m \rangle$. If I is trivial, then $IJ = I \cap J = \sqrt{I} = \langle 0 \rangle$, and $I + J = J$. Assuming neither I nor J are trivial we have,

- (i) $IJ = \langle t^{n+m} \rangle$
- (ii) $I \cap J = \langle t^{\max(n, m)} \rangle$
- (iii) $I + J = \langle t^{\min(n, m)} \rangle$
- (iv) $\sqrt{I} = \langle t \rangle$.

Universal Properties. Throughout this course we will see numerous constructions and operations (e.g., quotients, tensor products, localizations, completions, etc.) each of these will have an explicit construction where we describe the objects precisely as sets. However, there is an alternative perspective, which is just as useful, describing these objects via their *universal properties*, which roughly asks what makes each object special amongst all other such objects. The standard framework for a universal property is, “Object X is unique up to isomorphism among all objects having property P.”

Problem 3. Universal Property of Quotients.

Let R be a ring and $I \subseteq R$ an ideal. Let $\pi: R \rightarrow R/I$ be the quotient map. Suppose $\varphi: R \rightarrow S$ is a ring homomorphism such that $I \subseteq \ker(\varphi)$. Define a function

$$\tilde{\varphi}: R/I \rightarrow S \quad \text{by} \quad \tilde{\varphi}(r + I) = \varphi(r).$$

- (a) Prove that $\tilde{\varphi}$ is well-defined.
- (b) Prove that $\tilde{\varphi}$ is a ring homomorphism.
- (c) Prove that $\tilde{\varphi} \circ \pi = \varphi$.
- (d) Prove that $\tilde{\varphi}$ is the unique ring homomorphism $R/I \rightarrow S$ with the property that

$$\tilde{\varphi} \circ \pi = \varphi.$$
- (e) Explain in words what this says about the quotient ring R/I .

Solution.

- (a) Let $r + I = r' + I \in R/I$. Then, $r - r' \in I$.

$$\tilde{\varphi}(r + I) = \varphi(r) = \varphi(r + (r - r')) = \varphi(r') = \tilde{\varphi}(r' + I)$$

Hence, $\tilde{\varphi}$ is well-defined.

- (b) Let $\bar{a}, \bar{b} \in R/I$, we check if the three conditions hold.

Unital: Since φ is a ring homomorphism, we have

$$\tilde{\varphi}(1_{R/I}) = \varphi(1_R) = 1_S$$

Addition: Again, since φ is a ring homomorphism, we have

$$\tilde{\varphi}(\bar{a}) + \tilde{\varphi}(\bar{b}) = \varphi(a) + \varphi(b) = \varphi(a + b) = \tilde{\varphi}(\overline{a + b})$$

Multiplication: Again, φ is a ring homomorphism, we have

$$\tilde{\varphi}(\bar{a}) \cdot \tilde{\varphi}(\bar{b}) = \varphi(a) \cdot \varphi(b) = \varphi(a \cdot b) = \tilde{\varphi}(\overline{a \cdot b})$$

Hence, $\tilde{\varphi}$ is a ring homomorphism.

(c) Let $a \in R$. Then,

$$\tilde{\varphi} \circ \pi(a) = \tilde{\varphi}(\bar{a}) = \varphi(a)$$

Since a was an arbitrary element in R , this holds for all elements, and hence $\tilde{\varphi} \circ \pi = \varphi$.

(d) Assume there exists some other homomorphism $\sigma : R/I \rightarrow S$ such that $\sigma \circ \pi = \varphi$. Then for any $\bar{a} \in R/I$,

$$\sigma(\bar{a}) = \sigma \circ \pi(a) = \varphi(a) = \tilde{\varphi}(a)$$

(e) Every homomorphism φ such that $I \subseteq \ker(\varphi)$ can be written as $\tilde{\varphi} \circ \pi$ for some $\tilde{\varphi}$.

Problem 4. In this problem, we wish to consider the evaluation map:

$$\text{ev}_i: \mathbb{Z}[x] \rightarrow \mathbb{C}, \quad f(x) \mapsto f(i)$$

- (a) Prove that ev_i is a ring homomorphism.
- (b) Show that $\langle x^2 + 1 \rangle \subset \ker(\text{ev}_i)$.
- (c) Use Problem 3 to produce a ring homomorphism

$$\widetilde{\text{ev}}_i: \mathbb{Z}[x]/\langle x^2 + 1 \rangle \rightarrow \mathbb{C}.$$

- (d) What does $\widetilde{\text{ev}}_i$ do to the class of x in $\mathbb{Z}[x]/\langle x^2 + 1 \rangle$?
- (e) Prove that $\mathbb{Z}[x]/\langle x^2 + 1 \rangle$ is isomorphic to $\mathbb{Z}[i]$. Is $\langle x^2 + 1 \rangle$ prime and/or maximal?

Solution.

- (a) Let $f, g \in \mathbb{Z}[x]$, we check if the three conditions hold.

Unital: Let f_1 be the constant function 1. Then,

$$\text{ev}_i(f_1(x)) = f_1(i) = 1$$

Addition: Since $\mathbb{Z}[x]$ is a ring under pointwise addition and multiplication, we can write,

$$\text{ev}_i(f(x)) + \text{ev}_i(g(x)) = f(i) + g(i) = (f + g)(i) = \text{ev}_i((f + g)(x))$$

Multiplication: Since $\mathbb{Z}[x]$ is a ring under pointwise addition and multiplication, we can write,

$$\text{ev}_i(f(x)) \cdot \text{ev}_i(g(x)) = f(i) \cdot g(i) = (f \cdot g)(i) = \text{ev}_i((f \cdot g)(x))$$

Hence, ev_i is a ring homomorphism.

- (b) Note that $\text{ev}_i(x^2 + 1) = i^2 + 1 = -1 + 1 = 0$. For any $f(x)(x^2 + 1) \in \langle x^2 + 1 \rangle$ we have $\text{ev}_i(f(x)(x^2 + 1)) = f(i) \cdot 0 = 0 \in \ker(\text{ev}_i)$. Thus the ideal generated by $\langle x^2 + 1 \rangle$ under ev_i maps to 0 and hence is a subset of the kernel.
- (c) From Q3 we know that if $\langle x^2 + 1 \rangle \subseteq \ker(\text{ev}_i)$ there exists a unique homomorphism,

$$\widetilde{\text{ev}}_i: \mathbb{Z}[x]/\langle x^2 + 1 \rangle \rightarrow \mathbb{C}, \quad \widetilde{\text{ev}}_i(\overline{f(x)}) = f(i)$$

(d) $\widetilde{\text{ev}}_i(\bar{x}) = x(i) = i$.

(e) Define a map $\varphi : \mathbb{Z}[x] \rightarrow \mathbb{Z}[i]$, $\varphi(f(x)) \rightarrow f(i)$. We will show that $\ker(\varphi) = \langle x^2 + 1 \rangle$.
 $\langle x^2 + 1 \rangle \subseteq \ker(\varphi)$

Note,

$$\varphi(x^2 + 1) = i^2 + 1 = 0 \in \ker(\varphi)$$

$$\ker(\varphi) \subseteq \langle x^2 + 1 \rangle$$

Let $\varphi(f(x)) \in \ker(\varphi)$. If $f(i) = 0$, then $x^2 + 1 \mid f(x)$ in $\mathbb{Z}[x]$ (since $x^2 + 1$ is the minimal polynomial of i over $\mathbb{Q}$ and hence is primitive over $\mathbb{Z}$), and we have,

$$f(x) \in \langle x^2 + 1 \rangle \implies \ker(\varphi) \subseteq \langle x^2 + 1 \rangle$$

Then by the first isomorphism theorem,

$$\mathbb{Z}[x] / \langle x^2 + 1 \rangle \cong \text{im}(\varphi) = \mathbb{Z}[i]$$

Since $\mathbb{Z}[x]$ is an integral domain but not a field, $\langle x^2 + 1 \rangle$ is prime but not maximal.

Problem 5. Universal Property of Polynomial Rings.

The goal of this exercise is to prove the following characterization of polynomial rings. Let $\varphi: R \rightarrow S$ be a ring homomorphism. For any element $s \in S$ there is a unique ring homomorphism $\psi: R[x] \rightarrow S$ such that $\psi(x) = s$ and the following diagram commutes:

$$\begin{array}{ccc} R & \xrightarrow{\varphi} & S \\ \downarrow & \nearrow \psi & \\ R[x] & & \end{array} .$$

- (a) Prove there is at most one ring homomorphism $\psi: R[x] \rightarrow S$ with the above properties.
- (b) Construct such a homomorphism explicitly by setting

$$\psi\left(\sum_{i=0}^n a_i x^i\right) = \sum_{i=0}^n \varphi(a_i) s^i.$$

- (c) Conclude that to give a ring homomorphism $R[x] \rightarrow S$ is equivalent to giving:
- a ring homomorphism $R \rightarrow S$, and
 - an element of S .

Solution.

- (a) Suppose there exists another homomorphism $\sigma: R[x] \rightarrow S$ such that $\sigma(x) = \psi(x) = s$. Also, note that since the diagram commutes $\psi|_R = \sigma|_R = \varphi$.

Then for any $\sum_{i=0}^n a_i x^i \in R[x]$ we have,

$$\begin{aligned} \sigma\left(\sum_{i=0}^n a_i x^i\right) &= \sigma(a_1) \cdot \sigma(x) + \cdots + \sigma(a_n) \cdot \sigma(x^n) \\ &= \varphi(a_1) \cdot \psi(x) + \cdots + \varphi(a_n) \cdot \psi(x^n) \\ &= \psi(a_1) \cdot \psi(x) + \cdots + \psi(a_n) \cdot \psi(x^n) \\ &= \psi\left(\sum_{i=0}^n a_i x^i\right) \end{aligned}$$

Hence, $\sigma = \psi$.

(b) Let $\psi : R[x] \rightarrow S$ such that

$$\psi(x) = s, \quad \psi(a) = \varphi(a) \quad \forall a \in R$$

(c) Hence for any ring homomorphism $\sigma : R[x] \rightarrow S$ we have

$$\begin{array}{ccc} R & \xrightarrow{\varphi} & S \\ \downarrow & \nearrow \psi & \\ R[x] & & \end{array}$$

Problem 6. Evaluation maps.

Let R be a ring and let $a \in R$. Define the evaluation map

$$\text{ev}_a: R[x] \rightarrow R, \quad f(x) \mapsto f(a).$$

- (a) Prove that ev_a is a surjective ring homomorphism.
- (b) Prove that $\ker(\text{ev}_a) = \langle x - a \rangle$.
- (c) Use Problem 3 to show that ev_a factors through a ring homomorphism

$$R[x]/(x - a) \rightarrow R.$$

- (d) Explain what this means if R is a field or R is an integral domain.
- (e) Give an example of a ring R and prime/maximal ideal $I \subset R[x]$ such that $I \neq \ker(\text{ev}_a)$ for all $a \in R$.

Solution.

- (a) For all $r \in R$ consider the constant functions r represented by $f_r \in R[x]$. Then,

$$\text{ev}_a(f_r(x)) = f_r(a) = r$$

- (b) $\ker(\text{ev}_a) \subseteq \langle x - a \rangle$.

Note that for any $p(x) \in \ker(\text{ev}_a)$ we have

$$p(x) = (x - a)q(x) + r(x)$$

for some $q(x), r(x) \in R[x]$. Evaluating at ev_a we have,

$$\text{ev}_a(p(x)) = (a - a)q(a) + r(a) = 0 \implies r(a) = 0$$

That is

$$p(x) = (x - a)q(x) \implies p(x) \in \langle x - a \rangle$$

$$\langle x - a \rangle \subseteq \ker(\text{ev}_a).$$

Let $p \in \langle x - a \rangle$. Since $\langle x - a \rangle$ irreducible we have $p = f(x)(x - a)$, for some $f(x) \in R[x]$,

$$\text{ev}_a(p) = \text{ev}_a(f(x)(x - a)) = f(a)(a - a) = 0 \in \ker(\text{ev}_a)$$

Hence, $\ker(\text{ev}_a) = \langle x - a \rangle$.

- (c) Since $\ker(\text{ev}_a) \supseteq \langle x - a \rangle$ from problem 3 we know we have an induced ring homomorphism $\widetilde{\text{ev}}_a$ such that

$$\begin{array}{ccc} R[x] & \xrightarrow{\pi} & R[x]/\langle x - a \rangle \\ \text{ev}_a \downarrow & & \nwarrow \widetilde{\text{ev}}_a \\ R & & \end{array}$$

- (d) If R is a domain this means $\langle x - a \rangle$ is prime and also that $\widetilde{\text{ev}}_a = \text{id}$.

If R is a field $\langle x - a \rangle$ is maximal and $R \cong R[x]/\langle x - a \rangle$. Also since all domains are fields we have $\widetilde{\text{ev}}_a = \text{id}$

- (e) $R = \mathbb{C}, I = \langle 0 \rangle$.

Worksheet 1.3: Gröbner Bases Pt. I

Polynomial Division. Recall the following classical theorem:

Theorem 0.1. *Let $g \in \mathbb{K}[x]$ be a non-zero polynomial. Every polynomial $f \in \mathbb{K}[x]$ can be written uniquely as $f = qg + r$ where $q, r \in \mathbb{K}[x]$, and either $r = 0$ or $\deg(r) < \deg(g)$.*

While this theorem might seem innocuous when first learned as a teenager, it turns out to be very powerful when studying ideals and polynomials in $\mathbb{K}[x]$. The proof of this theorem also provides an algorithm for computing q and r , which when presented in pseudocode looks something like

Algorithm 1 Division Algorithm

Require: g, f

Ensure: q, r

```
 $q \leftarrow 0; r \leftarrow f$   
while  $r \neq 0$  and  $\text{LT}(g)$  divides  $\text{LT}(r)$  do  
     $q \leftarrow q + \text{LT}(r)/\text{LT}(g)$   
     $r \leftarrow r - (\text{LT}(r)/\text{LT}(g))g$   
end while
```

The goal of this worksheet is to begin exploring how we might extend this theorem to polynomial rings in more variables. Along the way we will see how these explorations will lead to both computational tools for studying the commutative algebra of polynomial rings over fields, but also gives us useful tools for proving theorems in this setting as well.

Problem 1. Use Theorem 1 to prove the following:

- (a) If $f \in \mathbb{K}[x]$ is a non-zero polynomial then f has at most $\deg(f)$ roots in $\mathbb{K}$.
- (b) If $g \in \mathbb{K}[x]$ is a non-zero polynomial then $f \in \langle g \rangle$ if and only if $r = 0$.

Solution.

- (a) Assume for the sake of contradiction that f has more than $n = \deg(f)$ roots, say m roots: $a_1, \dots, a_m$. Then by Theorem 1, dividing by $(x - a_i)$ for all $1 \leq i \leq m$ we get $f = c(x - a_1) \cdots (x - a_m)$, for some constant c and $r = 0$. But then we have m x terms, i.e., x^m is in f , but since $\deg(f) < m$ this is not possible. Hence a contradiction.

- (b) ($\implies$)

Assume $f \in \langle g \rangle$. Then there exists some $q \in \mathbb{K}[x]$ such that $f = qg$. Then,

$$f = qg + r \implies f - qg = r = 0$$

- ($\impliedby$)

Assume $r = 0$. Then by Theorem 1 we have,

$$f = qg + 0 = qg$$

That is there exists some $q \in \mathbb{K}[x]$ such that $f = qg$ and by definition of an ideal, $f \in \langle g \rangle$.

Problem 2. Implement the algorithm above in Macaulay2, and use your code to compute q and r for the following polynomials.

(a) $f = x^3 + x + 1, \quad g = x + 1.$

(d) $f = x^{100} - 1, \quad g = x^2 - 1.$

(b) $f = x^5 - 1, \quad g = x - 1.$

(e) $f = x^{1000} - 1, \quad g = x - 1.$

(c) $f = x^2 + 1, \quad g = x^3 + x.$

(f) $f = \sum_{i=0}^{500} x^i, \quad g = x^2 + x + 1.$

Solution.

(a) $(x^2 - x + 2, -1)$

(d) $\left(\sum_{k=0}^{49} x^{2k}, 0\right)$

(b) $(x^4 + x^3 + x^2 + 1, 0)$

(e) $\left(\sum_{k=0}^{99} x^k, 0\right)$

(c) $(0, x^2 + 1)$

(f) $\left(\sum_{k=0}^{166} x^{3k}, 0\right)$

Definition 0.2. A monomial ordering on $\mathbb{K}[x_1, \dots, x_n]$ is a relation $<$ on $\mathbb{Z}_{\geq 0}^n$ satisfying the following:

- (i) $<$ is a total ordering on $\mathbb{Z}_{\geq 0}^n$,
- (ii) if $\alpha < \beta$ and $\gamma \in \mathbb{Z}_{\geq 0}^n$ then $\alpha + \gamma < \beta + \gamma$, and
- (iii) $<$ is a well-ordering on $\mathbb{Z}_{\geq 0}^n$.

Recall that a *total ordering* means that of any two elements $\alpha, \beta \in \mathbb{Z}_{\geq 0}^n$ exactly one of the following statements is true: $\alpha < \beta$, $\alpha = \beta$, or $\beta < \alpha$. That $<$ is a *well-ordering* on $\mathbb{Z}_{\geq 0}^n$ means that every non-empty subset of $\mathbb{Z}_{\geq 0}^n$ has a smallest element with respect to $<$.

Once a monomial ordering has been fixed, every non-zero polynomial acquires a distinguished piece of data: its leading monomial, leading coefficient, and leading term. These are the ingredients that drive the multivariable division process just as degree and leading coefficient drive ordinary polynomial division.

Problem 3. Prove that a monomial ordering $<$ on $\mathbb{Z}_{\geq 0}^n$ is equivalent to a relation $<$ on the set of monomials in $K[x_1, \dots, x_n]$ satisfying analogous conditions to (i), (ii), and (iii). Explain why this justifies the moniker “monomial ordering”.

Solution.

There exists a bijection

$$\mathbb{Z}_{\geq 0}^n \ni \{\alpha = (a_1, \dots, a_n)\} \leftrightarrow \{x^\alpha = x_1^{a_1} \cdots x_n^{a_n}\} \in K[x_1, \dots, x_n]$$

that satisfies all the conditions because addition of vectors corresponds to addition of exponents during multiplication.

Problem 4. Prove or disprove that the following are monomial orderings:

(a) Lexicographic (Lex) Order: Given $\alpha = (\alpha_1, \dots, \alpha_n)$ and $\beta = (\beta_1, \dots, \beta_n)$ in $\mathbb{Z}_{\geq 0}^n$, we say $\beta <_{\text{lex}} \alpha$ if and only if the left-most non-zero entry of $\alpha - \beta$ is positive.

(b) Graded Lex Order: $\beta <_{\text{grlex}} \alpha$ if and only if

$$|\beta| = \sum_{i=1}^n \beta_i < |\alpha| = \sum_{i=1}^n \alpha_i$$

or $|\beta| = |\alpha|$ and $\beta <_{\text{lex}} \alpha$.

(c) Reverse Lex (RevLex) Order: $\beta <_{\text{revlex}} \alpha$ if and only if the right-most non-zero entry of $\alpha - \beta$ is negative.

(d) Graded RevLex Order: $\beta <_{\text{grevlex}} \alpha$ if and only if

$$|\beta| = \sum_{i=1}^n \beta_i < |\alpha| = \sum_{i=1}^n \alpha_i$$

or $|\beta| = |\alpha|$ and $\beta <_{\text{revlex}} \alpha$.

(e) Weight Order: Fix $w = (w_1, \dots, w_n) \in \mathbb{Z}^n$. Define $\beta <_w \alpha$ if and only if

$$\beta \cdot w = \sum_{i=1}^n \beta_i w_i < \alpha \cdot w = \sum_{i=1}^n \alpha_i w_i.$$

Solution.

(a) It is a monomial ordering. Let $\alpha = (\alpha_1, \dots, \alpha_n)$, $\beta = (\beta_1, \dots, \beta_n) \in \mathbb{Z}_{\geq 0}^n$.

(i) Total Ordering:

If $\alpha = \beta$ we're done. Otherwise, $\alpha - \beta$ has a leftmost non-zero entry. If this entry is positive, $\beta <_{\text{lex}} \alpha$, else $\alpha <_{\text{lex}} \beta$.

(ii) Let $\alpha <_{\text{lex}} \beta$, that is the leftmost non-zero entry of $\alpha - \beta$ is negative. Then for all $\gamma \in \mathbb{Z}_{\geq 0}^n$ we have,

$$(\alpha + \gamma) - (\beta + \gamma) = \alpha - \beta$$

That is the leftmost non-zero entry remains unchanged and hence $(\alpha + \gamma) <_{\text{lex}} (\beta + \gamma)$.

- (iii) Let $S \subseteq \mathbb{Z}_{\geq 0}^n$ be non-empty. Since $\mathbb{Z}_{\geq 0}$ is well-ordered we let $m_1 = \min\{\alpha_1 \mid \alpha \in S\}$. Then, among the elements with first coordinate m_1 we let $m_2 = \min\{\alpha_2 \mid \alpha \in S\}$ and so on for all n coordinates. This gives us,

$$m = (m_1, \dots, m_n) \in S$$

which is the lex-minimal element.

- (b) It is a monomial ordering. Let $\alpha = (\alpha_1, \dots, \alpha_n), \beta = (\beta_1, \dots, \beta_n) \in \mathbb{Z}_{\geq 0}^n$.

- (i) Total Ordering:

If $\alpha = \beta$ we're done. Else if $|\alpha| = |\beta|$, by part (a) we have an ordering. Otherwise, $|\alpha| \neq |\beta|$ has a leftmost non-zero entry. If $|\alpha| > |\beta|$, $\beta <_{\text{grlex}} \alpha$, else $\alpha <_{\text{grlex}} \beta$.

- (ii) Let $\alpha <_{\text{grlex}} \beta$, and $\gamma \in \mathbb{Z}_{\geq 0}^n$.

- If $|\alpha| < |\beta|$, we have

$$|\alpha| + |\gamma| < |\beta| + |\gamma| \implies |\alpha + \gamma| < |\beta + \gamma|$$

- If $|\alpha| = |\beta|$ and $\alpha <_{\text{lex}} \beta$ we have an ordering by part (a).

- (iii) Let $S \subseteq \mathbb{Z}_{\geq 0}^n$ be non-empty. Since all entries are non-negative there exists $m = \min\{|\alpha| \mid \alpha \in S\}$. Then, we take the lex-minimal element in the subset $H \subseteq S, H = \{|\alpha| = m \mid \alpha \in S\}$. This is the grlex-minimal element.

- (c) It is a monomial ordering. Let $\alpha = (\alpha_1, \dots, \alpha_n), \beta = (\beta_1, \dots, \beta_n) \in \mathbb{Z}_{\geq 0}^n$.

- (i) Total Ordering:

If $\alpha = \beta$ we're done. Otherwise, $\alpha - \beta$ has a rightmost non-zero entry. If this entry is negative, $\beta <_{\text{revlex}} \alpha$, else $\alpha <_{\text{revlex}} \beta$.

- (ii) Let $\alpha <_{\text{revlex}} \beta$, that is the leftmost non-zero entry of $\alpha - \beta$ is positive. Then for all $\gamma \in \mathbb{Z}_{\geq 0}^n$ we have,

$$(\alpha + \gamma) - (\beta + \gamma) = \alpha - \beta$$

That is the rightmost non-zero entry remains unchanged and hence $(\alpha + \gamma) <_{\text{revlex}} (\beta + \gamma)$.

- (iii) Let $S \subseteq \mathbb{Z}_{\geq 0}^n$ be non-empty. Since $\mathbb{Z}_{\geq 0}$ is well-ordered we let $m_n = \min\{\alpha_n \mid \alpha \in S\}$. Then, among the elements with first coordinate m_n we let $m_{n-1} = \min\{\alpha_{n-1} \mid \alpha \in S\}$ and so on for all n coordinates. This gives us,

$$m = (m_1, \dots, m_n) \in S$$

which is the revlex-minimal element.

- (d) It is a monomial ordering by (b) and (c).

- (e) Not a monomial ordering. For instance, if $w = (1, 1), \alpha = (0, 2), \beta = (2, 0)$ we have $\beta \cdot w = \alpha \cdot w$ but $\alpha \neq \beta$.

Problem 5. Leading Terms.

Fix a monomial ordering $<$ on $\mathbb{Z}_{\geq 0}^n$. Let $f = \sum_{\alpha} c_{\alpha} x^{\alpha} \in K[x_1, \dots, x_n]$ be a non-zero polynomial. Define the leading monomial $\text{LM}(f)$, leading coefficient $\text{LC}(f)$, and leading term $\text{LT}(f)$.

- (a) Compute $\text{LM}(f)$, $\text{LC}(f)$, and $\text{LT}(f)$ for

$$f = x^3 + 3x^2y + 5xy^3 - 7z^2 \in K[x, y, z]$$

with respect to Lex, Graded Lex, and Graded RevLex (assume $x > y > z$).

- (b) Prove that if $f, g \in K[x_1, \dots, x_n]$ are non-zero, then

$$\text{LM}(fg) = \text{LM}(f)\text{LM}(g), \quad \text{LT}(fg) = \text{LT}(f)\text{LT}(g).$$

- (c) Explain which property of a monomial ordering is used in part (b).

Solution.

- (a) • Lex:

$$\text{LM}(f) = x^3, \quad \text{LC}(f) = 1, \quad \text{LT}(f) = x^3$$

- Graded Lex:

$$\text{LM}(f) = xy^3, \quad \text{LC}(f) = 5, \quad \text{LT}(f) = 5xy^3$$

- Graded RevLex:

$$\text{LM}(f) = xy^3, \quad \text{LC}(f) = 5, \quad \text{LT}(f) = 5xy^3$$

- (b) Let $f = \text{LT}(f) + \sum_{\alpha < \text{LM}(f)} c_{\alpha} x^{\alpha}$, $g = \text{LT}(g) + \sum_{\beta < \text{LM}(g)} d_{\beta} x^{\beta} \in K[x_1, \dots, x_n]$ be non-zero. Then the product fg is,

$$fg = \sum_{\alpha, \beta} c_{\alpha} d_{\beta} x^{\alpha + \beta}$$

Note that $\text{LT}(f)\text{LT}(g) = c_0 d_0 x^{\alpha_0 + \beta_0}$ appears in fg . We know for all α s and β s, $\alpha < \alpha_0$ and $\beta < \beta_0$ so

$$\alpha + \beta < \alpha_0 + \beta_0$$

Hence $x^{\alpha + \beta} < x^{\alpha_0 + \beta_0}$, and $x^{\alpha + \beta} = \text{LM}(f)\text{LM}(g) = \text{LM}(fg)$. Moreover $\text{LT}(fg) = \text{LC}(fg)\text{LM}(fg)$, where $\text{LC}(fg) = \text{LC}(f)\text{LC}(g)$ by definition, and we showed $\text{LM}(fg) = \text{LM}(f)\text{LM}(g)$. Hence $\text{LT}(fg) = \text{LT}(f)\text{LT}(g)$.

- (c) (ii) property is used.

Problem 6. Let $G = (g_1, \dots, g_s)$ be an ordered list of non-zero polynomials in $K[x_1, \dots, x_n]$.

- (a) Write pseudocode for a division algorithm which takes as input f and G and returns polynomials $q_1, \dots, q_s, r$ such that

$$f = q_1g_1 + \dots + q_sg_s + r,$$

where no monomial in r is divisible by any $\text{LM}(g_1), \dots, \text{LM}(g_s)$.

- (b) In the algorithm above, what should happen if the leading monomial of the current polynomial is not divisible by any $\text{LM}(g_i)$?
- (c) Prove that your algorithm terminates. What plays the role of “degree” in the proof?

Solution.

- (a) **Require:** $f, g_1, \dots, g_s$

Ensure: $q_1, \dots, q_s, r$

$q_1, \dots, q_s \leftarrow 0; r \leftarrow 0; p \leftarrow f$

while $p \neq 0$ **do**

\leftarrow 0

for $i = 1, \dots, s$ **do**

if $\text{LT}(g_i)$ divides $\text{LT}(p)$ **then**

$q_i \leftarrow q_i + \text{LT}(p) / \text{LT}(g_i)$

$p \leftarrow p - (\text{LT}(p) / \text{LT}(g_i))g_i$

\leftarrow 1

break

end if

end for

if $\text{divides} = 0$ **then**

$r \leftarrow r + \text{LT}(p)$

$p \leftarrow p - \text{LT}(p)$

end if

end while

- (b) We add it to the remainder term r , and continue.

- (c) $\text{LM}(f)$ plays the role of degree. At every step, we update either $p \rightarrow p - \text{LT}(p)$ (if no g_i divides $\text{LM}(p)$) or $p \rightarrow \text{LT}(p) / \text{LT}(g_i) \cdot g_i$ (for some g_i dividing $\text{LM}(p)$). So the leading term strictly decreases. Hence, the program must terminate.

Problem 7. Work in $K[x, y]$ with Lex order and $x > y$. Let $f = xy^2 - x$, $g_1 = xy - 1$, and $g_2 = y^2 - 1$.

- (a) Divide f by the ordered list (g_1, g_2) .
- (b) Divide f by the ordered list (g_2, g_1) .
- (c) Show directly that $f \in \langle g_1, g_2 \rangle$.
- (d) What do your computations show about division in several variables? In particular, is the remainder uniquely determined by the ideal $\langle g_1, g_2 \rangle$?

Solution.

- (a) $q_1 = y, q_2 = 0, r = -x + y$
- (b) $q_1 = 0, q_2 = x, r = 0$
- (c) $x \in K[x, y]$ and $f = g_2 \cdot x$, hence $f \in \langle g_2 \rangle \subseteq \langle g_1, g_2 \rangle$.
- (d) Remainders are not unique.

Worksheet 2.1: Zariski Topology

Problem 1. Recall that the spectrum and maximal spectrum of a ring R are the sets:

$$\text{Spec}(R) := \{P \subset R \mid P \text{ is a prime ideal}\}$$

and

$$\text{mSpec}(R) := \{P \subset R \mid P \text{ is a maximal ideal}\}$$

Given any subset $\mathcal{S} \subset R$ define

$$\mathbb{V}(\mathcal{S}) := \{P \in \text{Spec}(R) \mid \mathcal{S} \subset P\}.$$

- (a) Describe $\text{Spec}(\mathbb{K})$ and $\text{mSpec}(\mathbb{K})$ when $\mathbb{K}$ is a field.
- (b) Describe both $\text{Spec}(\mathbb{K}[x]/\langle x^3 \rangle)$ and $\text{mSpec}(\mathbb{K}[x]/\langle x^3 \rangle)$ when $\mathbb{K}$ is a field.
- (c) For an arbitrary ring R , compute $\mathbb{V}(\{0\})$ and $\mathbb{V}(\{1\})$.
- (d) Explain the natural poset structure on $\text{Spec}(R)$. Describe the poset for $\text{Spec}(\mathbb{Z})$.

Solution.

- (a) Let $\mathbb{K}$ be a field. Then $\mathbb{K}$ only has 1 proper ideal: the ideal generated by 0. Note that this ideal is prime because $\mathbb{K}$ is an integral domain. Moreover it is also maximal because $\mathbb{K}/\langle 0 \rangle = \mathbb{K}$ is a field. So,

$$\text{Spec}(\mathbb{K}) = \text{mSpec}(\mathbb{K}) = \{\langle 0 \rangle\}$$

- (b) Note that the ideal generated by 0 is not prime in this ring because $x \cdot x^2 \in \langle 0 \rangle$ even though neither x nor x^2 are in $\langle x \rangle$. Then we get,

$$\text{Spec}(\mathbb{K}[x]/\langle x^3 \rangle) = \text{mSpec}(\mathbb{K}[x]/\langle x^3 \rangle) = \{\langle x \rangle\}$$

- (c) Let R be an arbitrary ring, then,

$$\mathbb{V}(\{0\}) = \text{Spec}(R)$$

since every ideal contains 0 and,

$$\mathbb{V}(\{1\}) = \emptyset$$

since 1 is not contained in any proper ideal of R .

- (d) The natural poset structure on $\text{Spec}(R)$ is given by inclusion of ideals, that is, for $P_1, P_2 \in \text{Spec}(R)$, we write $P_1 \leq P_2$ iff $P_1 \subseteq P_2$.

The poset for $\text{Spec}(\mathbb{Z})$ contains prime ideals in $\mathbb{Z}$. The zero ideal is contained in every other ideal and every other prime ideal is maximal.

A *topological space* is a set X together with a collection of subsets τ , which satisfies the following conditions: i) $\emptyset \in \tau$ and $X \in \tau$, ii) τ is closed under arbitrary unions, and iii) τ is closed under finite intersections. We call τ a *topology* on X , the elements of τ the open subsets of X , and a subset $Z \subset X$ is closed if and only if $X \setminus Z$ is open (i.e., $X \setminus Z \in \tau$). If $S \subset X$ is a subset, the *subspace topology* on S is the topology where the open sets of S are all sets of the form $S \cap U \subset S$ where $U \subset X$ is open in X . A map $f : X \rightarrow Y$ of topological spaces is *continuous* if and only if $f^{-1}(U) \subset X$ is open for all open subsets $U \subset Y$, i.e., the preimage of an open set is open.

Problem 2. The Zariski Topology.

We will now place a topology on the set $\text{Spec}(R)$, called the Zariski topology after Oscar Zariski.

- (a) Prove that if $\mathcal{S}, \mathcal{T} \subset R$ are arbitrary sets then $\mathbb{V}(\mathcal{S}) \cap \mathbb{V}(\mathcal{T}) = \mathbb{V}(\mathcal{S} \cup \mathcal{T})$. Explain why your argument extends to an *arbitrary* intersection.
- (b) Prove that if $\mathcal{S}, \mathcal{T} \subset R$ are arbitrary sets then $\mathbb{V}(\mathcal{S}) \cup \mathbb{V}(\mathcal{T}) = \mathbb{V}(\mathcal{S} \cdot \mathcal{T})$ where

$$\mathcal{S} \cdot \mathcal{T} := \{st \mid s \in \mathcal{S}, t \in \mathcal{T}\}.$$

- (c) Prove that the sets of the form $\mathbb{V}(\mathcal{S})$ are the closed sets of a topology on $\text{Spec}(R)$. We call this topology the Zariski topology on $\text{Spec}(R)$. (Hint: Rephrase the definition of a topology in terms of closed sets instead of open sets.)
- (d) Take a break and Google Oscar Zariski and read about his life.

Solution.

- (a) We prove this for arbitrary intersections. Let $\mathcal{S}_i, i \in \Lambda$ where Λ is some indexing set, be arbitrary subsets of R . Then by definition,

$$\bigcap_{i \in \Lambda} \mathbb{V}(\mathcal{S}_i) = \bigcap_{i \in \Lambda} \{[P] \in \text{Spec}(R), \mathcal{S}_i \subseteq P\} = \left\{ [P] \in \text{Spec}(R), \bigcup_{i \in \Lambda} \mathcal{S}_i \subseteq P \right\}$$

And

$$\mathbb{V}\left(\bigcup_{i \in \Lambda} \mathcal{S}_i\right) = \left\{ [P] \in \text{Spec}(R), \bigcup_{i \in \Lambda} \mathcal{S}_i \subseteq P \right\}$$

That is, $\bigcap_{i \in \Lambda} \mathbb{V}(\mathcal{S}_i) = \mathbb{V}\left(\bigcup_{i \in \Lambda} \mathcal{S}_i\right)$. Hence proved.

- (b) Let $\mathcal{S}, \mathcal{T}$ be arbitrary subsets of R . Then by definition,

$$\mathbb{V}(\mathcal{S} \cdot \mathcal{T}) = \{[P] \in \text{Spec}(R), \mathcal{S} \cdot \mathcal{T} \subseteq P\}$$

And

$$\mathbb{V}(\mathcal{S}) \cup \mathbb{V}(\mathcal{T}) = \{[P] \in \text{Spec}(R), \mathcal{S} \subseteq P\} \cup \{[P] \in \text{Spec}(R), \mathcal{T} \subseteq P\}$$

Note that $\mathcal{S} \cdot \mathcal{T} \subset \mathcal{S}, \mathcal{T}$. Then the containment $\mathbb{V}(\mathcal{S} \cdot \mathcal{T}) \subseteq \mathbb{V}(\mathcal{S}) \cup \mathbb{V}(\mathcal{T})$ is obvious.

To prove that $\mathbb{V}(\mathcal{S} \cdot \mathcal{T}) \supseteq \mathbb{V}(\mathcal{S}) \cup \mathbb{V}(\mathcal{T})$, it suffices to show that if $\mathcal{S} \cdot \mathcal{T} \subseteq P$ then either $\mathcal{S} \subseteq P$ or $\mathcal{T} \subseteq P$.

Assume $\mathcal{S} \cdot \mathcal{T} \subseteq P$. Let $s \in \mathcal{S}$ and $t \in \mathcal{T}$. Then, since P is a prime ideal and $st \in P$ then either $s \in P$ or $t \in P$. If both $s, t \in P$ we are done. Else wlog, assume that $s \notin P$. But note that $\forall t' \in \mathcal{T}, s \cdot t' \in P$. Hence, $t' \in P, \forall t' \in \mathcal{T}$, i.e. $\mathcal{T} \subseteq P$.

Hence, $\mathbb{V}(\mathcal{S}) \cup \mathbb{V}(\mathcal{T}) = \mathbb{V}(\mathcal{S} \cdot \mathcal{T})$.

(c) To prove that the sets of the form $\mathbb{V}(\mathcal{S})$ are the closed sets of the Zariski topology on $\text{Spec}(R)$ we must show that (i) $\emptyset, \text{Spec}(R)$ are closed sets, (ii) the finite union of closed sets is closed, and (iii) arbitrary intersection of closed sets is closed.

(i) Note that $\emptyset = \{\mathbb{V}(\{1\})\}$ since 1 is not contained in any proper ideal of R , and $\text{Spec}(R) = \{\mathbb{V}(\{0\})\}$ since 0 is contained in every ideal and thus in every prime ideal. Both $\mathbb{V}(\{1\})$ and $\mathbb{V}(\{0\})$ are closed sets, hence $\emptyset$ and $\text{Spec}(R)$ are closed.

(ii) From part (b) we know that the finite union of two closed sets is closed, and by induction we can extend this to a finite number of closed sets. Hence, the finite union of closed sets is closed.

(iii) From part (a) we know that any arbitrary intersection of closed sets is closed.

Hence, sets of the form $\mathbb{V}(\mathcal{S})$ are the closed sets of the Zariski topology on $\text{Spec}(R)$.

Problem 3. Describe the closed subsets of $\text{mSpec}(R) \subset \text{Spec}(R)$ in the subspace topology.

Solution.

Closed subsets of $\text{mSpec}(R)$ look like $\mathbb{V}(\mathcal{S}) \cap \text{mSpec}(R)$, for all $\mathbb{V}(\mathcal{S})$ in the Zariski Topology on $\text{Spec}(R)$. That is, the closed subsets of $\text{mSpec}(R)$ are maximal ideals M such that $\mathcal{S} \subseteq M$.

Problem 4. Let $\mathbb{K}$ be an algebraically closed field. This exercise will describe the Zariski topology on $\text{Spec}(\mathbb{K}[x])$.

- (a) Describe the points of $\text{Spec}(\mathbb{K}[x])$. (Hint: There are two types.)
- (b) Prove that if $S \subset \mathbb{K}[x]$ then $\mathbb{V}(S) = \mathbb{V}(\{f\})$ where f is the greatest common divisor of all elements of S .
- (c) Show that the subspace topology on $\text{mSpec}(\mathbb{K}[x])$ is the finite complement topology. (Recall the *finite complement topology* on a set X is defined by letting the open sets be the empty set and sets of the form $X \setminus T$ for $T \subset X$ a subset of finite cardinality.)

Solution.

- (a) Points of $\text{Spec}(\mathbb{K}[x]) = \{\langle 0 \rangle\} \cup \{\langle x - a \rangle, a \in \mathbb{K}\}$.
- (b) Note that since every element $s \in S$ is a multiple of the $\text{gcd}(S)$, the containment $\mathbb{V}(S) \subseteq \mathbb{V}(\{f\})$ is obvious. To show the opposite containment we use the following algorithm:

Require: A set S

Ensure: $\text{gcd}(S)$

Pick $a_1, a_2 \in S$; set $d \leftarrow \text{gcd}(a_1, a_2)$

while there exists $a \in S$ such that $d \nmid a$ **do**

 Pick such an $a \in S$

$d \leftarrow \text{gcd}(d, a)$

end while

return d

Then, there must exist a_i, a_j such that $\text{gcd}(a_i, a_j) = d$. Then by Bezout's Identity, we have $d \in \langle S \rangle \implies \langle d \rangle \subseteq \langle p \rangle$ for all prime ideals containing S .

Hence, $\mathbb{V}(S) = \mathbb{V}(\{f\})$.

- (c) Let C be a closed set in $\text{mSpec}(\mathbb{K}[x])$. Then we know $C = \mathbb{V}(I)$ for some $I \subseteq \mathbb{K}[x]$. Since $\mathbb{K}[x]$ is a PID every ideal is of the form $I = \langle f \rangle$ for some $f \in \mathbb{K}[x]$.
 - If $f = 0$ then $\mathbb{V}(f) = \text{mSpec}(\mathbb{K}[x])$.
 - If $f \neq 0$, since $\mathbb{K}$ is algebraically closed, $f = \prod_{i=1}^n (x_i - a_i)$ for finitely many factors. Moreover since $\mathbb{K}[x]$ is a PID over an algebraically closed field maximal ideals of $\mathbb{K}[x]$ correspond to irreducible polynomials, which are of the form $\langle x - a \rangle$. So, $\langle f \rangle \subseteq \langle x_i - a_i \rangle$ for its finitely many factors.

Hence, $\mathbb{V}(I)$ is finite and the closed sets are exactly $\text{mSpec}(\mathbb{K}[x])$ itself and finite subsets, which is exactly the finite complement topology.

Problem 5. Dense Points.

If X is a topological space we say that a set $S \subset X$ is *dense* if $U \cap S \neq \emptyset$ for every non-empty open subset $U \subset X$.

- (a) Let p be a point in a topological space X . Prove that $\{p\}$ is a dense subset of X if and only if the smallest closed subset which contains $\{p\}$ is X . We call such points *dense points* of X . (Note: no point in $\mathbb{R}^n$ or $\mathbb{C}^n$ with the Euclidean topology is dense in this sense.)
- (b) Given a prime ideal $P \subset R$ we will frequently write $[P]$ to denote the corresponding point in $\text{Spec}(R)$. Prove that a point $[P] \in \text{Spec}(R)$ is dense if and only if $V(P) = \text{Spec}(R)$.
- (c) Characterize all of the dense points in the following:
 - (i) $\text{Spec}(\mathbb{K}[x])$
 - (ii) $\text{Spec}(\mathbb{Z})$
 - (iii) $\text{Spec}(\mathbb{Z}[t])$
 - (iv) $\text{Spec}(\mathbb{K}[x]/\langle x^2 \rangle)$
 - (v) $\text{Spec}(\mathbb{K}[x, y])$
 - (vi) $\text{Spec}(\mathbb{Z}/\langle 6 \rangle)$.
 - (vii) $\text{Spec}(\mathbb{Z}/\langle 4 \rangle)$.
 - (viii) $\text{Spec}(\mathbb{Z}/\langle 12 \rangle)$.
- (d) Prove that if R is a domain then $\text{Spec}(R)$ has a unique dense point.
- (e) Repeat part 5(c) replacing Spec with mSpec in the subspace topology. Can $\text{mSpec}(R)$ ever have a dense point?
- (f) Consider the following two properties (which together are implied by the Hausdorff condition) for a topological space X : i) for every $p \in X$ the set $\{p\}$ is closed and ii) given two distinct points $p, q \in X$ there exist open subsets $U, V \subset X$ such that $p \in U$ and $q \in V$ and $U \cap V = \emptyset$. Prove that if X is a topological space with a dense point, and X is not a single point, then X fails both parts of this definition.

Solution.

(a) ($\implies$)

Assume that $\{p\}$ is a dense subset of X .

Then $U \cap \{p\} \neq \emptyset$ for all non-empty open subsets $U \subset X$. That is $\{p\} \not\subseteq X/U$ (since all non-empty proper closed subsets of X are complements of non-empty open subsets of X , $\{p\}$ is not contained in any proper closed subset of X). But $\{p\} \subset X$, hence smallest closed subset which contains $\{p\}$ is X .

($\impliedby$)

Assume that the smallest closed subset which contains $\{p\}$ is X .

Then for all proper closed subsets U' of X , $\{p\} \not\subseteq U'$ that is $X/U' = U \cap \{p\} \neq \emptyset$. Note that $\emptyset \neq U$ for any U' since U' is a proper subset of X . Therefore, for every non-empty open subset $U \subset X$, $\{p\} \cap U \neq \emptyset$, and hence $\{p\}$ is dense in X .

Hence proved that $\{p\}$ is a dense subset of X if and only if the smallest closed subset which contains $\{p\}$ is X .

(b) ($\implies$)

Assume $[P] \in \text{Spec}(R)$ is dense. Then the smallest closed set containing $[P]$ is the whole space, i.e. $\text{Spec}(R)$. That is $\mathbb{V}(P) = \text{Spec}(R)$.

($\impliedby$)

Assume $\mathbb{V}(P) = \text{Spec}(R)$. Then $P \subseteq T$ for all prime ideals $T \subset R$. That is, the smallest closed set containing P is the whole space ($\text{Spec}(R)$). Hence, $[P]$ is dense in $\text{Spec}(R)$.

Hence proved that a point $[P] \in \text{Spec}(R)$ is dense if and only if $\mathbb{V}(P) = \text{Spec}(R)$.

- (c) (i) Since $\mathbb{K}[x]$ is a PID, points of $\text{Spec}(\mathbb{K}[x])$ are $\langle 0 \rangle$ and maximal ideals $\langle f \rangle$ where f is a monic irreducible polynomial. From part (b) we know that dense points of $\text{Spec}(\mathbb{K}[x])$ are points $[P] \in \text{Spec}(\mathbb{K}[x])$ such that $\mathbb{V}(P) = \text{Spec}(\mathbb{K}[x])$. Since $\mathbb{V}(f) = \{\langle f \rangle\}$, $[f]$ is not dense in $\text{Spec}(\mathbb{K}[x])$. However since $\mathbb{V}(0) = \text{Spec}(\mathbb{K}[x])$, $[0]$ is dense in $\text{Spec}(\mathbb{K}[x])$.
- (ii) Since $\mathbb{Z}$ is a PID, points of $\text{Spec}(\mathbb{Z})$ are $\langle 0 \rangle$ and maximal ideals $\langle p \rangle$ where $p \in \mathbb{Z}$ prime. From part (b) we know that dense points of $\text{Spec}(\mathbb{Z})$ are points $[P] \in \text{Spec}(\mathbb{Z})$ such that $\mathbb{V}(P) = \text{Spec}(\mathbb{Z})$. Since $\mathbb{V}(p) = \langle p \rangle$, $[p]$ is not dense in $\text{Spec}(\mathbb{Z})$. However since $\mathbb{V}(0) = \text{Spec}(\mathbb{Z})$, $[0]$ is dense in $\text{Spec}(\mathbb{Z})$.
- (iii) From part (b) we know that dense points of $\text{Spec}(\mathbb{Z}[t])$ are points $[P] \in \text{Spec}(\mathbb{Z}[t])$ such that $\mathbb{V}(P) = \text{Spec}(\mathbb{Z}[t])$. Since $\mathbb{V}(0) = \text{Spec}(\mathbb{Z}[t])$, $[0]$ is dense in $\text{Spec}(\mathbb{Z}[t])$, and for every non-zero $[P] \in \text{Spec}(\mathbb{Z}[t])$, $\mathbb{V}(P)$ doesn't contain $[0]$ since $\langle P \rangle \not\subseteq \langle 0 \rangle$. Hence, $[P]$ is not dense.
- (iv) $\text{Spec}(\mathbb{K}[x]/\langle x^2 \rangle) = \{\langle x \rangle\}$ since $\langle 0 \rangle$ is not prime. Hence $[x]$ is trivially dense in $\text{Spec}(\mathbb{K}[x]/\langle x^2 \rangle)$.
- (v) From part (b) we know that dense points of $\text{Spec}(\mathbb{K}[x, y])$ are points $[P] \in \text{Spec}(\mathbb{K}[x, y])$ such that $\mathbb{V}(P) = \text{Spec}(\mathbb{K}[x, y])$. Since $\mathbb{V}(0) = \text{Spec}(\mathbb{K}[x, y])$, $[0]$ is dense, and for every non-zero $[P] \in \text{Spec}(\mathbb{K}[x, y])$, $\mathbb{V}(P)$ doesn't contain $[0]$ since $\langle P \rangle \not\subseteq \langle 0 \rangle$. Hence, $[P]$ is not dense.
- (vi) $\text{Spec}(\mathbb{Z}/\langle 6 \rangle) = \{\langle 2 \rangle, \langle 3 \rangle\}$. Note that they are both maximal since $\mathbb{Z}/\langle 6 \rangle/\langle 2 \rangle = \mathbb{Z}/\langle 2 \rangle$ and $\mathbb{Z}/\langle 6 \rangle/\langle 3 \rangle = \mathbb{Z}/\langle 3 \rangle$. From part (b) we know that dense points of $\text{Spec}(\mathbb{Z}/\langle 6 \rangle)$ are points $[P] \in \text{Spec}(\mathbb{Z}/\langle 6 \rangle)$ such that $\mathbb{V}(P) = \text{Spec}(\mathbb{Z}/\langle 6 \rangle)$. Since both ideals are maximal

and incomparable, $\mathbb{V}([2])$ doesn't contain $[3]$ and $\mathbb{V}([3])$ doesn't contain $[2]$. Hence, there are no dense points in $\text{Spec}(\mathbb{Z}/\langle 6 \rangle)$.

(vii) $\text{Spec}(\mathbb{Z}/\langle 4 \rangle) = \{\langle 2 \rangle\}$. Hence $[2]$ is trivially dense.

(viii) $\text{Spec}(\mathbb{Z}/\langle 12 \rangle) = \{\langle 2 \rangle, \langle 3 \rangle\}$. Note that they are both maximal since $\mathbb{Z}/\langle 12 \rangle / \langle 2 \rangle = \mathbb{Z}/\langle 2 \rangle$ and $\mathbb{Z}/\langle 12 \rangle / \langle 3 \rangle = \mathbb{Z}/\langle 3 \rangle$.

From part (b) we know that dense points of $\text{Spec}(\mathbb{Z}/\langle 12 \rangle)$ are points $[P] \in \text{Spec}(\mathbb{Z}/\langle 12 \rangle)$ such that $\mathbb{V}(P) = \text{Spec}(\mathbb{Z}/\langle 12 \rangle)$. Since both ideals are maximal and incomparable, $\mathbb{V}([2])$ doesn't contain $[3]$ and $\mathbb{V}([3])$ doesn't contain $[2]$. Hence, there are no dense points in $\text{Spec}(\mathbb{Z}/\langle 12 \rangle)$.

(d) Assume that R is an integral domain.

Existence: Note that $\langle 0 \rangle \subseteq R$ is a prime ideal, and also $\langle 0 \rangle \subseteq P$ for all prime ideals P in R . Also recall that the smallest closed set containing $[0]$ is $\mathbb{V}([0])$, and

$$\mathbb{V}([0]) = \mathbb{V}(\langle 0 \rangle) = \text{Spec}(R)$$

That is the smallest closed set containing $[0]$ is the whole space, hence $[0]$ is a dense point in $\text{Spec}(R)$.

Uniqueness: Suppose $[P] \in \text{Spec}(R)$ is also a dense point. Then by part(b), we know that $\mathbb{V}([P]) = \text{Spec}(R)$. That is $P \subseteq \langle 0 \rangle$, and $\langle 0 \rangle \subseteq P$ trivially. Hence, $[P] = [0]$.

Hence if R is a domain, $\text{Spec}(R)$ has a unique dense point.

- (e) (i) Since $\mathbb{K}[x]$ is a PID, points of $\text{mSpec}(\mathbb{K}[x])$ are maximal ideals $\langle f \rangle$ where f is a monic irreducible polynomial. Since $\mathbb{V}(f) = \langle f \rangle$, $[f]$ is not dense in $\text{mSpec}(\mathbb{K}[x])$.
- (ii) Since $\mathbb{Z}$ is a PID, points of $\text{mSpec}(\mathbb{Z})$ are maximal ideals $\langle p \rangle$ where $p \in \mathbb{Z}$ prime. Since $\mathbb{V}(p) = \langle p \rangle$, $[p]$ is not dense in $\text{Spec}(\mathbb{Z})$.
- (iii) For every non-zero $[P] \in \text{mSpec}(\mathbb{Z}[t])$, $\mathbb{V}(P) = \{[P]\}$. Hence, $[P]$ is not dense.
- (iv) $\text{mSpec}(\mathbb{K}[x]/\langle x^2 \rangle) = \{\langle x \rangle\}$. Hence $[x]$ is trivially dense in $\text{mSpec}(\mathbb{K}[x]/\langle x^2 \rangle)$.
- (v) For every maximal ideal $[P] \in \text{mSpec}(\mathbb{K}[x, y])$, $\mathbb{V}(P) = \{[P]\} \neq \text{mSpec}(\mathbb{K}[x, y])$. Hence, $[P]$ is not dense.
- (vi) $\text{mSpec}(\mathbb{Z}/\langle 6 \rangle) = \{\langle 2 \rangle, \langle 3 \rangle\}$. Since $\mathbb{V}([2])$ doesn't contain $[3]$ and $\mathbb{V}([3])$ doesn't contain $[2]$ there are no dense points in $\text{mSpec}(\mathbb{Z}/\langle 6 \rangle)$.
- (vii) $\text{mSpec}(\mathbb{Z}/\langle 4 \rangle) = \{\langle 2 \rangle\}$. Hence $[2]$ is trivially dense.
- (viii) $\text{mSpec}(\mathbb{Z}/\langle 12 \rangle) = \{\langle 2 \rangle, \langle 3 \rangle\}$. Since $\mathbb{V}([2])$ doesn't contain $[3]$ and $\mathbb{V}([3])$ doesn't contain $[2]$ there are no dense points in $\text{mSpec}(\mathbb{Z}/\langle 12 \rangle)$.
- $\text{mSpec}(R)$ only has dense points when it is a singleton set.

(f) Assume X is a topological space with a dense point p and X is not a single point.

Then by definition of denseness $p \in U$, for all non-empty open subsets $U \subset X$, i.e. we cannot find two distinct open subsets V, W such that $V \cap W = \emptyset$.

For the sake of contradiction assume $\{p\}$ is a closed set, then $X \setminus \{p\}$ is a non-empty open set (since X is not a single point) not containing p contradicting the definition of denseness. Hence, $\{p\}$ cannot be closed.

Problem 6. Let $\mathcal{S} \subset R$ be any subset. Prove that $\mathbb{V}(\mathcal{S}) = \mathbb{V}(\langle \mathcal{S} \rangle)$ where $\langle \mathcal{S} \rangle$ denotes the ideal generated by $\mathcal{S}$.

Solution.

Let I be an ideal containing $\mathcal{S}$, then one can see that $\langle \mathcal{S} \rangle \subseteq I$. And trivially $\mathcal{S} \subset \langle \mathcal{S} \rangle \subseteq I$. Hence,

$$\mathcal{S} \subseteq I \iff \langle \mathcal{S} \rangle \subseteq I$$

And it follows that $\mathbb{V}(\mathcal{S}) = \mathbb{V}(\langle \mathcal{S} \rangle)$.

Problem 7. In this exercise we wish to understand the Zariski topology on $\text{Spec}(\mathbb{C}[x, y])$. Fix a point $p = (a, b) \in \mathbb{C}^2$.

- (a) Use the *evaluation-at- p* map: $\text{ev} : \mathbb{C}[x, y] \rightarrow \mathbb{C}$ to prove that the set of polynomial functions vanishing at p is a maximal ideal $\mathfrak{m}_p \subset \mathbb{C}[x, y]$.
- (b) Find generators for the ideal $\mathfrak{m}_p$. Is this ideal principal?
- (c) Prove that the following map is injective:

$$\begin{array}{ccc} \mathbb{C}^2 & \xrightarrow{\eta} & \text{Spec}(\mathbb{C}[x, y]) \\ p & \longmapsto & \mathfrak{m}_p \end{array}.$$

- (d) Show that the map η from part 3c is not surjective by finding at least two distinct points in $\text{Spec}(\mathbb{C}[x, y])$ not in the image of η .
- (e) Describe the closed subsets of $\text{Spec}(\mathbb{C}[x, y])$. (Hint: They come in four types)

Solution.

- (a) We have, $\text{ev}_p : \mathbb{C}[x, y] \rightarrow \mathbb{C}$ where $f \mapsto f(a, b)$. If f vanishes at p then $f(a, b) = 0$, i.e. $f \in \ker(\text{ev}_p) = \mathfrak{m}_p$. The field homomorphism is surjective (constant functions), hence $\mathfrak{m}_p$ is maximal (First Isomorphism Theorem).
- (b) Note that $\mathfrak{m}_p \supseteq \langle x - a, y - b \rangle$ since both $x - a$ and $y - b$ vanish at (a, b) . Moreover since both $\langle x - a, y - b \rangle$ ($\mathbb{C}[x, y] / \langle x - a, y - b \rangle \cong \mathbb{C}$) and $\mathfrak{m}_p$ are maximal they must be equal.
- (c) Let $p = (a, b), q = (c, d) \in \mathbb{C}^2$. Assume $\eta(p) = \eta(q)$, then

$$\mathfrak{m}_p = \mathfrak{m}_q \implies x - a \in \langle x - c, y - d \rangle \implies c = a$$

similarly $y - b \in \langle x - c, y - d \rangle \implies b = d$. Hence $p = q$.

- (d) $\langle x \rangle, \langle y \rangle \in \text{Spec}(\mathbb{C}[x, y])$ but not in $\text{im}(\eta)$ since the latter only consists of maximal ideals.
- (e) The closed subsets of $\text{Spec}(\mathbb{C}[x, y])$ are of finite unions and arbitrary intersections of the following:

- $\mathbb{V}(\langle 1 \rangle) = \emptyset$
- $\mathbb{V}(\langle 0 \rangle) = \text{Spec}(\mathbb{C}[x, y])$

- $\mathbb{V}(\langle f \rangle) = \{[\mathfrak{p}] \in \text{Spec}(\mathbb{C}[x, y]), \langle f \rangle \subseteq \mathfrak{p}\}$, f is an irreducible polynomial.
- $\mathbb{V}(\mathfrak{m}_p) = \{\mathfrak{m}_p\}$, where $\mathfrak{m}_p$ is the maximal ideal that corresponds to the kernel of evaluation-at- p map.

A function $f : X \rightarrow Y$ of topological spaces is *continuous* if and only if $f^{-1}(U) \subset X$ is open for all open subsets $U \subset Y$, i.e., the preimage of an open is open. A *homeomorphism* $f : X \rightarrow Y$ is a continuous map of topological spaces with a continuous inverse $f^{-1} : Y \rightarrow X$. A useful fact is that a continuous map $f : X \rightarrow Y$ is a homeomorphism if and only if it is bijective and $f(U) \subset Y$ is open for every open subsets $U \subset X$.

Problem 8. Ring Maps Induce Continuous Maps.

Let $\phi : R \rightarrow S$ be a ring homomorphism. The goal of this exercise is to show ϕ induces a map $\phi^\# : \text{Spec}(S) \rightarrow \text{Spec}(R)$, which is continuous in the Zariski topology.

- (a) Prove that if $[P] \in \text{Spec}(S)$, then $[\phi^{-1}(P)] \in \text{Spec}(R)$.
- (b) Prove that the map below is continuous in the Zariski topology:

$$\begin{array}{ccc} \text{Spec}(S) & \xrightarrow{\phi^\#} & \text{Spec}(R) \\ [P] & \longmapsto & [\phi^{-1}(P)] \end{array}$$

- (c) For the ring map $\mathbb{Z} \rightarrow \mathbb{Z}/2\mathbb{Z}$ sending each integer to its residue class modulo 2, describe the induced map on spectra explicitly.
- (d) For the ring map $\mathbb{Z} \hookrightarrow \mathbb{Q}$, describe the induced map on spectra explicitly.
- (e) If $\phi : R \hookrightarrow S$ is an injective ring map, give a simple description of the induced map on spectra.

Solution.

- (a) We know that pre-images of ideals are ideals, so we only need to show that $\phi^{-1}(P)$ is prime. Let $ab \in P$, then $\phi^{-1}(ab) \in \phi^{-1}(P)$. Since P is prime either $a \in P$ or $b \in P$, and hence either $\phi^{-1}(a) \in \phi^{-1}(P)$ or $\phi^{-1}(b) \in \phi^{-1}(P)$. Hence $\phi^{-1}(P)$ is prime, and $[\phi^{-1}(P)] \in \text{Spec}(R)$.

- (b) We will show that pre-images of closed sets in $\text{Spec}(R)$ are closed. Let $\mathbb{V}(I)$ be a closed set for some ideal $I \subseteq R$. Then,

$$(\phi^\#)^{-1}(\mathbb{V}(I)) = \{\mathfrak{p} \mid I \subseteq \phi^{-1}(\mathfrak{p})\} = \{\mathfrak{p} \mid \langle \phi(I) \rangle \subseteq \mathfrak{p}\} = \mathbb{V}(\langle \phi(I) \rangle)$$

Which is a closed set in $\text{Spec}(S)$.

- (c) The map induced on spectra is $\text{Spec}(\mathbb{Z}/2\mathbb{Z}) \xrightarrow{\phi^\#} \text{Spec}(\mathbb{Z})$. $[0]$ is the only element in $\text{Spec}(\mathbb{Z}/2\mathbb{Z})$ and $\phi^\#([0]) = [2] \in \text{Spec}(\mathbb{Z})$.
- (d) The map induced on spectra is $\text{Spec}(\mathbb{Q}) \xrightarrow{\phi^\#} \text{Spec}(\mathbb{Z})$. $[0]$ is the only element in $\text{Spec}(\mathbb{Q})$ and $\phi^\#([0]) = [0] \in \text{Spec}(\mathbb{Z})$.
- (e) If $\phi : R \hookrightarrow S$ is injective, we can view R as a subring of S and $\phi^\#([\mathfrak{p}]) = [\phi^{-1}(\mathfrak{p})] = [\mathfrak{p} \cap R] \in \text{Spec}(R)$.

A *category* is a collection of objects together with, for each pair of objects, a set of morphisms between them, equipped with an associative composition law and an identity morphism for each object. A *functor* between categories $\mathbf{C}$ and $\mathbf{D}$ is a mapping that assigns to each object of $\mathbf{C}$ an object of $\mathbf{D}$ and to each morphism in $\mathbf{C}$ a morphism in $\mathbf{D}$, preserving identity morphisms and composition. A functor is *contravariant* if it reverses the order of composition and *covariant* if it preserves the order of composition.

Problem 9. Spec as a Functor.

In this problem we package together much of this worksheet to define the Spec functor. Consider the map:

$$\begin{array}{ccc} \mathbf{comRing} & \longrightarrow & \mathbf{Top} \\ R & \longmapsto & \mathrm{Spec}(R) \end{array}$$

from the category of commutative rings with unit with ring homomorphisms to the category of topological spaces with continuous maps.

- (a) Explain using Question 8 how this mapping defines a contravariant functor from **comRing** to **Top**. To check functoriality, verify

$$(\mathrm{Id}_R)^\# = \mathrm{Id}_{\mathrm{Spec}(R)} \quad \text{and} \quad (\psi \circ \phi)^\# = \phi^\# \circ \psi^\#$$

for ring maps $\phi : R \rightarrow S$ and $\psi : S \rightarrow T$. We call this the Spec functor.

- (b) Do isomorphic rings have Spectra that are homeomorphic topological spaces? Explain.
- (c) If two rings have homeomorphic spectra, are they necessarily isomorphic rings? Explain.

Solution.

- (a) We already showed that a homomorphism between rings $R \rightarrow S$ corresponds to a map between their spectra $\mathrm{Spec}(S) \rightarrow \mathrm{Spec}(R)$, and that this map is a continuous topological map. So it only remains to check that the functorial properties hold,

- Let $\mathrm{id}_R : R \rightarrow R$ be the identity map. Then for $[\mathfrak{p}] \in \mathrm{Spec}(R)$,

$$(\mathrm{id}_R)^\#([\mathfrak{p}]) = [\mathrm{id}_R^{-1}(\mathfrak{p})] = [\mathfrak{p}] = \mathrm{id}_{\mathrm{Spec}(R)}([\mathfrak{p}])$$

- Let $\phi : R \rightarrow S$, $\psi : S \rightarrow T$ be two ring homomorphisms. Then for $[\mathfrak{p}] \in \mathrm{Spec}(T)$,

$$\begin{aligned} (\psi \circ \phi)^\#([\mathfrak{p}]) &= [((\psi \circ \phi)^{-1}(\mathfrak{p}))] = [\phi^{-1}(\psi^{-1}(\mathfrak{p}))] \\ &= [\phi^{-1}(\psi^\#(\mathfrak{p}))] = [\phi^\#(\psi^\#(\mathfrak{p}))] \\ &= [\phi^\# \circ \psi^\#(\mathfrak{p})] \end{aligned}$$

Hence $(\mathrm{Id}_R)^\# = \mathrm{Id}_{\mathrm{Spec}(R)}$ and $(\psi \circ \phi)^\# = \phi^\# \circ \psi^\#$. The reversal of order makes this a covariant functor.

- (b) If $\phi : R \xrightarrow{\sim} S$ is a ring isomorphism then $\phi^{-1} : S \xrightarrow{\sim} R$ is also a ring isomorphism. Then,

$$(\phi^{-1} \circ \phi)^{\#} = \phi^{\#} \circ (\phi^{-1})^{\#} = (\text{Id}_R)^{\#} = \text{Id}_{\text{Spec}(R)}$$

$$(\phi \circ \phi^{-1})^{\#} = (\phi^{-1})^{\#} \circ \phi^{\#} = (\text{Id}_S)^{\#} = \text{Id}_{\text{Spec}(S)}$$

so $\phi^{\#}$ and $(\phi^{-1})^{\#}$ are inverse to each other. Since both $\phi^{\#}$ and $(\phi^{-1})^{\#}$ are continuous, $\phi^{\#} : \text{Spec}(S) \rightarrow \text{Spec}(R)$ is a homeomorphism. Hence isomorphic rings have homeomorphic spectra.

- (c) No, because we could have a homeomorphism between spectra of fields ($\text{Spec}(\mathbb{C}) \cong \text{Spec}(\mathbb{R})$) but that doesn't necessarily correspond to an isomorphism of fields ($\mathbb{C} \not\cong \mathbb{R}$).

Problem 10. Find a homeomorphism between $\text{Spec}(\mathbb{C}[x])$ and $\text{Spec}(\mathbb{C}[x, y]/\langle x - y \rangle)$. Construct the corresponding ring map explicitly in *Macaulay2* and compute its kernel.

Solution. Since $\mathbb{C}[x, y]/\langle x - y \rangle \cong \mathbb{C}[x]$ we send $\mathfrak{p} \subseteq \mathbb{C}[x]$ to $\bar{\mathfrak{p}} \subseteq \mathbb{C}[x, y]/\langle x - y \rangle$. And the corresponding ring map is $\mathbb{C}[x, y]/\langle x - y \rangle \rightarrow \mathbb{C}[x]$ with trivial kernel.

Problem 11. Closed Embeddings I.

Let $I \subset R$ be an ideal and $\pi : R \rightarrow R/I$ the natural quotient map. Consider the induced map on spectra

$$\operatorname{Spec}(R/I) \xrightarrow{\pi^\#} \operatorname{Spec}(R)$$

In this exercise we will show this map is a closed embedding. A *closed embedding* is a map of topological spaces whose image is closed and which is a homeomorphism onto its image.

- (a) Using the correspondence between ideals in R and R/I give an explicit description of $\pi^\#$.
- (b) Prove that the image of $\pi^\#$ is equal to $\mathbb{V}(I)$.
- (c) Explain why the ideal correspondence implies that $\pi^\#$ is injective.
- (d) Prove that $\pi^\#$ is a closed embedding onto $\mathbb{V}(I)$, in the subspace topology, by showing that if $J \subset R/I$ is an ideal then

$$\pi^\#(\mathbb{V}(J)) = \mathbb{V}(\pi^{-1}(J)).$$

Solution.

- (a) Ideals in R/I correspond to ideals in R containing I . And for any ideal $I \subseteq R$ prime ideal we have $\pi(I) = \bar{I}$. Let $\bar{\mathfrak{p}} \in \operatorname{Spec}(R/I)$. Then $\pi^\#(\bar{\mathfrak{p}}) = \pi^{-1}(\bar{\mathfrak{p}}) = \mathfrak{p}$.
- (b) The image of $\pi^\#$ by definition is

$$\operatorname{im}(\pi^\#) = \pi^\#(\operatorname{Spec}(R/I)) = \{\mathfrak{p} \in \operatorname{Spec}(R) \mid I \subseteq \pi(\mathfrak{p})\} = \mathbb{V}(I)$$

- (c) Suppose $\pi^\#(\bar{\mathfrak{p}}) = \pi^\#(\bar{\mathfrak{q}})$ then,

$$\pi^{-1}(\bar{\mathfrak{p}}) = \pi^{-1}(\bar{\mathfrak{q}}) \implies \mathfrak{p} = \mathfrak{q}$$

since π is surjective.

- (d) If $J \subseteq R/I$ is an ideal then $I \subseteq \pi^{-1}(J)$

$$\pi^\#(\mathbb{V}(J)) = \{\pi^\#(\bar{\mathfrak{p}}) \mid J \subseteq \bar{\mathfrak{p}}\} = \{\pi^{-1}(\bar{\mathfrak{p}}) \mid \pi^{-1}(J) \subseteq \pi^{-1}(\bar{\mathfrak{p}})\} = \mathbb{V}(\pi^{-1}(J))$$

Since $\pi^\#$ sends closed sets to closed sets and is a bijection onto $\mathbb{V}(I)$, it is a homeomorphism onto its image, and hence a closed embedding.

Worksheet 2.2: Algebraic Sets

Theorem 0.3 (Hilbert's Nullstellensatz I). *If $\mathbb{K}$ is algebraically closed, then every maximal ideal of $\mathbb{K}[x_1, \dots, x_n]$ is of the form $\langle x_1 - a_1, \dots, x_n - a_n \rangle$ for some point $(a_1, \dots, a_n) \in \mathbb{K}^n$.*

Note this is just the first of a series of theorems that will go by Hilbert's Nullstellensatz. Recall that a $\mathbb{K}$ -algebra is a ring R together with a specified ring homomorphism $\mathbb{K} \rightarrow R$. A morphism of $\mathbb{K}$ -algebras $\varphi : R \rightarrow S$ is a ring homomorphism that preserves scalars, i.e. $\varphi(c \cdot 1_R) = c \cdot 1_S$ for every $c \in \mathbb{K}$. In particular, $\mathbb{K}[x_1, \dots, x_n]$ is a $\mathbb{K}$ -algebra, every evaluation map $\text{ev}_p : \mathbb{K}[x_1, \dots, x_n] \rightarrow \mathbb{K}$ is a $\mathbb{K}$ -algebra homomorphism, and if $\mathfrak{m}$ is a maximal ideal then $R/\mathfrak{m}$ is naturally a field equipped with a $\mathbb{K}$ -algebra structure. When we say below that a residue field is “isomorphic to $\mathbb{K}$,” we mean *as a $\mathbb{K}$ -algebra*.

Problem 1. Evaluation Maps.

For a point $p = (a_1, \dots, a_n) \in \mathbb{K}^n$ the *evaluation-at- p* map is given by

$$\begin{array}{ccc} \mathbb{K}[x_1, \dots, x_n] & \xrightarrow{\text{ev}_p} & \mathbb{K} \\ f & \longmapsto & f(p) \end{array}.$$

- (a) Prove that ev_p is a ring homomorphism and find generators for $\mathfrak{m}_p := \ker(\text{ev}_p)$.
- (b) Prove that there is a bijection, induced by these evaluation maps, between $\mathbb{K}^n$ and the subset of $\text{Spec}(\mathbb{K}[x_1, \dots, x_n])$ consisting of maximal ideals whose residue field is isomorphic to $\mathbb{K}$ as a $\mathbb{K}$ -algebra.
- (c) Find a maximal ideal of $\mathbb{R}[x, y]$ that is not of the form $\mathfrak{m}_p$ for any $p \in \mathbb{R}^2$.
- (d) Restate Hilbert's Nullstellensatz I (above) in terms of $\mathfrak{m}_p$.

Solution.

- (a) To show that ev_p is a ring homomorphism, we need to show that for all $f_1, f_2 \in \mathbb{K}[x_1, \dots, x_n]$ the three conditions hold,

Additivity:

$$\text{ev}_p(f_1 + f_2) = (f_1 + f_2)(p) = f_1(p) + f_2(p) = \text{ev}_p(f_1) + \text{ev}_p(f_2)$$

Multiplicativity:

$$\text{ev}_p(f_1 \cdot f_2) = (f_1 \cdot f_2)(p) = f_1(p) \cdot f_2(p) = \text{ev}_p(f_1) \cdot \text{ev}_p(f_2)$$

And $\text{ev}_p(1) = 1$ since 1 is a constant function.

- (b) Consider the map

$$\phi : \mathbb{K}^n \longrightarrow \{\mathfrak{m} \in \text{mSpec}(\mathbb{K}[x_1, \dots, x_n]) : \mathbb{K}[x_1, \dots, x_n] / \mathfrak{m} \cong \mathbb{K}\}, \quad \phi(p) = \mathfrak{m}_p$$

To see that ϕ is injective, suppose $\mathfrak{m}_p = \mathfrak{m}_q$ for $p = (a_1, \dots, a_n)$ and $q = (b_1, \dots, b_n)$. By part (a),

$$\mathfrak{m}_p = (x_1 - a_1, \dots, x_n - a_n), \quad \mathfrak{m}_q = (x_1 - b_1, \dots, x_n - b_n).$$

Since $x_i - a_i \in \mathfrak{m}_q$, we have

$$(x_i - a_i)(q) = b_i - a_i = 0.$$

Hence $a_i = b_i$ for all i , so $p = q$.

To prove surjectivity, let $\mathfrak{m}$ be a maximal ideal such that

$$\mathbb{K}[x_1, \dots, x_n] / \mathfrak{m} \cong \mathbb{K}$$

as $\mathbb{K}$ -algebras. Let $\pi : \mathbb{K}[x_1, \dots, x_n] \rightarrow \mathbb{K}[x_1, \dots, x_n] / \mathfrak{m} \cong \mathbb{K}$ be the quotient map, and $a_i = \pi(x_i) \in \mathbb{K}$. Since π is a $\mathbb{K}$ -algebra homomorphism, for every polynomial $f \in \mathbb{K}[x_1, \dots, x_n]$ we have

$$\pi(f) = f(a_1, \dots, a_n).$$

Thus $\pi = \text{ev}_p$, where $p = (a_1, \dots, a_n)$, and therefore

$$\mathfrak{m} = \ker(\pi) = \ker(\text{ev}_p) = \mathfrak{m}_p.$$

Hence ϕ is surjective and therefore a bijection.

(c) Consider the ideal $\mathfrak{m} = (x^2 + 1, y) \subset \mathbb{R}[x, y]$. We have

$$\mathbb{R}[x, y] / \mathfrak{m} \cong \mathbb{R}[x] / (x^2 + 1) \cong \mathbb{C}.$$

Since $\mathbb{C}$ is a field, $\mathfrak{m}$ is maximal. But $\mathbb{C} \neq \mathbb{R}$ so $\mathfrak{m}$ is not of the form $\mathfrak{m}_p$ for any $p \in \mathbb{R}^2$.

(d) Hilbert's Nullstellensatz can be restated as saying that the radical of an ideal is the intersection of all maximal ideals of the form $\mathfrak{m}_p$ that contain it.

Problem 2. Algebraic Sets.

Let $\mathcal{S} \subset \mathbb{K}[x_1, \dots, x_n]$ be a subset. Define the corresponding *algebraic set* in $\mathbb{K}^n$ to be:

$$\mathbb{V}(\mathcal{S}) := \{p \in \mathbb{K}^n \mid f(p) = 0 \text{ for all } f \in \mathcal{S}\}.$$

- (a) Prove that $\mathbb{V}$ is order-reverse, i.e., if $\mathcal{S} \subset \mathcal{T}$ then $\mathbb{V}(\mathcal{T}) \subset \mathbb{V}(\mathcal{S})$.
- (b) Prove that $\mathbb{V}(\mathcal{S}) = \mathbb{V}(\langle \mathcal{S} \rangle) = \mathbb{V}(\sqrt{\langle \mathcal{S} \rangle})$.
- (c) Describe each of the following algebraic sets, and when possible draw a picture.
 - (i) $\mathcal{S} := \{x^2 - y\} \subset \mathbb{R}[x, y]$.
 - (ii) $\mathcal{S} := \{x^2 - xy\} \subset \mathbb{R}[x, y]$.
 - (iii) $\mathcal{S} := \{x^6 - 3x^5y + 3x^4y^2 - x^3y^3\} \subset \mathbb{R}[x, y]$.
 - (iv) $\mathcal{S} := \{x^4 - 2x^3y + x^2y^2\} \subset \mathbb{R}[x, y]$.
 - (v) $\mathcal{S} := \{x^2 + y^2 - 1\} \subset \mathbb{R}[x, y]$.
 - (vi) $\mathcal{S} := \{n(x^2 + y^2 - z^2) \mid n \in \mathbb{Z}_{\geq 0}\} \subset \mathbb{R}[x, y, z]$.
 - (vii) $\mathcal{S} := \{x^2 + 1\} \subset \mathbb{R}[x]$.
 - (viii) $\mathcal{S} := \{x^2 + 1\} \subset \mathbb{C}[x]$.
 - (ix) $\mathcal{S}$ is a collection of any linear polynomials in $\mathbb{K}[x_1, \dots, x_n]$.
- (d) Explain why $SL_n(\mathbb{K})$ has the structure of an algebraic set.
- (e) Explain why the set of $m \times n$ matrices of rank $< t$ is an algebraic set contained in $\mathbb{K}^{m \times n}$.

Solution.

- (a) Assume $\mathcal{S} \subset \mathcal{T}$. Let $p \in \mathbb{V}(\mathcal{T})$.

Then, for all $f \in \mathcal{T}$, $f(p) = 0$. But since $\mathcal{S} \subset \mathcal{T}$ this implies $g(p) = 0, \forall g \in \mathcal{S}$. Hence, $p \in \mathbb{V}(\mathcal{S})$. Since p was an arbitrary point in $\mathbb{V}(\mathcal{T})$ we conclude that $\mathbb{V}$ is order reversing.

- (b) By definition,

$$\mathcal{S} \subseteq \langle \mathcal{S} \rangle \subseteq \sqrt{\langle \mathcal{S} \rangle}.$$

And from part (a) we have,


$$\mathbb{V}(\mathcal{S}) \supseteq \mathbb{V}(\langle \mathcal{S} \rangle) \supseteq \mathbb{V}\left(\sqrt{\langle \mathcal{S} \rangle}\right).$$

To show $\mathbb{V}(\langle S \rangle) \subseteq \mathbb{V}(\sqrt{\langle S \rangle})$ we let $p \in \mathbb{V}(\langle S \rangle)$. That is for all $f \in \langle S \rangle \subseteq \sqrt{\langle S \rangle}$, $f(p) = 0$, i.e. $p \in \mathbb{V}(\sqrt{\langle S \rangle}) \implies \mathbb{V}(\langle S \rangle) \subseteq \mathbb{V}(\sqrt{\langle S \rangle})$.

To show that $\mathbb{V}(S) \subseteq \mathbb{V}(\langle S \rangle)$ we let $p \in \mathbb{V}(S)$. That is for all $f \in S$, $f(p) = 0$ but since $\langle S \rangle$ is the set of linear combinations of functions that evaluate to zero at p , they also evaluate to zero at p and $p \in \mathbb{V}(\langle S \rangle)$.


(c) (i) $S := \{x^2 - y\} \subset \mathbb{R}[x, y]$.

$$\mathbb{V}(S) = \{(x, y) \in \mathbb{R}^2 \mid x^2 = y\}$$



(ii) $S := \{x^2 - xy\} \subset \mathbb{R}[x, y]$.


$$\mathbb{V}(S) = \{(x, y) \in \mathbb{R}^2 \mid x = 0, x = y\}$$



(iii) $S := \{x^6 - 3x^5y + 3x^4y^2 - x^3y^3\} \subset \mathbb{R}[x, y]$.

Note that $x^6 - 3x^5y + 3x^4y^2 - x^3y^3 = x^3(x - y)^3$.


$$\mathbb{V}(S) = \{(x, y) \in \mathbb{R}^2 \mid x = 0, x = y\}$$



(iv) $S := \{x^4 - 2x^3y + x^2y^2\} \subset \mathbb{R}[x, y]$.


Note that $x^4 - 2x^3y + x^2y^2 = x^2(x - y)^2$.

$$\mathbb{V}(S) = \{(x, y) \in \mathbb{R}^2 \mid x = 0, x = y\}$$



(v) $\mathcal{S} := \{x^2 + y^2 - 1\} \subset \mathbb{R}[x, y]$.

$\mathbb{V}(\mathcal{S}) = \{(x, y) \in \mathbb{R}^2 \mid x^2 + y^2 = 1\}$



(vi) $\mathcal{S} := \{n(x^2 + y^2 - z^2) \mid n \in \mathbb{Z}_{\geq 0}\} \subset \mathbb{R}[x, y, z]$.


$\mathbb{V}(\mathcal{S}) = \{(x, y, z) \in \mathbb{R}^3 \mid x^2 + y^2 = z^2\}$

(vii) $\mathcal{S} := \{x^2 + 1\} \subset \mathbb{R}[x]$.

$\mathbb{V}(\mathcal{S}) = \emptyset$ since this equation has no solutions over $\mathbb{R}[x]$

(viii) $\mathcal{S} := \{x^2 + 1\} \subset \mathbb{C}[x]$.

$\mathbb{V}(\mathcal{S}) = \{(x, y) \in \mathbb{R}^2 \mid x^2 = y\}$



(ix) $\mathcal{S}$ is a collection of any linear polynomials in $\mathbb{K}[x_1, \dots, x_n]$.

$\mathbb{V}(\mathcal{S}) = \ker(A)$, where A is a $m \times n$ matrix where the i -th row is $(a_{i_1}, \dots, a_{i_n})$

(d) $SL_n(\mathbb{K})$ is the set of $n \times n$ matrices with determinant 1, and the determinant is a polynomial function on $\mathbb{K}^{n \times n}$. Therefore we can define the polynomial

$$f = \det(A) - 1 \in \mathbb{K}[x_1, \dots, x_{n \times n}]$$

where A is a $n \times n$ matrix, and $SL_n(\mathbb{K}) = \mathbb{V}(\{f\})$.

Problem 3. Defining Ideal of an Algebraic Set.

Let $V \subset \mathbb{K}^n$ be an algebraic set. The *defining ideal* of V is the ideal in $\mathbb{K}[x_1, \dots, x_n]$ given by

$$\mathbb{I}(V) := \{f \in \mathbb{K}[x_1, \dots, x_n] \mid f(p) = 0 \text{ for all } p \in V\}.$$

- (a) Prove that $\mathbb{I}(V)$ is in fact an ideal.
- (b) Prove that $\mathbb{I}(V)$ is radical.
- (c) If $V = \mathbb{V}(\mathcal{S})$ for some subset $\mathcal{S} \subset \mathbb{K}[x_1, \dots, x_n]$, show that $\langle \mathcal{S} \rangle \subset \sqrt{\langle \mathcal{S} \rangle} \subset \mathbb{I}(V)$.
- (d) Let $\mathbb{K} = \mathbb{R}$. Give an example to show that the inclusion $\sqrt{\langle \mathcal{S} \rangle} \subset \mathbb{I}(V)$ can be proper.
- (e) If $V = \mathbb{V}(\mathcal{S})$ for some subset $\mathcal{S} \subset \mathbb{K}[x_1, \dots, x_n]$, show that $\mathbb{V}(\mathcal{S}) = \mathbb{V}(\mathbb{I}(V))$.

Solution.

- (a) Let $g \in \mathbb{K}[x_1, \dots, x_n]$ and $f_1, f_2 \in \mathbb{I}(V)$. That is, $f_1(p) = f_2(p) = 0$.

Subgroup Under Addition: Note that,

$$(f_1 + f_2)(p) = f_1(p) + f_2(p) = 0$$

The zero polynomial $0(p) = 0$ acts as the identity, and

$$f_1(p) + (-f_1)(p) = 0(p) = 0$$

gives us the additive inverse.

Closed Under Multiplication: Notice,

$$(g \cdot f)(p) = g(p) \cdot f_1(p) = g(p) \cdot 0 = 0$$

Thus $g \cdot f \in \mathbb{I}(V)$ and $\mathbb{I}(V)$ is an ideal.

- (b) We want to show that $\mathbb{I}(V) = \sqrt{\mathbb{I}(V)}$.

Note that the containment $\mathbb{I}(V) \subseteq \sqrt{\mathbb{I}(V)}$ is obvious. To show the opposite containment take $f \in \sqrt{\mathbb{I}(V)}$. Then for some n we have,

$$f^n(p) = (f(p))^n = 0$$

But since $\mathbb{K}$ is a domain, this implies that $f(p) = 0$. Hence $f \in \mathbb{I}(V)$, $\mathbb{I}(V) \supseteq \sqrt{\mathbb{I}(V)}$, and $\mathbb{I}(V) = \sqrt{\mathbb{I}(V)}$.

- (c) Let $V = \mathbb{V}(\mathcal{S})$. We know that $\langle \mathcal{S} \rangle \subseteq \sqrt{\langle \mathcal{S} \rangle}$. So showing that $\sqrt{\langle \mathcal{S} \rangle} \subset \mathbb{I}(V)$ suffices. Note that $\mathbb{I}(V) = \mathbb{I}(\mathbb{V}(\mathcal{S}))$ which is the set of all functions that evaluate to 0 at p , for all $p \in \mathbb{V}(\mathcal{S})$. And $\mathbb{V}(\mathcal{S})$ is the set of all points p such that all functions $f \in \mathcal{S}$ evaluate to 0 at p . Let $f \in \sqrt{\langle \mathcal{S} \rangle}$. Then by definition for some n , $f^n \in \langle \mathcal{S} \rangle$. We can then write,

$$f^n = \sum_i h_i(p) g_i(p), \quad h_i \in \mathbb{K}[x_1, \dots, x_n], g_i \in \mathcal{S}$$

But then,

$$f^n(p) = \sum_i h_i(p) g_i(p) = \sum_i h_i(p) \cdot 0 = 0$$

Moreover since $\mathbb{K}$ is a domain, $f^n(p) = 0 \implies f(p) = 0 \implies f$ is a function that evaluates to 0 at p , hence $f \in \mathbb{I}(V)$.

- (d) Consider the set $\mathcal{S} = \{x^2 + 1\}$. We have,

$$\sqrt{\langle \mathcal{S} \rangle} = \sqrt{\langle x^2 + 1 \rangle}$$

And, because the polynomial has no solutions over $\mathbb{R}$, we have $V = \mathbb{V}(\mathcal{S}) = \emptyset$. But $\mathbb{I}(\mathbb{V}(\mathcal{S})) = \mathbb{R}[x]$. Also note that since $\langle \mathcal{S} \rangle$ is a prime ideal ($x^2 + 1$ is irreducible), we have $\langle x^2 + 1 \rangle = \sqrt{\langle x^2 + 1 \rangle} \subset \mathbb{R}[x]$. Hence, the containment $\sqrt{\langle \mathcal{S} \rangle} \subset \mathbb{I}(V)$ can be proper.

- (e) Let $V = \mathbb{V}(\mathcal{S})$.

$$\mathbb{V}(\mathcal{S}) \subseteq \mathbb{V}(\mathbb{I}(\mathbb{V}(\mathcal{S})))$$

This containment is obvious by construction. $\mathbb{I}(\mathbb{V}(\mathcal{S}))$ is the set of all functions that vanish at all points in $\mathbb{V}(\mathcal{S})$. And $\mathbb{V}(\mathbb{I}(\mathbb{V}(\mathcal{S})))$ is the set of all points that all functions in $\mathbb{I}(\mathbb{V}(\mathcal{S}))$ vanish at. So, $\mathbb{V}(\mathcal{S}) \subseteq \mathbb{V}(\mathbb{I}(\mathbb{V}(\mathcal{S})))$.

$$\mathbb{V}(\mathcal{S}) \supseteq \mathbb{V}(\mathbb{I}(\mathbb{V}(\mathcal{S})))$$

Note that by definition $\mathcal{S} \subseteq \mathbb{I}(\mathbb{V}(\mathcal{S}))$. Then by 2(a) $\mathbb{V}(\mathbb{I}(\mathbb{V}(\mathcal{S}))) \subseteq \mathbb{V}(\mathcal{S})$.

Hence, $\mathbb{V}(\mathcal{S}) = \mathbb{V}(\mathbb{I}(\mathbb{V}(\mathcal{S})))$.

So far you have built two order-reversing operations:

$$\mathcal{S} \longmapsto \mathbb{V}(\mathcal{S}) \quad \text{and} \quad V \longmapsto \mathbb{I}(V).$$

You have shown that passing from equations to a zero set only remembers the radical of the ideal they generate, and Question 4d shows that over a non-algebraically closed field this can lose information. The next theorem says that over an algebraically closed field this is the only ambiguity.

Theorem 0.4 (Hilbert's Nullstellensatz II). *If $\mathbb{K}$ is algebraically closed and $V = \mathbb{V}(\mathcal{S}) \subset \mathbb{K}^n$ is an algebraic set, for some subset $\mathcal{S} \subset \mathbb{K}[x_1, \dots, x_n]$, then $\mathbb{I}(V) = \sqrt{\langle \mathcal{S} \rangle}$.*

At this point the basic affine dictionary is taking shape; Question (4) asks you to make the correspondence between algebraic sets and radical ideals completely explicit. After that we turn to coordinate rings, which package an algebraic set V into algebra and will let us recover from $\mathbb{K}[V]$ its points, its topology, and ultimately its morphisms.

Problem 4. Let $\mathbb{K}$ be algebraically closed. Use Hilbert's Nullstellensatz II (above) to give an explicit inclusion-reversing bijection between algebraic sets in $\mathbb{K}^n$ and radical ideals.

Solution.

Let $\mathcal{S} \subset \mathbb{K}[x_1, \dots, x_n]$.

$$\mathbb{V}(\mathcal{S}) \xrightarrow{\mathbb{I}} \sqrt{\langle \mathcal{S} \rangle}$$

$$\mathbb{V}(\mathcal{S}) \xleftarrow{\mathbb{V}} \sqrt{\langle \mathcal{S} \rangle}$$

Problem 5. Coordinate Rings.

Let $V \subset \mathbb{K}^n$ be an algebraic set and $\mathcal{R} = \text{Hom}_{\text{Set}}(V, \mathbb{K})$ be the ring of all (set) functions from V to $\mathbb{K}$. In this question we study the restriction map from $\mathbb{K}[x_1, \dots, x_n]$ to $\mathcal{R}$ given by:

$$\begin{array}{ccc} \mathbb{K}[x_1, \dots, x_n] & \xrightarrow{\rho} & \mathcal{R} \\ f & \longmapsto & f|_V \end{array}.$$

Here we are viewing $f \in \mathbb{K}[x_1, \dots, x_n]$ as defining a function $\mathbb{K}^n \rightarrow \mathbb{K}$ given by $p \mapsto f(p)$. We call the image of ρ the *coordinate ring* of V and denote it $\mathbb{K}[V]$.

- (a) Prove that ρ is in fact a ring homomorphism.
- (b) Verify that each of the following functions lies in the coordinate ring of the given algebraic set.
 - (i) Let $V = \mathbb{V}(x^2 + y^2 - 1) \subset \mathbb{R}^2$ and consider the function $V \rightarrow \mathbb{R}$ given by $(x, y) \mapsto e^{x^2 + y^2}$.
 - (ii) Let $V = \mathbb{V}(xy - 1) \subset \mathbb{K}^2$ and consider the function $V \rightarrow \mathbb{K}$ given by $(x, y) \mapsto \frac{1}{x}$.
- (c) Identify the kernel of ρ and use it to give an explicit presentation for $\mathbb{K}[V]$ as a $\mathbb{K}$ -algebra.
- (d) The i -th coordinate of $\mathbb{K}^n$ is the linear functional $e_i^\vee : \mathbb{K}^n \rightarrow \mathbb{K}$ sending $(a_1, \dots, a_n) \mapsto a_i$. Explain why $e_i^\vee \in \mathbb{K}[\mathbb{K}^n]$, and under the presentation given in part (c), which polynomial corresponds to $e_i^\vee$.
- (e) Explain why $\mathbb{K}[V]$ is called the coordinate ring of V . Hint: Think about the previous part.
- (f) Describe the coordinate ring for the following algebraic sets:
 - (i) $\mathbb{V}(y - x^2) \subset \mathbb{K}^2$.
 - (ii) $\mathbb{V}(y^2, xy, x^{17}) \subset \mathbb{K}^2$.
 - (iii) $\mathbb{V}(x^6 - 3x^5y + 3x^4y^2 - x^3y^3) \subset \mathbb{K}^2$.
 - (iv) The set of 3×3 matrices over $\mathbb{F}_5$ of determinant 1 and trace 0. (Hint/Warning: There will likely be 9 more polynomials in the defining ideal than you expect, coming from the fact that finite fields have a special automorphism. This highlights another way we might get a counterexample for Question 4d.)

Solution.

- (a) Let $f_1, f_2 \in \mathbb{K}[x_1, \dots, x_n]$. Note that addition and multiplication is defined pointwise in R . To see if ρ is a ring homomorphism, we check if the three properties hold:

Addition:

$$\rho(f_1) + \rho(f_2)(p) = f_1|_V(p) + f_2|_V(p) = (f_1 + f_2)|_V(p) = \rho(f_1 + f_2)(p)$$

Multiplication:

$$\rho(f_1) \cdot \rho(f_2)(p) = f_1|_V(p) \cdot f_2|_V(p) = (f_1 \cdot f_2)|_V(p) = \rho(f_1 \cdot f_2)(p)$$

Unital:

$$\rho(1)(p) = 1|_V(p) = 1$$

- (b) (i) Let $f : (x, y) \mapsto e^{x^2+y^2}$. Note that on V , $x^2 + y^2 = 1$ then, $f|_V = e^{x^2+y^2} = e^1$. Which behaves like the constant polynomial function $g(x, y) = e$ and hence lies in the ring of coordinates of V .
- (ii) Let $f : (x, y) \mapsto \frac{1}{x}$. Note that on V , $\frac{1}{x} = y$, and $x \neq 0$ because $\mathbb{K}$ is a domain. Then, $f|_V = \frac{1}{x} = y$. Which behaves like the polynomial function $g(x, y) = y$ and hence lies in the ring of coordinates of V .

- (c) The kernel of ρ is $\mathbb{I}(V)$.

$$\mathbb{K}[V] \cong \mathbb{K}[x_1, \dots, x_n] / \mathbb{I}(V)$$

- (d) The coordinate ring of $\mathbb{K}^n$, $\mathbb{K}[\mathbb{K}^n]$ is the ring of all set functions $\mathbb{K}^n \rightarrow \mathbb{K}$.

$$e_i^\vee(x_1, \dots, x_n) = x_i$$

which corresponds to the polynomial function $g(x_1, \dots, x_n) = x_i$. Hence, $e_i^\vee \in \mathbb{K}[\mathbb{K}^n]$.

Under the presentation given in part (c) we have,

$$\mathbb{K}[\mathbb{K}^n] \cong \mathbb{K}[x_1, \dots, x_n] / \mathbb{I}(\mathbb{K}^n) = \mathbb{K}[x_1, \dots, x_n] / \{0\} = \mathbb{K}[x_1, \dots, x_n]$$

So, $e_i^\vee$ corresponds to the coset $[x_i] = x_i$.

- (e) $\mathbb{K}[V]$ is the coordinate ring of V because every polynomial is a combination of coordinate functions restricted to V . And $\mathbb{K}[V]$ is generated by these functions on V .

$$\mathbb{K}[V] = \mathbb{K}[e_1^\vee|_V, \dots, e_n^\vee|_V]$$

- (f) (i) Let $V = \mathbb{V}(y - x^2)$.

$$\mathbb{K}[V] \cong \mathbb{K}[x, y] / \mathbb{I}(y - x^2) = \mathbb{K}[x, x^2] = \mathbb{K}[x]$$

(ii) Let $V = \mathbb{V}(y^2, xy, x^{17})$.

Note that $\mathbb{I}(V) = \sqrt{\langle y^2, xy, x^{17} \rangle}$.

Since $\mathbb{K}$ is a field, if $(a, b) \in V$ then

$$a^{17} = 0 \implies a = 0$$

and

$$b^2 = 0 \implies b = 0$$

Then, $V = \{(0, 0)\}$. And $I(V) = I(\{(0, 0)\}) = \langle x, y \rangle$.

$$\mathbb{K}[V] \cong \mathbb{K}[x, y] / \mathbb{I}(\mathbb{V}(y^2, xy, x^{17})) = \mathbb{K}[x, y] / \langle x, y \rangle = \mathbb{K}$$

(iii) Let $V = \mathbb{V}(x^6 - 3x^5y + 3x^4y^2 + x^3y^3) = \mathbb{V}(x^3(x - y)^3)$.

Then if $(a, b) \in V$, either $a^3 = 0 \implies a = 0$ or $(x - y)^3 = 0 \implies x = y$.

Then $V = \{(0, a), (a, a) \mid a \in \mathbb{K}\} = \mathbb{V}(x) \cup \mathbb{V}(x - y)$.

$$\mathbb{I}(V) = \mathbb{I}(\mathbb{V}(x) \cup \mathbb{V}(x - y)) = \mathbb{I}(\mathbb{V}(x)) \cap \mathbb{I}(\mathbb{V}(x - y)) = \langle x \rangle \cap \langle x - y \rangle = \langle x(x - y) \rangle$$

And

$$\mathbb{K}[V] \cong \mathbb{K}[x, y] / \langle x(x - y) \rangle$$

(iv) Let $A = (x_{ij}) \in SL_3(\mathbb{F}_5)$, and $V = \mathbb{V}(\det(A) - 1, \text{tr}(A))$. Note that the trace formula is $x_{11} + x_{22} + x_{33}$, and we use fermat's little theorem to enforce staying in $\mathbb{F}_5$. Then the coordinate ring is,

$$\mathbb{K}[V] = \mathbb{F}_5[x_{11}, \dots, x_{33}] / \langle \det(A) - 1, x_{11} + x_{22} + x_{33}, x_{ij}^5 - x_{ij} \rangle$$

You should now think of $\mathbb{K}[V]$ as the algebraic avatar of V : its elements are precisely the polynomial functions on V , and the coordinate functions on $\mathbb{K}^n$ descend to generators of $\mathbb{K}[V]$. The next step is to recognize points of V purely algebraically, either as $\mathbb{K}$ -algebra maps $\mathbb{K}[V] \rightarrow \mathbb{K}$ or as maximal ideals whose residue fields are isomorphic to $\mathbb{K}$ as $\mathbb{K}$ -algebras.

Problem 6. Let R be a finitely generated $\mathbb{K}$ -algebra. Fix a presentation $R \cong \mathbb{K}[x_1, \dots, x_n]/I$. Describe a natural bijection between the following three sets:

- (i) points in $\mathbb{V}(I) \subset \mathbb{K}^n$,
- (ii) maximal ideals in R with residue field isomorphic to $\mathbb{K}$ as a $\mathbb{K}$ -algebra, and
- (iii) elements of $\text{Hom}_{\mathbb{K}\text{-alg}}(R, \mathbb{K})$, the set of morphisms from R to $\mathbb{K}$ in the category of $\mathbb{K}$ -algebras.

Solution.

Let $\mathbb{K}$ be algebraically closed.

Claim: Let $\pi : \mathbb{K}[x_1, \dots, x_n] \rightarrow \mathbb{K}[x_1, \dots, x_n]/I = R$, and let $J \subset R$ be maximal. We claim that $\pi^{-1}(J)$ is maximal in $\mathbb{K}[x_1, \dots, x_n]$.

Proof. We know that $\pi^{-1}(J)$ is an ideal. Let M be a maximal ideal containing $\pi^{-1}(J)$. Since π is surjective, we know that $\pi(M)$ is an ideal (worksheet 1.1, Q4.4), and $J \subseteq \pi(M)$. Then, since J is maximal, either $\pi(M) = J$ or $\pi(M)$ is the whole ring. $\pi(M) \neq R$ since ring homomorphisms are unital and $1 \notin M$. Hence, $M = \pi^{-1}(J)$ or $\pi^{-1}(J)$ is maximal. $\square$

(i) $\iff$ (ii)

Recall that $\mathbb{V}(I)$ contains points $p = (a_1, \dots, a_n)$ such that $f(p) = 0$ for all $f \in I$. Also,

$$\text{ev}_p : \mathbb{K}^n \rightarrow \mathbb{K}$$

By Hilbert's Nullstellensatz I, we know that maximal ideals of $\mathbb{K}[x_1, \dots, x_n]$ are of the form $\langle x_1 - a_1, \dots, x_n - a_n \rangle$, for some point $(a_1, \dots, a_n)$.

Problem 7. Zariski Topology on Algebraic Sets. Let $V \subseteq \mathbb{K}^n$ be an algebraic set. Define the *Zariski topology* on V by declaring the closed sets to be the subsets of V of the form $\mathbb{V}(I)$ for some ideal $I \subset \mathbb{K}[x_1, \dots, x_n]$. In other words, the closed sets are precisely the *algebraic subsets* of V .

- (a) Verify that this definition indeed determines a topology on V .
- (b) For $f \in \mathbb{K}[V]$, define

$$D(f) = \{p \in V \mid f(p) \neq 0\}.$$

Show that $\{D(f)\}$ form a basis for the Zariski topology on V .

- (c) For each point $p \in V$, let

$$\mathfrak{m}_p = \{f \in \mathbb{K}[V] \mid f(p) = 0\}$$

be the corresponding maximal ideal. Show that the map

$$\begin{aligned} V &\longrightarrow \text{Spec}(\mathbb{K}[V]) \\ p &\longmapsto \mathfrak{m}_p \end{aligned}$$

is a homeomorphism onto its image, which consists of those maximal ideals whose residue field is isomorphic to $\mathbb{K}$ as a $\mathbb{K}$ -algebra.

- (d) If $\mathbb{K}$ is algebraically closed how can the preceding part be rephrased in terms of $\text{mSpec}(\mathbb{K}[V])$?
- (e) Is the Zariski topology on V Hausdorff in general? If not, when is it?

Solution.

- (a) $V = \mathbb{V}(I)$ for some $I \subset \mathbb{K}[x_1, \dots, x_n]$. The closed sets of V are of the form $\mathbb{V}(J)$, where $I \subset J \subset \mathbb{K}[x_1, \dots, x_n]$. We need to check the following:
 - (i) $\emptyset$ and V are closed.
Note $\mathbb{V}(I) = V$, and $\mathbb{V}(\mathbb{K}[x_1, \dots, x_n]) = \emptyset$ is closed.
 - (ii) Finite union of closed sets is closed.
Let $\mathbb{V}(J), \mathbb{V}(K) \subseteq V$ be subsets. Then, $\mathbb{V}(I) \cup \mathbb{V}(J)$ is the set of points in $\mathbb{K}^n$ such that for all $p \in \mathbb{V}(I) \cup \mathbb{V}(J)$ either $f(p) = 0$ or $g(p) = 0$.
 - (iii) Arbitrary intersection of closed sets is closed.

Worksheet 3.1: Gröbner Bases Pt. II

Fix a field $\mathbb{K}$. Unless otherwise specified the ring throughout this worksheet will be $\mathbb{K}[x_1, \dots, x_n]$. In the previous worksheet we introduced monomial orderings and leading terms. In this worksheet we return to the multivariable division algorithm, and then use it to study monomial ideals, initial ideals, and Gröbner bases. The main goal is to push the story all the way to a major finiteness theorem: every ideal of $\mathbb{K}[x_1, \dots, x_n]$ is finitely generated.

Recall a *monomial ordering* on $\mathbb{K}[x_1, \dots, x_n]$ is a total well-ordering $<$ on $\mathbb{Z}_{\geq 0}^n$ such that if $\alpha < \beta$ then $\alpha + \gamma < \beta + \gamma$ for all $\gamma \in \mathbb{Z}_{\geq 0}^n$. Using the bijection between monomials in $\mathbb{K}[x_1, \dots, x_n]$ and elements of $\mathbb{Z}_{\geq 0}^n$ we think of a monomial ordering as giving us a way to compare monomials by saying $x^\alpha < x^\beta$ if and only if $\alpha < \beta$. In Worksheet 1.3 we saw several important monomial orders: lexicographic (lex) order, graded lexicographic (grlex) order, and graded reverse lexicographic (grevlex) order. Please review that worksheet for their definitions.

Returning to our goal of having a division algorithm in $\mathbb{K}[x_1, \dots, x_n]$ we are faced with two challenges compared to the familiar setting of $\mathbb{K}[x]$:

- (i) In several variables the notion of “leading term” is ambiguous. In one variable, the leading term is simply the term of highest degree. In several variables this no longer suffices: for instance, what should the leading term of $x^2 + xy + y^2$ be? All three terms have total degree 2.
- (ii) We want a division algorithm that can answer questions about ideals, i.e., membership, intersections, sums, products, radicals. In $\mathbb{K}[x]$ every ideal is principal, so dividing by a single generator is enough. For $n > 1$, however, $\mathbb{K}[x_1, \dots, x_n]$ is no longer a PID, so ideals generally require multiple generators. We therefore need a way to divide a polynomial by an entire *list* of polynomials.

As seen on Worksheet 1.3, once a monomial ordering has been fixed, we give a meaningful definition of leading term, leading monomial, and leading coefficient; solving issue (i). Note while we normally write $\text{LT}(f)$, $\text{LM}(f)$ and $\text{LC}(f)$ for these quantities; they do depend on the chosen order $<$, and so would more properly be called $\text{LT}_{<}(f)$, $\text{LM}_{<}(f)$ and $\text{LC}_{<}(f)$ to highlight this fact.

To resolve issue (ii), we generalize long division as follows. In single-variable polynomial long division, we repeatedly ask: “Does the leading term of the divisor divide the current leading term of the dividend?” If yes, we subtract off the appropriate multiple; if not, we are done. The multivariable algorithm works the same way, except that instead of checking a single divisor we scan through an *ordered* list of divisors $G = (g_1, \dots, g_s)$ and use the first one whose leading term divides the current leading term. More precisely, the algorithm maintains a “working polynomial” p (initialized to f), quotients $q_1, \dots, q_s$ (initialized to 0), and a remainder r (initialized to 0), then repeats:

- **Try to Divide.** Walk through the list $g_1, g_2, \dots, g_s$ in order. If you find some g_i whose leading term $\text{LT}(g_i)$ divides $\text{LT}(p)$, subtract the appropriate multiple: update $q_i \leftarrow q_i + \text{LT}(p)/\text{LT}(g_i)$ and $p \leftarrow p - (\text{LT}(p)/\text{LT}(g_i))g_i$. Then return to the start of the list and try again with the new p .
- **Move to Remainder.** If no g_i in the list has a leading term dividing $\text{LT}(p)$, then the current leading term of p cannot be reduced further. Move it to the remainder: $r \leftarrow r + \text{LT}(p)$ and $p \leftarrow p - \text{LT}(p)$.

We continue until the working polynomial p is equal to 0. Proving this algorithm terminates as expected, which you will do later in this worksheet, amounts essentially to proving the following theorem

Theorem 0.5. Fix a monomial ordering $<$ on $\mathbb{K}[x_1, \dots, x_n]$ and let $G = (g_1, \dots, g_s)$ be an ordered list of nonzero polynomials. If $f \in \mathbb{K}[x_1, \dots, x_n]$ then there exist $q_1, \dots, q_s, r \in \mathbb{K}[x_1, \dots, x_n]$ such that

$$f = q_1 g_1 + \dots + q_s g_s + r,$$

where no monomial appearing in r is divisible by any of $\text{LM}(g_1), \dots, \text{LM}(g_s)$.

Written in pseudocode the algorithm for multivariable polynomial division is shown below.

Algorithm 2 Division by an ordered list $G = (g_1, \dots, g_s)$

Require: $f, g_1, \dots, g_s$

Ensure: $q_1, \dots, q_s, r$

$q_1, \dots, q_s \leftarrow 0; r \leftarrow 0; p \leftarrow f$

while $p \neq 0$ **do**

\leftarrow 0

for $i = 1, \dots, s$ **do**

if $\text{LT}(g_i)$ divides $\text{LT}(p)$ **then**

$q_i \leftarrow q_i + \text{LT}(p)/\text{LT}(g_i)$

$p \leftarrow p - (\text{LT}(p)/\text{LT}(g_i))g_i$

 divides $\leftarrow 1$

break

end if

end for

if divides = 0 **then**

$r \leftarrow r + \text{LT}(p)$

$p \leftarrow p - \text{LT}(p)$

end if

end while

Problem 1. Review of Monomial Orders.

Work in $\mathbb{K}[x, y, z]$ and assume $x > y > z$.

- Order the monomials x^2y , xy^2 , xz^2 , y^3 , yz^4 from largest to smallest with respect to lex, graded lex, and graded reverse lex.
- Compute $\text{LM}(f)$, $\text{LC}(f)$, and $\text{LT}(f)$ for $f = 4x^2y - 3xy^2 + y^4 + 2z^5$ with respect to lex and grevlex.
- Prove that 1 is the smallest monomial for every monomial ordering.
- Deduce that every strictly decreasing sequence of monomials terminates.
- Explain why part (d) should make you optimistic that a division algorithm might terminate.

Solution.

- Lex: $x^2y, xy^2, xz^2, y^3, yz^4$
 - Graded Lex: $yz^4, x^2y, xy^2, xz^2, y^3$
 - Graded RevLex: $yz^4, x^2y, xy^2, y^3, xz^2$

(b) Lex:

$$\text{LM}(f) = x^2y, \quad \text{LC}(f) = 4, \quad \text{LT}(f) = 4x^2y$$

Grevlex:

$$\text{LM}(f) = z^5, \quad \text{LC}(f) = 2, \quad \text{LT}(f) = 2z^5$$

- The monomial 1 corresponds to $(0, \dots, 0)$. For the sake of contradiction assume there exist some $\alpha = (a_1, \dots, a_n)$ such that $\alpha \neq 1$ and $\alpha < 1$. Then, by property (ii) of monomial orders we have

$$\alpha + \alpha < 0 + \alpha$$

Applying (ii) again,

$$3\alpha < 2\alpha$$

And so on,

$$\dots < 4\alpha < 3\alpha < 2\alpha < \dots$$

But this violates the well-ordering property (iii) of monomial orderings, a contradiction! Hence, 1 must be the smallest monomial in every monomial ordering.

- Since we have a minimal element, every strictly decreasing sequence of monomials terminates.

- (e) In a division algorithm, at each step we reduce the polynomial f by replacing its leading term with a smaller combination of terms coming from the divisors. This produces a strictly decreasing sequence of leading monomials and by part (d) every such sequence must terminate. Hence, eventually the algorithm must terminate.

Problem 2. Examples of Multivariable Division.

For this question, we will work in $\mathbb{K}[x, y]$ considered with the lex order where $x > y$.

1. Divide $f = x^2y + xy + y - x - 2$ by the ordered list $G = (xy - 1, y - 1)$. Record the successive values of p , the quotients, and the remainder after each update.
2. Divide $f = xy^2 - x$ by the ordered list $G = (xy - 1, y^2 - 1)$. Record the successive values of p , the quotients, and the remainder after each update.
3. Repeat (b), but with the ordered list $(y^2 - 1, xy - 1)$, i.e. where the order of G is swapped.
4. Show directly that $xy^2 - x \in \langle xy - 1, y^2 - 1 \rangle$.
5. What do your computations show about division in several variables? In particular, what is always true when the remainder is 0, and what can go wrong when the remainder is non-zero?

Solution.

1. **Step 0:** $q_1 = 0, q_2 = 0, p = x^2y + xy + y - x - 2, r = 0$

Step 1: $q_1 = x, q_2 = 0, p = xy + y - 2, r = 0$

Step 2: $q_1 = xy, q_2 = 0, p = y - 1, r = 0$

Step 3: $q_1 = xy, q_2 = y, p = 0, r = 0$

2. **Step 0:** $q_1 = 0, q_2 = 0, p = xy^2 - x, r = 0$

Step 1: $q_1 = y, q_2 = 0, p = -x + y, r = 0$

Step 2: $q_1 = y, q_2 = 0, p = y, r = -x$

Step 3: $q_1 = y, q_2 = 0, p = 0, r = -x + y$

3. **Step 0:** $q_1 = 0, q_2 = 0, p = xy^2 - x, r = 0$

Step 1: $q_1 = x, q_2 = 0, p = 0, r = 0$

4. From part 3 we know that $xy^2 - x \in \langle y^2 - 1 \rangle \implies xy^2 - x \in \langle xy - 1, y^2 - 1 \rangle$.

5. If the remainder is zero the polynomial is in the ideal generated by the divisors. The converse is not true; that is if the remainder is non-zero, we can't conclude that the polynomial is not in the ideal.

Problem 3. Termination of Division Algorithm

Let $G = (g_1, \dots, g_s)$ be an ordered list of non-zero polynomials.

1. Show that throughout the algorithm one always has $f = q_1g_1 + \dots + q_sg_s + p + r$.
2. Show that after each pass through the `while` loop, the leading monomial of p strictly decreases with respect to the given monomial order.
3. Use Question 1(d) to prove that the algorithm terminates.
4. Prove that on termination the output satisfies $f = q_1g_1 + \dots + q_sg_s + r$, and no monomial appearing in r is divisible by any of $\text{LM}(g_1), \dots, \text{LM}(g_s)$.
5. Explain why a zero remainder certifies ideal membership: if dividing f by G produces remainder 0, then $f \in \langle g_1, \dots, g_s \rangle$.

Solution.

1. We will show this by induction on the number of steps.

Base Case: It holds initially since $p = f$ and $q_i = r = 0$.

Inductive Step: Assume that the invariant $f = q_1g_1 + \dots + q_sg_s + p + r$ is maintained for the first $n - 1$ steps, then at the n -th step, we either,

- Replace $q_i \leftarrow q_i + \frac{\text{LT}(p)}{\text{LT}(g_i)}$ and $p \leftarrow p - \frac{\text{LT}(p)}{\text{LT}(g_i)}g_i$, so the sum $q_ig_i + p$ is unchanged.
- Or replace $r \leftarrow r + \text{LT}(p)$ and $p \leftarrow p - \text{LT}(p)$, so $p + r$ is unchanged.

In both cases the invariant is preserved, thus we always have $f = q_1g_1 + \dots + q_sg_s + p + r$.

2. If a g_i divides the leading term, we multiply it with q_i and subtract the leading term, if no g_i divides the leading term, we subtract it from p (and add it to r), either ways each pass through the `while` loop removes the leading monomial and thus the leading monomial of p strictly decreases with respect to the given monomial order.
3. Since every strictly decreasing sequence of monomials terminates (Question 1(d)), p must also terminate, i.e. $p = 0$ eventually and the algorithm terminates.
4. From part 1 we know that we always have $f = q_1g_1 + \dots + q_sg_s + p + r$, since on termination we have $p = 0$ we get,

$$f = q_1g_1 + \dots + q_sg_s + r$$

Moreover, since the algorithm only adds $\text{LT}(p)$ to r when $\text{LT}(g_i) \nmid \text{LT}(p)$ for all i no monomial in r is divisible by any of $\text{LM}(g_1), \dots, \text{LM}(g_s)$.

5. If $r = 0$ we have $f = q_1g_1 + \cdots + q_sg_s$ which is precisely the condition for ideal membership.

Worksheet 3.2: The Nullstellensatz

Throughout this course, “ring” means *commutative* ring with unity. In this worksheet $\mathbb{K}$ will denote a field. In the previous worksheet you built the order-reversing maps

$$I \longmapsto \mathbb{V}(I) \quad \text{and} \quad V \longmapsto \mathbb{I}(V)$$


connecting ideals in $\mathbb{K}[x_1, \dots, x_n]$ with algebraic sets in $\mathbb{K}^n$. Our goal now is to prove that over an algebraically closed field these constructions *almost* recover each other exactly. The only restriction being that since $\mathbb{I}(V)$ is always a radical ideal, these functions are only bijections when restricted to the set of radical ideals in $\mathbb{K}[x_1, \dots, x_n]$. More precisely, our goal is to prove the first two versions of Hilbert’s Nullstellensatz under the condition that $\mathbb{K}$ is uncountable (i.e. $\mathbb{C}$). Although our work will also show the second version follows from the first for any algebraically closed field.

Theorem 0.6 (Hilbert’s Nullstellensatz I). *If $\mathbb{K}$ is algebraically closed, then every maximal ideal of $\mathbb{K}[x_1, \dots, x_n]$ is of the form $\langle x_1 - a_1, \dots, x_n - a_n \rangle$ for some point $(a_1, \dots, a_n) \in \mathbb{K}^n$.*

Theorem 0.7 (Hilbert’s Nullstellensatz II). *If $\mathbb{K}$ is algebraically closed and $V = \mathbb{V}(\mathcal{S}) \subset \mathbb{K}^n$ is an algebraic set, for some subset $\mathcal{S} \subset \mathbb{K}[x_1, \dots, x_n]$, then $\mathbb{I}(V) = \sqrt{\langle \mathcal{S} \rangle}$.*

For a point $p = (a_1, \dots, a_n) \in \mathbb{K}^n$, we will often write $\mathfrak{m}_p$ for the ideal $\langle x_1 - a_1, \dots, x_n - a_n \rangle \subset \mathbb{K}[x_1, \dots, x_n]$, which by the previous worksheet is the kernel of the evaluation-at- p map $\text{ev}_p : \mathbb{K}[x_1, \dots, x_n] \rightarrow \mathbb{K}$. An alternative statement of Hilbert’s Nullstellensatz I is to say there is a bijection between $\mathbb{K}^n$ and maximal ideals in $\mathbb{K}[x_1, \dots, x_n]$ given by $p \mapsto \mathfrak{m}_p$. We say that this statement can fail when $\mathbb{K}$ is not algebraically closed, e.g., $\langle x^2 + 1 \rangle \subset \mathbb{R}[x]$.

Since points are themselves algebraic sets, the second version of Hilbert’s Nullstellensatz can be thought of as a generalization of the first. In particular Nullstellensatz II can be thought of as saying that $\mathbb{V}$ and $\mathbb{I}$ are mutual inverses giving bijections



Somewhat amazingly, the entire difficulty of this theorem comes down to only one of the four inclusions needed. In particular, last week we showed that $\mathbb{V}(\mathbb{I}(V)) = V$ for any algebraic subset $V \subset \mathbb{K}^n$, regardless of the field. Similarly, we also proved that the inclusion $\sqrt{I} \subseteq \mathbb{I}(\mathbb{V}(I))$ holds over an arbitrary field. Thus, the hard part of the proof of Hilbert’s Nullstellensatz II is to show that

$$\mathbb{I}(\mathbb{V}(I)) \subseteq \sqrt{I}$$

for every ideal $I \subseteq \mathbb{K}[x_1, \dots, x_n]$. This can fail over non-algebraically closed fields, as the example $I = \langle x^2 + 1 \rangle \subset \mathbb{R}[x]$ shows.

The strategy of this worksheet is to isolate that difficulty: first review the geometry–algebra dictionary, and then prove the point Hilbert’s Nullstellensatz I for uncountable fields using field theory and a clever cardinality argument. We will then deduce what some call the weak Nullstellensatz:

Theorem 0.8 (Hilbert’s Weak Nullstellensatz). *If $\mathbb{K}$ is algebraically closed and $I \subset \mathbb{K}[x_1, \dots, x_n]$ is an ideal then $\mathbb{V}(I) = \emptyset$ if and only if $I = \langle 1 \rangle$.*

Note the difficult direction of the weak Nullstellensatz is the forward direction, the reverse is immediate. Intuitively this is saying that a system of polynomial equations has no solution over $\mathbb{K}$ if and only if the “obvious” obstruction – namely you can write 1 in terms of your polynomials – occurs. Finally we will use Rabinowitsch’s trick to obtain the full statement. Along the way we will also pause to recall some field theory language, since the key obstruction is the difference between algebraic and transcendental field extensions.

Problem 1. Warm-Up: Recalling the Correspondence.

Recall the two order-reversing maps

$$I \longmapsto \mathbb{V}(I) \quad \text{and} \quad V \longmapsto \mathbb{I}(V).$$

Before we move to proving the main theorems let us warm up with a few examples.

1. Give the definitions of $\mathbb{V}(I)$ and $\mathbb{I}(V)$.
2. If $V = \{p\} \subset \mathbb{K}^n$ for $p = (a_1, \dots, a_n)$, describe $\mathbb{I}(V)$ both intrinsically and by giving generators.
3. The ideals $\langle 0 \rangle$ and $\langle 1 \rangle$ are the extreme cases. Describe $\mathbb{V}(\langle 0 \rangle)$ and $\mathbb{V}(\langle 1 \rangle)$ as subsets of $\mathbb{K}^n$.
4. Over $\mathbb{R}$, describe $\mathbb{V}(\langle x, y \rangle)$ and $\mathbb{V}(\langle x^2 + y^2 \rangle) \subseteq \mathbb{R}^2$.
5. Over $\mathbb{F}_p$, describe $\mathbb{V}(\langle 0 \rangle)$ and $\mathbb{V}(\langle x^p - x, xy^p - x^p y \rangle) \subseteq \mathbb{F}_p^2$.

Solution.

1. For $I, V \in \mathbb{K}^n$,

$$\mathbb{V}(I) = \{p \in \mathbb{K}^n \mid f(p) = 0, \forall f \in I\}$$

$$\mathbb{I}(V) = \{f \in \mathbb{K}[x_1, \dots, x_n] \mid f(p) = 0, \forall p \in V\}$$

2. If $V = \{p\}$ then $\mathbb{I}(V)$ is the ideal generated by all functions that vanish on p . The generator is $\langle (x_1 - a_1), (x_2 - a_2), \dots, (x_n - a_n) \rangle$.
3. $\mathbb{V}(\langle 0 \rangle) = \mathbb{K}^n$ and $\mathbb{V}(\langle 1 \rangle) = \emptyset$.

4. Over $\mathbb{R}$,

$$\mathbb{V}(\langle x, y \rangle) = \{(0, 0)\}$$

And,

$$\mathbb{V}(\langle x^2 + y^2 \rangle) = \{(x^2, -x^2)\} = \{(0, 0)\}$$

5. Over $\mathbb{F}_p^2$,

$$\mathbb{V}(\langle 0 \rangle) = \mathbb{F}_p^2$$

And

$$\mathbb{V}(x^p - x, xy^p - x^p y) = \mathbb{F}_p^2$$

Problem 2. A Parameterized Example.

Consider the map

$$\phi : \overline{\mathbb{F}}_{17}^2 \longrightarrow \overline{\mathbb{F}}_{17}^3, \quad (u, v) \longmapsto (u^2, uv, v^2).$$

1. Show that every point in the image of ϕ satisfies the equation $xz - y^2 = 0$.
2. Let $(a, b, c) \in \overline{\mathbb{F}}_{17}^3$ satisfy $ac = b^2$. Show that if $a \neq 0$, then (a, b, c) lies in the image of ϕ . Handle the case $a = 0$ separately.
3. Deduce that the image of ϕ is equal to $\mathbb{V}(xz - y^2) \subseteq \overline{\mathbb{F}}_{17}^3$.
4. Show that $xz - y^2$ is irreducible in $\overline{\mathbb{F}}_{17}[x, y, z]$ and conclude that the corresponding radical ideal is $\langle xz - y^2 \rangle$.

Solution.

1. Let $p \in \text{im}(\phi)$, Then $p = (u^2, uv, v^2) = (x, y, z)$ for some $u, v \in \overline{\mathbb{F}}_{17}$. So,

$$xz - y^2 = u^2v^2 - (uv)^2 = 0$$

2. If $a \neq 0$, we have $(a, b, c) = (a, \sqrt{ac}, c)$ and since $\overline{\mathbb{F}}_{17}$ is algebraically closed let $u^2 = a$ and $v = b/u$ Then we have,

$$\phi(u, v) = (a, b, b^2/a) = (a, b, c)$$

If $a = 0$, then $b^2 = ac = 0$ so we are free to pick any $v \in \overline{\mathbb{F}}_{17}$ such that $v^2 = c$ and

$$\phi(0, v) = (0, 0, c) = (a, b, c)$$

3. Thus $\text{im}(\phi) = \{p \in \overline{\mathbb{F}}_{17}^3 \mid (xz - y^2)(p) = 0\} = \mathbb{V}(xz - y^2)$.
4. Note that $\overline{\mathbb{F}}_{17}[x, y, z] / \langle xz - y^2 \rangle \cong \overline{\mathbb{F}}_{17}[x, z, \sqrt{xz}]$ is an integral domain, so $xz - y^2$ is prime and hence also irreducible. Since it is prime, the corresponding radical ideal is $\langle xz - y^2 \rangle$.

Problem 3. Reduction to the Core Difficulty.

Recall what you proved last time about the interaction of $\mathbb{V}$ and $\mathbb{I}$ over an arbitrary field.

1. Show that if $V \subseteq \mathbb{K}^n$ is algebraic, then $\mathbb{V}(\mathbb{I}(V)) = V$.
2. Show that for every ideal $I \subseteq \mathbb{K}[x_1, \dots, x_n]$ one has $I \subseteq \sqrt{I} \subseteq \mathbb{I}(\mathbb{V}(I))$.
3. Show via an example – other than $\langle x^2 + 1 \rangle \subset \mathbb{R}[x]$ – that the inclusion in 2 can be proper when $\mathbb{K}$ is not algebraically closed
4. Explain why proving Hilbert's Nullstellensatz II reduces to showing that when $\mathbb{K}$ is algebraically closed, $\mathbb{I}(\mathbb{V}(I)) \subset \sqrt{I}$ for every ideal $I \subset \mathbb{K}[x_1, \dots, x_n]$.

Solution.

1. Let $V = \mathbb{V}(\mathcal{S})$.

$$\mathbb{V}(\mathcal{S}) \subseteq \mathbb{V}(\mathbb{I}(\mathbb{V}(\mathcal{S})))$$

This containment is obvious by construction. $\mathbb{I}(\mathbb{V}(\mathcal{S}))$ is the set of all functions that vanish at all points in $\mathbb{V}(\mathcal{S})$. And $\mathbb{V}(\mathbb{I}(\mathbb{V}(\mathcal{S})))$ is the set of all points that all functions in $\mathbb{I}(\mathbb{V}(\mathcal{S}))$ vanish at. So, $\mathbb{V}(\mathcal{S}) \subseteq \mathbb{V}(\mathbb{I}(\mathbb{V}(\mathcal{S})))$.

$$\mathbb{V}(\mathcal{S}) \supseteq \mathbb{V}(\mathbb{I}(\mathbb{V}(\mathcal{S})))$$

Note that by definition $\mathcal{S} \subseteq \mathbb{I}(\mathbb{V}(\mathcal{S}))$. Then by 2(a) $\mathbb{V}(\mathbb{I}(\mathbb{V}(\mathcal{S}))) \subseteq \mathbb{V}(\mathcal{S})$.

Hence, $\mathbb{V}(\mathcal{S}) = \mathbb{V}(\mathbb{I}(\mathbb{V}(\mathcal{S})))$.

2. Let $V = \mathbb{V}(I)$. We know that $\langle I \rangle \subseteq \sqrt{\langle I \rangle}$. So showing that $\sqrt{\langle I \rangle} \subseteq \mathbb{I}(V)$ suffices. Note that $\mathbb{I}(V) = \mathbb{I}(\mathbb{V}(I))$ which is the set of all functions that evaluate to 0 at p , for all $p \in \mathbb{V}(I)$. And $\mathbb{V}(I)$ is the set of all points p such that all functions $f \in I$ evaluate to 0 at p . Let $f \in \sqrt{\langle I \rangle}$. Then by definition for some n , $f^n \in I$. We can then write,

$$f^n = \sum_i h_i(p) g_i(p), \quad h_i \in \mathbb{K}[x_1, \dots, x_n], g_i \in I$$

But then,

$$f^n(p) = \sum_i h_i(p) g_i(p) = \sum_i h_i(p) \cdot 0 = 0$$

Moreover since $\mathbb{K}$ is a domain, $f^n(p) = 0 \implies f(p) = 0 \implies f$ is a function that evaluates to 0 at p , hence $f \in \mathbb{I}(V)$.

3. Let $I = \langle x^3 + 1 \rangle = \sqrt{I}$. And $\mathbb{I}(\mathbb{V}(I)) = \mathbb{I}(\{-1\}) = \langle x + 1 \rangle$.

4. The statement is for every ideal $I \subseteq \mathbb{K}^n$, $\mathbb{I}(\mathbb{V}(I)) = \sqrt{I}$. Since we have already shown one containment, it only remains to show that $\mathbb{I}(\mathbb{V}(I)) \subseteq \sqrt{I}$.

Before the next question, recall some field theory language. If $\mathbb{K}$ is a field and $\mathbb{E}$ is a field containing $\mathbb{K}$, we call $\mathbb{E}/\mathbb{K}$ a *field extension*. In this situation $\mathbb{E}$ is naturally a vector space over $\mathbb{K}$, and we write $[\mathbb{E} : \mathbb{K}] := \dim_{\mathbb{K}} \mathbb{E}$ for its dimension, called the *degree* of the extension. If $[\mathbb{E} : \mathbb{K}]$ is finite we say the extension is *finite*.

Fixing a field extension $\mathbb{E}/\mathbb{K}$, we say an element $\zeta \in \mathbb{E}$ is *algebraic* over $\mathbb{K}$ if there exists a nonzero polynomial $f \in \mathbb{K}[x]$ with $f(\zeta) = 0$; otherwise ζ is *transcendental* over $\mathbb{K}$. The extension $\mathbb{E}/\mathbb{K}$ is called *algebraic* if every element of $\mathbb{E}$ is algebraic over $\mathbb{K}$. Every finite extension is algebraic: if $[\mathbb{E} : \mathbb{K}] = d$, then for any $\zeta \in \mathbb{E}$ the $d + 1$ elements $\zeta^0, \zeta, \dots, \zeta^d$ are $\mathbb{K}$ -linearly dependent, giving a polynomial relation for ζ over $\mathbb{K}$.

More generally, elements $\zeta_1, \dots, \zeta_r \in \mathbb{E}$ are *algebraically independent* over $\mathbb{K}$ if no nonzero polynomial $f \in \mathbb{K}[x_1, \dots, x_r]$ satisfies $f(\zeta_1, \dots, \zeta_r) = 0$. In particular, a single element ζ is transcendental over $\mathbb{K}$ exactly when $\{\zeta\}$ is algebraically independent over $\mathbb{K}$. One helpful characterization of transcendental elements can be phrased in terms of our trusty friend evaluation maps.

Lemma 0.9. *Let $\mathbb{E}/\mathbb{K}$ be a field extension. An element $\zeta \in \mathbb{E}$ is transcendental over $\mathbb{K}$ if and only if the evaluation-at- ζ homomorphism below is injective:*

$$\begin{array}{ccc} \mathbb{K}[x] & \xrightarrow{\text{ev}_\zeta} & \mathbb{E} \\ f & \longmapsto & f(\zeta) \end{array} .$$

A useful consequence of this lemma is that if $\zeta \in \mathbb{E}$ is transcendental over $\mathbb{K}$, then the map ev_ζ identifies $\mathbb{K}[x]$ with the subring $\mathbb{K}[\zeta] \subseteq \mathbb{E}$. Passing to fields of fractions then identifies the rational function field $\mathbb{K}(x)$ with the subfield $\mathbb{K}(\zeta) \subseteq \mathbb{E}$. You will prove the converse in the exercises as well.

A field $\mathbb{K}$ is *algebraically closed* if every nonconstant polynomial in $\mathbb{K}[x]$ has a root in $\mathbb{K}$. An equivalent formulation is that $\mathbb{K}$ has no nontrivial algebraic extensions. That is, if $\mathbb{E}/\mathbb{K}$ is algebraic, then $\mathbb{E} = \mathbb{K}$. The field $\mathbb{C}$ of complex numbers is the prototypical example.

A $\mathbb{K}$ -*algebra* is a ring R equipped with a ring homomorphism $\alpha : \mathbb{K} \rightarrow R$. Given such a map, R becomes a $\mathbb{K}$ -vector space by defining $\lambda \cdot a := \alpha(\lambda)a$ for $\lambda \in \mathbb{K}$ and $a \in R$. This scalar multiplication is compatible with the ring multiplication in the sense that

$$\lambda \cdot (ab) = (\lambda \cdot a)b = a(\lambda \cdot b)$$

for all $a, b \in R$ and $\lambda \in \mathbb{K}$. Conversely, any ring R that is simultaneously a $\mathbb{K}$ -vector space satisfying the above compatibility condition is a $\mathbb{K}$ -algebra; the structure map is $\alpha(\lambda) = \lambda \cdot 1_R$. A *homomorphism of $\mathbb{K}$ -algebras* $\phi : R \rightarrow S$ is a ring homomorphism that also respects the $\mathbb{K}$ -action, i.e. $\phi(\lambda \cdot_R a) = \lambda \cdot_S \phi(a)$ for all $\lambda \in \mathbb{K}$ and $a \in R$. Equivalently, the

diagram

$$\begin{array}{ccc}
 & \mathbb{K} & \\
 \alpha_R \swarrow & & \searrow \alpha_S \\
 R & \xrightarrow{\phi} & S
 \end{array}$$

commutes, where α_S and α_R are the respective structure maps. In other words, ϕ is a ring homomorphism that is also $\mathbb{K}$ -linear. For example, $R = \mathbb{K}[x_1, \dots, x_n]$ is a $\mathbb{K}$ -algebra via the inclusion of constants, and any quotient R/I inherits a $\mathbb{K}$ -algebra structure from the surjection $\pi : R \twoheadrightarrow R/I$. If $I \subseteq \mathbb{K}[x_1, \dots, x_n]$ is an ideal, the quotient $\mathbb{K}[x_1, \dots, x_n]/I$ is a $\mathbb{K}$ -algebra. It is a field if and only if I is maximal. In that case, the quotient map π gives a field extension:

$$\mathbb{K} \longrightarrow \mathbb{K}[x_1, \dots, x_n] \xrightarrow{\pi} \mathbb{K}[x_1, \dots, x_n]/I$$

gives a field extension of $\mathbb{K}$. Every element of this quotient is a $\mathbb{K}$ -linear combination of the images of monomials $x_1^{a_1} \cdots x_n^{a_n}$, so properties of this extension (such as its dimension over $\mathbb{K}$) are controlled by the combinatorics of monomials together with the relations imposed by I .

The problem below uses all of these ideas to prove a beautiful dictionary between algebra and geometry: when $\mathbb{K}$ is both uncountable and algebraically closed (e.g. $\mathbb{K} = \mathbb{C}$), the maximal ideals of $\mathbb{K}[x_1, \dots, x_n]$ are in natural bijection with the points of $\mathbb{K}^n$.

Problem 4. Hilbert's Nullstellensatz I ($\mathbb{K}$ uncountable).

Let $R := \mathbb{K}[x_1, \dots, x_n]$ and $\mathfrak{m} \subset R$ be a maximal ideal. Write $L := R/\mathfrak{m}$ for the residue field, which contains $\mathbb{K}$ via the natural quotient map. The goal is to show that if $\mathbb{K}$ is uncountable and algebraically closed, then in fact $L \cong \mathbb{K}$.

1. Show that the monomials in R form a countable $\mathbb{K}$ -basis of R .
2. Deduce that L has at most countable dimension as a vector space over $\mathbb{K}$.
3. Let $\mathbb{E}/\mathbb{K}$ be a field extension and let $\zeta \in \mathbb{E}$. Explain why the following are equivalent:
 - (i) ζ is transcendental over $\mathbb{K}$;
 - (ii) $\{\zeta\}$ is algebraically independent over $\mathbb{K}$;
 - (iii) the evaluation map $\mathbb{K}[x] \rightarrow \mathbb{E}$ sending $f(x)$ to $f(\zeta)$ is injective.

In particular, if a field extension of $\mathbb{K}$ is not algebraic, then it contains a transcendental element.

4. Let $\mathbb{K}$ be any infinite field, and let $\zeta \in \mathbb{E}$ be transcendental over $\mathbb{K}$ for some field extension $\mathbb{E}/\mathbb{K}$. Prove that the set of elements in $\mathbb{E}$:

$$\left\{ \frac{1}{\zeta - \lambda} \mid \lambda \in \mathbb{K} \right\} \subseteq \mathbb{E}$$

is linearly independent over $\mathbb{K}$. (Hint: Suppose $\sum_{i=1}^m \frac{c_i}{\zeta - \lambda_i} = 0$ for distinct $\lambda_1, \dots, \lambda_m \in \mathbb{K}$ and coefficients $c_1, \dots, c_m \in \mathbb{K}$. Multiply both sides by $\prod_{j=1}^m (\zeta - \lambda_j)$ to obtain $\sum_{i=1}^m c_i \prod_{j \neq i} (\zeta - \lambda_j) = 0$. Now define $P(x) = \sum_{i=1}^m c_i \prod_{j \neq i} (x - \lambda_j) \in \mathbb{K}[x]$. Then $P(\zeta) = 0$. Since ζ is transcendental over $\mathbb{K}$, conclude that $P(x)$ must be the zero polynomial. Evaluate $P(x)$ at $x = \lambda_k$ for each k to show that $c_k = 0$.)

5. Now assume that $\mathbb{K}$ is uncountable. Using part 4, show that if a field extension $\mathbb{E}/\mathbb{K}$ contains a transcendental element, then $[\mathbb{E} : \mathbb{K}]$ is uncountable.
6. Assume $\mathbb{K}$ is uncountable. Show if $\mathfrak{m} \subseteq R$ is a maximal ideal, then the extension $\mathbb{K} \hookrightarrow R/\mathfrak{m} := L$ is algebraic.
7. Now assume that $\mathbb{K}$ is both uncountable and algebraically closed. Show that every algebraic extension of $\mathbb{K}$ is trivial, so $L \cong \mathbb{K}$ as $\mathbb{K}$ -algebras.
8. With the notation from part 7 let $\psi : R \rightarrow \mathbb{K}$ be the ring map given by the composition

$$R \xrightarrow{\pi} R/\mathfrak{m} = L \xrightarrow{\sim} \mathbb{K}$$

where π is the quotient map and the second map is the identification of L with $\mathbb{K}$ from part 7. Note $\ker(\psi) = \mathfrak{m}$. Define $a_i := \psi(x_i) \in \mathbb{K}$ for $i = 1, \dots, n$. Prove that $\mathfrak{m} = \langle x_1 - a_1, \dots, x_n - a_n \rangle$.

9. Deduce that when $\mathbb{K}$ is uncountable and algebraically closed, the assignment $p \mapsto \mathfrak{m}_p$ gives a bijection between points of $\mathbb{K}^n$ and maximal ideals of $\mathbb{K}[x_1, \dots, x_n]$.

Solution.

1. We know that $x_1^{a_1} \cdots x_n^{a_n}$ where $a_i \in \mathbb{Z}_{\geq 0}$ forms a $\mathbb{K}$ basis of R . It is countable because there is a bijection $x_1^{a_1} \cdots x_n^{a_n} \leftrightarrow \mathbb{Z}_{\geq 0}^n$, which is a finite product of countable sets and hence is countable.
2. We know that $\mathbb{K} \subseteq L \subseteq R$. Since R is countable dimensional over R , L must be at most countable dimensional over R .

3. (1 $\implies$ 2) by definition.

(2 $\implies$ 3)

If ζ is algebraically-independent over $\mathbb{K}$, it satisfies no non-zero polynomial over $\mathbb{K}[x]$, i.e. if $f(\zeta) = 0$ then $f = 0$, i.e. the evaluation at ζ function is injective.

(3 $\implies$ 1)

If the evaluation at ζ map is injective, then it satisfies no non-zero polynomial over $\mathbb{K}[x]$ and hence is transcendental over $\mathbb{K}$.

4. If $\mathbb{K}$ is uncountable and $\mathbb{E}/\mathbb{K}$ contains a transcendental element then by part 4 the set of elements

$$\left\{ \frac{1}{\zeta - \lambda} \mid \lambda \in \mathbb{K} \right\} \subseteq \mathbb{E}$$

is linearly independent and hence $\mathbb{E}$ has a uncountable $\mathbb{K}$ basis.

5. Recall that R has a countable $\mathbb{K}$ -basis, and $R/\mathfrak{m}$ is a field ($\mathfrak{m}$ maximal). Assume for the sake of contradiction that $L/\mathbb{K}$ is not algebraic. Then, $\exists t \in L$ such that

$$\left\{ \frac{1}{t - a} \mid a \in \mathbb{K} \right\} \subseteq L$$

are elements in L (since t is transcendental $t - a \neq 0$ and since L is a field the inverses exist.) But then by part 5 L has an uncountable number of independent elements over $\mathbb{K}$ which is a contradiction to R having a countable $\mathbb{K}$ -basis.

6. If $\mathbb{K}$ is algebraically closed, every polynomial with coefficients in $\mathbb{K}$ has a root in $\mathbb{K}$ and hence every algebraic extension of $\mathbb{K}$ is trivial. Thus $L \cong \mathbb{K}$ as $\mathbb{K}$ -algebras.
7. Since π is a surjection, $\ker(\pi) = \mathfrak{m}$. Since $\mathbb{K}[x_1, \dots, x_n] / \mathfrak{m} \cong \mathbb{K}$, we have $\psi(x_i) = a_i$ for some $a_i \in \mathbb{K}$. Then, $\langle x_1 - a_1, \dots, x_n - a_n \rangle \subset \mathfrak{m}$. Note that $\langle x_1 - a_1, \dots, x_n - a_n \rangle$ is maximal since $R / \langle x_1 - a_1, \dots, x_n - a_n \rangle$ is a field. Then, $\mathfrak{m}$ must be equal to $\langle x_1 - a_1, \dots, x_n - a_n \rangle$.
8. Thus when $\mathbb{K}$ is uncountable and algebraically closed the assignment $p \mapsto \mathfrak{m}_p$ gives a bijection between points of $\mathbb{K}^n$ and maximal ideals $\langle x_1 - a_1, \dots, x_n - a_n \rangle \subseteq \mathbb{K}[x_1, \dots, x_n]$.

We have now proven Hilbert's Nullstellensatz I under the additional assumption that $\mathbb{K}$ is uncountable, using virtually only field theory! Note this applies to $\mathbb{C}$ and $\mathbb{C}_p$ as these are both algebraically closed and uncountable, but does not apply to $\overline{\mathbb{Q}}$ and $\overline{\mathbb{F}_p}$ which while algebraically closed are countable.

In what follows we will *only* assume $\mathbb{K}$ is algebraically closed, and will use Hilbert's Nullstellensatz I to deduce the weak Nullstellensatz. After that we will use what has come to be known as *Rabinowitsch's trick* to upgrade the weak Nullstellensatz to the full statement of Hilbert's Nullstellensatz II.

Problem 5. Weak Nullstellensatz.

Let $\mathbb{K}$ be algebraically closed and let $I \subseteq \mathbb{K}[x_1, \dots, x_n]$ be an ideal.

1. Show that if I is proper, then I is contained in some maximal ideal $\mathfrak{m}$. (Hint: Use Zorn's Lemma.)
2. Assuming Hilbert's Nullstellensatz I, show that if I is proper then $\mathbb{V}(I) \neq \emptyset$. (Hint: $\mathbb{V}(\mathfrak{m}) \subset \mathbb{V}(I)$.)
3. Conclude that $\mathbb{V}(I) = \emptyset$ if and only if $I = \langle 1 \rangle$.

Solution.

1. Let Σ be the set of proper ideals of R containing I , ordered by inclusion. Then $\Sigma \neq \emptyset$ since $I \in \Sigma$. Let $\mathcal{C} \subseteq \Sigma$ be a chain and set $J^* = \bigcup_{J \in \mathcal{C}} J$. As a union of a chain of ideals, J^* is an ideal, and $I \subseteq J^*$. Moreover $1 \notin J$ for every $J \in \mathcal{C}$ (each is proper), so $1 \notin J^*$ and $J^* \neq R$. Hence $J^* \in \Sigma$ is an upper bound for $\mathcal{C}$. By Zorn's Lemma, Σ has a maximal element $\mathfrak{m}$.

Then it remains to check $\mathfrak{m}$ is a maximal ideal of R . It is proper since $\mathfrak{m} \in \Sigma$. If $\mathfrak{a}$ is an ideal with $\mathfrak{m} \subseteq \mathfrak{a} \subsetneq R$, then $\mathfrak{a} \in \Sigma$ and $\mathfrak{a} \supsetneq \mathfrak{m}$, so $\mathfrak{a} = \mathfrak{m}$ by maximality of $\mathfrak{m}$ in Σ . Thus no proper ideal strictly contains $\mathfrak{m}$, i.e. $\mathfrak{m}$ is maximal, and $I \subseteq \mathfrak{m}$.

2. Then $I \subseteq \mathfrak{m} \implies \mathbb{V}(\mathfrak{m}) \subseteq \mathbb{V}(I)$. $\mathbb{V}(\mathfrak{m}) = (a_1, \dots, a_n)$ where $\langle x_1 - a_1, \dots, x_n - a_n \rangle = \mathfrak{m}$. Hence $\mathbb{V}(I) \neq \emptyset$.
3. If $\mathbb{V}(I) = \emptyset$ then it is contained in no maximal ideal and hence we must have $I = \langle 1 \rangle$. And if $I = \langle 1 \rangle$ then $\mathbb{V}(I) = \emptyset$ (Problem 1.3).

Problem 6. Hilbert's Nullstellensatz II. (Rabinowitsch's Trick)

Let $\mathbb{K}$ be algebraically closed and $R := \mathbb{K}[x_1, \dots, x_n]$. Let $I = \langle f_1, \dots, f_t \rangle \subset R$ be an ideal. Choose any element $g \in \mathbb{I}(\mathbb{V}(I))$, we will show that some power of g lies in I . Note if $g = 0$ there is nothing to prove as $0 \in I$. Thus, we assume g is non-zero.

1. In the larger polynomial ring $R[z]$, define $J := \langle f_1, \dots, f_t, 1 - gz \rangle$. Show that $\mathbb{V}(J) = \emptyset$ without appealing to the weak Nullstellensatz. (Hint: If $(a_1, \dots, a_{n+1}) \in \mathbb{V}(J)$ then what do we know about $g(a_1, \dots, a_n)$ since $g \in \mathbb{I}(\mathbb{V}(I))$.)
2. Use Question 5 to conclude that $J = \langle 1 \rangle \subset R[z]$. Deduce there exist polynomials $h_1, \dots, h_t, h_{t+1} \in R[z]$ such that the following relation holds.

$$1 = h_1 f_1 + \dots + h_t f_t + h_{t+1}(1 - gz) \in R[z].$$

3. Consider the ring $R[1/g]$ consisting of expressions of the form p/g^k where $p \in R$ and $k \geq 0$, in other words, we enlarge R by formally allowing division by powers of g . Consider the map of $\mathbb{K}$ -algebras:

$$\begin{array}{ccc} R[z] & \xrightarrow{\psi} & R\left[\frac{1}{g}\right] \\ z & \longmapsto & \frac{1}{g} \end{array}$$

Explain why ψ is a well-defined $\mathbb{K}$ -algebra homomorphism. (Hint: What does it take to define a $\mathbb{K}$ -algebra map out of a polynomial ring $R[z]$?)

4. Prove that $\psi(1 - gz) = 0$ and deduce that $1 = \psi(h_1)f_1 + \dots + \psi(h_t)f_t$ in $R[1/g]$.
5. Explain why there exists $k \in \mathbb{N}$ such that $g^k \psi(h_i) \in R$ for every $i = 1, \dots, t$. Multiply the equation in part 4 by g^k to conclude that $g^k \in I$. (Hint: Each $\psi(h_i)$ is an element of $R[1/g]$, so it can be written as p_i/g^{k_i} for some $p_i \in R$ and $k_i \geq 0$.)
6. Conclude that $g \in \sqrt{I}$, and hence $\mathbb{I}(\mathbb{V}(I)) \subseteq \sqrt{I}$.
7. Combine this with Question 3 to finish the proof of Hilbert's Nullstellensatz II assuming Hilbert's Nullstellensatz I. (The latter we have proven over every uncountable algebraically closed field, in particular over $\mathbb{C}$.)

Solution.

1. Assume for contradiction $\mathbb{V}(J) \neq \emptyset$. Let $(a_1, \dots, a_{n+1}) \in \mathbb{V}(J)$ then

$$\begin{aligned} (1 - gz)(a_1, \dots, a_{n+1}) &= 1 - g(a_1, \dots, a_{n+1}) \cdot z(a_1, \dots, a_{n+1}) \\ \implies 1 &= g(a_1, \dots, a_{n+1}) \cdot z(a_1, \dots, a_{n+1}) \\ 1 &= 0 \cdot z(a_1, \dots, a_{n+1}) \end{aligned}$$

since $g(a_1, \dots, a_n) = 0$ since $I \subseteq J \implies \mathbb{V}(J) \subseteq \mathbb{V}(I)$, a contradiction. Hence $\mathbb{V}(J) = \emptyset$.

2. Then, by Question 5.3 we know that $J = \langle 1 \rangle$, i.e., by definition of an ideal there exist $h_1, \dots, h_{t+1} \in R[z]$ such that

$$h_1 f_1 + \dots + h_{t+1} (1 - gz) = 1$$

3. $R[z]$ is the polynomial ring in one variable over R , so for ψ to be well-defined we must send z to a unique element, which is $\frac{1}{g}$. To see that ψ is a homomorphism, consider

(a) for $p_i \in R$, we have

$$\psi \left(\sum_k p_k z^k \right) = \sum_k \frac{p_k}{g^k} = \sum_k \left(\psi(p_k z^k) \right)$$

(b) and

$$\psi \left(\prod_k p_k z^k \right) = \prod_k \frac{p_k}{g^k} = \prod_k \left(\psi(p_k z^k) \right)$$

4. Consider $\psi(1 - gz) = \psi(1) - \psi(g)\psi(z) = 1 - g \left(\frac{1}{g} \right) = 0$. Since all ring homomorphisms are unital we know that,

$$1 = \psi(h_1 f_1 + \dots + h_{t+1} (1 - gz)) = \psi(h_1) f_1 + \dots + \psi(h_{t+1}) (0) = \psi(h_1) f_1 + \dots + \psi(h_t) f_t$$

5. Since $h_i \in R[z]$ it might have monomials in z which under ψ get mapped to monomials in g^{-1} , $g \cdot g^{-1} = 1 \in R$ so we can cancel denominators by multiplying $\psi(h_i)$ by g^k where g^{-k} is the lowest degree of $g \in \psi(h_i)$. Hence we can express g^k as a finite sum of the generators of I ,

$$g^k = g^k \psi(h_1) f_1 + \dots + g^k \psi(h_t) f_t$$

6. Thus $g^k \in I$ and $g \in \sqrt{I}$. Since g was an arbitrary element in $\mathbb{I}(\mathbb{V}(I))$ this implies $\mathbb{I}(\mathbb{V}(I)) \subseteq \sqrt{I}$.
7. That gives us the last containment and hence we have proved Hilbert's Nullstellensatz II.

You have now proved Hilbert's Nullstellensatz for every *uncountable* algebraically closed field, including $\mathbb{C}$. The only remaining gap in the proof over an arbitrary algebraically closed field is the point case: one still needs to show that every maximal ideal is of the form $\mathfrak{m}_p$ without assuming uncountability. We will return to that later, after developing more tools.

For the remainder of the worksheet, assume Hilbert's Nullstellensatz I and II over all algebraically closed fields and use them freely. The point now is to unpack what the theorem says about coordinate rings, closed subsets, irreducibility, and functoriality. In particular, we would like a version of the Nullstellensatz that describes algebraic subsets not of just $\mathbb{K}^n$, but of any algebraic set $V \subset \mathbb{K}^n$. Towards this we will define and study relative versions of the $\mathbb{I}$ and $\mathbb{V}$ operators. In particular, for an algebraic subset $W \subset V$ and an ideal $J \subset \mathbb{K}[V]$ define

$$\mathbb{I}_V(W) := \{ \phi \in \mathbb{K}[V] \mid \phi(p) = 0 \text{ for all } p \in W \}$$

and

$$\mathbb{V}_V(J) := \{ p \in V \mid \phi(p) = 0 \text{ for all } \phi \in J \}.$$

Using these we how to prove the following relative version of the Nullstellensatz connecting algebraic subsets of V and radical ideals in $\mathbb{K}[V]$.

Theorem 0.10 (Hilbert's Nullstellensatz III). *Let $\mathbb{K}$ be an algebraically closed field. If $V \subset \mathbb{K}^n$ is an algebraic subset then the maps $\mathbb{I}_V$ and $\mathbb{V}_V$ induce inverse, inclusion-reversing bijections:*

$$\begin{array}{ccc} \left\{ \begin{array}{c} \text{algebraic subsets} \\ \text{in } V \end{array} \right\} & \begin{array}{c} \xrightarrow{\mathbb{I}_V} \\ \xleftarrow{\mathbb{V}_V} \end{array} & \left\{ \begin{array}{c} \text{radical ideals} \\ \text{in } \mathbb{K}[V] \end{array} \right\} \end{array}$$

Note that since the coordinate ring of $\mathbb{K}^n$ is canonically isomorphic to $\mathbb{K}[x_1, \dots, x_n]$ we could view Hilbert's Nullstellensatz II as a special case of this third version. However, the real difficulty in proving the third version boils down to applying the second. As you will see many of the following exercises boil down to the same ideas and arguments we used for $\mathbb{I}$ and $\mathbb{V}$, at times almost verbatim.

Problem 7. Properties of $\mathbb{I}_V$.

Assume $\mathbb{K}$ is algebraically closed, let $V \subseteq \mathbb{K}^n$ be an algebraic set with coordinate ring $\mathbb{K}[V]$. Let $W \subseteq V$ be an algebraic subset and $\rho_W : \mathbb{K}[V] \rightarrow \mathbb{K}[W]$ be the restriction homomorphism taking a function $\phi : V \rightarrow \mathbb{K}$ and restricting it to W to get a function $\phi|_W : W \rightarrow \mathbb{K}$. (If we wanted to feel fancy we could say $\phi|_W$ is the composition of $\phi : V \rightarrow \mathbb{K}$ with the inclusion $\iota : W \rightarrow V$, but in this setting that is needlessly complex.)

1. Show that $\mathbb{I}_V(W)$ is an ideal of $\mathbb{K}[V]$, and if $Z \subset W \subset V$ then $\mathbb{I}_V(W) \subset \mathbb{I}_V(Z)$.
2. Since V is an algebraic subset of itself, as a reality check what is $\mathbb{I}_V(V)$?
3. Prove that ρ_W is a surjective $\mathbb{K}$ -algebra homomorphism with kernel $\mathbb{I}_V(W)$.
4. Prove that ρ_W induces an $\mathbb{K}$ -algebra isomorphism $\mathbb{K}[V]/\mathbb{I}_V(W) \cong \mathbb{K}[W]$.
5. Show that if $W \subset V$ is an algebraic subset then the following diagram commutes:

$$\begin{array}{ccc} \mathbb{K}[x_1, \dots, x_n] & \xrightarrow{\rho} & \mathbb{K}[V] \\ & \searrow \rho' & \downarrow \rho_W \\ & & \mathbb{K}[W] \end{array}$$

where $\rho' : \mathbb{K}[x_1, \dots, x_n] \rightarrow \mathbb{K}[W]$ is the restriction of functions map defining $\mathbb{K}[W]$. Deduce that $\mathbb{I}_V(W) = \mathbb{I}(W)/\mathbb{I}(V)$.

6. Conclude that if $W \subset V$ is an algebraic subset then $\mathbb{I}_V(W)$ is a radical ideal of $\mathbb{K}[V]$.

Solution.

1. $\mathbb{K}[V] = \mathbb{K}[x_1, \dots, x_n]/\mathbb{I}(V)$. By definition,

$$\mathbb{I}_V(W) = \{\phi \in \mathbb{K}[V] \mid \phi(p) = 0, \forall p \in W\}$$

We check that $\mathbb{I}_V(W)$ is an ideal of $\mathbb{K}[V]$,

(a) **Zero:** $0(p) = 0$ for all p so $0 \in \mathbb{I}_V(W)$.

(b) **Additive Closure:** Let $\phi, \psi \in \mathbb{I}_V(W)$. For any $w \in W$,

$$(\phi + \psi)(w) = \phi(w) + \psi(w) = 0 + 0 = 0.$$

Hence $\phi + \psi \in \mathbb{I}_V(W)$.

(c) **Inverses:** If $\phi \in \mathbb{I}_V(W)$, then for any $w \in W$,

$$(-\phi)(w) = -\phi(w) = 0,$$

so $-\phi \in \mathbb{I}_V(W)$.

(d) **Closure under Multiplication from $\mathbb{K}[V]$:** Let $f \in \mathbb{K}[V]$. Then for every $w \in W$,

$$(f\phi)(w) = f(w)\phi(w) = f(w) \cdot 0 = 0.$$

Thus $f\phi \in \mathbb{I}_V(W)$.

Let $\phi \in \mathbb{I}_V(W)$. Since ϕ vanishes on all of W , it certainly vanishes on $Z \subseteq W$. Hence

$$\phi \in \mathbb{I}_V(W) \subseteq \mathbb{I}_V(Z).$$

2. Since V is an algebraic subset of itself,

$$\mathbb{I}_V(V) = \{\phi \in \mathbb{K}[V] \mid \phi(v) = 0, \forall v \in V\}.$$

The only regular function on V that vanishes at every point of V is the zero function. Hence

$$\mathbb{I}_V(V) = \{0\}.$$

3. To prove surjectivity, let $f \in \mathbb{K}[W]$. Since f is a function on W , there exists a polynomial $F \in \mathbb{K}[x_1, \dots, x_n]$ such that $f(w) = F(w)$ for all $w \in W$. Let $\phi \in \mathbb{K}[V]$ be the function on V represented by the same polynomial F . Then

$$\rho_W(\phi) = \phi|_W = f.$$

Hence ρ_W is surjective. The kernel of this homomorphism is

$$\ker(\rho_W) = \{\phi \in \mathbb{K}[V] \mid \rho_W(\phi) = 0\} = \{\phi \in \mathbb{K}[V] \mid \phi(w) = 0, \forall w \in W\} = \mathbb{I}_V(W)$$

4. By the First Isomorphism Theorem we have,

$$\mathbb{K}[V]/\mathbb{I}_V(W) \cong \mathbb{K}[W]$$

5. Let $f \in \mathbb{K}[x_1, \dots, x_n]$. Then

$$(\rho_W \circ \rho)(f) = \rho_W(f|_V) = (f|_V)|_W = f|_W = \rho'(f)$$

Hence the diagram commutes.

Note that

$$\rho^{-1}(\ker \rho_W) = \ker \rho' = \mathbb{I}(W)$$

And $\ker(\rho) = \mathbb{I}(V)$, so $\mathbb{I}_V(W) = \frac{\mathbb{I}(W)}{\mathbb{I}(V)}$.

6. $\mathbb{K}[V] = \mathbb{K}[x_1, \dots, x_n]/\mathbb{I}(V)$, and $\mathbb{I}_V(W) = \mathbb{I}(W)/\mathbb{I}(V)$. The a quotient of a ring by a radical ideal is reduced, so $\mathbb{I}(W)/\mathbb{I}(V)$ is radical because $\mathbb{I}(W)$ is radical.

Worksheet 4.1: Modules Pt. I

Throughout this course, “ring” means *commutative* ring with unity. In this worksheet $\mathbb{K}$ will denote a field. A R -module is an abelian group $(M, +)$ together with a scalar multiplication $R \times M \rightarrow M, (r, m) \mapsto rm$, satisfying the following:

- (i) Associativity: $(rs)m = r(sm)$ for all $r, s \in R$ and $m \in M$,
- (ii) Distribution over R : $(r + s)m = rm + sm$ for all $r, s \in R$ and $m \in M$,
- (iii) Distribution over M : $r(m + n) = rm + rn$ for all $r \in R$ and $m, n \in M$, and
- (iv) Identity: $1_R m = m$ for all $m \in M$.

Since M is an abelian group, it has a distinguished additive identity, which we normally denote by 0 or by 0_M if we need to distinguish it from 0_R . When the base ring is clear from context, we might simply say that M is a module, or we might say that M is a module over R . As we will see, modules are the analogue of vector spaces in which the scalars come from a ring instead of a field. They include ideals, quotient rings, abelian groups, vector spaces, and many other naturally occurring algebraic objects.

If M is an R -module, a subset $N \subseteq M$ is an R -submodule of M if and only if $rm + sn \in N$ for all $r, s \in R$ and $m, n \in N$. That is, a submodule is a subgroup of M such that the scalar multiplication of R on M restricts to a scalar multiplication on N .

Problem 1. Basic Identities. Let R be a ring and M an R -module. Prove the following identities.

1. $r0_M = 0_M$ for all $r \in R$.
2. $0_R m = 0_M$ for all $m \in M$.
3. $(-r)m = r(-m)$ for all $r \in R$ and all $m \in M$.

Solution.

1. Let $m \in M$.

$$r0_M = r(m - m) = rm - rm = 0_M$$

2. Let $r \in R$.

$$0_R m = (r - r)m = rm - rm = 0_M$$

- 3.

$$(-r)m = (0_R - r)m = 0_R m - rm$$

From (ii) we have that $0_R m = 0_M$,

$$\begin{aligned} (-r)m &= 0_R m - rm = 0_M - rm \\ &= 0_M r - rm = r(0_M - m) \\ &= r(-m) \end{aligned}$$

Problem 3. Submodules Generated by Elements.

Let M be an R -module. Given elements $m_1, \dots, m_t \in M$, the submodule of M generated by $m_1, \dots, m_t$ is the set

$$Rm_1 + Rm_2 + \cdots + Rm_t := \{r_1m_1 + \cdots + r_tm_t \mid r_1, \dots, r_t \in R\} \subseteq M.$$

We say that $m_1, \dots, m_t$ are *generators* for M if $Rm_1 + Rm_2 + \cdots + Rm_t = M$. A submodule $N \subseteq M$ is a *cyclic submodule* if $N = Rm$ for some $m \in M$.

1. Prove that $Rm_1 + Rm_2 + \cdots + Rm_t$ is an R -submodule of M .
2. Prove that $Rm_1 + Rm_2 + \cdots + Rm_t$ is the smallest R -submodule of M containing $m_1, \dots, m_t$.
3. Prove that a ring R is itself a cyclic R -module by finding a generator.
4. In the $\mathbb{Z}$ -module $\mathbb{Z}/12\mathbb{Z}$, compute the submodules generated by $\bar{8}$ and $\bar{6}$.
5. Let $R = \mathbb{K}[x]$ and $M = R/\langle x^3 \rangle$. Describe the submodules generated by $\bar{1}$, $\bar{x}$, and $\bar{x}^2$.

Solution.

Let $N = Rm_1 + Rm_2 + \cdots + Rm_t$.

1. Let $r, s \in R$ and $n, m \in N$. Then, by definition

$$m = r_1m_1 + \cdots + r_tm_t$$

and

$$n = s_1m_1 + \cdots + s_tm_t$$

with $r_1, \dots, r_t, s_1, \dots, s_t \in R$. Then by distributivity and associativity of R , and the abelian nature of M we have,

$$\begin{aligned} pm + qn &= p(r_1m_1 + \cdots + r_tm_t) + q(s_1m_1 + \cdots + s_tm_t) \\ &= ((pr_1)m_1 + \cdots + (pr_t)m_t) + ((qs_1)m_1 + \cdots + (qs_t)m_t) \\ &= ((pr_1 + qs_1)m_1 + \cdots + (pr_t + qs_t)m_t) \end{aligned}$$

but since R is closed under addition and multiplication $pr_1 + qs_1, \dots, pr_t + qs_t \in R$. Hence, $pm + qn \in N$ and N is a submodule.

2. Let L be the smallest R -submodule of M containing $m_1, \dots, m_t$. Then by minimality of L we have $L \subseteq N$. To show the reverse containment consider $m \in N$. By definition,

$$m = r_1m_1 + \cdots + r_tm_t$$

with $r_1, \dots, r_t \in R$. But since $m_1, \dots, m_t \in L$, all R -linear combinations of them is in L (by definition of a submodule), hence $m \in L$. Since m was an arbitrary element in N this implies that $N = L$, hence N is the smallest R -submodule of M containing $m_1, \dots, m_t$.

3. The ring R is a cyclic R -module with the generator 1_R .

By definition,

$$R \cdot 1_R = \{r \cdot 1_R \mid r \in R\}$$

But $r \cdot 1_R = r$ for all $r \in R$, i.e., $R \cdot 1_R = R$.

4. Consider $\mathbb{Z}/12\mathbb{Z} \cdot \bar{8}$,

$$\mathbb{Z}/12\mathbb{Z} \cdot \bar{8} = \{r \cdot \bar{8} \mid r \in \mathbb{Z}/12\mathbb{Z}\} = \{\bar{0}, \bar{8}, \bar{4}\}$$

Consider $\mathbb{Z}/12\mathbb{Z} \cdot \bar{6}$,

$$\mathbb{Z}/12\mathbb{Z} \cdot \bar{6} = \{r \cdot \bar{6} \mid r \in \mathbb{Z}/12\mathbb{Z}\} = \{\bar{0}, \bar{6}\}$$

5. $R \cdot \bar{1} = \{r \cdot \bar{1} = \bar{r} \mid r \in R\} = M$

$$R \cdot \bar{x} = \{r \cdot \bar{x} = \overline{rx} \mid r \in R\} = \{\overline{r_1x + r_2x^2} \mid r_1, r_2 \in R\} = \mathbb{K} \cdot \bar{x}$$

$$R \cdot \bar{x}^2 = \{r \cdot \bar{x}^2 = \overline{rx^2} \mid r \in R\} = \mathbb{K} \cdot \bar{x}^2$$

A *homomorphism* of R -modules is a group homomorphism $\phi : M \rightarrow N$ such that $\phi(rm) = r\phi(m)$ for all $r \in R$ and $m \in M$. Two R -modules M and N are *isomorphic* if and only if there exist R -module homomorphisms $\phi : M \rightarrow N$ and $\psi : N \rightarrow M$ such that $\phi \circ \psi = \text{Id}_N$ and $\psi \circ \phi = \text{Id}_M$. Equivalently, there is a bijective homomorphism of R -modules $\phi : M \rightarrow N$. We call such maps *isomorphisms* of R -modules. If $\phi : M \rightarrow N$ is a homomorphism of R -modules, there are two natural submodules associated to it: its *kernel* and its *image*:

$$\ker(\phi) := \{m \in M \mid \phi(m) = 0_N\} \subseteq M \quad \text{and} \quad \text{img}(\phi) := \{\phi(m) \mid m \in M\} \subseteq N.$$

One should readily check that both the kernel and image of a module homomorphism are submodules. As with linear maps, group homomorphisms, and ring homomorphisms, the kernel measures the failure of injectivity and the image measures the failure of surjectivity. Given a submodule $N \subseteq M$, the quotient abelian group M/N inherits an R -module structure via

$$r(m + N) := rm + N.$$

Well-definedness uses that N is a submodule, not merely a subgroup. The projection $\pi : M \rightarrow M/N$ is a homomorphism with $\ker(\pi) = N$. The quotient M/N is the module obtained by forcing $N = 0$. Precisely: if $f : M \rightarrow P$ is a homomorphism with $N \subseteq \ker(f)$, there is a unique $\bar{f} : M/N \rightarrow P$ with $f = \bar{f} \circ \pi$:

$$\begin{array}{ccc} M & \xrightarrow{f} & P \\ \pi \downarrow & \nearrow \bar{f} & \\ M/N & & \end{array}$$

Phrased differently, there is a natural isomorphism

$$\text{Hom}_R(M/N, P) \cong \{f \in \text{Hom}_R(M, P) \mid N \subseteq \ker(f)\};$$

That is, homomorphisms out of M/N are exactly homomorphisms out of M that kill N . This is usually summarized as the First Isomorphism Theorem.

Theorem 0.11 (First Isomorphism Theorem). *Every homomorphism $\phi : M \rightarrow N$ of R -modules induces an isomorphism*

$$M / \ker(\phi) \xrightarrow{\sim} \text{img}(\phi), \quad m + \ker(\phi) \mapsto \phi(m).$$

In other words, quotienting by the kernel accounts for all the redundancy: every surjection is a quotient map.

Theorem 0.12 (Third Isomorphism Theorem). *Let $N \subseteq M$ be a submodule and $\pi: M \twoheadrightarrow M/N$ the projection. There is an inclusion-preserving bijection*

$$\{\text{submodules of } M \text{ containing } N\} \xleftrightarrow{1:1} \{\text{submodules of } M/N\}$$

given by $L \mapsto L/N = \pi(L)$ with inverse $\bar{L} \mapsto \pi^{-1}(\bar{L})$. Moreover, for $N \subseteq L \subseteq M$:

$$M/L \cong (M/N)/(L/N).$$

The correspondence says that the submodule lattice of M/N is the interval $[N, M]$ in the submodule lattice of M . The supplementary isomorphism $(M/N)/(L/N) \cong M/L$ is sometimes called the *Third Isomorphism Theorem*: quotienting in two stages is the same as quotienting all at once.

Problem 4. Basic Examples of Module Homomorphisms.

1. If $\mathbb{K}$ is a field, explain why a homomorphism of $\mathbb{K}$ -modules is the same as a $\mathbb{K}$ -linear map.
2. Explain why a homomorphism of $\mathbb{Z}$ -modules is the same as a homomorphism of abelian groups.

Solution.

1. If $\mathbb{K}$ is a field, we know that a $\mathbb{K}$ -module is a $\mathbb{K}$ -vector space. Moreover, by definition, a homomorphism $\varphi : N \rightarrow M$ where N, M are $\mathbb{K}$ -modules ($\mathbb{K}$ -vector spaces) has the following property: for any $r, s \in \mathbb{K}$ and $a, b \in N$ we have,

$$\varphi(ra + sb) = \varphi(ra) + \varphi(sb) = r\varphi(a) + s\varphi(b)$$

which is the condition defining a $\mathbb{K}$ -linear map.

2. We proved in Problem 2 that $\mathbb{Z}$ -modules are the same as abelian groups, then any $\mathbb{Z}$ -module homomorphism must preserve the group operation and hence is the same as an abelian group homomorphism.

Problem 5. Cyclic Modules & Annihilators. Let M be an R -module. The *annihilator* of an element $m \in M$ is $\text{Ann}_R(m) := \{r \in R \mid rm = 0\}$.

1. Prove that $\text{Ann}_R(m)$ is an ideal of R .
2. What is $\text{Ann}_R(0_M)$?
3. Consider the multiplication by m function:

$$\phi_m : R \rightarrow M \quad \text{given by} \quad r \mapsto rm.$$

Prove that ϕ_m is a homomorphism of R -modules.

4. Show that $\ker(\phi_m) = \text{Ann}_R(m)$ and $\text{img}(\phi_m) = Rm$.
5. Prove that a submodule $N \subseteq M$ is cyclic if and only if $N \cong R/I$ for some ideal $I \subseteq R$.

Solution.

1. To show that $\text{Ann}_R(m)$ is an ideal of R we need to check that it is a subgroup under addition, and is closed under multiplication from elements in R .

(a) **Identity:** Note that $0 \in \text{Ann}_R(m)$ since $0 \cdot m = 0$.

(b) **Closure under Addition and Inverses:** If $a, b \in \text{Ann}_R(m)$, then

$$(a - b)m = am - bm = 0 - 0 = 0$$

so $a - b \in \text{Ann}_R(m)$.

Hence $\text{Ann}_R(m)$ is a subgroup of $(R, +)$.

(c) Let $r \in R$ and $a \in \text{Ann}_R(m)$. Then

$$(ra)m = r(am) = r \cdot 0 = 0$$

so $ra \in \text{Ann}_R(m)$.

Hence $\text{Ann}_R(m)$ is an ideal of R .

2. $\text{Ann}_R(0_M) = \{r \in R \mid r0_M = 0\} = R$. That is, $\text{Ann}_R(0_M)$ is the whole ring.
3. Let $a, b \in R$. Consider

$$\phi_m(a + b) = (a + b)m = am + bm = \phi_m(a) + \phi_m(b)$$

Hence ϕ_m is a group homomorphism. And, for any $r, a \in R$ we have,

$$\phi_m(ra) = (ra)m = r(am) = r\phi_m(a)$$

Hence, ϕ_m is a homomorphism of R -modules.

4. $\ker(\phi_m) = \{r \in R \mid rm = 0\}$ which is precisely the definition of $\text{Ann}_R(m)$.

And $\text{im}(\phi_m) = \{rm \mid r \in R\} = Rm$.

5. ($\implies$)

Assume submodule $N \subseteq M$ is cyclic. Then $N = R \cdot m$ for some $m \in M$. From part (4) we know that $\ker(\phi_m) = \text{Ann}_R(m)$ and $\text{im}(\phi_m) = Rm$, that is by the first isomorphism theorem,

$$N = Rm = \text{im}(\phi_m) \cong R / \ker(\phi_m) = R / \text{Ann}_R(m)$$

where we know that $\text{Ann}_R(m)$ is an ideal in R (part 1)

($\impliedby$)

Assume $N \cong R/I$ for some ideal $I \subseteq R$. Then there exists some isomorphism of R -modules $\varphi : R/I \rightarrow N$. Note that every element in R/I can be written as, $r(1 + I)$ for some $r \in R$ and therefore every element of N is of the form

$$\varphi(r(1 + I)) = r\varphi(1 + I)$$

Let $\varphi(1 + I) = n \in N$. That is every element is of the form rn for some $r \in R$, then $N \subseteq Rn$. Moreover since $n \in N$ and N is closed under scalar multiplication we have $Rn \subset N$. Hence, N is cyclic.

Let $\{M_i\}_{i \in I}$ be a family of R -modules indexed by a set I . The *direct product* is

$$\prod_{i \in I} M_i = \{(m_i)_{i \in I} \mid m_i \in M_i\},$$

with componentwise addition and scalar multiplication. The *direct sum* is the submodule

$$\bigoplus_{i \in I} M_i = \left\{ (m_i)_{i \in I} \in \prod_{i \in I} M_i \mid m_i = 0 \text{ for all but finitely many } i \right\}.$$

For each $j \in I$ there is a projection map π_j and an inclusion map ι_j :

$$\pi_j : \prod_{i \in I} M_i \rightarrow M_j, \quad ((m_i))_{i \in I} \mapsto m_j,$$

and

$$\iota_j : M_j \rightarrow \bigoplus_{i \in I} M_i, \quad m \mapsto (0, \dots, 0, m, 0, \dots).$$

The product is characterized by maps to the factors, while the direct sum is characterized by maps from the summands. These are the universal properties that make products and sums so useful.

Problem 6. Direct Sums and Products of Modules.

Let $\{M_i\}_{i \in I}$ be a family of R -modules indexed by a set I .

1. Prove that both $\prod_{i \in I} M_i$ and $\bigoplus_{i \in I} M_i$ are R -modules.
2. Explain why, if I is finite, $\prod_{i \in I} M_i = \bigoplus_{i \in I} M_i$.
3. Consider the $\mathbb{Z}$ -modules $M_n = \mathbb{Z}/n\mathbb{Z}$ for $n \geq 1$. Describe explicitly an element of $\prod_{n=1}^{\infty} M_n$ that is not an element of $\bigoplus_{n=1}^{\infty} M_n$.
4. Let $M_i = R$ for all $i \in \mathbb{N}$. Show that the submodule of $\prod_{i \in \mathbb{N}} R$ generated by the standard basis vectors $e_1, e_2, e_3, \dots$ is $\bigoplus_{i \in \mathbb{N}} R$, not the whole product, provided $R \neq 0$.
5. For two modules M and N , write down the inclusion maps ι_M, ι_N and projection maps π_M, π_N for $M \oplus N$. Verify the following identities

$$\pi_M \circ \iota_M = \text{Id}_M, \quad \pi_N \circ \iota_N = \text{Id}_N,$$

$$\pi_M \circ \iota_N = 0, \quad \pi_N \circ \iota_M = 0, \text{ and } \iota_M \circ \pi_M + \iota_N \circ \pi_N = \text{Id}_{M \oplus N}.$$

Solution.

1. Let $f, g, h \in \prod_{i \in I} M_i$ and $r, s \in R$. We can see that $\prod_{i \in I} M_i$ is an abelian group under componentwise addition because each M_i is an abelian group under addition by definition. Now we check the scalar multiplication:

(a) **Closure:** Since $r \cdot f(i) \in M_i$ for all i , $r \cdot f \in \prod_{i \in I} M_i$.

(b) **Associativity:** Since $(rs)f(i) = r(sf(i))$ for all i , $(rs)f = r(sf)$.

(c) **Distributivity over Module Addition:** Since for all i we have,

$$r(f(i) + g(i)) = r(f(i)) + r(g(i))$$

we get

$$r(f + g) = rf + rg$$

(d) **Distributivity over Ring Addition:** Since for all i we have,

$$(r + s)f(i) = r(f(i)) + s(f(i))$$

we get

$$(r + s)f = rf + sf$$

(e) **Unitality:** Since for all i we have $1_R \cdot f(i) = f(i)$ we get

$$1_R \cdot f = f$$

Since $\bigoplus_{i \in I} M_i \subseteq \prod_{i \in I} M_i$ it suffices to show that $\bigoplus_{i \in I} M_i$ is a submodule. Let $f, g \in \bigoplus_{i \in I} M_i$ and $r \in R$.

- (a) **Non-empty:** $0 \in \bigoplus_{i \in I} M_i$
 - (b) **Closure under Addition:** Since both f and g have only finitely many non-zero entries, their sum also has finitely many non-zero entries and hence $f + g \in \bigoplus_{i \in I} M_i$.
 - (c) **Closure under Scalar Multiplication:** Since both f has only finitely many non-zero entries, and $r \cdot 0_M = 0$ the product $r \cdot f$ also has finitely many non-zero entries and hence $r \cdot f \in \bigoplus_{i \in I} M_i$.
2. We know that $\bigoplus_{i \in I} M_i \subseteq \prod_{i \in I} M_i$, so we only need to show the opposite containment. But if I is finite, every element $f \in \prod_{i \in I} M_i$ only has finitely many non-zero entries and hence by definition $f \in \bigoplus_{i \in I} M_i$. Thus, $\bigoplus_{i \in I} M_i = \prod_{i \in I} M_i$.
 3. Note that $1 \in \mathbb{Z}/n\mathbb{Z}$ for all $n \geq 1$. Then the element $(1, 1, 1, \dots) \in \prod_{n=1}^{\infty} M_n$ but it is not in $\bigoplus_{n=1}^{\infty} M_n$ because it has infinitely many non-zero entries.
 4. An element of the submodule generated by $\{e_i\}$ is a finite R -linear combination, and hence has only finitely many non-zero entries and hence is in $\bigoplus_{i \in \mathbb{N}} R$. Since we are not allowed infinite sums elements like $(1, 1, 1, \dots) = \sum_{i=1}^{\infty} e_i$ are not in the generated submodule.
 5. We have,

$$\begin{aligned}\pi_M : M \bigoplus N &\rightarrow M, & \pi_M(m, n) &= m \\ \pi_N : M \bigoplus N &\rightarrow N, & \pi_N(m, n) &= n\end{aligned}$$

And,

$$\begin{aligned}\iota_M : M &\rightarrow M \bigoplus N, & \iota_M(m) &= (m, 0) \\ \iota_N : N &\rightarrow M \bigoplus N, & \iota_N(n) &= (0, n)\end{aligned}$$

Then,

$$\begin{aligned}\pi_M \circ \iota_M(m) &= \pi_M(0, m) = m = \text{id}_M(m), \\ \pi_N \circ \iota_N(n) &= \pi_N(0, n) = n = \text{id}_N(n), \\ \pi_M \circ \iota_N(n) &= \pi_M(0, n) = 0, \\ \pi_N \circ \iota_M(m) &= \pi_N(m, 0) = 0,\end{aligned}$$

$$(\iota_M \circ \pi_M + \iota_N \circ \pi_N)((m, n)) = \iota_M(m) + \iota_N(n) = (m, 0) + (0, n) = (m, n) = \text{Id}_{M \oplus N}(m, n).$$

Problem 7. Universal Properties of Products and Sums.

Let $\{M_i\}_{i \in I}$ be a family of R -modules indexed by I .

1. Let L be an R -module and suppose that for every $i \in I$ we are given a homomorphism $\phi_i : L \rightarrow M_i$. Prove that there exists a unique homomorphism $\phi : L \rightarrow \prod_{i \in I} M_i$ such that the following diagram commutes for all $i \in I$:

$$\begin{array}{ccc} L & \xrightarrow{\phi} & \prod_{i \in I} M_i \\ & \searrow \phi_i & \downarrow \pi_i \\ & & M_i \end{array} \quad .$$

2. Let L be an R -module and suppose that for every $i \in I$ we are given a homomorphism $\psi_i : M_i \rightarrow L$. Prove that there exists a unique homomorphism $\psi : \bigoplus_{i \in I} M_i \rightarrow L$ such that the following diagram commutes for all $i \in I$:

$$\begin{array}{ccc} \bigoplus_{i \in I} M_i & \xrightarrow{\psi} & L \\ \uparrow \iota_i & \nearrow \psi_i & \\ M_i & & \end{array} \quad .$$

3. Deduce natural isomorphisms of R -modules

$$\operatorname{Hom}_R \left(L, \prod_{i \in I} M_i \right) \cong \prod_{i \in I} \operatorname{Hom}_R(L, M_i)$$

and

$$\operatorname{Hom}_R \left(\bigoplus_{i \in I} M_i, L \right) \cong \prod_{i \in I} \operatorname{Hom}_R(M_i, L).$$

4. Explain why a family of maps $\psi_i : M_i \rightarrow L$ does not generally define a map $\prod_{i \in I} M_i \rightarrow L$. Use the case $M_i = L = \mathbb{Z}$ and $\psi_i = \operatorname{Id}_{\mathbb{Z}}$ to illustrate the problem.
5. When I is finite, explain why the two universal properties describe the same module. This is why finite direct sums are also finite direct products.

Solution.

1. **Existence:** We define $\phi : L \rightarrow \prod_{i \in I} M_i$ as follows

$$\phi(l) = (\phi_1(l), \dots, \phi_i(l), \dots)$$

This is a well-defined homomorphism because each ϕ_i is a homomorphism.

Uniqueness: Let $\psi : L \rightarrow \prod_{i \in I} M_i$ be another such homomorphism. Then we have $\pi_i \circ \psi(l) = \phi_i(l) = \pi_i \circ \phi(l)$ for all $i \in I$ and $l \in L$. Since π_i is well-defined, this means $\psi = \phi$.

2. **Existence:** We define $\psi : \bigoplus_{i \in I} M_i \rightarrow L$ as follows

$$\psi(f) = \sum_{i \in I} \psi_i(f(i))$$

Note that this sum is well-defined since for all $f \in \bigoplus_{i \in I} M_i$ only finitely many $f(i) \neq 0$. This is a homomorphism since,

$$\begin{aligned} \psi(f + g) &= \sum_{i \in I} \psi_i(f(i) + g(i)) \\ &= \sum_{i \in I} \psi_i(f(i)) + \psi_i(g(i)) \\ &= \sum_{i \in I} \psi_i(f(i)) + \sum_{i \in I} \psi_i(g(i)) \\ &= \psi(f) + \psi(g) \end{aligned}$$

And,

$$\begin{aligned} \psi(r \cdot f) &= \sum_{i \in I} \psi_i(r \cdot f(i)) \\ &= \sum_{i \in I} r \cdot \psi_i(f(i)) \\ &= r \sum_{i \in I} \psi_i(f(i)) \\ &= r \cdot \psi(f) \end{aligned}$$

Uniqueness: Let $\phi : \bigoplus_{i \in I} M_i \rightarrow L$ be another homomorphism such that $\phi \circ \iota_i = \psi_i$ for all $i \in I$. For any $f \in \bigoplus_{i \in I} M_i$, since f has only finitely many non-zero entries we can write $f = \sum_{i \in I} \iota_i(f(i))$, and so

$$\phi(f) = \sum_{i \in I} \phi(\iota_i(f(i))) = \sum_{i \in I} \psi_i(f(i)) = \psi(f)$$

Since f was arbitrary, $\phi = \psi$.

3. From (1) and (2) we can see that there exists a bijection between homomorphisms $\phi \leftrightarrow (\phi_i)$ and $\psi \leftrightarrow (\psi(i))$. Then we define homomorphisms:

$$\Phi : \text{Hom}_R \left(L, \prod_{i \in I} M_i \right) \rightarrow \prod_{i \in I} \text{Hom}_R(L, M_i), \quad \Phi(\phi) = (\pi_i \circ \phi)_{i \in I}$$

$$\Psi : \text{Hom}_R \left(\bigoplus_{i \in I} M_i, L \right) \rightarrow \prod_{i \in I} \text{Hom}_R(M_i, L), \quad \Psi(\phi) = (\psi \circ \iota_i)_{i \in I}$$

Since the universal properties say that every family (ϕ_i) and (ψ_i) come from unique ϕ and ψ respectively, the homomorphisms Φ and Ψ are bijective.

4. Let $M_i = L = \mathbb{Z}$ and $\psi_i = \text{id}_{\mathbb{Z}}$. Then $f = (1, 1, 1, \dots)$ is an element in $\prod_{i \in I} M_i$. Note that the number of non-zero entries in f can be infinite. Then, the map

$$\psi(f) = \sum_{i \in I} \text{id}_{\mathbb{Z}}(f(i)) = \sum_{i \in I} 1$$

is not defined because it describes an infinite sum. Hence, a family of maps $\psi_i : M_i \rightarrow L$ does not generally define a map $\prod_{i \in I} M_i \rightarrow L$.

5. This is clear from Problem 6 (2).

Over a field, every module (i.e. vector space) has a basis, and a linear map is determined by its values on a basis. Over a general ring, modules need not have bases at all—but those that do are of fundamental importance. The goal of this section is to set up the theory of free modules and to understand precisely how they differ from vector spaces.

Let M be an R -module. A subset $B \subseteq M$ *generates* (or *spans*) M if every element of M can be written as a finite R -linear combination of elements of B , i.e. if

$$M = \left\{ \sum_{i=1}^n r_i m_i \mid n \geq 0, r_i \in R, m_i \in B \right\}.$$

We say B is *linearly independent* if, whenever $m_1, \dots, m_n \in B$ are distinct and $\sum_{i=1}^n r_i m_i = 0$, we have $r_i = 0$ for all i . A *basis* for M is a subset that is both generating and linearly independent, or equivalently, a subset $\{b_s\}_{s \in S}$ such that every element of M can be written *uniquely* as a finite R -linear combination $\sum_s r_s b_s$.

Over a field, these notions interact exactly as in undergraduate linear algebra: every spanning set contains a basis, every linearly independent set extends to one, and all bases have the same cardinality. Over a general ring, the first two statements can fail; in particular, many modules do not admit bases at all. A module that does admit a basis is called *free*. Given a set S , the *free R -module on S* is

$$R^{(S)} := \bigoplus_{s \in S} R.$$

Concretely, $R^{(S)}$ consists of all functions $S \rightarrow R$ with finite support. It has a distinguished element e_s for each $s \in S$, where e_s has a 1 in the s -coordinate and 0 elsewhere. Every element of $R^{(S)}$ can then be written uniquely as a finite sum

$$\sum_{s \in S} r_s e_s,$$

so $\{e_s \mid s \in S\}$ is a basis, called the *standard basis*. When $S = \{1, \dots, n\}$ we write R^n instead of $R^{(S)}$.

The key feature of a free module is that a homomorphism out of it is completely determined by where the basis goes. Precisely: for every function of sets $\alpha: S \rightarrow M$ (where M is any R -module), there is a unique R -module homomorphism $\tilde{\alpha}: R^{(S)} \rightarrow M$ such that $\tilde{\alpha}(e_s) = \alpha(s)$ for all $s \in S$. In categorical language, the functor $S \mapsto R^{(S)}$ is left adjoint to the forgetful functor from R -modules to sets. This universal property characterizes $R^{(S)}$ up to unique isomorphism and is the reason free modules play a role analogous to that of vector spaces—they are the modules we can map out of “for free.”

Problem 8. Free Modules and Their Universal Property.

1. Verify directly that every element of $R^{(S)}$ can be written uniquely as a finite R -linear combination of the standard basis elements e_s .
2. Prove the universal property of $R^{(S)}$ stated above.
3. Deduce that there is a natural isomorphism

$$\operatorname{Hom}_R(R^{(S)}, M) \cong \prod_{s \in S} M.$$

Why is the right side a product rather than a direct sum?

4. Show that an R -module M is generated by $m_1, \dots, m_n$ if and only if there is a surjective homomorphism $R^n \twoheadrightarrow M$ sending e_i to m_i .
5. Conclude that every finitely generated R -module is a quotient of a finite free module.
6. More generally, show that every R -module is a quotient of a free R -module.
7. For $M = R/I$, write M as a quotient of a free module and identify the kernel.
8. For $R = \mathbb{K}[x]$ and $M = R/\langle x^3 \rangle$, describe M as a module with one generator and one relation.

Solution.

1. Let $f \in R^{(S)}$. Then by definition $f(s) \neq 0$ for only finitely many $s \in S$. So we can write $f = \sum_{s \in S} f(s) \cdot e_s = \sum_{s \in S} r_s \cdot e_s$ for some $r_s \in R$, which is a finite sum and hence every element of $R^{(S)}$ can be written uniquely as a finite R -linear combination of the standard basis elements.

Uniqueness: Suppose $f = \sum_{s \in S} r'_s e_s = \sum_{s \in S} r_s e_s$. Then comparing the s -th element gives us $r'_s e_s = r_s e_s \implies r'_s = r_s$. Hence the linear combination is unique.

2. **Existence:** Let $\alpha : S \rightarrow M$ be a set map. For each $s \in S$, define $\psi_s : R \rightarrow M$ by $\psi_s(r) = r \cdot \alpha(s)$. This is an R -module homomorphism since

$$\psi_s(r + r') = (r + r')\alpha(s) = r\alpha(s) + r'\alpha(s) = \psi_s(r) + \psi_s(r')$$

and $\psi_s(rr') = rr'\alpha(s) = r\psi_s(r')$. Since $R^{(S)} = \bigoplus_{s \in S} R$, by the universal property of the direct sum there exists a unique R -module homomorphism $\tilde{\alpha} : R^{(S)} \rightarrow M$ such

that $\tilde{\alpha} \circ \iota_s = \psi_s$ for all $s \in S$. In particular, $\tilde{\alpha}(e_s) = \psi_s(1_R) = \alpha(s)$, so the diagram

$$\begin{array}{ccc} R^{(S)} & \xrightarrow{\tilde{\alpha}} & M \\ \iota_s \uparrow & \nearrow \alpha & \\ S & & \end{array}$$

commutes for all $s \in S$.

Uniqueness: Any R -module homomorphism $\phi : R^{(S)} \rightarrow M$ satisfying $\phi(e_s) = \alpha(s)$ for all $s \in S$ must satisfy

$$\phi\left(\sum_{s \in S} r_s e_s\right) = \sum_{s \in S} r_s \phi(e_s) = \sum_{s \in S} r_s \alpha(s) = \tilde{\alpha}\left(\sum_{s \in S} r_s e_s\right).$$

Since every element of $R^{(S)}$ is a finite R -linear combination of the e_s , we conclude $\phi = \tilde{\alpha}$.

3. By the universal property, every homomorphism $\phi : R^{(S)} \rightarrow M$ is uniquely determined by $\phi(e_s) \in M$ for each $s \in S$. Then the map,

$$\text{Hom}_R(R^{(S)}, M) \rightarrow \prod_{s \in S} M, \quad \phi \mapsto (\phi(e_s))_{s \in S}$$

is a bijection. There is no finiteness restriction on how many $\phi(e_s) \neq 0$, hence the right-hand side must allow non-zero infinite tuples and hence is $\prod_{s \in S} M$.

4. ($\implies$)

Assume M is a R -module generated by $m_1, \dots, m_n$. Then from part (3) we know there exists a homomorphism $\phi : R^n \rightarrow M$ that sends $e_i \mapsto m_i$ for all $i \in \{1, \dots, n\}$. This homomorphism is surjective because all elements of M are finite linear combinations of m_i .

- ($\impliedby$)

Assume there exists a surjective homomorphism $\phi : R^n \rightarrow M$. Then for any $m \in M$, surjectivity gives some $x \in R^n$ with $\phi(x) = m$ and writing $x = \sum r_i e_i$ gives $m = \phi(\sum r_i e_i) = \sum r_i \phi(e_i)$. Hence, M is generated by $\{\phi(e_1), \dots, \phi(e_n)\}$.

5. Let M be a finitely-generated R -module. Then from part (5) we know there exists some surjective homomorphism $\phi : R^n \twoheadrightarrow M$ for some $n \in \mathbb{N}$. Then by the first isomorphism theorem we have an induced isomorphism,

$$\bar{\phi} : R^n / \ker(\phi) \cong M$$

6. Let M be a R -module, and define $S = M$. Then there exists a homomorphism $\phi : R^{(M)} \rightarrow M$ that sends $e_m \mapsto m$. Again, by the first isomorphism theorem there is an induced homomorphism $\bar{\phi}$,

$$\bar{\phi} : R^{(M)} / \ker(\phi) \cong M$$

7. Let $S = \{1\}$. Then we have $\phi : R^{(S)} \twoheadrightarrow R/I = M$ with $\ker(\phi) = I$.
8. M is generated by $\bar{1} \in M$ and obeys the relation $x^3 = 0$.

Problem 9. Free Modules Are Not Quite Vector Spaces.

1. Show that $\mathbb{Z}/n\mathbb{Z}$ is not a free $\mathbb{Z}$ -module for any $n > 1$.
2. More generally, if R is an integral domain and $0 \neq I \subsetneq R$ is an ideal, show that R/I is not a free R -module.
3. In the $\mathbb{Z}$ -module $\mathbb{Z}$, show that $\{2, 3\}$ is a generating set but not a basis.
4. In the $\mathbb{Z}$ -module $\mathbb{Z}$, show that $\{2\}$ is linearly independent but does not generate $\mathbb{Z}$.
5. Let $R = \mathbb{Z}/6\mathbb{Z}$. Show that the element $\bar{2} \in R$ is nonzero but not part of a basis for the free R -module R .
6. Explain why the previous examples show that bases over rings behave differently from bases over fields.

Solution.

1. Suppose for contradiction that $\mathbb{Z}/n\mathbb{Z}$ is a $\mathbb{Z}$ -module with basis B . Then B must be linearly independent, meaning whenever $m_1, \dots, m_k \in B$, are distinct and $\sum_{i=1}^k r_i m_i = 0$ we have $r_i = 0$ for all i . But for any $m \in B \subseteq \mathbb{Z}/n\mathbb{Z}$ we have $n \cdot m = 0$ in $\mathbb{Z}/n\mathbb{Z}$ with $n \neq 0 \in \mathbb{Z}$ contradicting linear independence. Hence $\mathbb{Z}/n\mathbb{Z}$ admits no basis and is not a free $\mathbb{Z}$ -module.
2. Suppose for contradiction that R/I is a free R -module with basis B . Since $I \neq 0$, pick some $0 \neq a \in I$. Then for any $m \in B \subseteq R/I$ we have $a \cdot m = 0 \in R/I$, with $a \neq 0$ thus contradicting linear independence. Hence R/I admits no basis and is not a free R -module.
3. $\{2, 3\}$ is a generating set since $3 - 2 = 1$ and 1 is a generator for $\mathbb{Z}$. But $\{2, 3\}$ is not a basis $3 \cdot 2 + (-2) \cdot 3 = 0$, where $3, -2 \neq 0$. Thus violating linear independence. Hence, $\{2, 3\}$ is not a basis.
4. $\{2\}$ is linearly independent because if $r \cdot 2 = 0$ then $r = 0$ since $\mathbb{Z}$ is an integral domain. But $\{2\}$ is not a generating set because no finite linear combination of $\{2\}$ gives us 1.
5. $\bar{2} \neq 0$ since $2 \not\equiv 0 \pmod{6}$, and $\bar{2} \cdot \bar{3} = 0$ and hence $\bar{2}$ fails linear independence and cannot be part of a basis.
6. Over a field every linearly-independent set extends to a basis and every spanning set contains a basis. The previous examples show that both properties fail over general rings.

Problem 10. Matrices and Maps Between Finite Free Modules.

1. Let $A = (a_{ij})$ be an $m \times n$ matrix with entries in R . Define a homomorphism

$$\phi_A : R^n \rightarrow R^m, \quad e_j \mapsto \sum_{i=1}^m a_{ij} e_i.$$

Prove that every homomorphism $R^n \rightarrow R^m$ arises uniquely in this way.

2. Under this correspondence, prove that composition of homomorphisms corresponds to multiplication of matrices.
3. When $R = \mathbb{K}$ is a field, explain how this recovers the usual matrix description of linear maps between finite-dimensional vector spaces.
4. Let $A = \begin{pmatrix} 2 & 0 \\ 0 & 3 \end{pmatrix}$ and view A as a homomorphism $\mathbb{Z}^2 \rightarrow \mathbb{Z}^2$. Describe $\text{coker}(A) = \mathbb{Z}^2 / \text{img}(A)$.
5. Let $R = \mathbb{K}[x]$ and consider multiplication by x as a map $R^2 \rightarrow R^2$. Write its matrix with respect to the standard basis, and compute its cokernel.

Solution.

1. **Existence:** By the universal property of free modules, a homomorphism $\phi : R^n \rightarrow R^m$ is determined by where it sends the basis vectors $e_1, \dots, e_n$. Then, for all j , $\phi(e_j) \in R^m$ can be written as

$$\phi(e_j) = \sum_{i=1}^m a_{ij} e_i$$

for some $a_{ij} \in R$. Then we define $A = (a_{ij})$ to be a $m \times n$ matrix. Then $\phi = \phi_A$ by construction.

Uniqueness: The matrix A is unique since each entry a_{ij} is uniquely determined as a linear combination of the basis elements e_i , hence no two matrices give the same homomorphism.

Hence every homomorphism $R^n \rightarrow R^m$ arises uniquely in this way.

2. Let $\phi : R^n \rightarrow R^p$, $\psi : R^p \rightarrow R^m$ be two homomorphisms with corresponding matrices $A = (a_{kj})$, $B = (b_{ik})$ respectively. Then for each basis vector $e_j \in R^n$,

$$(\psi \circ \phi)(e_j) = \psi \left(\sum_{k=1}^p a_{kj} e_k \right) = \sum_{k=1}^p a_{kj} \psi(e_k) = \sum_{k=1}^p a_{kj} \sum_{i=1}^m b_{ik} e_i = \sum_{i=1}^m \left(\sum_{k=1}^p b_{ik} a_{kj} \right) e_i$$

The coefficient of e_i is $\sum_{k=1}^p b_{ik}a_{kj}$, which is exactly the (i, j) -entry of the matrix product BA . Since $\psi \circ \phi$ is determined by its values on the basis vectors, we conclude that $\psi \circ \phi$ corresponds to BA .

3. Let $R = \mathbb{K}$ be a field. Then a homomorphism $\phi : \mathbb{K}^n \rightarrow \mathbb{K}^m$ is a $\mathbb{K}$ -linear map and the correspondence $\phi \rightarrow A$ is the usual representation of a linear map by its matrix with respect to the standard bases.
4. Note that $A(e_1) = (2, 0)$ and $A(e_2) = (0, 3)$. Then we have,

$$\text{img}(A) = \{(2a, 3b) \mid a, b \in \mathbb{Z}\} = 2\mathbb{Z} \oplus 3\mathbb{Z}$$

$$\text{Hence, } \text{coker}(A) = \mathbb{Z}^2 / \text{img}(A) = \mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/3\mathbb{Z}$$

5. Let A be the matrix that corresponds to multiplication by x . Then, $A = \begin{pmatrix} x & 0 \\ 0 & x \end{pmatrix}$, and $\text{coker}(A) = R^2 / \text{img}(A) = \mathbb{K} \oplus \mathbb{K}$.

For two R -modules M and N , we write $\text{Hom}_R(M, N)$ for the set of R -module homomorphisms $M \rightarrow N$. Since R is commutative, $\text{Hom}_R(M, N)$ is itself an R -module: addition and scalar multiplication are defined pointwise by

$$(\phi + \psi)(m) = \phi(m) + \psi(m), \quad (r\phi)(m) = r\phi(m).$$

This construction is one of the most important ways of building new modules from old ones. There is already a great deal to understand: Hom out of a free module is easy, Hom out of a quotient records annihilators, and Hom interacts cleanly with direct sums and products through their universal properties.

Problem 11. $\text{Hom}_R(M, N)$ as a Module.

1. Prove that $\text{Hom}_R(M, N)$ is an abelian group under pointwise addition. What is the zero element?
2. Prove that the formula $(r\phi)(m) = r\phi(m)$ makes $\text{Hom}_R(M, N)$ into an R -module.
3. Show that evaluation at $1 \in R$ defines an isomorphism of R -modules

$$\text{Hom}_R(R, M) \cong M.$$

4. More generally, prove that

$$\text{Hom}_R(R/I, M) \cong \{m \in M \mid Im = 0\}.$$

The isomorphism should again be given by evaluation at $\bar{1} \in R/I$.

5. Deduce that $\text{Hom}_R(R^n, M) \cong M^n$.
6. Prove that $\text{End}_R(R) \cong R$ as rings, where the ring structure on $\text{End}_R(R)$ is given by addition and composition.
7. If $I \subseteq R$ is an ideal, identify $\text{Hom}_R(R/I, R)$ in terms of an annihilator ideal of R .

Solution.

1. Let $\phi, \psi \in \text{Hom}_R(M, N)$. Then we define,

$$(\phi + \psi)(m) := \phi(m) + \psi(m) \quad \forall m \in M$$

Then since ϕ, ψ are homomorphisms and addition in N is commutative, for $m, m' \in M$ we have,

$$\begin{aligned} (\phi + \psi)(m + m') &= \phi(m + m') + \psi(m + m') \\ &= \phi(m) + \phi(m') + \psi(m) + \psi(m') \\ &= \phi(m) + \psi(m) + \phi(m') + \psi(m') \\ &= (\phi + \psi)(m) + (\phi + \psi)(m') \end{aligned}$$

Also note that associativity and commutativity are inherited from N , and the additive inverse of ϕ , $(-\phi)(m) := -\phi(m)$ is well-defined. The zero element is the zero map, $0(m) := 0_N$ for all $m \in M$,

$$(f + 0)(m) = f(m) + 0(m) = f(m) + 0_N = f(m)$$

Hence, $\text{Hom}_R(M, N)$ is an abelian group under pointwise addition.

2. We know that $\text{Hom}_R(M, N)$ is an abelian group, and the formula $(r\phi)(m) = r\phi(m)$ shows that $\text{Hom}_R(M, N)$ is closed under scalar multiplication,

$$(r\phi)(sm) = r\phi(sm) = rs\phi(m) = sr\phi(m) = s(r\phi)(m)$$

All other module properties follow, showing that it is a R -module.

3. Define a map $\eta : \text{Hom}_R(R, M) \rightarrow M$, where $\phi \mapsto \phi(1) = m$ for some $m \in M$. This map is injective since the homomorphism is completely determined by where 1 goes, so if $\phi(1) = \psi(1) \implies \phi = \psi$.

To show that this map is a surjection we must show that there exists a homomorphism $\phi_m \in \text{Hom}_R(R, M)$ such that $\phi_m(1) = m$ for all $m \in M$. So, we define

$$\phi_m(r) = rm \quad \forall r \in R$$

and show that ϕ_m is a homomorphism:

- $\phi_m(s \cdot r) = (s \cdot r)m = s(rm) = s\phi_m(r)$
- $\phi_m(s + r) = (s + r)m = sm + rm = \phi_m(s) + \phi_m(r)$

So for every $m \in M$ there exists a homomorphism ϕ_m that maps to it under η .

4. Similarly, we define a map $\eta : \text{Hom}_R(R/I, M) \rightarrow \{m \in M \mid Im = 0\}$, where $\phi \mapsto \phi(\bar{1}) = \phi(1 + I)$. This map is well-defined since for any $a \in I$ we have,

$$a\phi(1 + I) = \phi(a + aI) = \phi(\bar{0}) = 0$$

This map is injective since the homomorphism is completely determined by where $\bar{1}$ goes, so

$$\eta(\phi) = \eta(\psi) \implies \phi(1 + I) = \psi(1 + I) \implies \phi = \psi$$

To show that this map is a surjection we define

$$\phi_m(r + I) = rm \quad \forall r \in R$$

and show that ϕ_m is a well-defined homomorphism:

- Let $s + I = r + I$ then, $s - r \in I$ and

$$(s - r)m = 0 \implies sm = rm \implies \phi_m(s + I) = \phi_m(r + I)$$

- $\phi_m(s(r + I)) = \phi(sr + I) = srm = s\phi(r + I)$
- $\phi_m((s + r) + I) = (s + r)m = sm + rm = \phi_m(s) + \phi_m(r)$

So for every $m \in M$ such that $mI = 0$ there exists a homomorphism ϕ_m that maps to it under η .

- 5.

Worksheet 4.2: Localization

Throughout this course, “ring” means *commutative* ring with unity. In the previous worksheet we built the basic language of modules, quotient modules, free modules, Hom, and tensor products. The goal of this worksheet is to introduce one of the central operations in commutative algebra: *localization*. Informally, localization is the process of forcing a chosen collection of elements of a ring to become invertible. This construction simultaneously generalizes the passage from $\mathbb{Z}$ to $\mathbb{Q}$, the localization $\mathbb{Z}_{\langle p \rangle}$ at a prime number p , and the algebraic operation of restricting attention to a principal open subset $D(f)$ of $\text{Spec}(R)$.

Let R be a ring. A subset $U \subset R$ is called *multiplicatively closed* if $1 \in U$ and $u, t \in U$ implies $ut \in U$. Given a multiplicative set $U \subset R$, the localization of R at U is a ring $U^{-1}R$ together with a ring homomorphism $\eta : R \rightarrow U^{-1}R$ satisfying the following universal property: For all ring homomorphisms $\phi : R \rightarrow T$ such that every element of $\phi(U)$ is a unit in T there exists a unique homomorphism $\tilde{\phi} : U^{-1}R \rightarrow T$ such that the following diagram commutes:

$$\begin{array}{ccc} R & \xrightarrow{\eta} & U^{-1}R \\ \phi \downarrow & \nearrow \tilde{\phi} & \\ T & & \end{array} .$$

Roughly speaking we should think of $U^{-1}R$ as being the “smallest” ring where every element of U is a unit. As always the universal property guarantees that $U^{-1}R$ and the map η are unique up to unique isomorphism (assuming that they exist, which we will now discuss).

Informally, we think of elements of $U^{-1}R$ as obtained by formally inverting every element of U and imposing the necessary equivalence conditions to make it into a well-behaved ring. Elements of $U^{-1}R$ are formal fractions r/u where $r \in R$ and $u \in U$ subject to some equivalence relation. If R is an integral domain and $0 \notin U$, equality of fractions is the familiar cross-multiplication relation, i.e. $r_1/u_1 = r_2/u_2$ as elements in $U^{-1}R$ if and only if $r_1u_2 = r_2u_1$ as elements in R . In general, because zero divisors may be present, one must allow extra clearing denominators:

$$\frac{r_1}{u_1} = \frac{r_2}{u_2} \text{ in } U^{-1}R \iff t(r_1u_2 - r_2u_1) = 0 \text{ in } R \text{ for some } t \in U$$

Multiplication and addition in $U^{-1}R$ are then defined in the usual way for fractions:

$$\left(\frac{r_1}{u_1} \right) \left(\frac{r_2}{u_2} \right) = \frac{r_1r_2}{u_1u_2} \quad \text{and} \quad \frac{r_1}{u_1} + \frac{r_2}{u_2} = \frac{r_1u_2 + r_2u_1}{u_1u_2}$$

The additive identity is then the equivalence class of $0/1$ and the multiplicative identity is $1/1$. The function $\eta : R \rightarrow U^{-1}R$ is defined by $r \mapsto \frac{r}{1}$, which formally makes sense because

$1 \in U$. Although this construction is relatively easy to define, there is work involved in checking that the operations and η are well-defined, that $U^{-1}R$ is a ring, and that it satisfies the correct universal property. You will carefully go through this construction in the exercises.

A few warnings: when R is not a domain localization can behave somewhat strangely. For example, from our experience with fractions since childhood and its unassuming definition one might expect η to be injective; this is not the case. The element $\eta(r) = r/1$ can be equal to zero in $U^{-1}R$. By definition this occurs when $r/1 = 0/1$, which translates to $tr = 0$ in R for some element $t \in U$. Even worse, if $0 \in U$ then the ring $U^{-1}R$ becomes the zero ring.

Problem 1. Multiplicative Sets.

1. Verify using the definition directly which of the following subsets are multiplicatively closed:

(a) $R = \mathbb{Z}$ and $U = \{1, 3, 9, 27, \dots\}$.	(f) $R = \mathbb{R}[x, y]$ and $U = \{f \in R \mid f(1, 0) \neq 0\}$.
(b) $R = \mathbb{Z}$ and $U = \{1, 2, 3\}$.	(g) $R = \mathbb{R}[x, y] / \langle y^2 - x^3 + x \rangle$ and $U = R \setminus \langle \bar{x}, \bar{y} \rangle$.
(c) $R = \mathbb{Z}$ and $U = \mathbb{Z} \setminus \langle 2 \rangle$.	(h) $R = \mathbb{R}[x, y]$ and $U = \{f \in R \mid f(0, 0) \neq 0 \text{ and } f(1, 1) \neq 0\}$.
(d) $R = \mathbb{Z}$ and $U = \{n \geq 1\}$.	
(e) $R = \mathbb{R}[x, y]$ and $U = R \setminus \langle x, y \rangle$.	
2. Let $I \subset R$ be an ideal. Prove that $U = R \setminus I$ is multiplicatively closed if and only if I is prime.
3. Let $U \subset R$ be a multiplicatively closed set. If $\phi : R \rightarrow S$ is a ring homomorphism, is $\phi(U) \subset S$ a multiplicatively closed subset?

Solution.

1. (a) U is multiplicatively closed.
- (b) U is not multiplicatively closed; $2, 3 \in U$ but $2 \cdot 3 \notin U$.
- (c) U is multiplicatively closed.
- (d) U is multiplicatively closed.
- (e) $U = \mathbb{R}$ and hence is multiplicatively closed.
- (f) U is multiplicatively closed.
- (g) U is multiplicatively closed.

(h) U is multiplicatively closed.

2. Let $I \subset R$ ideal.

($\implies$)

Assume $U = R \setminus I$ is multiplicatively closed. Assume $a, b \notin I$. Then $a, b \in U$. Since U is multiplicatively closed, $ab \in U$, hence $ab \notin I$. That is if $a, b \notin I$ then $ab \notin I$. Thus, I is prime.

($\impliedby$)

Assume I is prime. Then, we know that if $a, b \notin I$, $ab \notin I$. Thus, $a, b, ab \in U$, and U is multiplicatively closed.

3. Let $U \subset R$ be multiplicatively closed. Let $\phi : R \rightarrow S$ be a ring homomorphism. Then for any $r, s \in U$ we have,

$$\phi(r) \cdot \phi(s) = \phi(rs) \in \phi(U)$$

Hence, $\phi(U) \subset S$ is multiplicatively closed.

Problem 2. First Examples.

Despite not having formally verified that our construction is well-defined, discuss ways to explicitly understand $U^{-1}R$ in each of the following cases.

1. $R = \mathbb{Z}$ and $U = \mathbb{Z} \setminus \{0\}$.
2. $R = \mathbb{Z}$ and $U = \{2^n \mid n \in \mathbb{Z}_{\geq 0}\}$.
3. $R = \mathbb{Z}$ and $U = \mathbb{Z} \setminus \langle 2 \rangle$.
4. $R = \mathbb{K}[x]$ and $U = \mathbb{K}[x] \setminus \{0\}$.
5. $R = \mathbb{K}[x]$ and $U = \{1, x, x^2, x^3, \dots\}$.
6. $R = \mathbb{Z}/12\mathbb{Z}$ and $U = \{\bar{1}, \bar{2}, \bar{4}, \bar{8}\}$.
7. $R = \mathbb{Z}/12\mathbb{Z}$ and $U = \{\bar{1}, \bar{3}, \bar{9}\}$.
8. $R = \mathbb{Z}[x]/\langle nx - 1 \rangle$ and $U = \{n^k \mid k \in \mathbb{Z}_{\geq 0}\}$ for a fixed non-zero integer n .
9. R an arbitrary ring and U a multiplicative closed set containing a nilpotent.

Solution.

1. $U^{-1}R = \mathbb{Q}$
2. $U^{-1}R = \{m/2^n \mid m \in \mathbb{Z}\}$. These are the dyadic integers.
3. $U^{-1}R = \{m/n \mid m \in \mathbb{Z}, n \text{ odd}\}$
4. $U^{-1}R = \mathbb{K}(x)$
5. $U^{-1}R = \mathbb{K}[x, x^{-1}]$
6. $U^{-1}R = \mathbb{Z}/3\mathbb{Z}$
7. $U^{-1}R = \mathbb{Z}/4\mathbb{Z}$
8. $U^{-1}R = \mathbb{Z} \left[\frac{1}{n} \right] = R$
9. If $u \in U$ is nilpotent, then for some $k \geq 1$ $u^k = 0 \in U$. Then for any two arbitrary fractions $a/b, c/d \in U$, pick $w = 0$. Then we have

$$w(ad - cd) = 0(ad - cd) = 0$$

that is, all fractions are equivalent and $U^{-1}R = \{0\}$.

Problem 3. Constructing $U^{-1}R$.

Let R be a ring and $U \subset R$ a multiplicatively closed set. Define a relation on $R \times U$ by saying $(r_1, u_1) \sim (r_2, u_2)$ if and only if $t(r_1u_2 - r_2u_1) = 0$ in R for some $t \in U$.

1. Prove that $\sim$ is an equivalence relation on $R \times U$.
2. Let $\frac{r}{u}$ denote the equivalence class of a pair (r, u) under the equivalence relation $\sim$ defined in the previous part. Prove that the operations:

$$\left(\frac{r_1}{u_1}\right) \left(\frac{r_2}{u_2}\right) = \frac{r_1r_2}{u_1u_2} \quad \text{and} \quad \frac{r_1}{u_1} + \frac{r_2}{u_2} = \frac{r_1u_2 + r_2u_1}{u_1u_2}$$

are well-defined. Hint: You must show that changing the representative of either input fraction does not change the resulting equivalence class.

3. Define $U^{-1}R$ to be the set of equivalence classes of $\sim$. Show that the operations above make $U^{-1}R$ into a ring where $\frac{0}{1}$ is the zero element and $\frac{1}{1}$ is the multiplicative identity.
4. Show that the function $\eta : R \rightarrow U^{-1}R$ given by $r \mapsto \frac{r}{1}$ is a well-defined ring homomorphism.
5. Prove that if $u \in U$ then $\eta(u)$ is a unit in $U^{-1}R$. Explicitly what is its inverse?
6. Where in this construction did you use that U was multiplicatively closed and contains 1?

Solution.

1. Let $(r_1, u_1) \sim (r_2, u_2) \sim (r_3, u_3) \in R \times U$.

(a) **Reflexivity:** $(r_1, u_1) \sim (r_1, u_1)$ since $1(r_1u_1 - r_1u_1) = 0$.

(b) **Symmetry:** If $(r_1, u_1) \sim (r_2, u_2)$ then there exists a $t \in U$ such that

$$t(r_1u_2 - r_2u_1) = 0 \implies tr_1u_2 = tr_2u_1 \implies t(r_2u_1 - r_1u_2) = 0$$

Hence, $(r_2, u_2) \sim (r_1, u_1)$.

(c) **Transitivity:** If $(r_1, u_1) \sim (r_2, u_2)$ and $(r_2, u_2) \sim (r_3, u_3)$ then there exist s, t such that

$$s(r_1u_2 - r_2u_1) = 0, \quad t(r_2u_3 - r_3u_2) = 0$$

Note that

$$stu_3(r_1u_2 - r_2u_1) = 0, \quad stu_1(r_2u_3 - r_3u_2) = 0$$

Adding,

$$0 = st(r_1u_2u_3 - r_2u_1u_3 + r_2u_3u_1 - r_3u_2u_1) = stu_2(r_1u_3 - r_3u_1)$$

Since $stu_2 \in U$ we have $(r_1, u_1) \sim (r_3, u_3)$.

Thus, $\sim$ is an equivalence relation on $R \times U$.

2. Let $(r, u) \sim (r', u')$ and $(s, t) \sim (s', t')$. Then there exist $p, q \in U$ such that

(a) We show that $\frac{rs}{ut} \sim \frac{r's'}{u't'}$:

$$p(ru' - r'u) = 0, \quad q(st' - s't) = 0$$

Multiplying,

$$pqst'(ru' - r'u) = 0, \quad pqr'u(st' - s't) = 0$$

Adding,

$$pq(st'ru' - st'r'u + r'ust' - r'us't) = pq(rsu't' - r's'ut) = 0$$

That is,

$$\left(\frac{r}{u}\right) \left(\frac{s}{t}\right) = \frac{rs}{ut} \sim \frac{r's'}{u't'} = \left(\frac{r'}{u'}\right) \left(\frac{s'}{t'}\right)$$

(b) We show that $\frac{rs}{ut} \sim \frac{r's'}{u't'}$

$$\begin{aligned} & pq((rt + su)(u't') - (r't' + s'u')(ut)) \\ &= pq(rtu't' + suu't' - r't'ut - s'u'ut) \\ &= pq(t't(ru' - r'u) + uu'(st' - s't)) \\ &= qt't \cdot p(ru' - r'u) + puu' \cdot q(st' - s't) \\ &= qt't \cdot 0 + puu' \cdot 0 = 0 \end{aligned}$$

3. Note that associativity, commutativity, and distributivity are inherited from R .

(a) **Additive Identity:**

$$\frac{r}{u} + \frac{0}{1} = \frac{r+0}{u} = \frac{r}{u}$$

(b) **Additive Inverses:**

$$\frac{r}{u} + \frac{-r}{u} = \frac{ru - ru}{u} = \frac{0}{1}$$

(c) **Multiplicative Identity:**

$$\frac{r}{u} \cdot \frac{1}{1} = \frac{r \cdot 1}{u \cdot 1} = \frac{r}{u}$$

Hence, $U^{-1}R$ is a ring.

4. Let $\eta : R \rightarrow U^{-1}R$, where $r \mapsto \frac{r}{1}$.

- $\eta(r) + \eta(s) = \frac{r}{1} + \frac{s}{1} = \frac{r+s}{1} = \eta(r+s)$
- $\eta(r) \cdot \eta(s) = \frac{r}{1} \cdot \frac{s}{1} = \frac{rs}{1} = \eta(rs)$

Hence, η is a ring homomorphism.

5. For any $u \in U$ we have,

$$\eta(u) \cdot \frac{1}{u} = \frac{u}{1} \cdot \frac{1}{u} = \frac{u}{u} \sim \frac{1}{1}$$

6. In proving transitivity and verifying that the operations are well-defined.

Problem 4. First Properties of Localization.

Let R be a ring and $U \subset R$ a multiplicatively closed set.

1. Prove that if $0 \in U$ then $U^{-1}R$ is the zero ring.
2. Prove that if R is an integral domain and $0 \notin U$ then $\frac{r_1}{u_1} = \frac{r_2}{u_2}$ in $U^{-1}R$ if and only if $r_1u_2 = r_2u_1$.
3. Let R be an integral domain and $U = R \setminus \{0\}$. Prove that $U^{-1}R$ is a field.
4. Find an example of a non-zero ring R for which $R \setminus \{0\}$ is not multiplicatively closed. What goes wrong with trying to form $(R \setminus \{0\})^{-1}R$?

Solution.

1. If $0 \in U$. Then for any two arbitrary fractions $a/b, c/d \in U$, pick $w = 0$. And we have

$$w(ad - cd) = 0(ad - cd) = 0$$

that is, all fractions are equivalent and $U^{-1}R = \{0\}$.

2. Assume R be an integral domain, and $0 \notin U$.

($\implies$)

Assume $\frac{r_1}{u_1} = \frac{r_2}{u_2}$ in $U^{-1}R$. Then there exists a $t \neq 0 \in U$ such that

$$t(r_1u_2 - u_1r_2) = 0$$

Since R is an integral domain this implies that

$$r_1u_2 - u_1r_2 = 0 \implies r_1u_2 = u_1r_2$$

($\impliedby$)

Assume $r_1u_2 = r_2u_1$. Then with $w = 1 \in U$,

$$r_1u_2 - r_2u_1 = 0 \implies \frac{r_1}{u_1} = \frac{r_2}{u_2}$$

Hence proved.

3. Let $\frac{r}{u} \approx \frac{0}{1} \in U^{-1}R$. From (2) we know that $r \neq 0$. Then $\frac{u}{r} \in U^{-1}R$ and hence, $U^{-1}R$ is a field.
4. Let $R = \mathbb{Z}/6\mathbb{Z}$. Then, we have $\frac{1}{2}, \frac{1}{3} \approx \frac{0}{1} \in (R \setminus \{0\})^{-1}R$. But $\frac{1}{2} \cdot \frac{1}{3} = \frac{1}{6} \sim \frac{1}{0}$ which is not defined.

Problem 5. The Kernel of the Localization Map.

Let R be a ring, $U \subset R$ be a multiplicatively closed set, and $\eta : R \rightarrow U^{-1}R$ the localization map as described in Problem 3.

1. Prove that η is injective if and only if every $u \in U$ is a nonzerodivisor on R , i.e. $ur = 0$ implies $r = 0$ for all $r \in R$.
2. Let $I = \{r \in R \mid ur = 0 \text{ for some } u \in U\}$. Prove that I is an ideal.
3. Show that there exists an injective ring homomorphism $R/I \rightarrow U^{-1}R$ such that the following diagram commutes:

$$\begin{array}{ccc} R & \xrightarrow{\pi} & R/I \\ & \searrow \eta & \downarrow \\ & & U^{-1}R \end{array} .$$

4. Prove that $\ker(\eta) = I$.
5. Revisit the examples in Problem 2. For which of them is the localization map $R \rightarrow U^{-1}R$ injective?

Solution.

1. ($\implies$)

Assume η is injective. Suppose $ur = 0$ for some $u \in U$. Then,

$$\eta(r) = \frac{r}{1} = \frac{ur}{u} = \frac{0}{u} \sim \frac{0}{1}$$

Since $\ker(\eta) = \{0\}$, this implies that either $r = 0$.

(We cannot have $u = 0$ since that would make $U^{-1}R$ into the zero ring, and η is injective.) Hence, U contains no zero-divisors on R .

($\impliedby$)

Assume U has no zero-divisors in R . An element r is in $\ker(\eta)$ iff $\eta(r) \sim \frac{0}{1}$. That is there exists $u \in U$ such that,

$$u(r - 0) = 0$$

But by assumption, u is not a zero-divisor and hence r must be 0. Since r was an arbitrary element, this implies that $\ker(\eta) = \{0\}$ and η is injective.

2. Since $I \subseteq R$ we only need to show that I is closed under addition and multiplication (from R).

(a) **Addition:** Let $r, s \in I$. Then there exist u, v such that $ru = 0, sv = 0$. Since U is multiplicatively closed, $uv \in U$. Then we have,

$$(r + s)uv = (ru)v + (sv)u = 0$$

Hence, $r + s \in I$.

(b) Let $r \in R$ and $s \in I$. Then there exist some $u \in U$ such that $su = 0$. Consider rs ,

$$(rs)u = r(su) = 0$$

Hence, $rs \in I$.

Thus I is an ideal.

3. Define $\phi : R/I \rightarrow U^{-1}R$, where $\bar{r} \mapsto \frac{r}{1}$. Now we check for the following:

(a) **Well-Definedness:** Let $\bar{r} = \bar{s}$. Then, $\bar{r} - \bar{s} \in I \implies \exists u \in U$ such that

$$u(r - s) = 0 \implies u(r \cdot 1 - s \cdot 1) \implies \frac{r}{1} \sim \frac{s}{1}$$

(b) **Ring Homomorphism:**

- $\phi(\bar{r} + \bar{s}) = \frac{r+s}{1} = \frac{r}{1} + \frac{s}{1} = \phi(\bar{r}) + \phi(\bar{s})$
- $\phi(\bar{r} \cdot \bar{s}) = \frac{r \cdot s}{1} = \frac{r}{1} \cdot \frac{s}{1} = \phi(\bar{r}) \cdot \phi(\bar{s})$

(c) **Injectivity:** Let $\bar{r} \in \ker(\phi)$. Then $\phi(\bar{r}) = \frac{r}{1} \sim \frac{0}{1}$. That is, there exists $u \in U$ such that $u(r) = 0$. Then $\bar{r} \in I$ and hence $\bar{r} = \bar{0} \in R/I$.

(d) **Commutates:** Let $r \in R$. Then we have,

$$\eta(r) = \frac{r}{1}, \quad \phi(\pi(r)) = \phi(\bar{r}) = \frac{r}{1}$$

Hence, there exists an injective ring homomorphism $R/I \rightarrow U^{-1}R$

4. $\ker(\eta) = \ker(\phi \circ \pi) = \pi^{-1}(\ker(\phi)) = \pi^{-1}(\langle 0 \rangle) = I$.

5. η is injective for subparts 1, 2, 3, 4, 5, 8.

Problem 6. Field of Fractions.

Let R be an integral domain. The *field of fractions* of R is the ring $U^{-1}R$ where $U = R \setminus \{0\}$. We denote the field of fractions of R by $\text{Frac}(R)$.

1. Show that $\text{Frac}(R)$ is a field containing R .
2. Describe $\text{Frac}(R)$ when $R = \mathbb{Z}$ and $R = \mathbb{K}[x]$.
3. If R is an integral domain and $V \subset R$ is any multiplicatively closed set not containing zero, explain how we can view $V^{-1}R$ as a subring of $\text{Frac}(R)$.

Solution.

1. From Question 4.3 we know that $\text{Frac}(R)$ is a field, and contains R since $\eta(r) = \frac{r}{1} \in \text{Frac}(R)$ for all $r \in R$.
2. $\text{Frac}(\mathbb{Z}) = \mathbb{Q}$, and $\text{Frac}(\mathbb{K}[x]) = \mathbb{K}(x)$.
3. Note that $V \subseteq U$. Then every element $a = \frac{r}{v} \in V^{-1}R$, is also an element of $\text{Frac}(R)$. Then we can define an embedding $\iota : V^{-1}R \rightarrow U^{-1}R$ where $\iota\left(\frac{r}{v}\right) = \left(\frac{r}{v}\right)$, showing that $V^{-1}R$ is a subring of $U^{-1}R = \text{Frac}(R)$.

Problem 7. Universal Property of Localization.

In this exercise we shall show that $U^{-1}R$ and η as constructed in Problem 3 satisfy the universal property of localization. Let R be a ring and $U \subset R$ a multiplicatively closed subset. Let $\phi : R \rightarrow T$ be a ring homomorphism such that $\phi(U) \subset T^\times$.

1. Define a function $\tilde{\phi} : U^{-1}R \rightarrow T$ by $\frac{r}{u} \mapsto \phi(r)\phi(u)^{-1}$. Explain why $\tilde{\phi}$ is well-defined and prove it is a ring homomorphism.
2. Show that $\tilde{\phi}$ makes the following diagram commute:

$$\begin{array}{ccc} R & \xrightarrow{\eta} & U^{-1}R \\ \phi \downarrow & \swarrow \tilde{\phi} & \\ T & & \end{array} .$$

3. Prove that $\tilde{\phi}$ is the unique ring homomorphism $U^{-1}R \rightarrow T$ making the diagram commute. Hint: If $\psi : U^{-1}R \rightarrow T$ is any such map, first determine $\psi\left(\frac{1}{u}\right)$ for $u \in U$, and then use the equality $\frac{r}{u} = \frac{r}{1} \frac{1}{u}$.

Solution.

1. Let $\frac{r}{u} \sim \frac{s}{v} \in U^{-1}R$, we want to show that $\phi(r)\phi(u)^{-1} = \phi(s)\phi(v)^{-1}$. Then there exists some $t \in U$ such that $t(rv - su) = 0$. Applying ϕ ,

$$\phi(t)(\phi(r)\phi(v) - \phi(s)\phi(u)) = \phi(0) = 0$$

Since $\phi(t) \in \phi(U) \subseteq T^\times$ we can multiply both sides by $\phi(t)^{-1}$,

$$\phi(r)\phi(v) = \phi(s)\phi(u) \implies \phi(r)\phi(u)^{-1} = \phi(s)\phi(v)^{-1}$$

Hence $\tilde{\phi}$ is well-defined.

Ring homomorphism:

• **Addition:**

$$\begin{aligned} \tilde{\phi}\left(\frac{r}{u} + \frac{s}{v}\right) &= \tilde{\phi}\left(\frac{rv + su}{uv}\right) \\ &= \phi(rv + su)\phi(uv)^{-1} \\ &= \phi(r)\phi(v)\phi(u)^{-1}\phi(v)^{-1} + \phi(s)\phi(u)\phi(u)^{-1}\phi(v)^{-1} \\ &= \phi(r)\phi(u)^{-1} + \phi(s)\phi(v)^{-1} \\ &= \tilde{\phi}\left(\frac{r}{u}\right) + \tilde{\phi}\left(\frac{s}{v}\right) \end{aligned}$$

• **Multiplication:**

$$\begin{aligned}
\tilde{\phi}\left(\frac{r}{u} \cdot \frac{s}{v}\right) &= \tilde{\phi}\left(\frac{rs}{uv}\right) \\
&= \phi(rs)\phi(uv)^{-1} \\
&= \phi(r)\phi(s)\phi(u)^{-1}\phi(v)^{-1} \\
&= \phi(r)\phi(u)^{-1} \cdot \phi(s)\phi(v)^{-1} \\
&= \tilde{\phi}\left(\frac{r}{u}\right) \cdot \tilde{\phi}\left(\frac{s}{v}\right)
\end{aligned}$$

Hence $\tilde{\phi}$ is a well-defined ring homomorphism.

2. Let $r \in R$. Then we have,

$$\tilde{\phi}(\eta(r)) = \tilde{\phi}\left(\frac{r}{1}\right) = \phi(r)\phi(1)^{-1} = \phi(r)$$

3. Let $\psi : U^{-1}R \rightarrow T$ be a ring homomorphism that commutes with the given diagram. Then since $\phi(u) = \psi(\eta(u)) = \psi\left(\frac{u}{1}\right)$, and $\psi(u)\psi(u^{-1}) = 1$ we must have $\psi\left(\frac{u}{1}\right) \cdot \psi\left(\frac{1}{u}\right) = \phi(u) \cdot \phi(u)^{-1}$.

That is for an arbitrary element $\frac{r}{u} \in U^{-1}R$ we have,

$$\psi\left(\frac{r}{u}\right) = \psi\left(\frac{r}{1} \cdot \frac{1}{u}\right) = \psi\left(\frac{r}{1}\right) \cdot \psi\left(\frac{1}{u}\right) = \phi(r)\phi(u)^{-1} = \tilde{\phi}\left(\frac{r}{u}\right)$$

Problem 8. Applications of the Universal Property of Localization.

Let R be a ring and $U \subset R$ a multiplicatively closed subset.

1. Use the universal property of localization to show that $U^{-1}R$ and η are unique up to unique isomorphism.
2. Suppose every element of U is already a unit in R . Prove that the localization map $\eta : R \rightarrow U^{-1}R$ is an isomorphism.
3. Suppose $V \subset R$ is another multiplicatively closed set and $U \subset V$. Use the universal property to construct a natural homomorphism $U^{-1}R \rightarrow V^{-1}R$.
4. Let $\phi : R \rightarrow S$ be a ring homomorphism such that $\phi(U) \subset S^\times$. Let $\tilde{\phi} : U^{-1}R \rightarrow S$ be the induced map. Prove that if ϕ is surjective, then $\tilde{\phi}$ is surjective. Prove that if ϕ is injective, then $\tilde{\phi}$ is injective.

Solution.

1. **TODO:**

2. If every element of U is a unit in R then there exist natural maps $\eta : R \rightarrow U^{-1}R$ and $\tilde{\phi} : U^{-1}R \rightarrow R$. We then have the following diagram,

$$\begin{array}{ccc} R & \xrightarrow{\eta} & U^{-1}R \\ \phi \downarrow & \swarrow \tilde{\phi} & \\ R & & \end{array} .$$

This forces $\phi = \text{id}_R$ hence showing that the composition $\tilde{\phi} \circ \eta = \text{id}_R$. Now consider the maps $\eta \circ \tilde{\phi}, \text{id}_{U^{-1}R} : U^{-1}R \rightarrow U^{-1}R$. They agree on composition with η , and hence by uniqueness of the map from $U^{-1}R \rightarrow T$, they must be the same map.

$$(\eta \circ \tilde{\phi}) \circ \eta = \eta \circ (\tilde{\phi} \circ \eta) = \eta \circ \text{id}_R = \eta = \text{id}_{U^{-1}R} \circ \eta$$

Hence $R \cong U^{-1}R$.

3. If $U \subset V$, then every element in U is a unit in $V^{-1}R$, then by the universal property, there exists a map $\tilde{\phi} : U^{-1}R \rightarrow V^{-1}R$.
4. Assume ϕ is surjective. Then for all $s \in S$ there exists some $r \in R$ such that $\phi(r) = s$. Then,

$$\tilde{\phi}(\eta(r)) = \tilde{\phi}\left(\frac{r}{1}\right) = \phi(r) = s$$

Thus $\tilde{\phi}$ must be surjective.

Assume ϕ is injective. Suppose $\tilde{\phi} \left(\begin{smallmatrix} r \\ 1 \end{smallmatrix} \right) = 0$, then $\tilde{\phi} \left(\begin{smallmatrix} r \\ 1 \end{smallmatrix} \right) = \phi(r)\phi(1)^{-1} = \phi(r) = 0$. But since ϕ is injective, this implies that $r = 0$. Thus, $\tilde{\phi}$ is injective.

The two most important examples of localization are: i) when U is equal to the complement of a prime ideal and ii) when $U = \{1, f, f^2, f^3, \dots\}$ for some non-zero element $f \in R$. Given a prime ideal $\mathfrak{p} \subset R$ you showed in Question 2 that $U = R \setminus \mathfrak{p}$ is a multiplicatively closed set. The ring $(R \setminus \mathfrak{p})^{-1}R$ occurs so frequently that we give it its own notation and name. The *localization of R at $\mathfrak{p}$* is the ring:

$$R_{\mathfrak{p}} := (R \setminus \mathfrak{p})^{-1}R.$$

The guiding feature of $R_{\mathfrak{p}}$ is that all elements outside of $\mathfrak{p}$ become invertible, while the elements in $\mathfrak{p}$ remain non-units. As we shall see shortly, this implies that $R_{\mathfrak{p}}$ has a unique maximal ideal, namely,

$$\mathfrak{p} R_{\mathfrak{p}} := \left\{ \frac{r}{u} \mid r \in \mathfrak{p}, u \in R \setminus \mathfrak{p} \right\}.$$

A ring with a unique maximal ideal is called a *local ring*. We often denote a local ring by $(R, \mathfrak{m})$, where $\mathfrak{m}$ is its unique maximal ideal. Local rings are algebraic analogues of rings of germs near a point.

Turning towards the second important example, fix a non-zero element $f \in R$ and consider the set $U = \{1, f, f^2, f^3, \dots\}$. Note we should assume f is not a nilpotent element, since otherwise as we saw in Question 2, the localization $U^{-1}R$ will be the zero ring. The localization $U^{-1}R$ consists of fractions of the form $\frac{r}{f^n}$ with $r \in R$ and $n \geq 0$. Every such fraction can be rewritten as $r \cdot \frac{1}{f^n} = r \cdot \left(\frac{1}{f}\right)^n$, so every element of $U^{-1}R$ is a polynomial expression in $\frac{1}{f}$ with coefficients in R . In other words, $U^{-1}R$ is generated, as a ring, by the image of R and the single new inverse $\frac{1}{f}$. We therefore write:

$$R\left[\frac{1}{f}\right] := \left\{1, f, f^2, \dots\right\}^{-1}R = \left\{\frac{r}{f^n} \mid r \in R, n \geq 0\right\}.$$

We will see shortly that we can make the idea that $R[\frac{1}{f}]$ is “polynomials in $\frac{1}{f}$ with coefficients in R ” precise by showing $R[\frac{1}{f}]$ is isomorphic to $R[x]/\langle xf - 1 \rangle$. Intuitively this makes sense since quotienting by $\langle xf - 1 \rangle$ means setting $xf = 1$, i.e., let the formal variable x equal the inverse of f . (Note we saw something similar to this when we used the Rabinowitsch trick on the Nullstellensatz worksheet.)

Problem 9. Localization at a Prime Ideal.

Let R be a ring and $\mathfrak{p} \subset R$ be a prime ideal.

1. Prove that an element $\frac{r}{u} \in R_{\mathfrak{p}}$ is a unit if and only if $r \notin \mathfrak{p}$.
2. Deduce that the non-units of $R_{\mathfrak{p}}$ are precisely the elements

$$\mathfrak{p} R_{\mathfrak{p}} := \left\{ \frac{r}{u} \mid r \in \mathfrak{p}, u \in R \setminus \mathfrak{p} \right\}.$$

3. Prove that $\mathfrak{p} R_{\mathfrak{p}}$ is an ideal in $R_{\mathfrak{p}}$.
4. Show that $\mathfrak{p} R_{\mathfrak{p}}$ is a maximal ideal in $R_{\mathfrak{p}}$. Hint: Show that any ideal strictly containing $\mathfrak{p} R_{\mathfrak{p}}$ must contain a unit.
5. Show that $R_{\mathfrak{p}}$ does not have any other maximal ideal, i.e. $(R_{\mathfrak{p}}, \mathfrak{p} R_{\mathfrak{p}})$ is a local ring.
6. More generally, show that a ring R is a local ring if and only if the set of non-units in R is an ideal.
7. Let $p \in \mathbb{Z}$ be a prime number. Describe the units and unique maximal ideal of $\mathbb{Z}_{\langle p \rangle}$.
8. Let $R = \mathcal{C}[x]$ and $\mathfrak{m} = \langle x - a \rangle$ for $a \in \mathcal{C}$. Which rational functions in $\mathcal{C}(x)$ lie in $R_{\mathfrak{m}}$? Which of them are units?

Solution.

1. ($\implies$)

Assume $\frac{r}{u} \in R_{\mathfrak{p}}$ is a unit. Assume for contradiction that $r \in \mathfrak{p}$, and let $\frac{s}{v}$ be the inverse of $\frac{r}{u}$. Then $\exists t \in U$ such that

$$t(rs - vu) = 0 \implies trs = tvu \implies tvu \in \mathfrak{p}$$

But $t, v, u \notin \mathfrak{p}$ by assumption and since $\mathfrak{p}$ is prime $tvu \notin \mathfrak{p}$, thus a contradiction.

Hence $r \notin \mathfrak{p}$.

($\impliedby$)

Assume $r \notin \mathfrak{p}$. Then $r \in (R \setminus \mathfrak{p})$. That is there exists $\frac{u}{r} \in R_{\mathfrak{p}}$ and thus $\frac{r}{u}$ is a unit.

2. Then we can conclude that the non-units of $R_{\mathfrak{p}}$ are the precisely elements $r \in \mathfrak{p}$, which is the definition of $\mathfrak{p} R_{\mathfrak{p}}$.

3. To show that $\mathfrak{p} R_{\mathfrak{p}}$ is an ideal we need to show that it is closed under addition and closed under multiplication from $R_{\mathfrak{p}}$. Let $\frac{r}{u}, \frac{s}{u} \in \mathfrak{p} R_{\mathfrak{p}}$, and $\frac{a}{v} \in R_{\mathfrak{p}}$, with $u, v \in R \setminus \mathfrak{p}$.

- $\frac{r}{u} + \frac{s}{u} = \frac{r+s}{u} \in \mathfrak{p} R_{\mathfrak{p}}$ since $\mathfrak{p}$ is closed under addition.
- $\frac{r}{u} \cdot \frac{a}{v} = \frac{ra}{uv} \in \mathfrak{p} R_{\mathfrak{p}}$ since both $R \setminus \mathfrak{p}$ and $\mathfrak{p}$ are closed under multiplication.

4. Assume for contradiction that there exists a maximal ideal M such that $\mathfrak{p} R_{\mathfrak{p}} \subset M \subset R_{\mathfrak{p}}$. Then there exists some element $\frac{a}{u} \in M$ that is not in $\mathfrak{p} R_{\mathfrak{p}}$. But from part (2) we know that if $\frac{a}{u} \in R_{\mathfrak{p}}$ but not in $\mathfrak{p} R_{\mathfrak{p}}$ it is a unit. So there exists some element $\frac{b}{v} \in R_{\mathfrak{p}}$ such that $\frac{a}{u} \cdot \frac{b}{v} = 1 \in M$ contradicting our assumption that M is a proper ideal.

Hence, $\mathfrak{p} R_{\mathfrak{p}}$ is maximal.

5. From part (2) we know that all elements not in $\mathfrak{p} R_{\mathfrak{p}}$ are units. Then by the argument in (4) is any ideal $I \subset R_{\mathfrak{p}}$ such that $I \neq \mathfrak{p} R_{\mathfrak{p}}$ contains 1 and hence is the whole ring. Hence, $(R_{\mathfrak{p}}, \mathfrak{p} R_{\mathfrak{p}})$ is a local ring.

6. So far we have shown that if the set of non-units is an ideal then R is a local ring, so we only need to show the opposite direction.

Assume R is a local ring with maximal ideal M . Let J be the set of non-units in R . The containment $M \subseteq J$ is obvious (because if not, we have $1 \in M$). To show the reverse containment, let $j \in J$. Consider the ideal $\langle j \rangle$. Then there must exist some maximal ideal in R containing $\langle j \rangle$. Since R is a local ring we must have $\langle j \rangle \subseteq M$. Since j was an arbitrary element in J , this implies that $J \subseteq M$. Thus, the set of non-units in R is an ideal.

7. The maximal ideal (and the set of units) of $\mathbb{Z}_{\langle p \rangle}$ is $\mathfrak{p} \mathbb{Z}_{\langle p \rangle} := \left\{ \frac{q}{u} \mid q \in \langle p \rangle, u \in \mathbb{Z} \setminus \langle p \rangle \right\}$.

8. All rational functions in $\mathbb{C}(x)$ that don't have f in the denominator lie in $R_{\mathfrak{m}}$ where $f \in \langle x - a \rangle$. The ones that don't have f in the numerator are units.

Problem 10. Inverting an Element.

Let R be a ring and let $f \in R$ be a nonzero, non-nilpotent element. Recall the ring $R\left[\frac{1}{f}\right]$ is defined to be the localization of R at $U = \{1, f, f^2, f^3, \dots\}$.

1. Show that the R -algebra homomorphism $\phi : R[x] \rightarrow R\left[\frac{1}{f}\right]$ determined by $\phi(r) = \frac{r}{1}$ for $r \in R$ and $\phi(x) = \frac{1}{f}$ is well-defined.
2. Prove that ϕ is surjective by explicitly finding an element that maps to $\frac{r}{f^k}$ for $r \in R$ and $k \in \mathbb{Z}_{\geq 0}$.
3. Show that $\ker(\phi) = \langle fx - 1 \rangle$.
4. Conclude that $R\left[\frac{1}{f}\right]$ is isomorphic to $R[x] / \langle fx - 1 \rangle$.
5. Give a different argument proving $R\left[\frac{1}{f}\right]$ and $R[x] / \langle fx - 1 \rangle$ are isomorphic using the universal property of localizations.
6. What changes if f is nilpotent? Show that both $R\left[\frac{1}{f}\right]$ and $R[x] / \langle fx - 1 \rangle$ are the zero ring.
7. Let $f, g \in R$. Use the universal property to prove that

$$\left(R\left[\frac{1}{f}\right]\right)\left[\frac{1}{g}\right] \cong R\left[\frac{1}{fg}\right].$$

Solution.

1. **TODO:**

2. For any element $\frac{r}{f^k} \in R\left[\frac{1}{f}\right]$ we have

$$\phi(rx^k) = \phi(r) \cdot \phi(x^k) = \frac{r}{1} \cdot \frac{1}{f^k} = \frac{r}{f^k}$$

Thus ϕ is surjective.

3. $\ker(\phi) \subseteq \langle fx - 1 \rangle$

Assume $a(x) \in \ker(\phi)$ non-zero polynomial. Then $a(x) = q(x)(fx - 1) + r(x)$ for

some $q(x), r(x) \in R[x]$. We know that $\phi(a(x)) = 0 = \phi(q(x)(fx - 1) + r(x))$. Then,

$$\begin{aligned}\phi(q(x)(fx - 1) + r(x)) &= 0 \\ \phi(q(x))(\phi(fx) - \phi(1)) + \phi(r(x)) &= 0 \\ \phi(q(x))\left(\frac{f}{1} \cdot \frac{1}{f} - 1\right) + \phi(r(x)) &= 0 \\ 0 + \phi(r(x)) &= 0\end{aligned}$$

Thus $\phi(r(x)) = 0$. But since we were dividing by a degree one polynomial, $\deg(r(x)) = 0$, i.e. r is a constant, that forces $r = 0$. Hence, $a(x) \in \langle fx - 1 \rangle$.

$$\langle fx - 1 \rangle \subseteq \ker(\phi)$$

Let $a \in \langle fx - 1 \rangle$. Then $a = g(fx - 1)$ for some $g \in R$. Consider $\phi(a)$.

$$\phi(a) = \phi(g(fx - 1)) = \phi(gfx) - \phi(g) = \frac{gf}{f} - \frac{g}{1} = 0$$

Since a was an arbitrary element, we have $\langle fx - 1 \rangle \subseteq \ker(\phi)$.

4. Then by the first isomorphism theorem we have,

$$R \left[\frac{1}{f} \right] \cong R[x] / \ker(\phi) = R[x] / \langle fx - 1 \rangle$$

5. We have $U = \{1, f, f^2, \dots\}$, and $\tilde{\phi}$ exists because $\phi(U) \subset (R[x] / \langle xf - 1 \rangle)^\times$ since f maps to a unit in that ring ($x \mapsto \frac{1}{f}$).

$$\begin{array}{ccc} R & \xrightarrow{\eta} & U^{-1}R[x] = R \left[\frac{1}{f} \right] \\ \phi \downarrow & \nwarrow \tilde{\phi} & \\ R[x] / \langle xf - 1 \rangle & & \end{array} .$$

Now, since $\langle xf - 1 \rangle = \ker(\eta)$ by the universal property of quotients we have the map $\tilde{\eta}$,

$$\begin{array}{ccc} R & \xrightarrow{\pi} & R[x] / \langle fx - 1 \rangle \\ \eta \downarrow & \nwarrow \tilde{\eta} & \\ R \left[\frac{1}{f} \right] & & \end{array}$$

Consider the following compositions:

- $\tilde{\eta} \circ \tilde{\phi} \left(\frac{r}{f^k} \right) = \tilde{\eta}(rx^k) = \frac{r}{f^k}$
- $\tilde{\phi} \circ \tilde{\eta}(rx^k) = \tilde{\phi} \left(\frac{r}{f^k} \right) = rx^k.$

Thus $R \left[\frac{1}{f} \right] \cong R[x] / \langle fx - 1 \rangle.$

6. If f is nilpotent, from Question 4(1) we know that $R \left[\frac{1}{f} \right] = \{0\}.$ By part (4), this implies that $R[x] / \langle fx - 1 \rangle = \{0\}.$

7. First consider the composition $R \rightarrow R[1/f] \rightarrow (R[1/f])[1/g].$

In $(R[1/f])[1/g],$ both f and g are units, so fg is a unit. Then by the universal property of localization (on $R[1/fg],$), there exists a unique map $\tilde{\phi} : R[1/fg] \rightarrow (R[1/f])[1/g].$

Now, note that in $R[1/fg], fg$ is a unit, so both f and g are units

$$f \cdot \frac{g}{fg} = 1, \quad g \cdot \frac{f}{fg} = 1$$

Then by the universal property of localization (on $(R[1/f])[1/g],$), there exists a unique map $\tilde{\psi} : (R[1/f])[1/g] \rightarrow R[1/fg].$ Both compositions $\tilde{\phi} \circ \tilde{\psi}$ and $\tilde{\psi} \circ \tilde{\phi}$ are the identity by uniqueness in the universal property, hence:

$$\left(R \left[\frac{1}{f} \right] \right) \left[\frac{1}{g} \right] \cong R \left[\frac{1}{fg} \right]$$

Given any ring homomorphism $\phi : R \rightarrow S$, there is a natural interplay between ideals of R and ideals of S . If $I \subset R$ is an ideal, its *extension* is the ideal of S generated by the image of I under ϕ :

$$\phi(I) \cdot S := \langle \phi(I) \rangle = \left\{ \sum_{i=1}^n s_i \phi(r_i) \mid r_i \in I, s_i \in S, n \geq 0 \right\}.$$

Note we must take the ideal generated by $\phi(I)$, not just the image of I itself, since as we previously saw $\phi(I)$ may not be an ideal of S .

Conversely, if $J \subseteq S$ is an ideal, its *contraction* is the preimage

$$\phi^{-1}(J) = \{r \in R \mid \phi(r) \in J\},$$

which is always an ideal of R . Note there are numerous different notations for extension and contraction of ideals: some (namely Atiyah–MacDonald) write I^e and J^c for the extension and contraction respectively, others write IS and $J \cap R$ (as a nod to what occurs when ϕ is an injection). We generally avoid these notations to emphasize that contraction and extension are dependent on the homomorphism ϕ .

Problem 11. Extension and Contraction of Ideals.

Let $\phi : R \rightarrow S$ be a homomorphism of rings. Let $I \subset R$ and $J \subset S$ be ideals.

1. Prove that $\phi^{-1}(J)$ is an ideal of R .
2. Give an explicit example showing that $\phi(I) \subset S$ need not be an ideal.
3. Prove that $I \subset \phi^{-1}(\phi(I)S)$.
4. Let $\phi : \mathbb{Z} \rightarrow \mathbb{Z}/\langle 6 \rangle$ be the natural quotient map and let $I = \langle 0 \rangle \subset \mathbb{Z}$. Explicitly describe $J = \phi(I)(\mathbb{Z}/\langle 6 \rangle)$ and $\phi^{-1}(J)$. Show that $I \subsetneq \phi^{-1}(J)$. Thus, the inclusion in the previous part can be strict.
5. Prove that $\phi(\phi^{-1}(J))S \subset J$.
6. Let $\phi : \mathbb{Z} \rightarrow \mathbb{Z}[x]$ be the inclusion map and let $J = \langle x \rangle \subset \mathbb{Z}[x]$. Explicitly describe $I = \phi^{-1}(J)$ and $\phi(I)\mathbb{Z}[x]$. Show that $\phi(I)\mathbb{Z}[x] \subsetneq J$. Thus, the inclusion in the previous part can be strict.
7. Prove that if $J \subset S$ is prime then the contraction $\phi^{-1}(J) \subset R$ is prime.
8. Give an example showing that if $J \subset S$ is maximal the contraction $\phi^{-1}(J) \subset R$ need not be maximal.
9. Give an example showing that if $I \subset R$ is prime then the extension $\phi(I)S \subset S$ need not be prime.
10. Let $\phi : R \hookrightarrow S$ be an injective ring homomorphism. Prove that

$$\phi^{-1}(J) = J \cap R \quad \text{and} \quad \phi(I)S = IS.$$

Solution.

1. We showed this in Worksheet 1.1 Question 4.1.
2. Let $R = \mathbb{Z}/4\mathbb{Z}$, $S = \mathbb{Z}$, $I = \langle 2 \rangle$ and ϕ be the inclusion map. Then we can see that $\phi(I) = \{0, 2\} \in S$ is not an ideal.
3. Let $a \in I$. Note that $\phi(a) \in \phi(I) \subseteq \phi(I)S$. Then we have,

$$\phi^{-1}(\phi(a)) = a \in \phi^{-1}(\phi(I)) \subseteq \phi^{-1}(\phi(I)S)$$

Since a was an arbitrary element, this implies that $I \subseteq \phi^{-1}(\phi(I)S)$.

4. We have $J = \phi(I)(\mathbb{Z}/\langle 6 \rangle) = \{\bar{0}\}$, and $\phi^{-1}(J) = \langle 6 \rangle$. Thus, there is a strict inclusion $I = \langle 0 \rangle \subsetneq \langle 6 \rangle = \phi^{-1}(J)$.

5. Let $I = \phi^{-1}(J)$, and let $a \in \phi(I)S$. Note that $s = \sum_i \phi(r_i)s_i$ where $r_i \in I$ and $s_i \in S$. Then $\phi(r_i) \in J$ and by definition of ideals, we have $s \in J$.
6. We have $I = \phi^{-1}J = \{r \in R \mid \phi(r) \in \langle x \rangle\} = \{0\}$, and $\phi(I)\mathbb{Z}[x] = \{0\}\mathbb{Z}[x] = \{0\}$. Thus, there is a strict inclusion $\phi(I)\mathbb{Z}[x] \subsetneq J$.
7. Let $ab \in \phi^{-1}(J)$. Then J is prime either $\phi(a) \in J$ or $\phi(b) \in J$. That is, either $a \in \phi^{-1}(J)$ or $b \in \phi^{-1}(J)$. Thus, $\phi^{-1}(J)$ is prime.
8. Let $R = \mathbb{Z}$, $S = \mathbb{Q}$, $J = \langle 0 \rangle$, and ϕ be the inclusion map. Then $\phi^{-1}(J) = \langle 0 \rangle$ which is not maximal in $\mathbb{Z}$.
9. Let $R = \mathbb{Z}$, $S = \mathbb{Z}/4\mathbb{Z}$, $I = \langle 0 \rangle$, and ϕ be the quotient map. Then $\phi(I)S = \langle 0 \rangle$ is not prime in $\mathbb{Z}/4\mathbb{Z}$ since $2 \times 2 = 0$.
10. Since ϕ is injective, R is isomorphic to a subring of S . Then $\phi^{-1}(J) = \{r \in R \mid r \in J\} = J \cap R$. Again since ϕ is injective $\phi(I) = I$ as a subset of S . So, $\phi(I)S = IS$.

As we saw in the previous exercise, in general, extension and contraction are far from being inverse operations: $\phi^{-1}(\phi(I)S)$ may strictly contain I , and $\phi(\phi^{-1}(J))S$ may be strictly smaller than J . In the case of localization, extension and contraction along the canonical localization map $\eta : R \rightarrow U^{-1}R$ sending $r \mapsto r/1$ have very nice forms. For an ideal $I \subset R$ the extension along η is

$$\eta(I) (U^{-1}R) = U^{-1}I := \left\{ \frac{r}{u} \mid r \in I, u \in U \right\} \subset U^{-1}R.$$

For an ideal $J \subset U^{-1}R$ the contraction along η is:

$$\eta^{-1}(J) = \left\{ r \in R \mid \frac{r}{1} \in J \right\}.$$

Although the extension–contraction correspondence remains lossy for arbitrary ideals, it becomes perfectly behaved for prime ideals. If $\mathfrak{q} \subset U^{-1}R$ is a proper ideal, then no element of U can lie in its contraction. (This is because $\frac{u}{1}$ is a unit in $U^{-1}R$ for every $u \in U$.) Thus every prime ideal of $U^{-1}R$ contracts to a prime ideal of R which is disjoint from U . Conversely, if $\mathfrak{p} \subset R$ is a prime ideal disjoint from U , then its extension

$$U^{-1}\mathfrak{p} = \left\{ \frac{r}{u} \mid r \in \mathfrak{p}, u \in U \right\}$$

is a prime ideal of $U^{-1}R$. In fact, extension and contraction give an order-preserving bijection

$$\left\{ \text{prime ideals of } U^{-1}R \right\} \longleftrightarrow \left\{ \mathfrak{p} \in \text{Spec}(R) \mid \mathfrak{p} \cap U = \emptyset \right\}.$$

Under this correspondence, a prime ideal $\mathfrak{q} \subset U^{-1}R$ is sent to its contraction $\eta^{-1}(\mathfrak{q})$, and a prime ideal $\mathfrak{p} \subset R$ disjoint from U is sent to its extension $U^{-1}\mathfrak{p}$.

This is especially important in the two basic examples above. If $U = R \setminus \mathfrak{p}$, then the prime ideals of $R_{\mathfrak{p}}$ correspond exactly to the prime ideals of R contained in $\mathfrak{p}$. If $U = \{1, f, f^2, \dots\}$, then the prime ideals of $R[\frac{1}{f}]$ correspond exactly to the prime ideals of R which do not contain f .

Problem 12. Ideals and Localization.

Let R be a ring, let $U \subset R$ be a multiplicatively closed set, and let $\eta : R \rightarrow U^{-1}R$ be the localization map.

1. Let $I \subset R$ be an ideal. Prove that $U^{-1}I := \{\frac{r}{u} \mid r \in I, u \in U\}$ is an ideal of $U^{-1}R$.
2. Prove that $U^{-1}I$ is the extension of I along the localization map $\eta : R \rightarrow U^{-1}R$.
3. Prove that $U^{-1}I = U^{-1}R$ if and only if $I \cap U \neq \emptyset$.
4. Prove that $\frac{r}{u} \in U^{-1}I$ if and only if $tr \in I$ for some $t \in U$.
5. Deduce that the contraction of $U^{-1}I$ back to R is

$$\eta^{-1}(U^{-1}I) = \{r \in R \mid tr \in I \text{ for some } t \in U\}.$$

6. Let $R = \mathbb{Z}$ and $U = \{1, 2, 4, 8, \dots\}$. If $I = \langle 6 \rangle$, compute $U^{-1}I \subset U^{-1}\mathbb{Z}$ and its contraction back to $\mathbb{Z}$. Show that $I \subsetneq \eta^{-1}(U^{-1}I)$.
7. Let $R = \mathbb{Z}$ and $U = \mathbb{Z} \setminus \langle 2 \rangle$. If $I = \langle 6 \rangle$, compute $U^{-1}I \subset \mathbb{Z}_{\langle 2 \rangle}$ and its contraction back to $\mathbb{Z}$.
8. Let $R = \mathbb{K}[x, y]$ and $U = \{1, x, x^2, \dots\}$. For each of the ideals $\langle x \rangle$, $\langle y \rangle$, and $\langle x, y \rangle$, describe its extension to $R[\frac{1}{x}]$. Which of these ideals become the whole ring after localization?

Solution.

1. $U^{-1}I$ is a subset of $U^{-1}R$ so we only need to show closure under addition and closure under multiplication from $U^{-1}R$ to prove that it is an ideal. Let $\frac{r}{u}, \frac{s}{v} \in U^{-1}I$ and $\frac{a}{w} \in U^{-1}R$.

- $\frac{r}{u} + \frac{s}{v} = \frac{rv+su}{uv} \in U^{-1}I$ since I is an ideal.
- $\frac{r}{u} \cdot \frac{a}{w} = \frac{ra}{uw} \in U^{-1}I$ since I is an ideal.

Hence, $U^{-1}I$ is an ideal.

2. $\eta(I)U^{-1}R = \left\{ \sum_{i=1}^n \frac{r_i}{u_i} \cdot \frac{a_i}{v_i} \mid r_i \in R, u_i, v_i \in U, a_i \in I, n \geq 0 \right\}$. But since I is an ideal and U is multiplicatively closed we have $r_i a_i = s_i \in I$ and $u_i v_i = w_i \in U$ for all i , giving us,

$$\eta(I)U^{-1}R = \left\{ \sum_{i=1}^n \frac{s_i}{w_i} \mid w_i \in U, s_i \in I, n \geq 0 \right\}$$

which is the definition of $U^{-1}I$.

3. ($\implies$)

Assume $U^{-1}I = U^{-1}R$. Then $\eta(I)$ must have a unit, i.e. $I \cap U \neq \emptyset$.

($\impliedby$)

Assume $I \cap U \neq \emptyset$. Let $a \in I \cap U$. $\eta(a)$ is a unit in $U^{-1}R$. That is $\frac{a}{a} = \frac{1}{1} \in U^{-1}I$ and thus $U^{-1}I = U^{-1}R$.

4. ($\implies$)

Assume $\frac{r}{u} \in U^{-1}I$. Then, $\frac{r \cdot u}{u \cdot 1} = \frac{r}{1} \in U^{-1}I$, since this is the image of $\eta(r)$, we must have $r \in I$, with $t = 1$.

($\impliedby$)

Assume $tr \in I$ for some $t \in U$. Since U is multiplicatively closed $ut \in U$, and we have $\frac{tr}{ut} = \frac{r}{u} \in U^{-1}R$.

5. Therefore we can conclude that the contraction of $U^{-1}I$ is

$$\eta^{-1}(U^{-1}I) = \{r \in R \mid tr \in I \text{ for some } t \in U\}.$$

6. $U^{-1}I = \langle 6 \rangle U^{-1}\mathbb{Z} = \left\{ \frac{3n}{2^k} \mid n \in \mathbb{Z}, k \geq 0 \right\}$

And $\eta^{-1}(U^{-1}I) = \{r \in \mathbb{Z} \mid \exists 2^k \in U \text{ such that } 2^k r \in \langle 6 \rangle\}$. Then $2^k r = 6n$ for some n . Then, we have $r = \frac{3n}{2^{k-1}}$ and thus $3 \mid r \implies r \in \langle 3 \rangle = \eta^{-1}(U^{-1}I)$.

Now note that $3 \in \langle 3 \rangle$ but not in $\langle 6 \rangle$.

7. $U^{-1}I = \langle 6 \rangle U^{-1}\mathbb{Z} = \left\{ \frac{6n}{k} \mid n \in \mathbb{Z}, k \text{ odd} \right\} = \langle 2 \rangle U^{-1}\mathbb{Z}$ since 3 is invertible.

And $\eta^{-1}(U^{-1}I) = \{r \in \mathbb{Z} \mid \exists k \in U \text{ such that } kr \in \langle 6 \rangle\}$. This means that $2 \mid r$, then we have $\langle 2 \rangle = \eta^{-1}(U^{-1}I)$.

8. • $\langle x \rangle \cap U \neq \emptyset$.

Then by part (3) $\langle x \rangle (U^{-1}R) = U^{-1}R = R \left[\frac{1}{x} \right]$

• $\langle y \rangle \cap U = \emptyset$.

$$U^{-1}\langle y \rangle = \left\{ \frac{r}{u} \mid r \in \langle y \rangle, u \in U \right\} = \left\{ \frac{ay}{x^n} \mid ay \in \langle y \rangle, n \geq 0 \right\} = y \cdot R \left[\frac{1}{x} \right].$$

• $\langle x, y \rangle \cap U \neq \emptyset$.

$$\text{Then by part (3) } \langle x, y \rangle (U^{-1}R) = U^{-1}R = R \left[\frac{1}{x} \right]$$

Problem 13. Prime Ideal Correspondence for Localization.

Let R be a ring, let $U \subset R$ be a multiplicatively closed set, and let $\eta : R \rightarrow U^{-1}R$ be the localization map.

1. Let $\mathfrak{q} \subset U^{-1}R$ be a prime ideal. Prove that $\eta^{-1}(\mathfrak{q})$ is a prime ideal of R and that $\eta^{-1}(\mathfrak{q}) \cap U = \emptyset$.
2. Let $\mathfrak{p} \subset R$ be a prime ideal such that $\mathfrak{p} \cap U = \emptyset$. Prove that $U^{-1}\mathfrak{p}$ is a proper prime ideal of $U^{-1}R$. Hint: Use the criterion from the previous exercise for when a fraction belongs to $U^{-1}\mathfrak{p}$.
3. Let $\mathfrak{p} \subset R$ be a prime ideal such that $\mathfrak{p} \cap U = \emptyset$. Prove that the contraction of $U^{-1}\mathfrak{p}$ is $\mathfrak{p}$.
4. Let $\mathfrak{q} \subset U^{-1}R$ be a prime ideal. Prove that the extension of its contraction is again $\mathfrak{q}$, i.e.

$$U^{-1}\eta^{-1}(\mathfrak{q}) = \mathfrak{q}.$$

Hint: If $\frac{r}{u} \in U^{-1}R$, then $\frac{r}{u} \in \mathfrak{q}$ if and only if $\frac{r}{1} \in \mathfrak{q}$, since $\frac{u}{1}$ is a unit.

5. Conclude that extension and contraction give an order-preserving bijection

$$\text{Spec}(U^{-1}R) \longleftrightarrow \{\mathfrak{p} \in \text{Spec}(R) \mid \mathfrak{p} \cap U = \emptyset\}.$$

6. Specialize to the case $U = R \setminus \mathfrak{p}$. Prove that the prime ideals of $R_{\mathfrak{p}}$ correspond to the prime ideals of R contained in $\mathfrak{p}$.
7. Specialize to the case $U = \{1, f, f^2, \dots\}$. Prove that the prime ideals of $R[\frac{1}{f}]$ correspond to the prime ideals of R which do not contain f .
8. Describe the prime ideals of $\mathbb{Z}_{\langle p \rangle}$ for a prime number p .

Solution.

1. We already showed that pre-image of a prime ideal is prime. So we only need to show that $\eta^{-1}(\mathfrak{q}) \cap U = \emptyset$. Suppose not, then the image $\mathfrak{q}$ has a unit, contradicting our assumption that $\mathfrak{q}$ is a prime ideal. Thus, $\mathfrak{q} \cap U$ must be trivial.
2. Let $\mathfrak{p} \cap U = \emptyset$ for $\mathfrak{p}$ prime, then we have,

$$U^{-1}\mathfrak{p} = \left\{ \frac{r}{u} \mid r \in \mathfrak{p}, u \in U \right\}$$

Assume $\frac{r \cdot s}{u \cdot v} \in U^{-1}\mathfrak{p}$, then by definition either $r \in \mathfrak{p}$ or $s \in \mathfrak{p}$. That is either $\frac{r}{u} \in U^{-1}\mathfrak{p}$ or $\frac{s}{v} \in U^{-1}\mathfrak{p}$. Thus, $U^{-1}\mathfrak{p}$ is a prime ideal in U .

3. Let $\mathfrak{p} \cap U = \emptyset$ for $\mathfrak{p}$ prime. Consider $\eta^{-1}(U^{-1}\mathfrak{p})$. From Problem 12.5 we know that this is the set,

$$\eta^{-1}(U^{-1}\mathfrak{p}) = \{r \in R \mid tr \in \mathfrak{p} \text{ for some } t \in U\}$$

But since $\mathfrak{p} \cap U = \emptyset$ by assumption and $\mathfrak{p}$ is prime this is just the set,

$$\eta^{-1}(U^{-1}\mathfrak{p}) = \{r \in R \mid r \in \mathfrak{p}\} = \mathfrak{p}$$

4. $\mathfrak{q} \subseteq U^{-1}\eta^{-1}(\mathfrak{q})$

Let $\frac{r}{u} \in \mathfrak{q}$. Then $\frac{r}{1} \in \mathfrak{q} \implies r \in \eta^{-1}(\mathfrak{q})$. Hence, $\frac{r}{u} \in U^{-1}\eta^{-1}(\mathfrak{q})$.

$$\mathfrak{q} \supseteq U^{-1}\eta^{-1}(\mathfrak{q})$$

Let $\frac{r}{u} \in U^{-1}\eta^{-1}(\mathfrak{q})$. Then $r \in \eta^{-1}(\mathfrak{q}) \implies \frac{r}{1} \in \mathfrak{q}$.

5. Thus we can conclude that prime ideals in $U^{-1}R$ are in direct correspondce with prime ideals in R that have trivial intersection with U . Giving us an order preserving bijection:

$$\text{Spec}(U^{-1}R) \longleftrightarrow \{\mathfrak{p} \in \text{Spec}(R) \mid \mathfrak{p} \cap U = \emptyset\}.$$

6. In the case $U = R \setminus \mathfrak{p}$, the only ideals that have non-trivial intersection with U are inside $\mathfrak{p}$. Thus our bijection becomes,

$$\text{Spec}(R_{\mathfrak{p}}) \longleftrightarrow \{\mathfrak{q} \in \text{Spec}(R) \mid \mathfrak{q} \subseteq \mathfrak{p}\}.$$

7. In the case $U = \{1, f, f^2, \dots\}$, the ideals that have non-trivial intersection with U are ideals that do not contain f . Thus our bijection becomes,

$$\text{Spec}\left(R \left[\frac{1}{f}\right]\right) \longleftrightarrow \{\mathfrak{p} \in \text{Spec}(R) \mid f \notin \mathfrak{p}\}.$$

8. From (6) $\text{Spec}(\mathbb{Z}_{\langle p \rangle}) = \{[0], [\mathfrak{p}]\}$.

Localization is not limited to rings: we can also localize modules in essentially the same way. Let R be a ring and let M be an R -module. If $U \subset R$ is a multiplicatively closed subset, we want to build a new module $U^{-1}M$ by formally dividing elements of M by the elements of U . The elements of $U^{-1}M$ are fractions $\frac{m}{u}$, where $m \in M$ and $u \in U$. So the numerators come from the module M , while the denominators come from the set U which lives in the ring. Of course, we must introduce a criterion of when two fractions $\frac{m_1}{u_1}$ and $\frac{m_2}{u_2}$ are equal. Just as when localizing a ring we say

$$\frac{m_1}{u_1} = \frac{m_2}{u_2} \quad \text{in } U^{-1}M \quad \Longleftrightarrow \quad t(u_2m_1 - u_1m_2) = 0 \quad \text{in } M \text{ for some } t \in U$$

The factor of $t \in U$ is needed in order to ensure this is a transitive relation. Define addition on $U^{-1}M$ by

$$\frac{m_1}{u_1} + \frac{m_2}{u_2} = \frac{m_1u_2 + m_2u_1}{u_1u_2}$$

which makes $U^{-1}M$ into an abelian group where the identity is $0/1$. The abelian group $U^{-1}M$ has the structure of both an R -module and a $U^{-1}R$ -module. The R -module structure and the $U^{-1}R$ -module structure are given by

$$r \left(\frac{m}{u} \right) = \frac{rm}{u} \quad \text{and} \quad \left(\frac{r}{u_1} \right) \left(\frac{m}{u_2} \right) = \frac{rm}{u_1u_2}$$

respectively. These module structures are compatible: the R -module structure on $U^{-1}M$ is the one obtained from the $U^{-1}R$ -module structure by restriction of scalars along the localization map $\eta : R \rightarrow U^{-1}R$.

In general, if $\phi : R \rightarrow S$ is any ring homomorphism and N is an S -module, then N becomes an R -module by defining $rn := \phi(r)n$ for $r \in R$ and $n \in N$. This construction is called *restriction of scalars*. One checks immediately that this gives N a well-defined R -module structure, since ϕ preserves addition, multiplication, and the identity. In our situation, the localization map $\eta : R \rightarrow U^{-1}R$ is defined by $\eta(r) = r/1$. Since $U^{-1}M$ is a $U^{-1}R$ -module, restriction of scalars along η gives $U^{-1}M$ an R -module structure via

$$r \cdot \frac{m}{u} = \eta(r) \cdot \frac{m}{u} = \frac{r}{1} \cdot \frac{m}{u} = \frac{rm}{u},$$

which is exactly the R -module structure we defined above. So there is really only one module structure at play: the $U^{-1}R$ -module structure is the fundamental one, as the R -module structure is inherited from it.

Just as the localization of a ring is characterized by a universal property, so is the localization of a module. There is a natural R -module homomorphism $\eta_M : M \rightarrow U^{-1}M$ given by $m \mapsto \frac{m}{1}$, which we call the *localization map*. Note that the target $U^{-1}M$ carries a $U^{-1}R$ -module structure, and in particular every element of U acts invertibly on $U^{-1}M$: for

any $u \in U$, multiplication by $u/1 \in U^{-1}R$ is an isomorphism with inverse given by multiplication by $1/u$. The pair $U^{-1}M$ and η_M satisfies the following universal property: Let N be any $U^{-1}R$ -module, viewed as an R -module by restriction of scalars along $\eta : R \rightarrow U^{-1}R$. Then for every R -module homomorphism $\phi : M \rightarrow N$, there exists a unique $U^{-1}R$ -module homomorphism $\tilde{\phi} : U^{-1}M \rightarrow N$ such that the following diagram commutes:

$$\begin{array}{ccc} M & \xrightarrow{\eta_M} & U^{-1}M \\ \phi \downarrow & \nwarrow \tilde{\phi} & \\ N & & \end{array} .$$

Problem 14. Constructing $U^{-1}M$.

Let R be a ring, $U \subset R$ a multiplicatively closed set, and M be an R -module. Define a relation on $M \times U$ by $(m_1, u_1) \sim (m_2, u_2)$ if and only if $t(m_1u_2 - m_2u_1) = 0$ in M for some $t \in U$.

1. Prove that $\sim$ is an equivalence relation on $M \times U$.
2. Let $\frac{m}{u}$ denote the equivalence class of a pair (m, u) under the equivalence relation $\sim$ defined in the previous part. Prove that the operation:

$$\frac{m_1}{u_1} + \frac{m_2}{u_2} = \frac{m_1u_2 + m_2u_1}{u_1u_2}$$

is well-defined. Show that this makes $U^{-1}M$ an abelian group.

3. Show that the operations

$$r \left(\frac{m}{u} \right) = \frac{rm}{u} \quad \text{and} \quad \left(\frac{r}{u_1} \right) \left(\frac{m}{u_2} \right) = \frac{rm}{u_1u_2}$$

are well-defined. Deduce that $U^{-1}M$ has the structure of both an R -module and a $U^{-1}R$ -module.

4. Let $\eta_M : M \rightarrow U^{-1}M$ be the function $m \mapsto \frac{m}{1}$. Show that η_M is an R -module homomorphism.
5. Prove that $\eta_M(m) = 0$ if and only if $um = 0$ for some $u \in U$. Deduce that η_M is injective if and only if no element of U acts as a zero-divisor on M .

Solution.

1. Let $(m_1, u_1) \sim (m_2, u_2) \sim (m_3, u_3) \in M \times U$.

(a) **Reflexivity:** $(m_1, u_1) \sim (m_1, u_1)$ since $1(m_1u_1 - m_1u_1) = 0$.

(b) **Symmetry:** If $(m_1, u_1) \sim (m_2, u_2)$ then there exists a $t \in U$ such that

$$t(m_1u_2 - m_2u_1) = 0 \implies tm_1u_2 = tm_2u_1 \implies t(m_2u_1 - m_1u_2) = 0$$

Hence, $(m_2, u_2) \sim (m_1, u_1)$.

(c) **Transitivity:** If $(m_1, u_1) \sim (m_2, u_2)$ and $(m_2, u_2) \sim (m_3, u_3)$ then there exist s, t such that

$$s(m_1u_2 - m_2u_1) = 0, \quad t(m_2u_3 - m_3u_2) = 0$$

Note that

$$stu_3(m_1u_2 - m_2u_1) = 0, \quad stu_1(m_2u_3 - m_3u_2) = 0$$

Adding,

$$0 = st(m_1u_2u_3 - m_2u_1u_3 + m_2u_3u_1 - m_3u_2u_1) = stu_2(m_1u_3 - m_3u_1)$$

Since $stu_2 \in U$ we have $(m_1, u_1) \sim (m_3, u_3)$.

Thus, $\sim$ is an equivalence relation on $R \times U$.

2. Well-definedness:

Let $(m_1, u_1) \sim (m'_1, u'_1)$ and $(m_2, u_2) \sim (m'_2, u'_2)$. Then there exist s and t such that,

$$s(m_1u'_1 - m'_1u_1) = 0, \quad t(m_2u'_2 - m'_2u_2) = 0$$

We show that $\frac{m_1u_2 + m_2u_1}{u_1u_2} \sim \frac{m'_1u'_2 + m'_2u'_1}{u'_1u'_2}$:

$$\begin{aligned} & st((m_1u_2 + m_2u_1)(u'_1u'_2) - (m'_1u'_2 + m'_2u'_1)(u_1u_2)) \\ &= st(m_1u_2u'_1u'_2 + m_2u_1u'_1u'_2 - m'_1u'_2u_1u_2 - m'_2u'_1u_1u_2) \\ &= st(u_2u'_2(m_1u'_1 - m'_1u_1) + u_1u'_1(m_2u'_2 - m'_2u_2)) \\ &= tu_2u'_2 \cdot s(m_1u'_1 - m'_1u_1) + su_1u'_1 \cdot t(m_2u'_2 - m'_2u_2) \\ &= tu_2u'_2 \cdot 0 + su_1u'_1 \cdot 0 = 0 \end{aligned}$$

Identity:

$$\frac{m}{u} + \frac{0}{1} = \frac{m+0}{u} = \frac{m}{u}$$

And note that associativity, commutativity, and distributivity are inherited from R .

Thus $U^{-1}M$ is an abelian group.

3. Let $(m, u) \sim (m', u')$. Then there exist $s \in U$ such that

$$s(mu' - m'u) = 0$$

Then,

$$s(rmu' - rm'u) = sr(mu' - m'u) = 0$$

Thus $U^{-1}M$ has the structure of a R -module.

Let $(m, u) \sim (m', u')$. Then there exist $s \in U$ such that

$$s(mu' - m'u) = 0$$

Then,

$$s(rmu' - rm'u) = sr(mu' - m'u) = 0$$

Thus $U^{-1}M$ has the structure of a R -module.

Let $m_1, m_2 \in M$, $u_1, u_2 \in U$, $r, r' \in R$.

First assume $\frac{m}{u_2} \sim \frac{m'}{u_2'}$, so $\exists t \in U$ with $t(mu_2' - m'u_2) = 0$. We have:

$$t(rmu_1u_2' - rm'u_1u_2) = u_1t(rmu_2' - rm'u_2) = ru_1t(mu_2' - m'u_2) = 0$$

Now assume $\frac{r}{u_1} \sim \frac{r'}{u_1'}$, so $\exists s \in U$ with $s(ru_1' - r'u_1) = 0$. We have:

$$s(rmu_1'u_2 - r'mu_1u_2) = mu_2 \cdot s(ru_1' - r'u_1) = 0$$

Thus $U^{-1}M$ has the structure of a $U^{-1}R$ -module.

4. Let $m, n \in M$ and $r \in R$.

- $\eta_M(mn) = \frac{mn}{1} = \frac{m}{1} \cdot \frac{n}{1} = \eta_M(m)\eta_M(n)$
- $\eta_M(m+n) = \frac{m+n}{1} = \frac{m}{1} + \frac{n}{1} = \eta_M(m) + \eta_M(n)$
- $r\eta_M(m) = r \cdot \frac{m}{1} = \frac{rm}{1} = \eta_M(rm)$

Thus η_M is a R -module homomorphism.

5. ($\implies$)

Assume $\eta_M(m) = 0$. Then we have,

$$\eta_M(m) = \frac{m}{1} \sim \frac{0}{1} \implies \exists u \in U \text{ such that } u(m - 0) = 0 \implies um = 0$$

($\impliedby$)

Assume $um = 0$ for some $u \in U$. Then $\eta_M(m) = \frac{m}{1}$, we want to show that $\eta_M(um) \sim \frac{0}{1}$. But we have,

$$u(m - 0) = um - 0 = 0$$

Thus $\eta_M(m) = 0$.

Thus, η_M is injective if and only if no element of U acts as a zero-divisor on M .

Problem 15. Examples of Localizing Modules.

Discuss ways to explicitly understand $U^{-1}M$ in each of the following cases. Describe the natural map $M \rightarrow U^{-1}M$ and determine whether it is injective.

1. $R = \mathbb{Z}$, $U = \mathbb{Z} \setminus \{0\}$, and $M = \mathbb{Z}$.
2. $R = \mathbb{Z}$, $U = \{1, 2, 4, 8, \dots\}$, and $M = \mathbb{Z}$.
3. $R = \mathbb{Z}$, $U = \{1, 2, 4, 8, \dots\}$, and $M = \mathbb{Z}/\langle 6 \rangle$.
4. $R = \mathbb{Z}$, $U = \mathbb{Z} \setminus \langle 2 \rangle$, and $M = \mathbb{Z}/\langle 6 \rangle$.
5. $R = \mathbb{K}[x]$, $U = \{1, x, x^2, \dots\}$, and $M = R/\langle x^3 \rangle$.
6. $R = \mathbb{K}[x]$, $U = R \setminus \langle x \rangle$, and $M = R/\langle x^3 \rangle$.
7. $R = \mathbb{K}[x, y]$, $U = \{1, y, y^2, \dots\}$, and $M = R/\langle xy \rangle$.
8. R is an arbitrary ring, M is an R -module, and U contains an element u such that $uM = 0$.

Solution.

1. $U^{-1}M = U^{-1}R = \mathbb{Q}$, and η_M is injective.
2. $U^{-1}M = U^{-1}R = \left\{ \frac{r}{2^k} \mid r \in \mathbb{Z}, k \geq 0 \right\}$, and η_M is injective.
3. $U^{-1}M \cong \mathbb{Z}/2\mathbb{Z}$, and η_M not injective since $2 \in U$.
4. $U^{-1}M \cong \mathbb{Z}/3\mathbb{Z}$, and η_M not injective since $3 \in U$.
5. $U^{-1}M = \{0\}$, and η_M not injective since $x, x^2, x^3 \in U$.
6. $U^{-1}M = M$, and η_M is injective.
7. $U^{-1}M = \mathbb{K}[x, y]$, and η_M **TODO:**
8. $U^{-1}M = \{0\}$, and η_M not injective.

Problem 16. Universal Property of $U^{-1}M$.

Let R be a ring, let $U \subset R$ be a multiplicatively closed set, and let M be an R -module. Let N be a $U^{-1}R$ -module, which we view as an R -module via the localization map $\eta : R \rightarrow U^{-1}R$. Let $\psi : M \rightarrow N$ be an R -module homomorphism.

1. Define a function $\tilde{\psi} : U^{-1}M \rightarrow N$ by $\tilde{\psi}(\frac{m}{u}) = \frac{1}{u}\psi(m)$. Prove that $\tilde{\psi}$ is well-defined.
2. Prove that $\tilde{\psi}$ is a homomorphism of $U^{-1}R$ -modules.
3. Show that $\tilde{\psi}$ makes the following diagram commute:

$$\begin{array}{ccc} M & \xrightarrow{\eta_M} & U^{-1}M \\ \psi \downarrow & \swarrow \tilde{\psi} & \\ N & & \end{array}$$

4. Prove that $\tilde{\psi}$ is the unique $U^{-1}R$ -module homomorphism making the diagram commute.
5. Deduce that $U^{-1}M$ is unique up to unique isomorphism as a $U^{-1}R$ -module satisfying this universal property.

Solution.

1. Let $\frac{m}{u} \sim \frac{n}{s}$. Then there exists a $t \in U$ such that $t(ms - nu) = 0$. Then we have,

$$\psi(t(ms - nu)) = t(\psi(m)s - \psi(n)u) = 0$$

Multiplying by $\frac{1}{tsu}$ gives us,

$$\frac{1}{u}\psi(m) - \frac{1}{s}\psi(n) = 0 \implies \tilde{\psi}(m) = \tilde{\psi}(n)$$

2. Let $\frac{m}{u}, \frac{n}{u} \in U^{-1}M$ and $\frac{r}{t} \in U^{-1}R$.

- $\tilde{\psi}(\frac{mn}{su}) = \frac{1}{su}\psi(mn) = \frac{1}{u}\psi(m)\frac{1}{s}\psi(n)$
- $\tilde{\psi}(\frac{ms+nu}{su}) = \frac{1}{su}\psi(ms+nu) = \frac{1}{u}\psi(m) + \frac{1}{s}\psi(n) = \tilde{\psi}(\frac{m}{u}) + \tilde{\psi}(\frac{n}{s})$
- $\frac{r}{t}\tilde{\psi}(\frac{m}{u}) = \frac{r}{t} \cdot \frac{1}{u}\psi(m) = \frac{r}{tu}\psi(m) = \frac{1}{tu}\psi(rm) = \tilde{\psi}(\frac{rm}{tu})$

Thus $\tilde{\psi}$ is a $U^{-1}R$ -module homomorphism.

3. We need to show that $\tilde{\psi} \circ \eta_M = \psi$. Let $m \in M$. Then,

$$(\tilde{\psi} \circ \eta_M)(m) = \tilde{\psi}(\eta_M(m)) = \tilde{\psi}\left(\frac{m}{1}\right) = \frac{1}{1}\psi(m) = \psi(m)$$

Thus the diagram commutes.

4. Assume another $U^{-1}R$ -module homomorphism exists such that $\phi \circ \eta_M = \psi$. Then, under it we must have $\phi(\frac{m}{1}) = \psi(m)$, and since it is a $U^{-1}R$ -module homomorphism $\phi(\frac{m}{u}) = \phi(\frac{1}{u} \cdot \frac{m}{1}) = \frac{1}{u}\psi(m)$. So $\phi = \tilde{\psi}$, proving uniqueness.

5. **TODO:**

Problem 17. $U^{-1}M$ as a Tensor Product.

Let R be a ring, $U \subset R$ be a multiplicatively closed set, and let M be an R -module. We view $U^{-1}R$ as an R -module via the localization map $\eta : R \rightarrow U^{-1}R$.

1. Show that the function given below is a well-defined bilinear map of R -modules:

$$U^{-1}R \times M \xrightarrow{\bar{\alpha}} U^{-1}M$$

$$\left(\frac{r}{u}, m\right) \longmapsto \frac{rm}{u}$$

2. Use the universal property of tensor products to construct an R -module homomorphism $\alpha : U^{-1}R \otimes_R M \rightarrow U^{-1}M$. Find a formula for $\alpha\left(\frac{r}{u} \otimes m\right)$.
3. Show that the function β given below is a well-defined R -module homomorphism:

$$U^{-1}M \xrightarrow{\beta} U^{-1}R \otimes_R M$$

$$\frac{m}{u} \longmapsto \frac{1}{u} \otimes m$$

4. Prove that α and β are mutual inverses. Conclude that $U^{-1}M \cong U^{-1}R \otimes_R M$ as R -modules.
5. Deduce that $U^{-1}(R^n) \cong (U^{-1}R)^n$ as $U^{-1}R$ -modules.
6. If $f : M \rightarrow N$ is an R -module homomorphism, define a natural $U^{-1}R$ -module homomorphism $U^{-1}f : U^{-1}M \rightarrow U^{-1}N$. What is its formula on fractions?

Solution.

1. Suppose $\frac{r}{u} \sim \frac{s}{v}$. Then there exists $t \in U$ such that $t(rv - su) = 0$. Then,

$$\bar{\alpha}\left(\frac{r}{u}, m\right) = \frac{rm}{u}, \text{ and } \bar{\alpha}\left(\frac{s}{v}, m\right) = \frac{sm}{v}$$

And we have,

$$t(rmv - smu) = mt(rv - su) = m \cdot 0 = 0$$

Thus $\frac{rm}{u} \sim \frac{sm}{v}$, and the map is well-defined.

Now we show bilinearity:

- $\frac{s}{v} \bar{\alpha}\left(\frac{r}{u}, m\right) = \frac{s}{v} \cdot \frac{rm}{u} = \frac{sr m}{vu} = \bar{\alpha}\left(\frac{s}{v} \frac{r}{u}, m\right)$
- $\bar{\alpha}\left(\frac{s}{v} + \frac{r}{u}, m\right) = \left(\frac{s}{v} + \frac{r}{u}\right) \frac{m}{1} = \frac{sm}{v} + \frac{rm}{u} = \bar{\alpha}\left(\frac{s}{v}, m\right) + \bar{\alpha}\left(\frac{r}{u}, m\right)$

- $n \cdot \bar{\alpha} \left(\frac{r}{u}, m \right) = n \cdot \frac{rm}{u} = \frac{nr m}{u} = \bar{\alpha} \left(\frac{r}{u}, nm \right)$
- $\bar{\alpha} \left(\frac{r}{u}, n + m \right) = \frac{r}{u} \left(\frac{n+m}{1} \right) = \frac{rn}{u} + \frac{rm}{u} = \bar{\alpha} \left(\frac{r}{u}, n \right) + \bar{\alpha} \left(\frac{r}{u}, m \right)$

Thus $\bar{\alpha}$ is a well-defined bilinear map of R -modules.

2. By the for $U^{-1}R \times M$ there exists a map $\alpha : U^{-1}R \otimes_R M \rightarrow U^{-1}M$ where $\alpha \left(\frac{r}{u} \otimes m \right) = \frac{rm}{u}$.

$$\begin{array}{ccc} U^{-1}R \times M & \xrightarrow{\tau} & U^{-1}R \otimes_R M \\ & \searrow \bar{\alpha} & \swarrow \alpha \\ & U^{-1}M & \end{array}$$

3. • Suppose $\frac{m}{u} \sim \frac{n}{v}$. Then there exists $t \in U$ such that $t(mv - nu) = 0$. Then,

$$\beta \left(\frac{m}{u} \right) = \frac{1}{u} \otimes_R m, \text{ and } \beta \left(\frac{n}{v} \right) = \frac{1}{v} \otimes_R n$$

Then,

$$\frac{1}{u} \otimes_R m = \frac{tv}{utv} \otimes_R m = \frac{1}{utv} \otimes_R tmv = \frac{1}{utv} \otimes_R nt u = \frac{tu}{utv} \otimes_R n = \frac{1}{v} \otimes_R n$$

Hence the map is well-defined.

- $r \cdot \beta \left(\frac{m}{u} \right) = r \left(\frac{1}{u} \otimes_R m \right) = \frac{1}{u} \otimes_R rm = \beta \left(\frac{rm}{u} \right)$
- $\beta \left(\frac{m}{u} + \frac{n}{v} \right) = \beta \left(\frac{mv+nu}{uv} \right) = \frac{1}{uv} \otimes_R mv + nu = \frac{1}{uv} \otimes_R mv + \frac{1}{uv} \otimes_R nu = \beta \left(\frac{m}{u} \right) + \beta \left(\frac{n}{v} \right)$

Thus β is a well-defined R -module homomorphism.

4. • $(\alpha \circ \beta) \left(\frac{m}{u} \right) = \alpha \left(\frac{1}{u} \otimes_R m \right) = \frac{m}{u} = \text{id} \left(\frac{m}{u} \right)$
- $(\beta \circ \alpha) \left(\frac{r}{u} \otimes_R m \right) = \beta \left(\frac{rm}{u} \right) = \frac{r}{u} \otimes_R m = \text{id} \left(\frac{r}{u} \otimes_R m \right)$

Thus α and β are inverses of each other and we have a R -module isomorphism $U^{-1}M \cong U^{-1}R \otimes_R M$.

5. Let $M = R^n$. Then from part 4 we have $U^{-1}(R^n) \cong U^{-1}R \otimes_R R^n$. And from Worksheet 4.1 Problem 15 we have,

$$U^{-1}R \otimes_R R^n = (U^{-1}R \otimes_R R)^n = (U^{-1}R)^n$$

So,

$$U^{-1}(R^n) \cong (U^{-1}R)^n$$

6. We define $U^{-1}f : U^{-1}M \rightarrow U^{-1}N$ as $U^{-1}f \left(\frac{m}{u} \right) = \frac{f(m)}{u}$.

Throughout this course, “ring” means a *commutative* ring with unity. Recall that, given an ideal $I \subset R$, we define the vanishing set of I to be

$$\mathbb{V}(I) := \{[P] \in \operatorname{Spec}(R) \mid I \subset P\}.$$

The *Zariski topology* on $\operatorname{Spec}(R)$ is then defined by declaring the closed sets to be the subsets of the form $\mathbb{V}(I)$ as I ranges over all ideals of R . In Worksheet 2.1 we verified that this is indeed a topology and studied it in detail. For example, we showed that any ring map $\phi : R \rightarrow S$ induces a continuous map of topological spaces $\phi^\# : \operatorname{Spec}(S) \rightarrow \operatorname{Spec}(R)$ given by $[P] \mapsto [\phi^{-1}(P)]$. (More precisely, Spec is a contravariant functor from the category of commutative rings to the category of topological spaces.) Further, we showed that the natural quotient map $\pi : R \rightarrow R/I$ induces a closed embedding $\pi^\# : \operatorname{Spec}(R/I) \rightarrow \operatorname{Spec}(R)$ whose image is $\mathbb{V}(I)$.

Now that we have localization in our toolbox, we return to studying the Zariski topology on $\operatorname{Spec}(R)$. The main slogan is that localizing a ring lets us study the part of $\operatorname{Spec}(R)$ where a given “function” is nonzero. We have not yet defined functions on $\operatorname{Spec}(R)$, but in many familiar examples elements of R really are functions—for example, when R is the coordinate ring of an algebraic set. Recall the following theorem from the last worksheet.

Theorem 0.13. *Let R be a ring and let $U \subset R$ be a multiplicatively closed set. The localization map $\eta : R \rightarrow U^{-1}R$ induces an inclusion-preserving bijection:*

$$\begin{array}{ccc} \left\{ \begin{array}{c} \text{prime ideals in} \\ U^{-1}R \end{array} \right\} & \xrightarrow{\sim} & \left\{ \begin{array}{c} \text{prime ideals in } R \\ \text{disjoint from } U \end{array} \right\} \\ P & \longmapsto & \eta^{-1}(P) \end{array}$$

We begin by using this theorem in the case that $U = \{1, f, f^2, f^3, \dots\}$ for an element $f \in R$ to study what are called the basic open sets of $\operatorname{Spec}(R)$. A *basic open subset* – also called a *distinguished open subset* – of $\operatorname{Spec}(R)$ is a set of the form

$$D(f) := \{[P] \in \operatorname{Spec}(R) \mid f \notin P\} \subset \operatorname{Spec}(R)$$

for some element $f \in R$. We will show that these subsets form a nice basis for the Zariski topology on $\operatorname{Spec}(R)$. A *basis* for a topological space X is a collection $\mathcal{B}$ of open subsets of X such that for every open set $U \subset X$ and every point $x \in U$, there exists some $B \in \mathcal{B}$ with $x \in B \subset U$. Equivalently, every open subset of X can be written as a union of elements of $\mathcal{B}$.

Problem 1. Basic Open Sets.

Let R be a ring and $f, g \in R$.

1. Prove directly from the definition of the Zariski topology that $D(f)$ is an open set.
2. Compute $D(0)$ and $D(1)$.
3. Prove that $D(f) = D(f^n)$ for all $n \geq 1$.
4. Prove that $D(f) \cap D(g) = D(fg)$. Use induction to find a formula for the intersection of finitely many basic opens.
5. Prove that $D(f) \subset D(g)$ if and only if $f \in \sqrt{\langle g \rangle}$. Hint: Take complements to rewrite $D(f) \subset D(g)$ as $\mathbb{V}(g) \subset \mathbb{V}(f)$, and use the fact that $\mathbb{V}(I) \subset \mathbb{V}(J)$ if and only if $\sqrt{J} \subset \sqrt{I}$.
6. Prove that $D(f) = D(g)$ if and only if $\sqrt{\langle f \rangle} = \sqrt{\langle g \rangle}$.
7. Let $U \subset \text{Spec}(R)$ be an open subset and let $[\mathfrak{p}] \in U$. Show that there exists $f \in R$ such that $[\mathfrak{p}] \in D(f) \subset U$. Hint: Since U is open, write $\text{Spec}(R) \setminus U = \mathbb{V}(I)$ for some ideal I . Then use the fact that $[\mathfrak{p}] \notin \mathbb{V}(I)$ to choose $f \in I \setminus \mathfrak{p}$.
8. Deduce that the basic open sets form a basis for the Zariski topology on $\text{Spec}(R)$.

Solution.

1. $D(f)$ is the set of all prime ideals where $f \neq 0$, i.e. where f doesn't vanish. So, $\text{Spec}(R) \setminus D(f) = \mathbb{V}(f)$. Which by definition is a closed set, thus $D(f)$ is open.
2. $D(0) = \emptyset$, and $D(1) = \text{Spec}(R)$ since 0 is in every prime ideal, and all prime ideals are proper.
3. $D(f^n) \subseteq D(f)$ since P is prime and $f^n \notin P$ we must have $f \notin P$. And $D(f) \subseteq D(f^n)$ since if $f \notin P$ and P is prime, $f^n \notin P$. Thus $D(f) = D(f^n)$ for all $n \geq 1$.
4. $D(f) \cap D(g) = \{[P] \in \text{Spec}(R) \mid f, g \notin P\}$. Since P is prime, this implies that $fg \notin P$. Thus we have,

$$D(f) \cap D(g) = \{[P] \in \text{Spec}(R) \mid f, g \notin P\} = \{[P] \in \text{Spec}(R) \mid fg \notin P\} = D(fg)$$

5. ($\implies$)

Assume $D(f) \subset D(g)$.

Then we have $\mathbb{V}(g) \subset \mathbb{V}(f) \implies \sqrt{\langle f \rangle} \subset \sqrt{\langle g \rangle} \implies f \in \sqrt{\langle g \rangle}$.

($\Leftarrow$)

Assume $f \in \sqrt{\langle g \rangle}$. Then $f^n \in \langle g \rangle$ for some n . We first show that $\sqrt{\langle f \rangle} \subset \sqrt{\langle g \rangle}$. Let $h \in \sqrt{\langle f \rangle}$, then $h^m = rf \in \langle f \rangle$ for some $m \in \mathbb{N}$ and some $r \in R$. Then, we have $h^{nm} = r^n f^n \in \langle g \rangle$. Thus, $h \in \sqrt{\langle g \rangle}$ and $\sqrt{\langle f \rangle} \subset \sqrt{\langle g \rangle}$.

But that implies $\mathbb{V}(g) \subset \mathbb{V}(f) \implies D(f) \subset D(g)$.

6. ($\implies$)

Assume $D(f) = D(g)$. Then we have $\mathbb{V}(g) = \mathbb{V}(f) \implies \sqrt{\langle f \rangle} = \sqrt{\langle g \rangle}$.

($\Leftarrow$)

Assume $\sqrt{\langle f \rangle} = \sqrt{\langle g \rangle}$. Then we have $\mathbb{V}(g) = \mathbb{V}(f) \implies D(f) = D(g)$.

7. Let $U \in \text{Spec}(R)$ open set. Then its complement $\text{Spec}(R) \setminus U = \mathbb{V}(I)$ is a closed set for some $I \subset R$. Since $[\mathfrak{p}] \in U$ we know that $[\mathfrak{p}] \notin \mathbb{V}(I)$, then we have $I \not\subset \mathfrak{p}$. So there exists some $f \in I$ such that $f \notin \mathfrak{p}$ giving us $[\mathfrak{p}] \in D(f) \subset U$.
8. Since $[\mathfrak{p}]$ and U were arbitrary, by part (7) the basic open sets satisfy the basis criterion and hence form a basis.

Problem 2. Examples of Basic Opens.

Explicitly describe the basic open set $D(f) \subset \text{Spec}(R)$ in each of the following cases.

- | | |
|--|--|
| 1. $R = \mathbb{K}[x]$ and $f = x$. | 5. $R = \mathbb{K}[x, y] / \langle xy \rangle$ and $f = \bar{y}$. |
| 2. $R = \mathbb{K}[x]$ and $f = x - a$ for $a \in \mathbb{K}$. | 6. $R = \mathbb{Z} / \langle 12 \rangle$ and $f = \bar{2}$. |
| 3. $R = \mathbb{Z}$ and $f = p$ for a prime number p . | 7. $R = \mathbb{Z} / \langle 12 \rangle$ and $f = \bar{3}$. |
| 4. $R = \mathbb{K}[x, y] / \langle xy \rangle$ and $f = \bar{x}$. | 8. $R = \mathbb{Z} / \langle 12 \rangle$ and $f = \bar{6}$. |

Solution.

1. $D(x) = \text{Spec}(R) \setminus \langle x \rangle = \{ \langle 0 \rangle \cup \langle p(x) \rangle \mid p \neq x, p \text{ is irreducible} \}.$
2. $D(x - a) = \text{Spec}(R) \setminus \langle x - a \rangle = \{ \langle 0 \rangle \cup \langle p(x) \rangle \mid p \neq x - a, p \text{ is irreducible} \}.$
3. $D(p) = \text{Spec}(\mathbb{Z}) \setminus \langle p \rangle = \{ [\mathfrak{p}] \in \text{Spec}(\mathbb{Z}), \mathfrak{p} \neq \langle p \rangle \}.$
4. $D(\bar{x}) = \{ \langle \bar{y}, q(x) \rangle, q \text{ is irreducible or zero} \}.$
5. $D(\bar{y}) = \{ \langle \bar{x}, q(y) \rangle, q \text{ is irreducible or zero} \}.$
6. $D(\bar{2}) = \{ \langle \bar{3} \rangle \}.$
7. $D(\bar{3}) = \{ \langle \bar{2} \rangle \}.$
8. $D(\bar{6}) = \emptyset.$

Problem 3. Induced Map by Localization.

Let $U \subset R$ be a multiplicatively closed set and let $\eta : R \rightarrow U^{-1}R$ be the localization map. In this exercise we will study the induced map $\eta^\# : \text{Spec}(U^{-1}R) \rightarrow \text{Spec}(R)$.

1. Use the correspondence between prime ideals in $U^{-1}R$ and R given in Theorem 0.13 to prove that $\eta^\#$ is injective and the image of $\eta^\#$ is

$$X_U := \{[\mathfrak{p}] \in \text{Spec}(R) \mid \mathfrak{p} \cap U = \emptyset\} \subset \text{Spec}(R).$$

2. Let $R = \mathbb{Z}$ and $U = \mathbb{Z} \setminus \langle p \rangle$ for a prime $p \in \mathbb{Z}$. Show that $X_U = \{\langle 0 \rangle, \langle p \rangle\}$. Prove that this is not open in $\text{Spec}(\mathbb{Z})$.
3. Let $R = \mathbb{Z}$ and $U = \{2^i 3^j \mid i, j \in \mathbb{Z}_{\geq 0}\}$. Show that $X_U = \{\langle 0 \rangle\} \cup \{\langle p \rangle \mid p \text{ prime}, p \neq 2, 3\}$. Show that this is an open subset of $\text{Spec}(\mathbb{Z})$.
4. Describe X_U when $R = \mathbb{C}[x, y]$ and $U = \{(xy)^n \mid n \in \mathbb{Z}_{\geq 0}\}$.
5. Prove that $X_U = \bigcap_{f \in U} D(f)$.
6. Deduce that if U is a multiplicative set generated by finitely many elements then X_U is open.
7. Fix an element $\frac{r}{u} \in U^{-1}R$. Prove that $\eta^\#(D(\frac{r}{u})) = D(r) \cap X_U$. Hint: In $U^{-1}R$, the element u is a unit, so $D(\frac{r}{u}) = D(\frac{r}{1})$. Then use part (1) to identify the image of $\eta^\#$.
8. Prove that $\eta^\# : \text{Spec}(U^{-1}R) \rightarrow \text{Spec}(R)$ is a homeomorphism onto its image X_U (considered with the subspace topology).
9. Specializing to the case that $U = \{1, f, f^2, f^3, \dots\}$, prove that $\eta^\# : \text{Spec}(R[\frac{1}{f}]) \rightarrow \text{Spec}(R)$ is a homeomorphism onto the basic open subset $D(f)$.

Solution.

1. By Theorem 0.1 the map $\mathfrak{q} \rightarrow \eta^{-1}(\mathfrak{q})$ gives a bijection between prime ideals in $U^{-1}R$ to prime ideals disjoint from U in R . That is,

$$\text{im}(\eta^\#) = X_U := \{[\mathfrak{p}] \in \text{Spec}(R) \mid \mathfrak{p} \cap U = \emptyset\} \subset \text{Spec}(R).$$

and $\eta^\#$ is injective since the bijection is injective.

2. We have, by construction,

$$X_U = \{[\mathfrak{p}] \in \text{Spec}(R) \mid \mathfrak{p} \cap U = \emptyset\} = \{\langle p \rangle, \langle 0 \rangle\}$$

Assume X_U is open in $\text{Spec}(\mathbb{Z})$. Since distinguished open sets form a basis for the Zariski topology, there exists some $D(f) \subseteq X_U$ such that $\langle 0 \rangle \subseteq D(f)$ which means $f \neq 0$. So we must have $f = n$, for some $n \in \mathbb{Z}$. Since n is divisible by only a finite number of primes, we have $D(n) = \{[P] \in \text{Spec}(\mathbb{Z}) \mid P \nmid n\}$ which is an infinite set, and thus $D(f) \not\subseteq X_U$. Thus, a contradiction and X_U cannot be a closed set.

3. By definition,

$$\begin{aligned} X_U &= \{[p] \in \text{Spec}(\mathbb{Z}) \mid p \cap U = \emptyset\} \\ &= \{[p] \in \text{Spec}(\mathbb{Z}) \mid p \neq 2, 3\} \\ &= \{\langle 0 \rangle\} \cup \{\langle p \rangle \mid p \text{ prime}, p \neq 2, 3\} \end{aligned}$$

Since X_U is the complement of a closed set in $\text{Spec}(\mathbb{Z})$, namely, $V(\langle 6 \rangle) = \{\langle 2 \rangle, \langle 3 \rangle\}$, it is an open set of $\text{Spec}(\mathbb{Z})$.

4. By definition,

$$\begin{aligned} X_U &= \{[p] \in \text{Spec}(\mathbb{C}[x, y]) \mid p \cap U = \emptyset\} \\ &= \{[p] \in \text{Spec}(\mathbb{C}[x, y]) \mid p \neq x, y\} \\ &= \{\langle 0 \rangle\} \end{aligned}$$

5. $X_U \supseteq \cap_{f \in U} D(f)$ is clear by definition. To see the other containment, let $[p] \in \cap_{f \in U} D(f)$. Then $f \notin p$ for all f . Hence, $p \cap U = \emptyset$ and thus $[p] \in X_U$.

6. In this case, X_U is a finite intersection of open sets and thus is open.

7. We know that $D(\frac{r}{u}) = D(\frac{r}{1})$. For each prime $[p] \in D(\frac{r}{1})$,

$$\frac{r}{1} = \eta(r) \notin p \iff r \notin \eta^\#(p)$$

8. From part 1 we know that $\eta^\#$ is injective, thus it is a bijection onto its image. Let $D_u^r \subseteq \text{Spec}(U^{-1}R)$ be a basic open set. Then by part 7,

$$\eta^\#(D(\frac{r}{u})) = D(r) \cap X_U$$

which is open in the subspace topology on X_U .

9. If $U = \{1, f, f^2, \dots\}$, then

$$X_U = \cap_{f \in U} D(f) = D(f)$$

as $D(f)$ contains $D(f^n)$ for all n . Hence, $\eta(\text{Spec}(f[\frac{1}{f}])) \cong D(f)$.

Let X be a topological space. An *open cover* of X is a collection $\{U_i\}_{i \in \Lambda}$ of open subsets of X such that $X = \bigcup_{i \in \Lambda} U_i$. A *finite subcover* is a finite subcollection $\{U_{i_1}, \dots, U_{i_n}\}$ that still covers X , i.e. $X = U_{i_1} \cup \dots \cup U_{i_n}$.

With these terms in hand, recall that X is *quasi-compact* if every open cover admits a finite subcover. More generally, a subset $Y \subset X$ is *quasi-compact* if every cover of Y by open subsets of X admits a finite subcover; equivalently, Y is quasi-compact in the subspace topology. In the Hausdorff setting this is simply compactness; the prefix “quasi-” is standard in algebraic geometry because Zariski topologies are virtually never Hausdorff, and one wants to keep the two notions distinct.

Every affine spectrum $\text{Spec } R$ is quasi-compact. The proof is a clean illustration of how the basic opens $D(f)$ serve as a bridge between topology and commutative algebra. If $\{D(f_i)\}_{i \in \Lambda}$ covers $\text{Spec}(R)$, then no prime ideal contains every f_i , so the f_i generate the unit ideal. But any expression $1 = \sum a_i f_i$ involves only finitely many terms, so a finite subfamily already generates the unit ideal and the corresponding $D(f_i)$ ’s already cover. The argument generalizes immediately: any open subset that can be written as a finite union of basic opens is itself quasi-compact.

This kind of translation—reformulating a covering condition as an ideal-membership statement, then exploiting the finiteness built into ring equations—is one of the workhorses of scheme theory.

Problem 4. Quasi-compactness of the Zariski Topology.

Let R be a ring.

1. Let $\Lambda \subset R$ be a collection of elements of R . Prove that $\{D(f)\}_{f \in \Lambda}$ is an open cover of $\text{Spec}(R)$ if and only if $\langle f \mid f \in \Lambda \rangle$ is the unit ideal.
2. Let $\Lambda \subset R$ be a collection of elements of R . Explain why if $\langle f \mid f \in \Lambda \rangle$ is the unit ideal then there exist finitely many elements $f_1, \dots, f_n \in \Lambda$ such that $\langle f_1, \dots, f_n \rangle$ is also the unit ideal.
3. Deduce that every cover of $\text{Spec}(R)$ by basic open subsets has a finite subcover.
4. Use the fact that the $D(f)$ form a basis to prove that $\text{Spec}(R)$ is quasi-compact. Hint: Refine each open set in the cover by basic open neighborhoods using the basis property from Problem 1.
5. Prove that $\mathbb{V}(I)$ is quasi-compact for every ideal $I \subseteq R$. Hint: Use the homeomorphism $\text{Spec}(R/I) \cong \mathbb{V}(I)$ from Worksheet 2.1.
6. Let $I \subset R$ be an ideal and set $U := \text{Spec}(R) \setminus \mathbb{V}(I)$.

(a) Show that $U = \bigcup_{f \in I} D(f)$.

(b) Show that if U is quasi-compact, then $U = D(f_1) \cup \dots \cup D(f_n)$ for some $f_1, \dots, f_n \in I$.

(c) Deduce that U is quasi-compact if and only if $\sqrt{I} = \sqrt{\langle f_1, \dots, f_n \rangle}$ for some $f_1, \dots, f_n \in I$.

Equivalently, U is quasi-compact if and only if $\sqrt{I}$ is the radical of a finitely generated ideal.

7. Deduce that if R is Noetherian, every open subset of $\text{Spec}(R)$ is quasi-compact. Hint: First show that $\text{Spec}(R) \setminus \mathbb{V}(I) = \bigcup_{f \in I} D(f)$.

Solution.

1. ($\implies$)

Assume $\{D(f)\}_{f \in \Lambda}$ is an open cover of $\text{Spec}(R)$. Then $\langle f \mid f \in \Lambda \rangle \not\subseteq \mathfrak{p}$ for any prime ideal. In particular, $\langle f \mid f \in \Lambda \rangle$ is not contained in any maximal ideal, and hence must be the unit ideal.

($\impliedby$)

Assume $\langle f \mid f \in \Lambda \rangle$ is the unit ideal. Then it is not contained in any prime ideal. In particular, there exists at least one $f \in \Lambda$ such that $f \notin \mathfrak{p}$ for all $\mathfrak{p}$. Hence, $\{D(f)\}_{f \in \Lambda}$

is a open cover of $\text{Spec}(R)$.

2. If $\langle f \mid f \in \Lambda \rangle$ is the unit ideal, then $1 = \sum_i a_i f_i \in \langle f \mid f \in \Lambda \rangle$ is a finite sum. Then, these (finite) f_i s generate the unit ideal.
3. Thus, the corresponding finitely many $D(f_i)$ s generate a finite subcover for any given cover of $\text{Spec}(R)$ by basic open sets.
4. Every open set can be written as a union of the basic open subsets and hence every open cover is a cover of $\text{Spec}(R)$ by basic open subsets and hence has a finite subcover. Thus, $\text{Spec}(R)$ is quasi-compact.
5. Recall that, $\mathbb{V}(I) \cong \text{Spec}(R/I)$. Since R/I is a ring, and we just showed that the spectrum of a ring is quasi-compact, $\text{Spec}(R/I)$ is quasi-compact.
6. (a) $U = \text{Spec}(R) \setminus \mathbb{V}(I) = \text{Spec}(R) \setminus \text{Spec}(R/I)$. We know that ideals in $\text{Spec}(R/I)$ correspond to ideals in $\text{Spec}(R)$ that contain I . So, U is the set of ideals in $\text{Spec}(R)$ that don't contain I . Hence,

$$U = \text{Spec}(R) \setminus \text{Spec}(R/I) = \cup_{f \in I} D(f)$$

- (b) If U is quasi-compact, then any cover of U has a finite-subcover. Since $\cup_{f \in I} D(f)$ is a cover, it also admits a finite subcover,

$$U = D(f_1) \cup \dots \cup D(f_n)$$

- (c) ($\implies$)

Assume U is quasi-compact. Then,

$$\begin{aligned} U &= D(f_1) \cup \dots \cup D(f_n) \\ &= \text{Spec}(R) \setminus (\mathbb{V}(f_1) \cap \dots \cap \mathbb{V}(f_n)) \\ &= \text{Spec}(R) \setminus \mathbb{V}(\langle f_1, \dots, f_n \rangle) \\ &= \text{Spec}(R) \setminus \mathbb{V}(\sqrt{\langle f_1, \dots, f_n \rangle}) \\ &= \text{Spec}(R) \setminus \mathbb{V}(I) = \text{Spec}(R) \setminus \mathbb{V}(\sqrt{I}) \\ &\implies \sqrt{I} = \sqrt{\langle f_1, \dots, f_n \rangle} \end{aligned}$$

- ($\impliedby$)

Assume $\sqrt{I} = \sqrt{\langle f_1, \dots, f_n \rangle}$ for some $f_1, \dots, f_n \in I$. Then,

$$\begin{aligned} U &= \operatorname{Spec}(R) \setminus \mathbb{V}(I) = \operatorname{Spec}(R) \setminus \mathbb{V}(\sqrt{I}) \\ &= \operatorname{Spec}(R) \setminus \mathbb{V}(\sqrt{\langle f_1, \dots, f_n \rangle}) \\ &= \operatorname{Spec}(R) \setminus (\mathbb{V}(f_1) \cap \dots \cap \mathbb{V}(f_n)) \\ &= \operatorname{Spec}(R) \setminus \mathbb{V}(\langle f_1, \dots, f_n \rangle) \\ &= D(f_1) \cup \dots \cup D(f_n) \end{aligned}$$

7. If R is Noetherian, every ideal $\mathbb{V}(I)$ is finitely-generated. Then by part 6.b, every open subset of $\operatorname{Spec}(R)$ is quasi-compact.

Problem 5. Counterexamples to Quasi-compactness of Subsets.

In the previous problem we saw that $\text{Spec}(R)$ and every closed subset $V(I) \subset \text{Spec}(R)$ are quasi-compact in the Zariski topology. We also saw that when R is Noetherian every open subset of $\text{Spec}(R)$ is quasi-compact. In this exercise we explore some pathologies for open subsets when R is not Noetherian.

1. Let $R = \mathbb{K}[x_1, x_2, \dots]$ and $I = \langle x_1, x_2, \dots \rangle$. Consider the open subset $U := \text{Spec}(R) \setminus V(I)$. Prove that $\{D(x_i)\}_{i=1}^\infty$ is an open cover of U , which has no finite subcover. Hence an open subset of $\text{Spec}(R)$ need not be quasi-compact. Hint: If you try to cover U by only $D(x_1), \dots, D(x_n)$, consider the prime ideal $\langle x_1, \dots, x_n \rangle$.
2. Explain why each basic open set $D(x_i)$ is quasi-compact. Deduce that an arbitrary union of quasi-compact open subsets of $\text{Spec}(R)$ need not be quasi-compact.

Solution.

1. For the sake of contradiction, assume there exists a finite subcover of $\{D(x_i)\}_{i=1}^\infty$ that covers U , say $D(x_1) \cup \dots \cup D(x_n)$. Then, the ideal generated by $\langle x_1, \dots, x_n \rangle \subseteq I$ is in U . Which is a contradiction. Hence, no finite subcover exists.
2. We know that $D(x_i) \cong \text{Spec}\left(R\left[\frac{1}{x_i}\right]\right)$, since the spectrum of any ring is quasi-compact $D(x_i)$ is quasi compact. And in part (1) we see that arbitrary unions of quasi-compact open subsets of $\text{Spec}(R)$ are not quasi-compact, however finite unions are.

Problem 6. Checking Properties on a Finite Cover.

Let R be a ring. Suppose that $f_1, \dots, f_n \in R$ are such that $\{D(f_i)\}_{i=1}^n$ is a finite open cover of $\text{Spec}(R)$, i.e., $\text{Spec}(R) = D(f_1) \cup D(f_2) \cup \dots \cup D(f_n)$. We will see that certain ring-theoretic properties of R can be checked locally by looking at the rings $R[\frac{1}{f_i}]$.

1. Let $r \in R$. Prove that if $r/1 = 0$ in $R[\frac{1}{f_i}]$ for all i then $r = 0$ in R . Hint: In $R[\frac{1}{f_i}]$, the equality $r/1 = 0$ means that $f_i^{n_i}r = 0$ for some $n_i \geq 0$.
2. Deduce that if $R[\frac{1}{f_i}] = 0$ for all i then $R = 0$.
3. Prove that if $R[\frac{1}{f_i}]$ is reduced for all i , then R is reduced.

Solution.

1. Since $\{D(f_i)\}_{i=1}^n$ covers $\text{Spec}(R)$, we know $\langle f_1, \dots, f_n \rangle = R$. And for any prime $\mathfrak{p}$ we have $f_i \in \mathfrak{p} \iff f_i^{n_i} \in \mathfrak{p}$, so $D(f_i) = D(f_i^{n_i})$.

Hence $\{D(f_i^{n_i})\}_{i=1}^n$ also covers $\text{Spec}(R)$, and we have

$$1 = \sum_{i=1}^n b_i f_i^{n_i} \implies r = 1 \cdot r = \sum_{i=1}^n b_i f_i^{n_i} r = \sum_{i=1}^n b_i \cdot 0 = 0.$$

Thus, $r = 0$ in R .

2. If $R[\frac{1}{f_i}] = 0$ for all i then each f_i is nilpotent in R . But $D(f_i)_{1 \leq i \leq n}$ covers R . Since $f_i \in \sqrt{\langle 0 \rangle} \subseteq \mathfrak{p}$ for all primes, $D(f_i) = \emptyset$ for all i . Thus, $\text{Spec}(R) = \emptyset \implies R = 0$.
3. Let $r \in R$ such that $r^m = 0$. Then for all i , in $R[\frac{1}{f_i}]$, we have $\frac{r^m}{1} = 0$. Since this ring is reduced, this implies $\frac{r}{1} = 0 \implies r = 0$ (by part 1).

An element $r \in R$ is *nilpotent* if $r^n = 0$ for some integer $n \geq 1$. If $I \subset R$ is an ideal, the *radical* of I is

$$\sqrt{I} := \{r \in R \mid r^n \in I \text{ for some } n \geq 1\},$$

which we previously checked is an ideal of R containing I and $\sqrt{\sqrt{I}} = \sqrt{I}$. An ideal $I \subset R$ is *radical* if and only if $I = \sqrt{I}$. We saw that radical ideals play a crucial role in the study of algebraic sets as $\mathbb{I}(V)$ is always radical and Hilbert's Nullstellensatz gave an inclusion-reversing bijection between radical ideals in $\mathbb{K}[x_1, \dots, x_n]$ and algebraic subsets of $\mathbb{K}^n$ when $\mathbb{K}$ is an algebraically closed field. These facts highlight the following motto: *(classical) geometry does not see nilpotents*. (Here classical means anything one encounters pre-scheme theory; topology, classical algebraic geometry, complex geometry, etc.)

We shall see that the invisibility of nilpotents to geometry also applies to spectra of rings. The *nilradical* of R is the set of all nilpotent elements of R , equivalently the radical of the zero ideal:

$$\text{Nil}(R) := \sqrt{\langle 0 \rangle} = \{r \in R \mid r^n = 0 \text{ for some } n \geq 1\}.$$

If $f \in \text{Nil}(R)$ and $f^n = 0$, then $f^n \in I$ for every ideal $I \subset R$. In particular, if $P \subset R$ is a prime ideal, then $f^n = 0 \in P$, which implies that $f \in P$. Thus the nilradical of R is contained in every prime ideal of R . In the exercises below you will prove the reverse inclusion, giving the fundamental characterization

$$\text{Nil}(R) = \bigcap_{\mathfrak{p} \in \text{Spec}(R)} \mathfrak{p}.$$

Since prime ideals of $R/\text{Nil}(R)$ are in bijection with prime ideals of R containing $\text{Nil}(R)$, the equality above implies that prime ideals of R are in bijection with prime ideals of $R/\text{Nil}(R)$. Geometrically, this means that the quotient map $\pi : R \rightarrow R/\text{Nil}(R)$ induces a map of topological spaces

$$\text{Spec}(R/\text{Nil}(R)) \xrightarrow{\pi^\#} \text{Spec}(R)$$

that is bijective and continuous. We will see shortly that this map is actually a homeomorphism; hence, topologically $\text{Spec}(R)$ and $\text{Spec}(R/\text{Nil}(R))$ are indistinguishable. This is another instance of the motto: *geometry does not see nilpotents*. A ring R is called *reduced* if $\text{Nil}(R) = \langle 0 \rangle$, i.e., if the only nilpotent element is 0. We often call $R/\text{Nil}(R)$ the *reduced ring* associated to R and denote it $R_{\text{red}} := R/\text{Nil}(R)$.

More generally, the nilradical characterization extends to arbitrary ideals. One checks that $\text{Nil}(R/I) = \sqrt{I}/I$, since the elements of R/I whose powers lie in I are exactly the images of elements of $\sqrt{I}$. Applying the nilradical formula in the ring R/I and pulling back to R gives

$$\sqrt{I} = \bigcap_{\substack{\mathfrak{p} \in \text{Spec}(R) \\ \mathfrak{p} \supseteq I}} \mathfrak{p}.$$

This has immediate geometric consequences. For example, this gives another proof that $\mathbb{V}(I) = \mathbb{V}(\sqrt{I})$. Thus the Zariski-closed subsets of $\text{Spec}(R)$ determined by I and $\sqrt{I}$ are identical. This interacts cleanly with quotient rings; in particular, an argument like the one above shows that $\text{Spec}(R/I)$ and $\text{Spec}(R/\sqrt{I})$ are homeomorphic.

Problem 7. Nilradicals and Reduced Rings.

Let R be a ring.

1. Explain why $\text{Nil}(R)$ is an ideal and $\text{Nil}(R) \subset \bigcap_{\mathfrak{p} \in \text{Spec}(R)} \mathfrak{p}$.
2. Prove that if $f \in R$ is not a nilpotent then $R[\frac{1}{f}] \neq 0$ and hence contains a prime ideal. Hint: If $R[\frac{1}{f}] = 0$, then $1 = 0$ in $R[\frac{1}{f}]$. What does that say about some power of f ?
3. Use the correspondence between prime ideals in $R[\frac{1}{f}]$ and R not containing f to show there is a prime ideal $\mathfrak{p} \subset R$ such that $f \notin \mathfrak{p}$.
4. Deduce the theorem

$$\text{Nil}(R) = \bigcap_{\mathfrak{p} \in \text{Spec}(R)} \mathfrak{p}.$$

5. Define R_{red} to be equal to $R/\text{Nil}(R)$. Prove that R_{red} is reduced.
6. Use the above characterization to prove that $D(f) = \emptyset$ if and only if f is a nilpotent element.
7. Show that $D(f) = \text{Spec}(R)$ if and only if f is a unit in R .
8. Let $\pi : R \rightarrow R_{\text{red}}$ be the natural quotient map. Prove that the induced map $\pi^\# : \text{Spec}(R_{\text{red}}) \rightarrow \text{Spec}(R)$ is a homeomorphism. (Hint: We already know it is continuous. Use part 4 to show it is a bijection. Finally, show that the map is closed by proving that for every ideal $I \supseteq \text{Nil}(R)$, the closed set $\mathbb{V}(I/\text{Nil}(R)) \subseteq \text{Spec}(R_{\text{red}})$ maps to $\mathbb{V}(I) \subseteq \text{Spec}(R)$.)
9. Compute R_{red} for each of the following rings:

(a) $\mathbb{K}[\epsilon]/\langle \epsilon^2 \rangle,$

(c) $\mathbb{K}[x, y]/\langle x^2, xy \rangle,$

(b) $\mathbb{K}[x]/\langle x^3 \rangle,$

(d) $\mathbb{Z}/12\mathbb{Z}.$

Solution.

1. We only need to show that $\text{Nil}(R)$ is closed under addition and multiplication from the ring.
 - (a) Let $a, b \in \text{Nil}(R)$ such that $a^n = b^m = 0$. Wlog assume $m > n$, then by the binomial theorem $(a + b)^{m+n} = 0$.
 - (b) Let $r \in R$, and $a \in \text{Nil}(R)$ such that $a^n = 0$. Then, $(ra)^n = r^n a^n = 0$.

Hence, $\text{Nil}(R)$ is an ideal. Let $a \in \text{Nil}(R)$ then $a^n = 0 \in \mathfrak{p}$ then by primeness of $\mathfrak{p}$ $a \in \mathfrak{p}$ for all $\mathfrak{p}$, hence $\text{Nil}(R) \subset \bigcap_{\mathfrak{p} \in \text{Spec}(R)} \mathfrak{p}$.

2. We will prove this by contrapositive. Assume $R[\frac{1}{f}] = 0$. Then $\frac{1}{1} = \frac{0}{1}$, and there exists f^n such that $f^n \cdot 1 = 0 \implies f^n = 0 \implies f$ is a nilpotent.
3. Prime ideals in $R[\frac{1}{f}]$ correspond to prime ideals in R not containing f . Since a non-zero ring has at least one maximal (and hence prime) ideal there is at least one $\mathfrak{p} \subset R$ such that $f \notin \mathfrak{p}$.
4. We only need to show the containment $\bigcap_{\mathfrak{p} \in \text{Spec}(R)} \mathfrak{p} \subseteq \text{Nil}(R)$. Let $f \in \bigcap_{\mathfrak{p} \in \text{Spec}(R)} \mathfrak{p}$ then $f \in \mathfrak{p}$ for all $\mathfrak{p}$. Hence $R[\frac{1}{f}] = 0$. By part (2), this implies that f is nilpotent. Hence,

$$\text{Nil}(R) = \bigcap_{\mathfrak{p} \in \text{Spec}(R)} \mathfrak{p}.$$

5. Suppose $\bar{a} \in R_{\text{red}} = R/\text{Nil}(R)$ is nilpotent, say $\bar{a}^k = 0$ in R_{red} . Then $a^k \in \text{Nil}(R)$, so $(a^k)^\ell = 0$ in R for some ℓ , i.e. $a^{k\ell} = 0$. Thus $a \in \text{Nil}(R)$, which means $\bar{a} = 0$ in R_{red} . Hence the only nilpotent of R_{red} is 0, so R_{red} is reduced.
6. By Problem 3.9, we know that $\eta^\# : \text{Spec}(R[\frac{1}{f}]) \rightarrow D(f)$ is a homeomorphism. If f is nilpotent, then $\text{Spec}(R[\frac{1}{f}]) = \emptyset$ and thus $D(f) = \emptyset$. Similarly, if $D(f) = \emptyset$ then $R[\frac{1}{f}] = 0$, then f is a nilpotent.
7. If f is unit, then $f \notin \mathfrak{p}$ for all $\mathfrak{p}$ hence $D(f) = \text{Spec}(R)$. And if $D(f) = \text{Spec}(R)$ then $f \notin \bigcup_{\mathfrak{p} \in \text{Spec}(R)} \mathfrak{p} \implies f$ is not in any maximal ideal and hence is a unit.
8. We know that $\pi^\#$ is continuous. It is a bijection because ideals in $\text{Spec}(R_{\text{red}})$ correspond to ideals in R containing $\text{Nil}(R)$, by part 4 this is every prime ideal. To see that this map is closed, let $\mathbb{V}(I_{\text{red}})$ be a closed set in R_{red} . Then $\pi^\#(\mathbb{V}(I_{\text{red}})) = \{\mathfrak{p} \mid \pi^{-1}(I_{\text{red}}) \subseteq \mathfrak{p}\} = \{\mathfrak{p} \mid I \subseteq \mathfrak{p}\} = \mathbb{V}(I)$, which is closed in $\text{Spec}(R)$.

9. (a) $\mathbb{K}$ (c) $\mathbb{K}[y]$
(b) $\mathbb{K}$ (d) $\mathbb{Z}/6\mathbb{Z}$

Given a ring homomorphism $\phi : R \rightarrow S$, we know that it induces a continuous map $\phi^\# : \text{Spec}(S) \rightarrow \text{Spec}(R)$ of topological spaces. A natural geometric question is: what is the fibre of this map over a point $[\mathfrak{p}] \in \text{Spec}(R)$? Recall that if $f : X \rightarrow Y$ is a map of topological spaces, then the fibre over a point $y \in Y$ is just the preimage $f^{-1}(\{y\}) = \{x \in X \mid f(x) = y\} \subset X$, considered with the subspace topology. In algebraic geometry, authors often write X_y for the fibre over y when the map f is clear. Returning to the induced map $\phi^\#$, since we know that $\phi^\#([\mathfrak{q}]) = [\phi^{-1}(\mathfrak{q})]$, we can directly describe the fibre as

$$(\phi^\#)^{-1}([\mathfrak{p}]) = \{[\mathfrak{q}] \in \text{Spec}(S) \mid \phi^{-1}(\mathfrak{q}) = \mathfrak{p}\} \subset \text{Spec}(S).$$

However, we would like a better, more algebraic description of this set. Answering this involves two conceptual steps: localization and reduction to the residue field. It is worth pausing to try to get an intuitive understanding of each step. As a roadmap: to compute the fibre over $[\mathfrak{p}]$, we first localize at $\mathfrak{p}$, then quotient by $\mathfrak{p} R_{\mathfrak{p}}$; this is what leads naturally to the fibre ring $\kappa(\mathfrak{p}) \otimes_R S$.

Let $f : X \rightarrow Y$ be any continuous map of topological spaces. The fibre over a point $y \in Y$ is local on Y in the sense that it really only depends on the behavior of f at the point y , or perhaps in an open neighborhood of y . More precisely, if $U \subset Y$ contains y and $V = f^{-1}(U)$, then

$$f^{-1}(\{y\}) = (f|_V)^{-1}(\{y\}),$$

where $f|_V : V \rightarrow U$ is the restriction of f . If we wished to be fancy, we might draw a commutative diagram of topological spaces

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ \uparrow \iota & & \uparrow \\ V & \xrightarrow{f|_V} & U \end{array}$$

where ι is the inclusion of V into X . Since $V = f^{-1}(U)$ and $y \in U$, every point of the fibre over y already lies in V , so the displayed equality holds. In other words, restricting f to V does not change the fibre over y ; more generally, we may replace X by any subset that still contains the fibre over y .

Returning to the algebraic situation $\phi^\# : \text{Spec}(S) \rightarrow \text{Spec}(R)$, we happen to know from the previous exercise on localization a very nice subset of $\text{Spec}(R)$ containing the point $[\mathfrak{p}]$, namely

$$[\mathfrak{p}] \in X_U \subset \text{Spec}(R) \quad \text{where} \quad U = R \setminus \mathfrak{p}.$$

Even better, the localization map $\eta : R \rightarrow R_{\mathfrak{p}}$ induces a homeomorphism from $\text{Spec}(R_{\mathfrak{p}})$ onto X_U . We could thus hope to compute the fibre over $[\mathfrak{p}]$ by trying to factor $\phi^\#$ through

$\eta^\#$, i.e. to find a commutative diagram of topological spaces as shown on the left below for some subset $V \subset \text{Spec}(S)$.

$$\begin{array}{ccc} \text{Spec}(S) & \xrightarrow{\phi^\#} & \text{Spec}(R) \\ \uparrow \text{---} & & \uparrow \eta^\# \\ V & \text{-----} & \text{Spec}(R_{\mathfrak{p}}) \end{array} \quad \begin{array}{ccc} R & \xrightarrow{\phi} & S \\ \downarrow \eta & & \downarrow \text{---} \\ R_{\mathfrak{p}} & \text{-----} & ?? \end{array} .$$

It turns out we can find such a subset using the universal property of localization. To see why, let us just hope that V can be realized as the spectrum of some ring and that the dashed maps arise from ring homomorphisms. This means we are trying to complete the diagram of rings above on the right. Recall that the image of a multiplicatively closed set is again multiplicatively closed, so we may localize S at

$$T := \phi(R \setminus \mathfrak{p}) \subset S.$$

Let $\eta_S : S \rightarrow T^{-1}S$ be the localization map. We now have a diagram in which the diagonal map is the composition of ϕ and η_S :

$$\begin{array}{ccc} R & \xrightarrow{\phi} & S \\ \downarrow \eta & \searrow & \downarrow \eta_S \\ R_{\mathfrak{p}} & \xrightarrow{\tilde{\phi}} & T^{-1}S \end{array}$$

where the diagonal map is just the composition of ϕ and η_S . Every element of $R \setminus \mathfrak{p}$ maps to a unit in $T^{-1}S$ under the diagonal map, so the universal property of localization gives a unique ring map $\tilde{\phi} : R_{\mathfrak{p}} \rightarrow T^{-1}S$ making the diagram commute. Passing to spectra gives a commutative diagram

$$\begin{array}{ccc} \text{Spec}(S) & \xrightarrow{\phi^\#} & \text{Spec}(R) \\ \uparrow \eta_S^\# & & \uparrow \eta^\# \\ \text{Spec}(T^{-1}S) & \xrightarrow{\tilde{\phi}^\#} & \text{Spec}(R_{\mathfrak{p}}) \end{array}$$

and hence the fibre of $\phi^\#$ over $[\mathfrak{p}]$ is naturally identified with the fibre of $\tilde{\phi}^\#$ over the point $[\mathfrak{p} R_{\mathfrak{p}}]$. Since $\eta^\#$ and $\eta_S^\#$ are homeomorphisms onto their images, this reduces the computation of the fibre of $\phi^\#$ over $[\mathfrak{p}]$ to the computation of the fibre of $\tilde{\phi}^\#$ over $[\mathfrak{p} R_{\mathfrak{p}}]$.

Now we would like to understand this latter fibre algebraically. Since $R_{\mathfrak{p}}$ is a local ring with unique maximal ideal $\mathfrak{p} R_{\mathfrak{p}}$, a prime ideal $\mathfrak{q} \subset T^{-1}S$ lies over $\mathfrak{p} R_{\mathfrak{p}}$ if and only if it contains the extended ideal $\mathfrak{p}(T^{-1}S)$. Thus the fibre is exactly the closed subset

$$\mathbb{V}(\mathfrak{p}(T^{-1}S)) \subset \operatorname{Spec}(T^{-1}S),$$

so

$$(\phi^{\#})^{-1}([\mathfrak{p}]) \cong \operatorname{Spec}(T^{-1}S / \mathfrak{p}(T^{-1}S)).$$

At first glance this quotient ring looks a bit mysterious, but tensor products package it very nicely. In fact there is a natural isomorphism $R_{\mathfrak{p}} \otimes_R S \cong T^{-1}S$. The *residue field* at $\mathfrak{p}$ is $\kappa(\mathfrak{p}) := R_{\mathfrak{p}} / \mathfrak{p} R_{\mathfrak{p}}$, which one checks is isomorphic to $\operatorname{Frac}(R / \mathfrak{p})$. (Recall $R / \mathfrak{p}$ is a domain since $\mathfrak{p}$ is a prime ideal.) There is then an isomorphism

$$\kappa(\mathfrak{p}) \otimes_R S \cong (R_{\mathfrak{p}} \otimes_R S) / \mathfrak{p}(R_{\mathfrak{p}} \otimes_R S) \cong T^{-1}S / \mathfrak{p}(T^{-1}S).$$

Therefore the fibre of $\phi^{\#}$ over $[\mathfrak{p}]$ is naturally homeomorphic to $\operatorname{Spec}(\kappa(\mathfrak{p}) \otimes_R S)$. The ring $\kappa(\mathfrak{p}) \otimes_R S$ is called the *fibre ring* of ϕ at $[\mathfrak{p}]$. The next exercise asks you to prove this carefully.

Problem 8. Residue Field.

Let R be a ring. Given a prime ideal $\mathfrak{p} \subset R$, the residue field of R at $\mathfrak{p}$ is the ring $R_{\mathfrak{p}} / \mathfrak{p} R_{\mathfrak{p}}$, which we denote by $\kappa(\mathfrak{p})$.

1. Explain why $\mathfrak{p} R_{\mathfrak{p}}$ is the unique maximal ideal of $R_{\mathfrak{p}}$. Deduce that $\kappa(\mathfrak{p})$ is a field.
2. Show that the composite map

$$R \xrightarrow{\eta} R_{\mathfrak{p}} \xrightarrow{\pi} \kappa(\mathfrak{p})$$

has kernel $\mathfrak{p}$. Deduce that it induces an injective ring map $R / \mathfrak{p} \hookrightarrow \kappa(\mathfrak{p})$.

3. Prove that every nonzero element of $R / \mathfrak{p}$ becomes invertible in $\kappa(\mathfrak{p})$. Conclude that $\kappa(\mathfrak{p}) \cong \text{Frac}(R / \mathfrak{p})$.
4. Use part (3) to compute the residue field $\kappa(\mathfrak{p})$ in each of the following cases. In each case it may be helpful to first identify the quotient ring $R / \mathfrak{p}$:

(a) $R = \mathbb{Z}$ and $\mathfrak{p} = \langle p \rangle$ for a prime p . (b) $R = \mathbb{Z}$ and $\mathfrak{p} = \langle 0 \rangle$. (c) $R = \mathbb{K}[x]$ and $\mathfrak{p} = \langle x - a \rangle$ for $a \in \mathbb{K}$. (d) $R = \mathbb{K}[x]$ and $\mathfrak{p} = \langle 0 \rangle$. (e) $R = \mathbb{R}[x]$ and $\mathfrak{p} = \langle x^2 + 1 \rangle$. (f) $R = \mathbb{K}[x, y]$ and $\mathfrak{p} = \langle y - x^2 \rangle$.	(g) $R = \mathbb{K}[x, y]$ and $\mathfrak{p} = \langle x, y \rangle$. (h) $R = \mathbb{Z}[x]$ and $\mathfrak{p} = \langle p, x \rangle$ for a prime p . (i) $R = \mathbb{Z}[i]$ and $\mathfrak{p} = \langle 1 + i \rangle$. (j) $R = \mathbb{Z}[\sqrt{-5}]$ and $\mathfrak{p} = \langle 2, 1 + \sqrt{-5} \rangle$.
--	--

Solution.

1. Prime ideals of $R_{\mathfrak{p}}$ correspond to prime ideals $\mathfrak{q} \subseteq R$ such that $\mathfrak{q} \subseteq \mathfrak{p}$, since $\mathfrak{p}$ is the largest such ideal it is maximal. $\kappa(\mathfrak{p})$ is a field since quotienting by a maximal ideal gives us a field.
2. Consider $(\pi \circ \eta)^{-1}(0)$.

$$\eta^{-1}(\pi^{-1}(0)) = \eta^{-1}(\mathfrak{p} R_{\mathfrak{p}}) = \mathfrak{p}$$

by primeness. Hence, $\mathfrak{p} = \ker(\pi \circ \eta)$. Then by the first isomorphism theorem,

$$R / \mathfrak{p} \hookrightarrow \kappa(\mathfrak{p})$$

3. Let $0 \neq f \in R / \mathfrak{p}$. Then $\eta(\pi(f)) \neq 0$. Since $\kappa(\mathfrak{p})$ is a field, every element is a unit and hence $\eta(\pi(f))$ is invertible. Thus, $\text{Frac}(R / \mathfrak{p}) \subseteq \kappa(\mathfrak{p})$.

Every element of $\kappa(\mathfrak{p}) = R_{\mathfrak{p}} / \mathfrak{p} R_{\mathfrak{p}}$ is the class of some a/s with $a \in R$, $s \in R \setminus \mathfrak{p}$, and this class equals $\iota(\bar{a}) \iota(\bar{s})^{-1}$ with $\bar{s} \neq 0$ (as $s \notin \mathfrak{p}$). That is exactly the image of $\bar{a}/\bar{s} \in \text{Frac}(R/\mathfrak{p})$. Hence there is an isomorphism, $\kappa(\mathfrak{p}) \cong \text{Frac}(R/\mathfrak{p})$.

4. (a) $R/\mathfrak{p} = \mathbb{Z}/\mathfrak{p}\mathbb{Z} = \mathbb{F}_{\mathfrak{p}} = \kappa(\mathfrak{p})$. (g) $R/\mathfrak{p} = \mathbb{K} = \kappa(\mathfrak{p})$.
- (b) $R/\mathfrak{p} = \mathbb{Z}$ and $\kappa(\mathfrak{p}) = \text{Frac}(\mathbb{Z}) = \mathbb{Q}$. (h) $R/\mathfrak{p} = \mathbb{Z}/\mathfrak{p}\mathbb{Z} = \mathbb{F}_{\mathfrak{p}} = \kappa(\mathfrak{p})$.
- (c) $R/\mathfrak{p} = \mathbb{K}[a] = \mathbb{K} = \kappa(\mathfrak{p})$.
- (d) $R/\mathfrak{p} = \mathbb{K}[x]$ and $\kappa(\mathfrak{p}) = \mathbb{K}(x)$. (i) $R/\mathfrak{p} = \mathbb{Z}/2\mathbb{Z} = \mathbb{F}_2 = \kappa(\mathfrak{p})$.
- (e) $R/\mathfrak{p} = \mathbb{R}[i] = \mathbb{C} = \kappa(\mathfrak{p})$. (j) $R/\mathfrak{p} = \mathbb{Z}/2\mathbb{Z}[\sqrt{-5}]/\langle 1 + \sqrt{5} \rangle = \mathbb{F}_2 = \kappa(\mathfrak{p})$.
- (f) $R/\mathfrak{p} = \mathbb{K}[x]$ and $\kappa(\mathfrak{p}) = \mathbb{K}(x)$.

Problem 9. Fibre Ring.

Let $\phi : R \rightarrow S$ be a ring map and fix a point $[\mathfrak{p}] \in \text{Spec}(R)$. Set $U := R \setminus \mathfrak{p}$ and $T := \phi(U) \subset S$. Let $\tilde{\phi} : R_{\mathfrak{p}} \rightarrow T^{-1}S$ be the map induced by the universal property of localization discussed above.

1. Prove that there is a natural isomorphism

$$R_{\mathfrak{p}} \otimes_R S \xrightarrow{\sim} T^{-1}S$$

$$\frac{r}{u} \otimes s \longmapsto \frac{\phi(r)s}{\phi(u)}.$$

2. Tensor the exact sequence

$$\mathfrak{p} R_{\mathfrak{p}} \rightarrow R_{\mathfrak{p}} \rightarrow \kappa(\mathfrak{p}) \rightarrow 0$$

with S over R to obtain an exact sequence

$$(\mathfrak{p} R_{\mathfrak{p}}) \otimes_R S \rightarrow R_{\mathfrak{p}} \otimes_R S \rightarrow \kappa(\mathfrak{p}) \otimes_R S \rightarrow 0.$$

Hint: View S as an R -module via ϕ , and use the right exactness of $- \otimes_R S$, which will be proved in Worksheet 5-1.

3. Show that the image of

$$(\mathfrak{p} R_{\mathfrak{p}}) \otimes_R S \rightarrow R_{\mathfrak{p}} \otimes_R S$$

is exactly the ideal $\mathfrak{p}(R_{\mathfrak{p}} \otimes_R S)$.

4. Deduce that there are natural isomorphisms

$$\kappa(\mathfrak{p}) \otimes_R S \cong (R_{\mathfrak{p}} \otimes_R S) / \mathfrak{p}(R_{\mathfrak{p}} \otimes_R S) \cong T^{-1}S / \mathfrak{p}(T^{-1}S).$$

Solution.

1. From the universal property of localization applied twice we have,

$$\begin{array}{ccc} R & \xrightarrow{\phi} & S \\ \eta \downarrow & \searrow f & \downarrow \\ R_{\mathfrak{p}} & \xrightarrow[\tilde{\phi}]{} & T^{-1}S \end{array}$$

Tensoring by $\otimes_R S$ gives us,

$$\begin{array}{ccc} R \otimes_R S & \xrightarrow{\phi \otimes_R \text{id}_S} & S \\ \eta \otimes_R \text{id}_S \downarrow & \searrow f \otimes_R \text{id}_S & \downarrow \\ R_{\mathfrak{p}} \otimes_R S & \xrightarrow[\tilde{\phi} \otimes_R \text{id}_S]{} & T^{-1}S \end{array}$$

Let $\tilde{\phi} \otimes_R \text{id}_S = \psi$.

ψ is injective: We show this for simple tensors. Let $\frac{\phi(r)s}{\phi(u)} = 0$. Then, there exists $t \in T$ such that $t\phi(r)s = 0$. Let $t = \phi(u)$ for some $u \in U$. Then, $\phi(vr)s = 0$. That is, in $R_{\mathfrak{p}} \otimes_R S$,

$$\frac{r}{u} \otimes s = \frac{vr}{vu} \otimes s = \frac{1}{vu} \otimes \phi(vr)s.$$

Since $\phi(vr)s = 0$,

$$\frac{1}{vu} \otimes \phi(vr)s = \frac{1}{vu} \otimes 0 = 0.$$

Therefore

$$\frac{r}{u} \otimes s = 0.$$

Thus $\ker(\psi) = 0$, and so ψ is injective.

ψ is surjective: An element in $T^{-1}S$ is of the form $\frac{s}{t}$, where $t = \phi(u)$ for some $u \in U$. Then,

$$\psi \left(\frac{1}{u} \otimes s \right) = \frac{1}{t} \otimes S = \frac{s}{t}$$

Hence ψ is surjective, and thus $R_{\mathfrak{p}} \otimes_R S \cong T^{-1}S$.

2. Since tensoring is right-exact, and the sequence

$$\mathfrak{p} R_{\mathfrak{p}} \longrightarrow R_{\mathfrak{p}} \longrightarrow \kappa(\mathfrak{p}) \longrightarrow 0$$

is exact, the obtained sequence is also exact,

$$(\mathfrak{p} R_{\mathfrak{p}}) \otimes_R S \xrightarrow{f} R_{\mathfrak{p}} \otimes_R S \xrightarrow{g} \kappa(\mathfrak{p}) \otimes_R S \longrightarrow 0.$$

3. Since the sequence is exact at $R_{\mathfrak{p}} \otimes_R S$, we have $\ker(g) = \text{im}(f)$. An element of $(\mathfrak{p} R_{\mathfrak{p}}) \otimes_R S$ is of the form $\frac{p}{u} \otimes s$ for $p \in \mathfrak{p}$ and $u \in R \setminus \mathfrak{p}$. Then,

$$\iota \otimes \text{id}_S \left(\frac{p}{u} \otimes s \right) = \frac{p}{u} \otimes s = p \left(\frac{1}{u} \otimes s \right) \in \mathfrak{p}(R_{\mathfrak{p}} \otimes_R S)$$

So we have $\text{im}(f) \subseteq \mathfrak{p}(R_{\mathfrak{p}} \otimes_R S)$. To see the other containment, let note that an element of $\mathfrak{p}(R_{\mathfrak{p}} \otimes_R S)$ is of the form $p \left(\frac{r}{u} \otimes s \right)$ where $p \in \mathfrak{p}$ and $\frac{r}{u} \otimes s$ is a simple tensor in $R_{\mathfrak{p}} \otimes_R S$. Since $pr \in \mathfrak{p}$, the tensor $p \left(\frac{r}{u} \otimes s \right) \in f \left(\frac{pr}{u} \otimes s \right) \in \text{im}(f)$.

Hence, $\text{im}(f) = \mathfrak{p}(R_{\mathfrak{p}} \otimes_R S)$.

4. By the First Isomorphism Theorem we have,

$$\kappa(\mathfrak{p}) \otimes_R S \cong R_{\mathfrak{p}} \otimes_R S / \ker(g)$$

and exactness gives us $\ker(g) = \text{im}(f)$, so,

$$\kappa(\mathfrak{p}) \otimes_R S \cong R_{\mathfrak{p}} \otimes_R S / \text{im}(f) = R_{\mathfrak{p}} \otimes_R S / \mathfrak{p}(R_{\mathfrak{p}} \otimes_R S)$$

Since we have an isomorphism $R_{\mathfrak{p}} \otimes_R S \cong T^{-1}S$, we also have

$$\kappa(\mathfrak{p}) \otimes_R S \cong R_{\mathfrak{p}} \otimes_R S / \mathfrak{p}(R_{\mathfrak{p}} \otimes_R S) \cong T^{-1}S / \mathfrak{p}(T^{-1}S)$$

Problem 10. Fibres of Spec Maps.

Let $\phi : R \rightarrow S$ be a ring map and fix a point $[\mathfrak{p}] \in \text{Spec}(R)$. Set $U := R \setminus \mathfrak{p}$ and $T := \phi(U) \subset S$. Let $\tilde{\phi} : R_{\mathfrak{p}} \rightarrow T^{-1}S$ be the map induced by the universal property of localization discussed above.

1. Show that the fibre $(\phi^{\#})^{-1}([\mathfrak{p}])$ is naturally homeomorphic to the fibre of $\tilde{\phi}^{\#} : \text{Spec}(T^{-1}S) \rightarrow \text{Spec}(R_{\mathfrak{p}})$ over the point $[\mathfrak{p} R_{\mathfrak{p}}]$.
2. Let $\mathfrak{q} \in \text{Spec}(T^{-1}S)$. Prove that $\tilde{\phi}^{-1}(\mathfrak{q}) = \mathfrak{p} R_{\mathfrak{p}}$ if and only if $\mathfrak{p}(T^{-1}S) \subseteq \mathfrak{q}$. Hint: Use that $R_{\mathfrak{p}}$ is a local ring with unique maximal ideal $\mathfrak{p} R_{\mathfrak{p}}$.

3. Deduce that

$$(\phi^{\#})^{-1}([\mathfrak{p}]) \cong \mathbb{V}(\mathfrak{p}(T^{-1}S)) \subset \text{Spec}(T^{-1}S),$$

and hence that there is a homeomorphism of topological spaces $(\phi^{\#})^{-1}([\mathfrak{p}]) \cong \text{Spec}(T^{-1}S / \mathfrak{p}(T^{-1}S))$.

4. Use Exercise 9 to conclude that $(\phi^{\#})^{-1}([\mathfrak{p}]) \cong \text{Spec}(\kappa(\mathfrak{p}) \otimes_R S)$.
5. Deduce that the fibre over $[\mathfrak{p}]$ is empty if and only if $\kappa(\mathfrak{p}) \otimes_R S = 0$.

Solution.

1. We have,

$$\begin{array}{ccc} R & \xrightarrow{\phi} & S \\ \eta \downarrow & \searrow f & \downarrow \eta_S \\ R_{\mathfrak{p}} & \xrightarrow[\tilde{\phi}]{} & T^{-1}S \end{array}$$

And the corresponding maps induced by Spec

$$\begin{array}{ccc} \text{Spec}(R) & \xleftarrow{\phi^{\#}} & \text{Spec}(S) \\ \eta^{\#} \uparrow & \nwarrow f^{\#} & \uparrow \eta_S^{\#} \\ \text{Spec}(R_{\mathfrak{p}}) & \xleftarrow[\tilde{\phi}^{\#}]{} & \text{Spec}(T^{-1}S) \end{array}$$

We know that both localization maps are homeomorphisms onto their image and also that $\mathfrak{p} R_{\mathfrak{p}} \in \text{Spec}(R_{\mathfrak{p}})$ corresponds to $\mathfrak{p} \in \text{Spec}(R)$, so the preimage of $\mathfrak{p} R_{\mathfrak{p}} \in \text{Spec}(R_{\mathfrak{p}})$ corresponds to the preimage of $\mathfrak{p} \in \text{Spec}(R)$ since the composition $\phi^{\#} \circ \eta_S^{\#} = \eta^{\#} \circ \tilde{\phi}^{\#}$. That is,

$$\mathfrak{q} \in (\tilde{\phi}^{\#})^{-1}([\mathfrak{p} R_{\mathfrak{p}}]) \iff \tilde{\phi}^{\#}(\mathfrak{q}) = [\mathfrak{p} R_{\mathfrak{p}}] \iff \eta^{\#}(\tilde{\phi}^{\#}(\mathfrak{q})) = [\mathfrak{p}] \iff \phi^{\#}(\eta_S^{\#}(\mathfrak{q})) = [\mathfrak{p}]$$

2. ($\implies$)

Assume $\tilde{\phi}^{-1}(\mathfrak{q}) = \mathfrak{p} R_{\mathfrak{p}}$. Since $\mathfrak{p}(T^{-1}S)$ is the extension of the contraction $\tilde{\phi}^{-1}(\mathfrak{q})$ we have $\mathfrak{p}(T^{-1}S) \subseteq \mathfrak{q}$.

($\impliedby$)

Assume $\mathfrak{p}(T^{-1}S) \subseteq \mathfrak{q}$. Then,

$$\tilde{\phi}^{-1}(\mathfrak{p}(T^{-1}S)) \subseteq \tilde{\phi}^{-1}(\mathfrak{q})$$

We know they are both ideals since pre-images of ideals are ideals. Since $R_{\mathfrak{p}}$ is a local ring with maximal ideal $\mathfrak{p} R_{\mathfrak{p}}$, the only proper ideal containing $\mathfrak{p} R_{\mathfrak{p}}$ is $\mathfrak{p} R_{\mathfrak{p}}$. So $\tilde{\phi}^{-1}(\mathfrak{q}) = \mathfrak{p} R_{\mathfrak{p}}$.

3. From parts 1 and 2 we have,

$$\begin{aligned} (\phi^{\#})^{-1}([\mathfrak{p}]) &\cong (\tilde{\phi}^{\#})^{-1}([\mathfrak{p} R_{\mathfrak{p}}]) \\ &= \{[\mathfrak{q}] \in \operatorname{Spec}(T^{-1}S) \mid \mathfrak{p}(T^{-1}S) \subseteq \mathfrak{q}\} \\ &= \mathbb{V}(\mathfrak{p} T^{-1}S) \cong \operatorname{Spec}(T^{-1}S / \mathfrak{p}(T^{-1}S)) \end{aligned}$$

4. From problem 9 we know that,

$$\kappa(\mathfrak{p}) \otimes_R S \cong T^{-1}S / \mathfrak{p}(T^{-1}S) \implies \operatorname{Spec}(\kappa(\mathfrak{p}) \otimes_R S) \cong (\phi^{\#})^{-1}([\mathfrak{p}])$$

5. Thus, the fibre over $[\mathfrak{p}]$ is empty if and only if $\kappa(\mathfrak{p}) \otimes_R S = 0$.

Problem 11. Computing Fibres of Spec Maps.

In this problem, we use the previous exercise to compute several examples:

1. Let $\phi : \mathbb{Z} \rightarrow \mathbb{Z}[x]$ be the inclusion. Compute the fibre of the induced map $\phi^\# : \text{Spec}(\mathbb{Z}[x]) \rightarrow \text{Spec}(\mathbb{Z})$ over the generic point $\langle 0 \rangle$ and over a closed point $\langle p \rangle$ for $p \in \mathbb{Z}$ prime.
2. Fix an integer $n \geq 2$, and consider the localization map $\eta : \mathbb{Z} \rightarrow \mathbb{Z}[\frac{1}{n}]$. For which points of $\text{Spec}(\mathbb{Z})$ is the fibre with respect to $\eta^\#$ empty? Compute the nonempty fibres.
3. Let $\mathbb{K}$ be a field and consider the map

$$\mathbb{K}[t] \longrightarrow \mathbb{K}[x, y], \quad t \longmapsto x.$$

Compute the fibre of the induced map over $\langle 0 \rangle \in \text{Spec}(\mathbb{K}[t])$ and the fibre over $\langle t - a \rangle$ for $a \in \mathbb{K}$.

4. Consider the inclusion of $\mathbb{C}$ -algebras $\mathbb{C}[t] \rightarrow \mathbb{C}[x, y, t] / \langle xy - t \rangle$ given by $t \mapsto t$.
 - (a) Compute the fibre over the generic point $\langle 0 \rangle$.
 - (b) Compute the fibre over $\langle t - a \rangle$ for $a \neq 0$.
 - (c) Compute the fibre over $\langle t \rangle$.
 - (d) Draw the corresponding map on closed points and compare it with the fibres above.

Solution.

1. We know that $(\phi^\#)^{-1}([q]) \cong \text{Spec}(\kappa(\mathfrak{p}) \otimes_R S)$. So in particular we have,

$$(\phi^\#)^{-1}([0]) \cong \text{Spec}(\kappa(0) \otimes_{\mathbb{Z}} \mathbb{Z}[x]) = \text{Spec}(\mathbb{Q} \otimes_{\mathbb{Z}} \mathbb{Z}[x]) \cong \text{Spec}(\mathbb{Q}[x])$$

And,

$$(\phi^\#)^{-1}([p]) \cong \text{Spec}(\kappa(\mathfrak{p}) \otimes_{\mathbb{Z}} \mathbb{Z}[x]) = \text{Spec}(\mathbb{F}_p \otimes_{\mathbb{Z}} \mathbb{Z}[x]) \cong \text{Spec}(\mathbb{F}_p[x])$$

2. Let $n \geq 2$. We know that ideals $\mathfrak{q}$ in $\text{Spec}(\mathbb{Z}[\frac{1}{n}])$ are in bijection with ideals $\mathfrak{p}$ in $\text{Spec}(\mathbb{Z})$ such that $n \notin \mathfrak{p}$. So the fibre over all $\mathfrak{p} \in \mathbb{Z}$ such that $\mathfrak{p} \mid n$ is empty. And the non-empty fibres correspond to $\mathfrak{p} \in \mathbb{Z}$ such that $\mathfrak{p} \nmid n$, so

$$(\phi^\#)^{-1}([p]) \cong \text{Spec} \left(\kappa(\mathfrak{p}) \otimes_{\mathbb{Z}} \mathbb{Z} \left[\frac{1}{n} \right] \right) = \text{Spec}(\mathbb{F}_p)$$

And for $\langle 0 \rangle$ we have,

$$(\phi^\#)^{-1}([0]) \cong \operatorname{Spec} \left(\kappa(0) \otimes_{\mathbb{Z}} \mathbb{Z} \left[\frac{1}{n} \right] \right) = \operatorname{Spec}(\mathbb{Q})$$

3. We have,

$$(\phi^\#)^{-1}([0]) \cong \operatorname{Spec} \left(\kappa(0) \otimes_{\mathbb{K}[t]} \mathbb{K}[x, y] \right) = \operatorname{Spec} \left(\mathbb{K}(t) \otimes_{\mathbb{K}[t]} \mathbb{K}[x, y] \right) \cong \operatorname{Spec}(\mathbb{K}[y](x))$$

And,

$$\begin{aligned} (\phi^\#)^{-1}([t - a]) &\cong \operatorname{Spec} \left(\kappa(a) \otimes_{\mathbb{K}[t]} \mathbb{K}[x, y] / \langle x - a \rangle \right) = \operatorname{Spec} \left(\mathbb{K} \otimes_{\mathbb{K}[t]} \mathbb{K}[x, y] \right) \cong \\ &\operatorname{Spec}(\mathbb{K}[x, y] / \langle x - a \rangle) \cong \operatorname{Spec}(\mathbb{K}[y]) \end{aligned}$$

4. (a)

$$\begin{aligned} (\phi^\#)^{-1}([0]) &\cong \operatorname{Spec} \left((\kappa(0) \otimes_{\mathbb{C}[t]} \mathbb{C}[x, y, t]) / \langle xy - t \rangle \right) \\ &\cong \operatorname{Spec} \left((\mathbb{C}(t) \otimes_{\mathbb{C}[t]} \mathbb{C}[x, y, t]) / \langle xy - t \rangle \right) \\ &\cong \operatorname{Spec}(\mathbb{C}[x, y](t) / \langle xy - t \rangle) \cong \operatorname{Spec}(\mathbb{C}(t)[x, x^{-1}]) \end{aligned}$$

(b)

$$\begin{aligned} (\phi^\#)^{-1}([t - a]) &\cong \operatorname{Spec} \left((\kappa(a) \otimes_{\mathbb{C}[t]} \mathbb{C}[x, y, t]) / \langle xy - t \rangle \right) \\ &\cong \operatorname{Spec} \left((\mathbb{C} \otimes_{\mathbb{C}[t]} \mathbb{C}[x, y]) / \langle xy - a \rangle \right) \\ &\cong \operatorname{Spec}(\mathbb{C}[x, y] / \langle xy - a \rangle) \cong \operatorname{Spec}(\mathbb{C}[x, x^{-1}]) \end{aligned}$$

(c)

$$\begin{aligned} (\phi^\#)^{-1}([t]) &\cong \operatorname{Spec} \left((\kappa(t) \otimes_{\mathbb{C}[t]} \mathbb{C}[x, y, t]) / \langle xy - t \rangle \right) \\ &\cong \operatorname{Spec} \left((\mathbb{C}(t) \otimes_{\mathbb{C}[t]} \mathbb{C}[x, y]) / \langle xy - t \rangle \right) \\ &\cong \operatorname{Spec}((\mathbb{C}[x, y]) / \langle xy \rangle) \end{aligned}$$

Throughout this course, “ring” means *commutative* ring with unity. In the previous worksheets we introduced modules, homomorphisms, quotient modules, free modules, Hom, tensor products, and localization. The goal of this worksheet is to organize these constructions around one central idea: *exactness*. Exactness is a way to say that one map accounts for exactly the elements killed by the next map. It packages kernels, images, quotient modules, presentations, and many theorems into one flexible language.

Let R be a ring. A *sequence* of R -modules is $\dots, M_0, M_1, M_2, \dots$ together with R -module homomorphisms $\phi_i : M_i \rightarrow M_{i-1}$. We often arrange a sequence into a diagram like

$$\dots \xrightarrow{\phi_{i+1}} M_i \xrightarrow{\phi_i} M_{i-1} \xrightarrow{\phi_{i-1}} M_{i-2} \longrightarrow \dots$$

We are most interested in sequences when the maps are related in some nice way that allows us to compare the image of one map with the kernel of the next. A *chain complex*, or simply a *complex*, of R -modules is a sequence in which $\phi_{i-1} \circ \phi_i = 0$ for every i . Equivalently, $\text{img}(\phi_i) \subset \ker(\phi_{i-1})$ for all i . We often write $(M_\bullet, \phi_\bullet)$ for a complex and call the maps ϕ_i *differentials*. A complex is *exact at* M_i if $\text{img}(\phi_{i+1}) = \ker(\phi_i)$. One way to measure the failure, or success, of a complex to be exact at M_i is by looking at the *i-th homology group* of $(M_\bullet, \phi_\bullet)$, which is defined to be the quotient R -module:

$$H_i(M_\bullet, \phi_\bullet) := \frac{\ker(\phi_i)}{\text{img}(\phi_{i+1})}.$$

The definition of a complex guarantees this quotient makes sense, since $\text{img}(\phi_{i+1}) \subset \ker(\phi_i)$. Equivalently, a complex is exact at M_i if and only if $H_i(M_\bullet, \phi_\bullet) = 0$. (Here 0 denotes the zero R -module.) A complex is *exact* if it is exact at every module where this condition makes sense. Thus every exact sequence is a complex, but a complex need not be exact.

Complexes appear throughout commutative algebra and other areas of mathematics. Perhaps the most important special case of a complex is a short exact sequence. A *short sequence* is a complex of the form

$$0 \longrightarrow A \xrightarrow{i} B \xrightarrow{p} C \longrightarrow 0.$$

It is called a *short exact sequence* if it is exact at A , B , and C . Strictly speaking, a complex extends infinitely in both directions, so the diagram above is shorthand for

$$\dots \longrightarrow 0 \longrightarrow A \xrightarrow{i} B \xrightarrow{p} C \longrightarrow 0 \longrightarrow \dots,$$

where all modules outside the displayed range are zero. We will follow the standard convention of suppressing these trailing and leading zero modules. Saying that this sequence is exact is the same as saying that i is injective, p is surjective, and $\text{img}(i) = \ker(p)$. Thus a short exact sequence says that A is being identified with a submodule of B , and that the quotient of B by this submodule is isomorphic to C .

Problem 1. First Encounters with Exactness.

Let A, B, C be R -modules.

1. Prove that a sequence

$$0 \longrightarrow A \xrightarrow{i} B$$

is exact at A if and only if i is injective.

2. Prove that a sequence

$$B \xrightarrow{p} C \longrightarrow 0$$

is exact at C if and only if p is surjective.

3. Prove carefully that

$$0 \longrightarrow A \xrightarrow{i} B \xrightarrow{p} C \longrightarrow 0$$

is exact if and only if i is injective, p is surjective, and $\text{img}(i) = \ker(p)$.

4. If

$$A \xrightarrow{f} B \xrightarrow{g} C$$

is exact at B , prove that $g \circ f = 0$.

5. Give an example of maps of R -modules

$$A \xrightarrow{f} B \xrightarrow{g} C$$

such that $g \circ f = 0$ but the sequence is not exact at B .

6. Consider the complex below. Compute the homology at B in each of the following cases:

$$A \xrightarrow{f} B \xrightarrow{g} C$$

- (a) $f = 0$ and $g = 0$.
- (b) f is injective and $g = 0$.
- (c) g is injective.

Solution.

1. ($\implies$)

Assume the sequence is exact at A . Then image of the 0 map is equal to $\ker(i)$. But the image of 0 under any map is 0. So we have $\ker(i) = \{0\} \implies i$ is injective.

($\Leftarrow$)

Assume i is injective. Then $\ker(i) = 0 = \text{im}(0)$, thus making the sequence exact at A .

2. ($\implies$)

Assume the sequence is exact at C . Then $\text{im}(p) = C$, thus p is surjective.

($\Leftarrow$)

Assume p is surjective. Then $\text{im}(p) = C$ which is the kernel of the map from $C \rightarrow 0$ thus making the sequence exact at C .

3. ($\implies$)

Assume the sequence is exact. Then it is exact at A and C . From parts (1) and (2) this implies that i is injective, p is surjective. Since it is also exact at B we must also have $\ker(p) = \text{im}(i)$.

($\Leftarrow$)

Assume i is injective, p is surjective, and $\text{im}(i) = \ker(p)$. Then by parts (1), (2), the sequence is exact at A and C . And since $\text{im}(i) = \ker(p)$, the sequence is exact at B by definition. Thus the sequence is exact.

4. Let $a \in A$. Then $f(a) \in \text{im}(f)$, and since the sequence is exact we have $\text{im}(f) = \ker(g)$. So, $f(a) \in \ker(g) \implies g(f(a)) = 0$. Since a was arbitrary this implies that $g \circ f = 0$.

5. Consider the $\mathbb{Z}$ -modules $A = C = \mathbb{Z}$, $B = \mathbb{Z}^2$ and the morphisms $f = 0$ and $g = \pi$. Then, we have $g \circ f = 0$ but $\ker(g) \subset \text{im}(f)$, so the sequence is not exact.

6. (a) Note that $\text{im}(f) = \{0\}$, $\ker(g) = B$. Then,

$$H = \ker(g) / \text{im}(f) = B / \{0\} = B$$

(b) Note that $\text{im}(f) \cong A$, $\ker(g) = B$. Then,

$$H = \ker(g) / \text{im}(f) \cong B / A$$

(c) Note that $\ker(g) = \{0\}$. Then,

$$H = \ker(g) / \text{im}(f) = \{0\} / \text{im}(f) = \{0\}$$

Problem 2. Examples of Short Exact Sequences.

Let R be a ring. In this problem we will see that a number of constructions we are quite familiar with can be framed in terms of short exact sequences.

1. Let $I \subset R$ be an ideal. Construct a short exact sequence

$$0 \longrightarrow I \xrightarrow{\iota} R \xrightarrow{\pi} R/I \longrightarrow 0$$

where π is the natural quotient map and ι is the inclusion I into R .

2. Let $N \subset M$ be a submodule. Prove that the sequence

$$0 \longrightarrow N \xrightarrow{\iota} M \xrightarrow{\pi} M/N \longrightarrow 0$$

is exact, where ι is the inclusion of N into M .

3. Let $I \subset J \subset R$ be ideals. Construct a short exact sequence

$$0 \longrightarrow J/I \longrightarrow R/I \longrightarrow R/J \longrightarrow 0.$$

Hint: Define $J/I \rightarrow R/I$ by $j + I \mapsto j + I$ and define $R/I \rightarrow R/J$ by $r + I \mapsto r + J$. Check well-definedness before checking exactness.

4. For a positive integer n , prove that multiplication by n gives

$$0 \longrightarrow \mathbb{Z} \xrightarrow{\times n} \mathbb{Z} \longrightarrow \mathbb{Z}/n\mathbb{Z} \longrightarrow 0$$

which is a short exact sequence of $\mathbb{Z}$ -modules.

Solution.

1.

2. Note that $\text{im}(\iota) \cong N$ and $\ker(\pi) = N$. That is, $\text{im}(\iota) = \ker(\pi)$. Moreover since ι is injective and π is surjective the sequence is exact (using Problem 1.3).

3. We define the maps ι and π as follows:

- $\iota : J/I \rightarrow R/I$ where $\iota(j + I) = j + I$.
- $\pi : R/I \rightarrow R/J$ where $\pi(r + I) = r + J$.

$$0 \longrightarrow J/I \xrightarrow{\iota} R/I \xrightarrow{\pi} R/J \longrightarrow 0.$$

The sequence is exact because ι is injective, π is surjective, and $\text{img}(\iota) \cong J/I = \ker(\pi)$ by the first and third isomorphism theorems.

4. The map $\times n$ is injective because $\ker(\times n) = \{0\}$, $\mathbb{Z} \rightarrow \mathbb{Z}/n\mathbb{Z}$ is surjective because it is the natural quotient map, and $\text{im}(\times n) = \langle n \rangle = \ker(\mathbb{Z} \rightarrow \mathbb{Z}/n\mathbb{Z})$, thus the sequence is exact.

Problem 3. Kernels, Cokernels, and Images.

Let $\phi : A \rightarrow B$ be a homomorphism of R -modules. The *cokernel* of ϕ is the R -module $\text{coker}(\phi) := B / \text{img}(\phi)$.

1. Prove that the sequence

$$0 \longrightarrow \ker(\phi) \xrightarrow{\iota} A \xrightarrow{\phi} B$$

where ι is the inclusion of $\ker(\phi)$ into A is exact.

2. Suppose

$$0 \longrightarrow A \xrightarrow{f} B \xrightarrow{g} C$$

is an exact sequence of R -modules. Prove that f induces an isomorphism $A \cong \ker(g)$.

3. Prove that the sequence

$$A \xrightarrow{\phi} B \xrightarrow{\pi} \text{coker}(\phi) \longrightarrow 0$$

where π is the natural quotient map is exact.

4. Suppose

$$A \xrightarrow{f} B \xrightarrow{g} C \longrightarrow 0$$

is an exact sequence of R -modules. Prove that g induces an isomorphism $\text{coker}(f) \cong C$.

5. Deduce there is an exact sequence of R -modules

$$0 \longrightarrow \ker(\phi) \xrightarrow{\iota} A \xrightarrow{\phi} B \xrightarrow{\pi} \text{coker}(\phi) \longrightarrow 0 .$$

6. Prove that ϕ induces an isomorphism $A / \ker(\phi) \cong \text{img}(\phi)$, and deduce that the exact sequence in part (5) factors through $\text{img}(\phi)$ as two short exact sequences

$$0 \longrightarrow \ker(\phi) \xrightarrow{\iota} A \xrightarrow{\bar{\phi}} \text{img}(\phi) \longrightarrow 0$$

and

$$0 \longrightarrow \text{img}(\phi) \xrightarrow{j} B \xrightarrow{\pi} \text{coker}(\phi) \longrightarrow 0$$

where $\bar{\phi}$ is ϕ with codomain restricted to $\text{img}(\phi)$ and j is the inclusion.

Solution.

1. • $\text{im}(0 \rightarrow \ker(\phi)) = 0 = \ker(\iota)$ (inclusion map)
 • $\text{im}(\iota) = \ker(\phi)$

Thus, the sequence is exact.

2. If the sequence is exact, then we have $\text{im}(0 \rightarrow A) = \{0\} = \ker(f)$ and $\text{im}(f) = \ker(g)$. Now we define $\psi : A \rightarrow \ker(g)$ where $\psi(a) = f(a)$. This is well-defined since $\text{im}(f) = \ker(g)$.

- ψ is injective since $\psi(a) = 0 \iff f(a) = 0$ that is $\ker(\psi) = \ker(f)$ but $\ker(f) = \{0\}$.
- ψ is surjective; since $\text{im}(f) = \ker(g)$ we have for any $b \in \ker(g)$ we have $b = f(a) = \psi(a)$ for some $a \in A$.

Thus we have a bijection $\psi : A \rightarrow \ker(g)$ giving us $A \cong \ker(g)$.

3. • $\text{im}(\phi) = \ker(\pi)$ (by definition)
 • $\text{im}(\pi) = \text{coker}(\phi) = \ker(\text{coker}(\phi) \rightarrow 0)$

Thus, the sequence is exact.

4. If the sequence is exact, then we have $\text{im}(f) = \ker(g)$ and $\text{im}(g) = \ker(C \rightarrow 0) = C$. By definition $\text{coker}(f) = B / \text{im}(f)$ Now we define a map $\psi : B / \text{im}(f) \rightarrow C$ by $\psi(\bar{b}) = g(b)$. Then,

- ψ is well-defined since if $\bar{b} = \bar{b}'$ then $b - b' \in \text{im}(f) = \ker(g)$, so $g(b) = g(b')$.
- ψ is injective since if $\psi(\bar{b}) = g(b) = 0 \implies b \in \text{im}(f) \implies \bar{b} = 0$.
- ψ is surjective since $\text{im}(g) = C$.

Thus we have a bijection $\psi : B / \text{im}(f) \rightarrow C$ giving us $\text{coker}(f) \cong C$.

5. From parts (1) and (3) we can conclude that the sequence is exact.

6. We define $\psi : A / \ker(\phi) \rightarrow \text{img}(\phi)$ by $\psi(\bar{a}) = \phi(a)$. Then,

- ψ is well-defined since if $\bar{a} = \bar{a}'$ then $a - a' \in \ker(\phi) \implies \psi(a - a') = 0 \implies \psi(a) = \psi(a')$.
- ψ is injective since if $\psi(\bar{a}) = \phi(a) = 0 \implies a \in \ker(\phi) \implies \bar{a} = 0$.
- ψ is surjective since $\text{im}(\psi) = \text{im}(\phi)$.

Thus we have a bijection $\psi : A / \ker(\phi) \rightarrow \text{im}(\phi)$ giving us $A / \ker(\phi) \cong \text{im}(\phi)$.

Then we have note that the following sequences are exact:

$$0 \longrightarrow \ker(\phi) \xrightarrow{\iota} A \xrightarrow{\bar{\phi}} \text{img}(\phi) \longrightarrow 0$$

- At $\ker(\phi)$: since the inclusion map is injective we have $\text{im}(0 \rightarrow \ker(\phi)) = 0 = \ker(\iota)$.
- At A : $\text{im}(\iota) \cong \ker(\phi) = \ker(\bar{\phi})$
- At $\text{img}(\phi)$: $\bar{\phi}$ is surjective, and $\ker(\text{im}(\phi) \rightarrow 0) = \text{im}(\phi) = \text{im}(\bar{\phi})$.

And

$$0 \longrightarrow \text{img}(\phi) \xrightarrow{j} B \xrightarrow{\pi} \text{coker}(\phi) \longrightarrow 0$$

- At $\text{img}(\phi)$: since the inclusion map j is injective we have $\text{im}(0 \rightarrow \text{im}(\phi)) = 0 = \ker(j)$.
- At B : $\text{im}(j) \cong \text{im}(\phi) = \ker(\pi)$ since $\text{coker}(\phi) = B / \text{im}(\phi)$
- At $\text{coker}(\phi)$: π is surjective by definition, and $\ker(\text{coker}(\phi) \rightarrow 0) = \text{coker}(\phi) = \text{im}(\pi)$.

Thus, the exact sequence in part (5) factors through $\text{img}(\phi)$.

Problem 4. Presentations of Modules.

Let R be a ring. A *presentation* of an R -module M is an exact sequence of the form

$$F_1 \xrightarrow{\varphi} F_0 \longrightarrow M \longrightarrow 0$$

where F_0 and F_1 are free modules.

1. Explain why a presentation of M as above says that M is obtained from the generators of F_0 by imposing the relations coming from $\varphi(F_1)$. Hint: Use exactness to identify M with $F_0 / \text{img}(\varphi)$.
2. Let $R = \mathbb{K}[x]$ and $M = R / \langle x^3 \rangle$. Write a presentation of M using finite free R -modules.
3. Let $R = \mathbb{K}[x, y]$ and $M = R / \langle x^2, xy, y^2 \rangle$. Write a presentation of M using one free generator and three relations.
4. Let A be an $m \times n$ matrix with entries in R , viewed as a map $R^n \rightarrow R^m$. Interpret the sequence

$$R^n \xrightarrow{A} R^m \longrightarrow \text{coker}(A) \longrightarrow 0$$

as a presentation of $\text{coker}(A)$.

Solution.

1. From problem 3.4 we know that $\text{coker}(\varphi) = F_0 / \text{im}(\varphi) \cong M$. And $F_0 / \text{im}(\varphi)$ is determined by what quotienting by $\text{im}(\varphi)$ does to F_0 's distinguished elements. Since the sequence is exact at F_0 , we have $\ker(F_0 \rightarrow M) = \text{im}(\varphi) = \varphi(F_1)$, so these are precisely the relations between the generators. Thus, $M = F_0 / \text{im}(\varphi) = F_0 / \varphi(F_1)$.

2.

$$\mathbb{K}[x] \xrightarrow{\varphi} \mathbb{K}[x] \longrightarrow \mathbb{K}[x] / \langle x^3 \rangle \longrightarrow 0$$

where $\varphi(e_1) = x^3$.

3.

$$(\mathbb{K}[x])^3 \xrightarrow{\varphi} \mathbb{K}[x] \longrightarrow \mathbb{K}[x] / \langle x^2, xy, y^2 \rangle \longrightarrow 0$$

where $\varphi(e_1) = x^2$, $\varphi(e_2) = xy$, $\varphi(e_3) = y^2$.

4. We have $\text{coker}(A) = R^m / \text{im}(A)$, which is a module with m -generators and n relations (the n columns of A give n relations).

We just saw that a short exact sequence $0 \rightarrow A \xrightarrow{i} B \xrightarrow{p} C \rightarrow 0$ of R -modules says that B has a submodule isomorphic to A (namely $\text{img } i$) whose quotient is isomorphic to C (namely $B/\text{img } i \cong C$). One might hope that knowing A and C determines B up to isomorphism, and in the simplest cases it does: $B \cong A \oplus C$. When this happens – and when the isomorphism is compatible with the maps i and p – the sequence is called *split*. More precisely, a short exact sequence is *split* (also called *split exact*) if there exists an isomorphism of R -modules $h : B \rightarrow A \oplus C$ such that the following diagram commutes

$$\begin{array}{ccccccccc} 0 & \longrightarrow & A & \xrightarrow{i} & B & \xrightarrow{p} & C & \longrightarrow & 0 \\ & & \parallel & & \downarrow h & & \parallel & & \\ 0 & \longrightarrow & A & \xrightarrow{\iota} & A \oplus C & \xrightarrow{\pi} & C & \longrightarrow & 0 \end{array}$$

where $\iota(a) = (a, 0)$ is the natural inclusion and $\pi(a, c) = c$ is the natural projection. Saying that the diagram commutes means that the two squares commute. This is the generic situation for vector spaces: every subspace has a complement, so every short exact sequence of vector spaces splits. Over more general rings, however, splitting is the exception rather than the rule.

We can characterize when a short exact sequence splits in terms of the existence of either a section of p or a retraction of i . A *section* of $p : B \rightarrow C$ is an R -module homomorphism $s : C \rightarrow B$ such that $p \circ s = \text{Id}_C$. A *retraction* of $i : A \rightarrow B$ is an R -module homomorphism $r : B \rightarrow A$ such that $r \circ i = \text{Id}_A$. Note that in general s is not an inverse to p , and r is not an inverse to i ; the compositions in the other order need not be the identity.

Lemma 0.14 (Splitting Lemma). *Let R be a ring, and consider a short exact sequence of R -modules*

$$0 \longrightarrow A \xrightarrow{i} B \xrightarrow{p} C \longrightarrow 0.$$

The following are equivalent:

- (i) *The sequence splits, i.e. $B \cong A \oplus C$ in a way compatible with i and p .*
- (ii) *The map p admits a section $s : C \rightarrow B$.*
- (iii) *The map i admits a retraction $r : B \rightarrow A$.*

Condition (ii) is often called *right split*, while condition (iii) is called *left split*. Thus, for R -modules, “right split,” “left split,” and “split” are all equivalent. Warning: This equivalence is special to the setting of R -modules (more generally, abelian categories). For example, in the category of non-abelian groups, a sequence may admit a section without being a direct product: such extensions correspond to semidirect products. Fortunately, none of these issues arise for R -modules.

Problem 5. Split Short Exact Sequences.

Let

$$0 \longrightarrow A \xrightarrow{i} B \xrightarrow{p} C \longrightarrow 0$$

be a short exact sequence of R -modules.

1. Suppose there exists a homomorphism $s : C \rightarrow B$ such that $p \circ s = \text{Id}_C$. Prove that the map defined below is an isomorphism of R -modules:

$$\Phi : A \oplus C \longrightarrow B, \quad (a, c) \longmapsto i(a) + s(c).$$

2. Conversely, suppose there is an isomorphism $B \cong A \oplus C$ under which i becomes the standard inclusion $A \rightarrow A \oplus C$ and p becomes the standard projection $A \oplus C \rightarrow C$. Construct a homomorphism $s : C \rightarrow B$ with $p \circ s = \text{Id}_C$.
3. Suppose there exists a homomorphism $r : B \rightarrow A$ such that $r \circ i = \text{Id}_A$.

- (a) Prove that the map

$$B \longrightarrow A \oplus C, \quad b \longmapsto (r(b), p(b))$$

is injective.

- (b) Prove that it is surjective. Hint: Given $(a, c) \in A \oplus C$, choose $b_0 \in B$ with $p(b_0) = c$ and modify b_0 by an element of $i(A)$.
 - (c) Conclude that $B \cong A \oplus C$.
4. Conversely, suppose there is an isomorphism $B \cong A \oplus C$ under which i becomes the standard inclusion $a \mapsto (a, 0)$ and p becomes the standard projection $(a, c) \mapsto c$. Construct a homomorphism $r : B \rightarrow A$ with $r \circ i = \text{Id}_A$.

Solution.

1. Φ is an R -module homomorphism:

- $\Phi((a + b, c + d)) = i(a + b) + s(c + d) = i(a) + i(b) + s(c) + s(d) = \Phi(a, c) + \Phi(b, d)$.
- $\Phi(r(a, c)) = i(ra) + s(rc) = ri(a) + rs(c) = r(i(a) + s(c)) = r(\Phi(a, c))$.

Φ is injective:

Suppose $\Phi((a, c)) = 0$, then $i(a) + s(c) = 0$. We apply p to both sides. $p(\Phi(a, c)) = p(i(a) + s(c)) = p(i(a)) + c$. But $i(a) \in \ker(p)$, so we have $p(i(a)) + c = 0 + c = 0$. Hence, $c = 0$. Substituting back, $i(a) + s(0) = 0 \implies i(a) = 0 \implies a = 0$ since i is injective. Thus, if $\Phi((a, c)) = 0$ then $(a, c) = (0, 0)$.

Φ is surjective:

Let $b \in B$. Then we have, $p(b) = c$ for some $c \in C$. Now consider $p(b - s(c))$,

$$p(b - s(c)) = p(b) - p(s(c)) = c - c = 0$$

That is $b - s(c) \in \ker(p) = \text{im}(A)$, i.e. there exists a $a \in A$ such that $i(a) = b - s(c)$. Upon re-arranging we get,

$$b = i(a) + s(c) = \Phi((a, c))$$

Thus the map Φ is an isomorphism, and $A \oplus C \cong B$.

2. Let $\phi : B \rightarrow A \oplus C$ be the isomorphism, and let $j : C \rightarrow A \oplus C$ be the standard inclusion. Then we define $s : C \rightarrow B$ as $s(c) = \phi^{-1}(j(c)) = \phi^{-1}((0, c))$ for all $c \in C$. Then

$$p \circ s = p \circ \phi^{-1} \circ j = \pi \circ j = \text{id}_C$$

3. (a) Let ϕ denote the map from $B \rightarrow A \oplus C$. Then, if $\phi(b) = (0, 0) \implies (r(b), p(b)) = (0, 0)$.

- $p(b) = 0 \implies b \in \ker(p) = \text{im}(i)$. So $b = i(a)$ for some a .
- $r(b) = 0 \implies r(i(a)) = 0$ but $r \circ i = \text{id}_A$ so $a = 0$.

Thus the map ϕ is injective.

- (b) Let $(a, c) \in A \oplus C$ be an arbitrary element. Since p is surjective we have $c = p(b)$ for some $b \in B$. Let $b' = b - i(r(b)) + i(a)$. Then $\phi(b') = (r(b'), p(b'))$. Where,

- $r(b') = r(b) - r(i(r(b))) + r(i(a)) = r(b) - r(b) + a = a$
- $p(b') = p(b) - p(i(r(b))) + p(i(a)) = c + 0 + 0 + c$ since $\text{im}(i) = \ker(p)$.

Thus the map ϕ is surjective.

- (c) Therefore we conclude $B \cong A \oplus C$.

4. Let $\phi : B \rightarrow A \oplus C$ be the isomorphism. Then we define $r = \pi_A \circ \phi$, where $\pi_A : A \oplus C \rightarrow A$ be the standard projection to A . Then,

$$r \circ i = \pi_A \circ \phi \circ i = \pi_A \circ \iota = \text{id}_A$$

Problem 6. Examples of Split & Non-Split Exact Sequences.

1. Prove that every short exact sequence of vector spaces over a field $\mathbb{K}$ splits.
Hint: Choose a basis for the first term, extend it to a basis for the middle term, and identify the remaining basis vectors with a basis for the third term.
2. Explain why the Rank-Nullity theorem for finite dimensional vector spaces over $\mathbb{K}$ is a consequence of the splitting theorem.
3. Prove that the sequence

$$0 \longrightarrow \mathbb{Z} \xrightarrow{\cdot 2} \mathbb{Z} \longrightarrow \mathbb{Z}/2\mathbb{Z} \longrightarrow 0$$

does not split as a sequence of $\mathbb{Z}$ -modules.

4. Let p be a prime number. Prove that the sequence of $\mathbb{Z}$ -modules:

$$0 \longrightarrow \mathbb{Z}/p\mathbb{Z} \longrightarrow \mathbb{Z}/p^2\mathbb{Z} \longrightarrow \mathbb{Z}/p\mathbb{Z} \longrightarrow 0,$$

where the first map sends $\bar{1} \mapsto \bar{p}$ and the second is reduction modulo p , does not split. Hint: A section would send $\bar{1} \in \mathbb{Z}/p\mathbb{Z}$ to an element of $\mathbb{Z}/p^2\mathbb{Z}$ whose reduction modulo p is $\bar{1}$. What is the order of such an element?

5. Let $R = \mathbb{K}[x]$. Prove that the sequence below is exact, but does not split as a sequence of R -modules:

$$0 \longrightarrow \langle x \rangle \longrightarrow \mathbb{K}[x] \longrightarrow \mathbb{K} \longrightarrow 0.$$

Hint: If a section $s : \mathbb{K} \rightarrow \mathbb{K}[x]$ existed as a $\mathbb{K}[x]$ -module map, what could $s(1)$ be? Use the fact that $x \cdot 1 = 0$ in $\mathbb{K} \cong \mathbb{K}[x]/\langle x \rangle$.

6. Let R be a ring and let $e \in R$ be an idempotent, meaning $e^2 = e$. Prove that

$$0 \longrightarrow eR \hookrightarrow R \twoheadrightarrow R/eR \longrightarrow 0$$

splits. Then prove that the map $R \rightarrow (1-e)R$ given by $r \mapsto (1-e)r$ induces an isomorphism $R/eR \cong (1-e)R$.

Solution.

1.

We now introduce just enough categorical language to talk cleanly about Hom, tensor product, and localization. A *category* is a collection of objects together with, for each pair of objects, a set of morphisms between them, equipped with an associative composition law and an identity morphism for each object. The quintessential category, both in this class and arguably beyond, is the category of R -modules, denoted $R - \mathbf{Mod}$, whose objects are R -modules and whose morphisms are R -module homomorphisms.

A *functor* between categories $\mathbf{C}$ and $\mathbf{D}$ is a mapping that assigns to each object of $\mathbf{C}$ an object of $\mathbf{D}$ and to each morphism in $\mathbf{C}$ a morphism in $\mathbf{D}$, preserving identity morphisms and composition. A functor is contravariant if it reverses the order of composition and covariant if it preserves the order of composition. More precisely, a *covariant functor* $F : \mathbf{C} \rightarrow \mathbf{D}$ is the data of

1. for every object X of $\mathbf{C}$, $F(X)$ is an object of $\mathbf{D}$,
2. for every morphism $f : X \rightarrow Y$ in $\mathbf{C}$, $F(f) : F(X) \rightarrow F(Y)$ is a morphism in $\mathbf{D}$,

which satisfies the following conditions:

- (i) $F(\text{Id}_X) = \text{Id}_{F(X)}$ for every object X of $\mathbf{C}$,
- (ii) $F(g \circ f) = F(g) \circ F(f)$ for all composable morphisms $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ in $\mathbf{C}$.

Similarly a *contravariant functor* $F : \mathbf{C} \rightarrow \mathbf{D}$ is the data of:

1. for every object X of $\mathbf{C}$, $F(X)$ is an object of $\mathbf{D}$,
2. for every morphism $f : X \rightarrow Y$ in $\mathbf{C}$, $F(f) : F(Y) \rightarrow F(X)$ is a morphism in $\mathbf{D}$,

which satisfies the following conditions:

- (i) $F(\text{Id}_X) = \text{Id}_{F(X)}$ for every object X of $\mathbf{C}$,
- (ii) $F(g \circ f) = F(f) \circ F(g)$ for all composable morphisms $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ in $\mathbf{C}$.

We have already constructed several operations that take modules as input and produce new modules. For example, after fixing an R -module T , the assignments

$$M \longmapsto \text{Hom}_R(T, M), \quad M \longmapsto \text{Hom}_R(M, T), \quad M \longmapsto M \otimes_R T$$

each send an object of $R - \mathbf{Mod}$ to another object of $R - \mathbf{Mod}$. Similarly, if $U \subset R$ is a multiplicatively closed set, localization gives an assignment

$$M \longmapsto U^{-1}M,$$

which sends an R -module to a module over the localized ring $U^{-1}R$. We would like to view each of these constructions as defining functors:

$$R\text{-}\mathbf{Mod} \xrightarrow{\mathrm{Hom}_R(T,-)} R\text{-}\mathbf{Mod} \qquad R\text{-}\mathbf{Mod} \xrightarrow{\mathrm{Hom}_R(-,T)} R\text{-}\mathbf{Mod}$$

$$R\text{-}\mathbf{Mod} \xrightarrow{-\otimes_R T} R\text{-}\mathbf{Mod} \qquad R\text{-}\mathbf{Mod} \xrightarrow{U^{-1}-} U^{-1}R\text{-}\mathbf{Mod}$$

However, a functor is not merely a rule on objects. To view these constructions functorially, we must also understand what they do to morphisms. Thus, given an R -module homomorphism $f : M \rightarrow N$, we must define induced maps between the corresponding Hom modules, tensor products, and localizations, and then check that these induced maps respect identities and composition. The following exercise asks you to make these constructions explicit and verify that they really do define functors.

Problem 7. Hom, Tensor, and Localization as Functors.

Throughout let R be a ring and fix an R -module T .

1. Let T be a fixed R -module. If $f : M \rightarrow N$ is a homomorphism, define

$$f_* : \text{Hom}_R(T, M) \rightarrow \text{Hom}_R(T, N), \quad \phi \mapsto f \circ \phi.$$

Prove that $\text{Hom}_R(T, -)$ is a covariant functor from $R - \mathbf{Mod}$ to $R - \mathbf{Mod}$.

2. If $f : M \rightarrow N$ is a homomorphism, define

$$f^* : \text{Hom}_R(N, T) \rightarrow \text{Hom}_R(M, T), \quad \psi \mapsto \psi \circ f.$$

Prove that $\text{Hom}_R(-, T)$ is a contravariant functor from $R - \mathbf{Mod}$ to $R - \mathbf{Mod}$.

3. If $f : M \rightarrow N$ is a homomorphism, use the universal property of tensor products to construct

$$f \otimes \text{Id}_T : M \otimes_R T \rightarrow N \otimes_R T, \quad m \otimes t \mapsto f(m) \otimes t.$$

Prove that $- \otimes_R T$ is a covariant functor from $R - \mathbf{Mod}$ to $R - \mathbf{Mod}$.

4. Let $U \subset R$ be a multiplicatively closed set. If $f : M \rightarrow N$ is a homomorphism, define

$$U^{-1}f : U^{-1}M \rightarrow U^{-1}N, \quad \frac{m}{u} \mapsto \frac{f(m)}{u}.$$

Prove that this is well-defined and that $U^{-1}(-)$ is a covariant functor from $R - \mathbf{Mod}$ to $U^{-1}R - \mathbf{Mod}$.

Solution.

1. • We know that $\text{Hom}_R(T, -)$ is a $R - \mathbf{Mod}$. That is, for every R -module M we have $\text{Hom}_R(T, M)$ another R -module.
• Let $f : M \rightarrow N, g : N \rightarrow O$ be R -module homomorphisms. Then, we need to show that f_* and g_* are R -module homomorphisms.

$$- r \cdot f_*(\phi) = r \cdot f \circ \phi = f \circ (r\phi) = f_*(r\phi)$$

$$- f_*(\phi + \psi) = f \circ (\phi + \psi) = f \circ \phi + f \circ \psi = f_*(\phi) + f_*(\psi)$$

A similar proof works to show g_* is a R -module homomorphism.

- $(g \circ f)_*(\phi) = g \circ f \circ \phi = g_*(f \circ \phi) = g_*(f_*(\phi)) = (g_* \circ f_*)(\phi)$
• Let $h : M \rightarrow M$ be a R -module homomorphism, where $h = \text{id}_M$. Then

$$h_* = h \circ \phi = \text{id}_M \circ \phi = \text{id}_{\text{Hom}_R(T, M)}$$

Thus, $\text{Hom}_R(T, -)$ is a covariant functor from $R - \mathbf{Mod}$ to $R - \mathbf{Mod}$.

2. • We know that $\text{Hom}_R(-, T)$ is a $R - \mathbf{Mod}$. That is, for every R -module M we have $\text{Hom}_R(M, T)$ another R -module.
- Let $f : M \rightarrow N, g : N \rightarrow O$ be R -module homomorphisms. Then, we need to show that f^* and g^* are R -module homomorphisms.

$$- r \cdot f^*(\phi) = r \cdot \phi \circ f = (r\phi) \circ f = f^*(r\phi)$$

$$- f^*(\phi + \psi) = (\phi + \psi) \circ f = \phi \circ f + \psi \circ f = f^*(\phi) + f^*(\psi)$$

A similar proof works to show g^* is a R -module homomorphism.

- $(g \circ f)^*(\phi) = \phi \circ g \circ f = f^*(\phi \circ g) = f^*(g^*(\phi)) = (f^* \circ g^*)(\phi)$
- Let $h : M \rightarrow M$ be a R -module homomorphism, where $h = \text{id}_M$. Then

$$h_* = \phi \circ h = \phi \circ \text{id}_M = \text{id}_{\text{Hom}_R(M, T)}$$

Thus, $\text{Hom}_R(T, -)$ is a contravariant functor from $R - \mathbf{Mod}$ to $R - \mathbf{Mod}$.

3. • We know that for any R -module $M, M \otimes_R T$ is a R -module.
- Let $f : M \rightarrow N, g : N \rightarrow O$ be R -module homomorphisms. Then, we need to show that $f \otimes \text{Id}_T$ and $g \otimes \text{Id}_T$ are R -module homomorphisms.

$$- r \cdot f \otimes \text{Id}_T(m \otimes t) = r \cdot f(m) \otimes t = f(rm) \otimes t = f \otimes \text{Id}_T(rm \otimes t)$$

-

$$\begin{aligned} f \otimes \text{Id}_T((m + n) \otimes t) &= f(m + n) \otimes t \\ &= f(m) \otimes t + f(n) \otimes t \\ &= f \otimes \text{Id}_T(m \otimes t) + f \otimes \text{Id}_T(n \otimes t) \end{aligned}$$

A similar proof works to show $g \otimes \text{Id}_T$ is a R -module homomorphism.

•

$$\begin{aligned} (g \circ f) \otimes \text{Id}_T(m \otimes t) &= (g \circ f)(m) \otimes t = g(f(m)) \otimes t \\ &= g \otimes \text{Id}_T(f(m) \otimes t) \\ &= (g \otimes \text{Id}_T) \circ (f \otimes \text{Id}_T)(m \otimes t) \end{aligned}$$

- Let $h : M \rightarrow M$ be a R -module homomorphism, where $h = \text{Id}_M$. Then

$$h \otimes \text{Id}_T(m \otimes t) = h(m) \otimes t = m \otimes t = (\text{Id}_M \otimes \text{Id}_T)(m \otimes t)$$

Thus, $- \otimes_R T$ is a covariant functor from $R - \mathbf{Mod}$ to $R - \mathbf{Mod}$.

4. • We know that $U^{-1}(-)$ is a $R - \mathbf{Mod}$. That is, for every R -module M we have $U^{-1}M$ another R -module.

- Let $f : M \rightarrow N, g : N \rightarrow O$ be R -module homomorphisms. Then, we need to show that $U^{-1}f$ and $U^{-1}g$ are R -module homomorphisms.

$$\begin{aligned}
- r \cdot U^{-1}f\left(\frac{m}{u}\right) &= \frac{rf(m)}{u} = \frac{f(rm)}{u} = U^{-1}f\left(\frac{rm}{u}\right) \\
- U^{-1}f\left(\frac{m}{u}\right) + U^{-1}f\left(\frac{n}{s}\right) &= \frac{f(m)}{u} + \frac{f(n)}{s} = \frac{f(ms)+f(nu)}{us} = U^{-1}f\left(\frac{ms+nu}{us}\right) = \\
&U^{-1}f\left(\frac{m}{u} + \frac{n}{s}\right)
\end{aligned}$$

A similar proof works to show g^* is a R -module homomorphism.

- $(U^{-1}g \circ U^{-1}f)\left(\frac{m}{u}\right) = U^{-1}g\left(U^{-1}f\left(\frac{m}{u}\right)\right) = \frac{g(f(m))}{u} = U^{-1}(f \circ g)\left(\frac{m}{u}\right)$
- Let $h : M \rightarrow M$ be a R -module homomorphism, where $h = \text{id}_M$. Then

$$U^{-1}h\left(\frac{m}{u}\right) = \frac{h(m)}{u} = \frac{m}{u}$$

Thus, $U^{-1}(-)$ is a covariant functor from $R - \mathbf{Mod}$ to $R - \mathbf{Mod}$.

The functors constructed above let us ask a new version of the exactness question. Instead of asking whether a given sequence of modules is exact, we can ask whether a construction preserves exactness. Roughly speaking, a functor is exact if applying it to an exact sequence does not introduce any new homology: the kernel of one map is still the image of the previous. This is a very strong condition, and many of the most important functors in algebra preserve only part of an exact sequence.

For the functors in this worksheet, applying the functor to a sequence again produces a sequence: these functors are additive, so they send zero maps to zero maps, and they respect composition. Let $F : R\text{-}\mathbf{Mod} \rightarrow S\text{-}\mathbf{Mod}$ be one of the covariant functors under consideration, where S is some ring. We say that F is *exact* if, whenever

$$M' \xrightarrow{f} M \xrightarrow{g} M''$$

is exact at M , the transformed sequence

$$F(M') \xrightarrow{F(f)} F(M) \xrightarrow{F(g)} F(M'')$$

is exact at $F(M)$. Equivalently, F sends exact sequences to exact sequences. In particular, an exact covariant functor sends every short exact sequence

$$0 \longrightarrow A \xrightarrow{i} B \xrightarrow{p} C \longrightarrow 0$$

to a short exact sequence

$$0 \longrightarrow F(A) \xrightarrow{F(i)} F(B) \xrightarrow{F(p)} F(C) \longrightarrow 0.$$

Thus exactness means that $F(i)$ remains injective, $F(p)$ remains surjective, and $\text{img}(F(i)) = \ker(F(p))$. A covariant functor F is called *left exact* if every exact sequence

$$0 \longrightarrow A \xrightarrow{i} B \xrightarrow{p} C$$

is sent to a sequence

$$0 \longrightarrow F(A) \xrightarrow{F(i)} F(B) \xrightarrow{F(p)} F(C)$$

which is exact at $F(A)$ and $F(B)$. So a left exact functor preserves the injectivity at the left end and preserves the kernel condition in the middle, but it may fail to make $F(p)$ surjective. Dually, F is called *right exact* if every exact sequence

$$A \xrightarrow{i} B \xrightarrow{p} C \longrightarrow 0$$

is sent to a sequence

$$F(A) \xrightarrow{F(i)} F(B) \xrightarrow{F(p)} F(C) \longrightarrow 0$$

which is exact at $F(B)$ and $F(C)$. Thus a right exact functor preserves the quotient at the right end, but it may fail to preserve injectivity at the left end. A functor that is both left exact and right exact is exact. One can also define exactness, left exactness, and right exactness for contravariant functors; in that case, the arrows are reversed.

The next exercises focus on three fundamental examples. The Hom functors are naturally left exact. For $\text{Hom}_R(T, -)$, the reason is that a map $T \rightarrow B$ whose composite with $B \rightarrow C$ is zero is exactly a map whose image lands in the submodule $A \subseteq B$. For the contravariant functor $\text{Hom}_R(-, T)$, the same idea appears in quotient form: a map $B \rightarrow T$ that kills A factors uniquely through $C = B/A$. In both cases, surjectivity can fail; a map into or out of one end of a short exact sequence need not lift or extend across the middle term.

Tensor product behaves in the opposite direction. The functor $- \otimes_R T$ is always right exact: it preserves cokernels and therefore preserves quotient presentations. This is why tensor product is so useful for computing modules from generators and relations. But tensor product need not be left exact; after tensoring, an injective map can acquire a kernel. The modules T for which tensoring with T is exact are precisely the flat modules, which will be introduced below.

Localization is the especially well-behaved example in commutative algebra. Since $U^{-1}M \cong U^{-1}R \otimes_R M$, localization is right exact for the same reason tensor product is right exact. What makes localization better than an arbitrary tensor product is that it is also left exact: if a fraction becomes zero after applying a localized map, then after clearing denominators its numerator already came from the previous module. Thus localization preserves all exact sequences, and this exactness is one of the main reasons localization is such a powerful tool.

Problem 8. Left Exactness of $\text{Hom}_R(T, -)$.

Let T be an R -module and let

$$0 \longrightarrow A \xrightarrow{i} B \xrightarrow{p} C \longrightarrow 0$$

be a short exact sequence.

1. Define the maps i_* and p_* by postcomposition, and prove that $p_* \circ i_* = 0$.
2. Prove that $i_* : \text{Hom}_R(T, A) \rightarrow \text{Hom}_R(T, B)$ is injective.
3. Suppose $\phi : T \rightarrow B$ satisfies $p \circ \phi = 0$. Prove that ϕ factors uniquely through $i : A \rightarrow B$. In other words, prove that there is a unique homomorphism $\tilde{\phi} : T \rightarrow A$ such that $\phi = i \circ \tilde{\phi}$.
4. Conclude that the sequence below is exact

$$0 \longrightarrow \text{Hom}_R(T, A) \xrightarrow{i_*} \text{Hom}_R(T, B) \xrightarrow{p_*} \text{Hom}_R(T, C).$$

5. Suppose the original short exact sequence splits. Prove that $p_* : \text{Hom}_R(T, B) \rightarrow \text{Hom}_R(T, C)$ is surjective. Thus $\text{Hom}_R(T, -)$ sends split short exact sequences to short exact sequences.
6. Let $R = \mathbb{Z}$ and apply $\text{Hom}_{\mathbb{Z}}(\mathbb{Z}/2\mathbb{Z}, -)$ to

$$0 \longrightarrow \mathbb{Z} \xrightarrow{\cdot 2} \mathbb{Z} \longrightarrow \mathbb{Z}/2\mathbb{Z} \longrightarrow 0$$

Compute the resulting sequence and explain why $\text{Hom}_{\mathbb{Z}}(\mathbb{Z}/2\mathbb{Z}, -)$ is not right exact.

Solution.

1. We define $i_*(\phi) = \phi \circ i$ and $p_*(\psi) = \psi \circ p$ for $\phi \in \text{Hom}_R(T, A)$ and $\psi \in \text{Hom}_R(T, B)$. Let $i(a) = b$. Then,

$$p_* \circ i_*(\phi) = \psi \circ p \circ \phi \circ i$$

1. **Left Exactness of $\text{Hom}_R(-, T)$.** Let T be an R -module and consider the short exact sequence below:

$$0 \longrightarrow A \xrightarrow{i} B \xrightarrow{p} C \longrightarrow 0.$$

- (a) Using precomposition define the maps as below. Prove that $i^* \circ p^* = 0$:

$$\text{Hom}_R(C, T) \xrightarrow{p^*} \text{Hom}_R(B, T) \xrightarrow{i^*} \text{Hom}_R(A, T) :$$

- (b) Prove that p^* is injective.
- (c) Suppose $\psi : B \rightarrow T$ satisfies $\psi \circ i = 0$. Prove that ψ factors uniquely through $p : B \rightarrow C$. In other words, prove that there is a unique homomorphism $\tilde{\psi} : C \rightarrow T$ such that $\psi = \tilde{\psi} \circ p$.
- (d) Conclude that the sequence below is exact:

$$0 \longrightarrow \text{Hom}_R(C, T) \xleftarrow{p^*} \text{Hom}_R(B, T) \xrightarrow{i^*} \text{Hom}_R(A, T).$$

- (e) Suppose the original short exact sequence splits. Prove that $i^* : \text{Hom}_R(B, T) \rightarrow \text{Hom}_R(A, T)$ is surjective. Thus $\text{Hom}_R(-, T)$ sends split short exact sequences to short exact sequences.
- (f) Let $R = \mathbb{Z}$ and apply $\text{Hom}_{\mathbb{Z}}(-, \mathbb{Z})$ to

$$0 \longrightarrow \mathbb{Z} \xrightarrow{\cdot 2} \mathbb{Z} \longrightarrow \mathbb{Z}/2\mathbb{Z} \longrightarrow 0$$

Compute the resulting sequence and explain why $\text{Hom}_{\mathbb{Z}}(-, \mathbb{Z})$ is not exact.

2. Hom Computations from Exactness.

- (a) Let M be an R -module and let $I \subseteq R$ be an ideal. Use the short exact sequence

$$0 \longrightarrow I \longrightarrow R \longrightarrow R/I \longrightarrow 0$$

together with $\text{Hom}_R(-, M)$ to recover the description

$$\text{Hom}_R(R/I, M) \cong \{m \in M \mid Im = 0\}.$$

Hint: Identify $\text{Hom}_R(R, M)$ with M by evaluating a homomorphism at $1 \in R$. Under this identification, what does the condition that a map $R \rightarrow M$ vanish on I become?

- (b) Let N be an abelian group and n a positive integer. Apply $\text{Hom}_{\mathbb{Z}}(-, N)$ to

$$0 \longrightarrow \mathbb{Z} \xrightarrow{\cdot n} \mathbb{Z} \longrightarrow \mathbb{Z}/n\mathbb{Z} \longrightarrow 0$$

Show that there is an exact sequence

$$0 \longrightarrow \text{Hom}_{\mathbb{Z}}(\mathbb{Z}/n\mathbb{Z}, N) \longrightarrow N \xrightarrow{\cdot n} N$$

Deduce that $\text{Hom}_{\mathbb{Z}}(\mathbb{Z}/n\mathbb{Z}, N)$ is naturally isomorphic to the subgroup of n -torsion elements of N .

- (c) For positive integers m, n , use the previous part to compute $\text{Hom}_{\mathbb{Z}}(\mathbb{Z}/m\mathbb{Z}, \mathbb{Z}/n\mathbb{Z})$.

- (d) Let $R = \mathbb{K}[x]$ and $M = R/\langle x^b \rangle$. Compute $\text{Hom}_R(R/\langle x^a \rangle, M)$ for positive integers a and b .
- (e) Let A be an $m \times n$ matrix with entries in R , let $M = \text{coker}(A)$, and let N be an R -module. Use the presentation

$$R^n \xrightarrow{A} R^m \twoheadrightarrow M \longrightarrow 0$$

to describe $\text{Hom}_R(M, N)$ as the kernel of an explicit map $N^m \rightarrow N^n$. Hint: Identify $\text{Hom}_R(R^m, N)$ with N^m and $\text{Hom}_R(R^n, N)$ with N^n . The induced map is precomposition with A ; in coordinates it is given by the transpose matrix A^t acting on column vectors, with the usual convention.

Problem 11. Right Exactness of Tensor Product.

Let T be an R -module and let the below be an exact sequence:

$$A \xrightarrow{f} B \xrightarrow{g} C \longrightarrow 0$$

1. Prove that $g \otimes \text{Id}_T : B \otimes_R T \rightarrow C \otimes_R T$ is surjective.
2. Construct a bilinear map $C \times T \rightarrow (B \otimes_R T) / \langle f(a) \otimes t \mid a \in A, t \in T \rangle$ by choosing a representative $b \in B$ of $c \in C$ and sending (c, t) to the class of $b \otimes t$. Check that this is well-defined.
3. Since $C \cong B / \text{img}(f)$, use the universal property of quotients and tensor products to show that

$$C \otimes_R T \cong (B \otimes_R T) / \langle f(a) \otimes t \mid a \in A, t \in T \rangle.$$

4. Prove that $\ker(g \otimes \text{Id}_T) = \text{img}(f \otimes \text{Id}_T)$.
5. Conclude that the sequence below is exact:

$$A \otimes_R T \longrightarrow B \otimes_R T \longrightarrow C \otimes_R T \longrightarrow 0.$$

6. Suppose the original short exact sequence splits. Prove that $g \otimes \text{Id}_T$ is injective. Thus $- \otimes_R T$ sends split short exact sequences to short exact sequences.

Solution.

1. We know that for all $c \in C$, there exists $b \in B$ such that $g(b) = c$. Now, let $c \otimes_R t \in C \otimes_R T$. Then we have,

$$g \otimes_R \text{Id}_T(b \otimes_R t) = g(b) \otimes_R t = c \otimes_R t$$

Hence the map $g \otimes_R \text{Id}_T$ is surjective.

2. Let $\phi : C \times T \rightarrow (B \otimes_R T) / \langle f(a) \otimes t \rangle$, where $(c, t) \mapsto \overline{b \otimes t}$ where $g(b) = c$. To check well-definedness let $g(b) = g(b')$. Then we have $g(b - b') = 0$, i.e. $b - b' \in \ker(g) = \text{img}(f)$. So $b - b' = f(a)$ for some $a \in A$.

$$(b - b') \otimes t = f(a) \otimes t \in \langle f(a) \otimes t \rangle \implies b \otimes t = b' \otimes t$$

- 3.

Problem 12. Tensor Product Need Not Be Left Exact.

The next example shows that tensor product can turn a nonzero injective map into the zero map.

1. Tensor the short exact sequence

$$0 \longrightarrow \mathbb{Z} \xrightarrow{\cdot 2} \mathbb{Z} \longrightarrow \mathbb{Z}/2\mathbb{Z} \longrightarrow 0$$

with $\mathbb{Z}/2\mathbb{Z}$. Compute the resulting sequence explicitly.

2. Explain why the map $\mathbb{Z} \otimes_{\mathbb{Z}} \mathbb{Z}/2\mathbb{Z} \rightarrow \mathbb{Z} \otimes_{\mathbb{Z}} \mathbb{Z}/2\mathbb{Z}$ induced by multiplication by 2 is the zero map.
3. Conclude that $- \otimes_{\mathbb{Z}} \mathbb{Z}/2\mathbb{Z}$ is not left exact.
4. Let n be a positive integer. Compute $(\mathbb{Z}/n\mathbb{Z}) \otimes_{\mathbb{Z}} (\mathbb{Z}/2\mathbb{Z})$ and use this to predict what happens when one tensors the exact sequence below with $\mathbb{Z}/2\mathbb{Z}$:

$$0 \longrightarrow \mathbb{Z} \xrightarrow{\cdot n} \mathbb{Z} \longrightarrow \mathbb{Z}/n\mathbb{Z} \longrightarrow 0.$$

Solution.

1. We get,

$$0 \longrightarrow \mathbb{Z}/2\mathbb{Z} \xrightarrow{\cdot 2 \otimes \text{id}_{\mathbb{Z}/2\mathbb{Z}}} \mathbb{Z}/2\mathbb{Z} \longrightarrow \mathbb{Z}/2\mathbb{Z} \longrightarrow 0$$

Since $\mathbb{Z} \otimes_{\mathbb{Z}} \mathbb{Z}/2\mathbb{Z} \cong \mathbb{Z}/2\mathbb{Z}$ and $\mathbb{Z}/\mathbb{Z} \otimes_{\mathbb{Z}} \mathbb{Z}/2\mathbb{Z} \cong \mathbb{Z}/2\mathbb{Z}$.

2. Let $a \otimes_{\mathbb{Z}} b \in \mathbb{Z} \otimes_{\mathbb{Z}} \mathbb{Z}/2\mathbb{Z}$. Then,

$$\cdot 2 \otimes_{\mathbb{Z}} \text{id}_{\mathbb{Z}/2\mathbb{Z}}(a \otimes_{\mathbb{Z}} b) = 2a \otimes_{\mathbb{Z}} b = a \otimes_{\mathbb{Z}} 2b = a \otimes_{\mathbb{Z}} 0 = 0(a \otimes_{\mathbb{Z}} b) = 0$$

Hence the map induced by multiplication by 2 is the zero map.

3. Since the map induced by multiplication by 2 is the zero map, it is not injective and hence $- \otimes_{\mathbb{Z}} \mathbb{Z}/2\mathbb{Z}$ is not left exact.
4. $(\mathbb{Z}/n\mathbb{Z}) \otimes_{\mathbb{Z}} (\mathbb{Z}/2\mathbb{Z}) \cong \mathbb{Z}/(n\mathbb{Z} + 2\mathbb{Z}) = \mathbb{Z}/(\gcd(n, 2)\mathbb{Z})$. That is if n is even, we have $(\mathbb{Z}/n\mathbb{Z}) \otimes_{\mathbb{Z}} (\mathbb{Z}/2\mathbb{Z}) \cong \mathbb{Z}/2\mathbb{Z}$ and the map induced by multiplication by n is the zero map and the sequence is not exact at the first term.

$$\mathbb{Z}/2\mathbb{Z} \xrightarrow{0} \mathbb{Z}/2\mathbb{Z} \longrightarrow \mathbb{Z}/2\mathbb{Z} \longrightarrow 0$$

If n is odd, we have $(\mathbb{Z}/n\mathbb{Z}) \otimes_{\mathbb{Z}} (\mathbb{Z}/2\mathbb{Z}) \cong 0$, the map induced by multiplication by n remains injective, and the sequence is exact.

$$\mathbb{Z}/2\mathbb{Z} \xrightarrow{\cdot n} \mathbb{Z}/2\mathbb{Z} \longrightarrow 0 \longrightarrow 0$$

Problem 13. Flat Modules.

An R -module T is called *flat* if the functor $- \otimes_R T$ is exact, or equivalently if tensoring with T preserves injective maps.

1. Explain why right exactness of tensor product implies that T is flat if and only if, for every injection $A \hookrightarrow B$, the induced map $A \otimes_R T \rightarrow B \otimes_R T$ is injective.
2. Prove that R is flat as an R -module.
3. Prove that a finite free module R^n is flat.
4. Prove that every free R -module is flat. Hint: Tensor product commutes with direct sums.

Solution.

1. By right exactness of tensor products we have two of the three sufficient conditions for exactness. The third condition requires injective maps to map to injective maps. Thus, T is flat iff for every injection $A \hookrightarrow B$, the induced map $A \otimes_R T \rightarrow B \otimes_R T$ is injective.
2. We need to show that tensoring with R preserves exactness, but since $A \otimes_R R \cong A$ gives us back the original sequence. Hence the functor $\otimes_R R$ is exact and R is flat.
3. For any exact sequence $0 \rightarrow A \rightarrow B$, we need to show $0 \rightarrow A \otimes_R R^n \rightarrow B \otimes_R R^n$ is exact. But $A \otimes_R R^n \cong A^n$, this becomes $0 \rightarrow A^n \rightarrow B^n$, and the map $A^n \rightarrow B^n$ is just $i^{\oplus n}$ (applying i componentwise), which is injective since i is injective. Hence a finite free module R^n is flat.
4. We know $A \otimes R^{(I)} \cong \bigoplus_{i \in I} A$, then the map $\bigoplus_{i \in I} A \rightarrow \bigoplus_{i \in I} B$ is injective because $A \rightarrow B$ is injective componentwise.

Localization is the most important exact functor in commutative algebra. Let $U \subset R$ be a multiplicatively closed set. The localization construction sends an R -module M to a $U^{-1}R$ -module $U^{-1}M$, and sends a homomorphism $f : M \rightarrow N$ to the homomorphism

$$U^{-1}M \xrightarrow{U^{-1}f} U^{-1}N \quad \frac{m}{u} \mapsto \frac{f(m)}{u}.$$

We already saw that $U^{-1}M \cong U^{-1}R \otimes_R M$. Since tensor product is right exact, localization is automatically right exact. The special feature of localization is that it is also left exact. The reason is the denominator-clearing argument: if a fraction maps to zero after localization, then some element of U already kills the relevant numerator.

Problem 14. Exactness of Localization.

Let $U \subset R$ be a multiplicatively closed set and let the sequence below be exact at M :

$$M' \xrightarrow{f} M \xrightarrow{g} M''.$$

1. Prove that

$$U^{-1}M' \xrightarrow{U^{-1}f} U^{-1}M \xrightarrow{U^{-1}g} U^{-1}M''$$

is a sequence of $U^{-1}R$ -modules whose consecutive composition is zero.

2. Prove that $\text{img}(U^{-1}f) \subseteq \ker(U^{-1}g)$.
3. Let $m/u \in U^{-1}M$ satisfy $(U^{-1}g)(m/u) = 0$. Unpack the definition of equality to zero in $U^{-1}M''$ to show that there exists $t \in U$ such that $g(tm) = 0$.
4. Use exactness of the original sequence at M to find $m' \in M'$ with $f(m') = tm$.
5. Prove that $\frac{m}{u} = \frac{f(m')}{tu}$ in $U^{-1}M$, and conclude that $m/u \in \text{img}(U^{-1}f)$.
6. Deduce that localization preserves exactness of arbitrary exact sequences.

Solution.

1. Let $\frac{m'}{u} \in U^{-1}M'$. Then the composition $U^{-1}g \circ U^{-1}f$ gives us,

$$U^{-1}g \circ U^{-1}f \left(\frac{m'}{u} \right) = \frac{g(f(m'))}{u} = \frac{0}{u}$$

since $g(f(m)) = 0$. Thus the consecutive composition is zero.

2. Let $\frac{m}{u} \in \text{img}(U^{-1}f)$. Then we know that $m \in \text{img}(f) = \ker(g)$. That is $g(m) = 0$ and thus $\frac{m}{u} \in \ker(U^{-1}g)$.

3. If $U^{-1}g \left(\frac{m}{u} \right) = \frac{g(m)}{u} = 0$. Then there exists $t \in U$ such that

$$t(g(m) \cdot 1 - 0 \cdot u) = t \cdot g(m) = g(tm) = 0$$

4. Since the original sequence is exact at M we have $\text{im}(f) = \ker(g)$, since $tm \in \ker(g)$ we must have some $m' \in M'$ such that $f(m') = tm$.

5. Then, $\frac{f(m')}{tu} = \frac{tm}{tu} = \frac{m}{u} \in U^{-1}M$, since $tu \in U$, we have $\frac{m}{u} \in \text{img}(U^{-1}f)$.

6. Since $\frac{m}{u}$ is an arbitrary element in $\ker(U^{-1}g)$, we have $\ker(U^{-1}g) \subseteq \text{img}(U^{-1}f)$. And part (2) gives us $\text{img}(U^{-1}f) \subseteq \ker(U^{-1}g)$, so $\ker(U^{-1}g) = \text{img}(U^{-1}f)$, and the sequence is exact at M . Thus localization preserves exactness of arbitrary exact sequences.

Problem 15. Localization Preserves Short Exact Sequences.

Let

$$0 \longrightarrow A \xrightarrow{i} B \xrightarrow{p} C \longrightarrow 0$$

be a short exact sequence and let $U \subset R$ be a multiplicatively closed set.

1. Prove directly that $U^{-1}i : U^{-1}A \rightarrow U^{-1}B$ is injective.
2. Prove directly that $U^{-1}p : U^{-1}B \rightarrow U^{-1}C$ is surjective.
3. Use the previous exercise to conclude that

$$0 \longrightarrow U^{-1}A \longrightarrow U^{-1}B \longrightarrow U^{-1}C \longrightarrow 0$$

is a short exact sequence of $U^{-1}R$ -modules.

4. Combine this with the isomorphism $U^{-1}M \cong U^{-1}R \otimes_R M$ to prove that $U^{-1}R$ is a flat R -module.

Solution.

1. Let $U^{-1}i\left(\frac{a}{u}\right) = \frac{i(a)}{u} = \left(\frac{0}{1}\right)$. Then there exists a $t \in U$ such that

$$t(i(a) \cdot 1 - u \cdot 0) = ti(a) = i(ta) = 0$$

Since i is injective, this implies $ta = 0$. But then, $t(a \cdot 1 - u \cdot 0) = ta = 0$, so $\frac{a}{u} \sim \frac{0}{1} \in U^{-1}A$. Hence $U^{-1}i$ is injective.

2. Since p is surjective, for every $c \in C$ there exists some $b \in B$ such that $p(b) = c$. Then, for some $\frac{c}{u} \in U^{-1}C$ we have, $U^{-1}p\left(\frac{b}{u}\right) = \frac{p(b)}{u} = \frac{c}{u}$. Thus $U^{-1}p$ is surjective.
3. Since $U^{-1}i$ is injective, $U^{-1}p$ is surjective, and $\ker(U^{-1}p) = \text{img}(U^{-1}i)$ by Problem 14, the sequence is exact at every term and hence is a short exact sequence of $U^{-1}R$ modules.
4. Since $U^{-1}M \cong U^{-1}R \otimes_R M$ we have the resulting exact sequence,

$$0 \longrightarrow U^{-1}R \otimes_R A \longrightarrow U^{-1}R \otimes_R B \longrightarrow U^{-1}R \otimes_R C \longrightarrow 0$$

Thus $U^{-1}R \otimes_R -$ preserves exact sequences and hence $U^{-1}R$ is a flat R -module.

Problem 16. Computing Localized Exact Sequences.

1. Let $I \subseteq R$ be an ideal. Use exactness of localization to prove that $U^{-1}(R/I) \cong U^{-1}R/IU^{-1}R$.
2. Localize the short exact sequence at $U = \{1, 2, 2^2, 2^3, \dots\}$. Identify each term explicitly

$$0 \longrightarrow \mathbb{Z} \xrightarrow{\cdot 6} \mathbb{Z} \longrightarrow \mathbb{Z}/6\mathbb{Z} \longrightarrow 0.$$

3. Localize the same sequence at $V = \mathbb{Z} \setminus \langle 2 \rangle$. Identify $V^{-1}(\mathbb{Z}/6\mathbb{Z})$.
4. Let $R = \mathbb{K}[x]$ and $M = R/\langle x^2(x-1) \rangle$. Compute $M[1/x]$ using the localized presentation

$$R[1/x] \xrightarrow{\cdot x^2(x-1)} R[1/x] \longrightarrow M[1/x] \longrightarrow 0$$

5. Explain why localizing $R/\langle x^2(x-1) \rangle$ at the prime ideal $\langle x \rangle$ “keeps” the $x = 0$ component and “removes” the $x = 1$ component.
6. With R and M as in the previous part, compute $M_{\langle x \rangle}$. Hint: What happens to $x - 1$ in $R_{\langle x \rangle}$?
7. Let $R = \mathbb{K}[x, y]$ and $M = R/\langle xy \rangle$. Compute $M[1/x]$ and $M[1/y]$.

Solution.

1. Using previous exercises, we get the exact sequence,

$$0 \longrightarrow U^{-1}I \xrightarrow{U^{-1}\iota} U^{-1}R \xrightarrow{U^{-1}\pi} U^{-1}(R/I) \longrightarrow 0$$

Then by the first isomorphism theorem we have,

$$U^{-1}(R/I) \cong U^{-1}R / \ker(U^{-1}\pi) = U^{-1}R / \text{img}(U^{-1}\iota) \cong U^{-1}R / U^{-1}I = U^{-1}R / IU^{-1}R$$

2. Localizing at U gives us,

$$0 \longrightarrow \mathbb{Z} \begin{bmatrix} 1 \\ 2 \end{bmatrix} \xrightarrow{\cdot 6} \mathbb{Z} \begin{bmatrix} 1 \\ 2 \end{bmatrix} \longrightarrow \mathbb{Z}/3\mathbb{Z} \longrightarrow 0.$$

Since $U^{-1}\mathbb{Z}$ gives us $\mathbb{Z} \left[\frac{1}{2} \right]$ and since $\bar{2}$ is a unit in $\mathbb{Z}/3\mathbb{Z}$,

$$\begin{aligned} U^{-1}(\mathbb{Z}/6\mathbb{Z}) &\cong U^{-1}\mathbb{Z}/(6U^{-1}\mathbb{Z}) \\ &= \mathbb{Z} \left[\frac{1}{2} \right] / \left(6\mathbb{Z} \left[\frac{1}{2} \right] \right) \\ &= \mathbb{Z} \left[\frac{1}{2} \right] / \left(3\mathbb{Z} \left[\frac{1}{2} \right] \right) \\ &\cong U^{-1}(\mathbb{Z}/3\mathbb{Z}) = \mathbb{Z}/3\mathbb{Z} \end{aligned}$$

3. Since $\frac{3}{1} \in U^{-1}(\mathbb{Z}/6\mathbb{Z})$ is a unit, we have $\frac{2}{1} = \frac{3 \cdot 2}{3} = \frac{0}{1} \in U^{-1}(\mathbb{Z}/6\mathbb{Z})$. Hence, $U^{-1}(\mathbb{Z}/6\mathbb{Z}) \cong \mathbb{Z}/2\mathbb{Z}$.
4. Note that in $R \left[\frac{1}{x} \right] = \mathbb{K}[x, x^{-1}]$, x is a unit. Then

$$\text{img}(\cdot x^2(x-1)) = \langle x^2(x-1) \rangle = \langle x-1 \rangle$$

So, $M \left[\frac{1}{x} \right] = R \left[\frac{1}{x} \right] / \langle x-1 \rangle = \mathbb{K}$.

5. Localizing at $\langle x \rangle$ means inverting $U = \mathbb{R} \setminus \langle x \rangle$, i.e. all polynomials that do not vanish at $x = 0$. Since $x-1 \neq 0$ when $x = 0$ it gets inverted killing the $x = 1$ component. But since $x \notin U$, we preserve the $x = 0$ component.
6. Since $\langle x-1 \rangle$ is now a unit we get $M_{\langle x \rangle} \cong R / \langle x^2 \rangle$.
7. We have $R \left[\frac{1}{x} \right] = \mathbb{K}[x, y, x^{-1}]$. In $M \left[\frac{1}{x} \right]$, x is a unit, so $\langle xy \rangle = \langle y \rangle$. Then $M \left[\frac{1}{x} \right] = \mathbb{K}[x, y, x^{-1}] / \langle y \rangle \cong \mathbb{K}[x, x^{-1}]$. Similarly, in $M \left[\frac{1}{y} \right]$, y is a unit, so $\langle xy \rangle = \langle x \rangle$, and $M \left[\frac{1}{y} \right] = \mathbb{K}[x, y, y^{-1}] / \langle x \rangle \cong \mathbb{K}[y, y^{-1}]$.

Worksheet 7.1: Integral Extension Pt. I

Throughout this course, “ring” means *commutative* ring with unity and all ring homomorphisms preserve unity. In the previous worksheets we have repeatedly moved between rings, modules, quotients, localization, and spectra. The goal of this worksheet is to introduce a language that lets us treat many ring maps as geometric families: the language of *R-algebras*. Once this language is in place, we will distinguish two finiteness notions that sound similar but behave very differently: finite generation as an algebra and finite generation as a module.

The main bridge between these two notions is integrality. Integral elements are elements of an extension ring that satisfy monic polynomial equations over the base. The monic condition is what lets one convert algebraic equations into module-theoretic finiteness. This is one of the basic mechanisms behind many finiteness theorems in commutative algebra and algebraic geometry.

Let R be a ring. An *R-algebra* is a ring S together with a specified ring homomorphism $\alpha : R \rightarrow S$. Note we do not require that α be injective, however, the case when α is an inclusion, i.e., $\mathbb{K}$ into $\mathbb{K}[x]$, is generally the motivating example. We often think of an *R-algebra* structure on a ring S as giving us a way to multiply elements of S by elements of R , i.e., $r \cdot s := \alpha(r)s$. Thus every *R-algebra* S is naturally an *R-module*. The key difference between an *R-module* M and an *R-algebra* S is that for module we may only multiply elements of M by elements of R , but for an *R-algebra* every element of S may be multiplied together. The key analogy might be: *R-modules* are to vector spaces, as *R-algebras* are to rings.

An *R-algebra* homomorphism $\phi : S \rightarrow T$ is a ring homomorphism compatible with the structure maps from R , meaning that the diagram

$$\begin{array}{ccc} S & \xrightarrow{\phi} & T \\ & \searrow \alpha_S & \swarrow \alpha_T \\ & T & \end{array}$$

commutes. Equivalently, $\phi(\alpha_S(r)) = \alpha_T(r)$ for every $r \in R$. An *R-subalgebra* of S is a subring $S' \subset S$, which contains $\alpha(R)$. Given any set $\mathcal{S} \subset S$ the *R-algebra generated by $\mathcal{S}$* is the smallest *R-subalgebra* of S containing $\mathcal{S}$. We often write $R[\mathcal{S}]$ to denote *R-subalgebra* generated by $\mathcal{S}$, because it can explicitly be written down as the set of polynomials in the elements in $\mathcal{S}$ with coefficients in R .

Problem 1. First Encounters with R -algebras.

Let $\alpha : R \rightarrow S$ be an R -algebra.

1. Verify that the rule $r \cdot s = \alpha(r)s$ makes S into an R -module.
2. Give an example where α is not injective. In your example, identify a nonzero element of R that acts as zero on S .
3. Show that every quotient R/I is naturally an R -algebra.
4. Show that if S and T are R -algebras and $\phi : S \rightarrow T$ is an R -algebra homomorphism, then ϕ is R -linear for the underlying R -module structures.
5. Let $\mathbb{K} \subseteq \mathbb{E}$ be a field extension. Explain why $\mathbb{E}$ is a $\mathbb{K}$ -algebra. What is the underlying $\mathbb{K}$ -module structure?

Solution.

1. We verify the axioms. For all $r, p \in R, s, t \in S$ we have:

$$(a) \quad r(s + t) = \alpha(r)(s + t) = \alpha(r)s + \alpha(r)t = rs + rt$$

$$(b) \quad (r + p)t = \alpha(r + p)t = \alpha(r)t + \alpha(p)t = rt + pt$$

$$(c) \quad (rp)t = \alpha(rp)t = \alpha(r)\alpha(p)t = \alpha(r)(\alpha(p)t) = r(pt)$$

Thus the rule $r \cdot s = \alpha(r)s$ makes S into a R -module.

2. $\alpha : \mathbb{Z} \rightarrow \mathbb{Z}/2\mathbb{Z}$.

2 is a non-zero element in $\mathbb{Z}$ but acts as zero on $\mathbb{Z}/2\mathbb{Z}$.

3. The natural projection $\pi : R \rightarrow R/I$ is a ring homomorphism, and the property we proved in part (1) works with substituting $\alpha = \pi$.
4. Let $\alpha : R \rightarrow S$ and $\beta : R \rightarrow T$. Then, for $r, s \in R$,

$$\phi(r \cdot s) = \phi(\alpha(r)s) = \phi(\alpha(r))\phi(s) = \beta(r)\phi(s) = r\phi(s)$$

Hence, ϕ is R -linear.

5. $\mathbb{E}$ is a $\mathbb{K}$ -algebra with the inclusion map $\iota : \mathbb{K} \rightarrow \mathbb{E}, \iota(k) = k$ as the ring homomorphism. Moreover multiplication in $\mathbb{E}$ is $\mathbb{K}$ -linear and,

$$k \cdot e = \iota(k)e = ke$$

Problem 2. Polynomial Rings and Generation of R -algebras.

Let S be an R -algebra and let $s_1, \dots, s_n \in S$.

1. Construct an R -algebra homomorphism

$$\text{ev}_{s_1, \dots, s_n} : R[x_1, \dots, x_n] \longrightarrow S, \quad f(x_1, \dots, x_n) \longmapsto f(s_1, \dots, s_n).$$

2. Show that the image of this map is the smallest R -subalgebra of S containing $s_1, \dots, s_n$. We denote this image by $R[s_1, \dots, s_n]$.
3. Prove that $S = R[s_1, \dots, s_n]$ if and only if $\text{ev}_{s_1, \dots, s_n}$ is surjective.
4. Deduce that S is generated as an R -algebra by n elements if and only if $S \cong R[x_1, \dots, x_n]/I$ as an R -algebra for some ideal $I \subset R[x_1, \dots, x_n]$.

Solution.

1. We define $\text{ev}_{s_1, \dots, s_n} : R[x_1, \dots, x_n] \longrightarrow S$ by,

$$\text{ev}_{s_1, \dots, s_n}(f(x_1, \dots, x_n)) = \text{ev}_{s_1, \dots, s_n}\left(\sum_i r_i x_i\right) = f(s_1, \dots, s_n) = \sum_i r_i(s_i) \in S$$

This is a homomorphism since both

$$(f + g)(s_1, \dots, s_n) = \sum_i (r_i + r'_i)s_i = \sum_i r_i s_i + \sum_i r'_i s_i = f(s_1, \dots, s_n) + g(s_1, \dots, s_n)$$

and,

$$(fg)(s_1, \dots, s_n) = \sum_i (r_i \cdot r'_i)s_i = f(s_1, \dots, s_n) \cdot g(s_1, \dots, s_n)$$

hold.

2. We know that $\text{im}(\text{ev}_{s_1, \dots, s_n}) = R[s_1, \dots, s_n]$. Let $T \subseteq S$ be a R -subalgebra with $s_1, \dots, s_n \in T$. Note that a general element of $R[s_1, \dots, s_n]$ has the form

$$f(s_1, \dots, s_n) = \sum_i r_i s_1^{i_1} \cdots s_n^{i_n}$$

But since T is a R -subalgebra the sum $\sum_i r_i s_1^{i_1} \cdots s_n^{i_n} \in T$ since T is closed under addition and multiplication and $r_i \in T$ for all $r_i \in R$. Thus, $R[s_1, \dots, s_n] \subseteq T$. Since $R[s_1, \dots, s_n]$ is contained in every R -subalgebra of S containing $s_1, \dots, s_n$ and is itself such a subalgebra it must be the smallest one.

3. We know that $\text{im}(\text{ev}_{s_1, \dots, s_n}) = R[s_1, \dots, s_n] \subseteq S$. If $S = R[s_1, \dots, s_n]$ then

$$S = \text{im}(\text{ev}_{s_1, \dots, s_n})$$

and $\text{ev}_{s_1, \dots, s_n}$ are surjective.

4. ($\implies$)

Assume S is a R -algebra generated by $s_1, \dots, s_n \in S$. Then we have $S = R[x_1, \dots, x_n] = \text{im}(\text{ev}_{s_1, \dots, s_n})$ by part 3. Then by the first isomorphism theorem,

$$S = \text{im}(\text{ev}_{s_1, \dots, s_n}) = R[x_1, \dots, x_n]/I$$

for some $I \subseteq R[x_1, \dots, x_n]$.

($\impliedby$)

Assume $S \cong R[x_1, \dots, x_n]/I$. Then there exists an isomorphism $\phi : R[x_1, \dots, x_n]/I \rightarrow S$, where $s_i = \phi(x_i + I)$ for $1 \leq i \leq n$. Then every element of S can be written as

$$\phi(f + I) = \phi\left(\sum_i r_i x_1^{i_1} \cdots x_n^{i_n} + I\right) = r_i s_1^{i_1} \cdots s_n^{i_n} = f(s_1, \dots, s_n)$$

so every element of S is a polynomial expression in $s_1, \dots, s_n$ with coefficients in R . Hence $S = R[s_1, \dots, s_n]$ is an R -algebra generated by n elements.

Problem 3. Examples of R -algebras.

In each case, identify a set of algebra generators over the indicated base ring. Try to choose a minimal-looking set, though you do not need to prove minimality.

- | | |
|--|--|
| 1. $R[x]/\langle x^2 \rangle$ over R . | 5. $\mathbb{Z}[i]$ over $\mathbb{Z}$. |
| 2. $R[x, y]/\langle xy \rangle$ over R . | 6. $\mathbb{K}[x, y]/\langle y^2 - x^3 \rangle$ over $\mathbb{K}[x]$. |
| 3. R/I over R . | 7. $\mathbb{K}(t)[[t]]$ over the $\mathbb{K}(t)$. |
| 4. $\mathbb{K}[t, t^{-1}]$ over $\mathbb{K}$. | 8. $\mathbb{Z}[\sqrt{n}; n \in \mathbb{Z}_{\geq 1}]$ over $\mathbb{Z}$. |

Solution.

- | | |
|---------------------------|--|
| 1. $\{\bar{x}\}$ | 5. $\{i\}$ |
| 2. $\{\bar{x}, \bar{y}\}$ | 6. $\{\bar{y}\}$ |
| 3. $\{\}$ | 7. TODO: |
| 4. $\{t, t^{-1}\}$ | 8. $\{\sqrt{p} \mid p \text{ prime}\}$ |

As discussed above an R -algebra S is also always an R -module with the same R -multiplication. This leads to two different finiteness conditions depending on whether we wish to view S as an R -module or as an R -algebra. Let S be an R -algebra.

Definition 0.15. We say that S is algebra-finite over R if S is finitely generated as an R -algebra.

Definition 0.16. We say that S is module-finite over R if S is finitely generated as an R -module.

The condition S be algebra-finite over R is sometimes called being *finite type* over R , i.e. we would say S is finite type over R if and only if S is algebra-finite over R . Similarly, many authors will use the phrase S is *finite* over R or S is a *finite R -algebra* to mean S is a R -algebra that is module-finite. In this worksheet we will attempt to restrict to using *algebra-finite* and *module-finite* to hopefully minimize confusion.

The key point is that module-finite is much stronger than algebra-finite. A module-finite algebra is controlled by finitely many R -linear combinations. An algebra-finite algebra is controlled by finitely many elements and all polynomial expressions in them.

Problem 4. Module-Finite Implies Algebra-Finite.

Let S be an R -algebra and suppose that S is generated as an R -module by $m_1, \dots, m_n \in S$.

1. Show that the R -subalgebra $R[m_1, \dots, m_n]$ contains every R -linear combination of $m_1, \dots, m_n$.
2. Deduce that $S = R[m_1, \dots, m_n]$.
3. Conclude that every module-finite R -algebra is algebra-finite over R .

Solution.

1. An R -linear combination is of the form $r_1 m_1 + \dots + r_n m_n$ for all $r_i \in R$. But this is the polynomial $r_1 x_1 + \dots + r_n x_n$ evaluated at $m_1, \dots, m_n$ which is an element of $\text{im}(ev_{m_1, \dots, m_n}) = R[m_1, \dots, m_n]$.
2. The containment $S \supseteq R[m_1, \dots, m_n]$ is obvious because the latter is a subalgebra of S and part 1 gives us the reverse containment, hence we have $S = R[m_1, \dots, m_n]$.
3. Thus every module-finite R -algebra generated by n elements can be written in the form $R[m_1, \dots, m_n]$ (by part 2) and thus is algebra-finite over R .

Problem 5. Algebra-Finite Need Not Imply Module-Finite.

Assume R is nonzero.

1. Show that $R[x]$ is algebra-finite over R .
2. Suppose $p_1, \dots, p_n \in R[x]$. Let d be the maximum of their degrees. Show that every R -linear combination of the p_i has degree at most d , unless it is zero.
3. Deduce that $R[x]$ is not module-finite over R . Hint: Is x^{d+1} in the R -span of $p_1, \dots, p_n$?
4. Specialize to $R = \mathbb{K}$ a field and restate the result in the language of vector spaces.

Solution.

1. $R[x]$ is generated by $\{x\}$ as an algebra and hence is algebra-finite.
2. Every R -linear combination of the p_i is of the form

$$r_1 p_1 + \cdots + r_n p_n, \quad r_i \in R$$

since $\deg(r_i) = 0$ the degree of the equation is the maximum of the degrees of the p_i which is d .

3. Assume $R[x]$ is module-finite over R . Then there exists a finite set of generating elements $p_1, \dots, p_n \in R[x]$. Let d be the maximum of their degrees. Now, $x^{d+1} \in R[x]$ with degree $d + 1$ but by part 2 we know that any R linear combination of the p_i has degree at most d — a contradiction. Hence, $R[x]$ is not module finite over R .
4. If $R = \mathbb{K}$ then $R[x]$ is an infinite dimensional vector space.

Problem 6. Permanence of Algebra-Finiteness.

The goal of this exercise is to explore how being algebra-finite plays with many natural algebraic operations: quotienting, tensoring, localizing, and composing the underlying morphisms giving the algebra structure. Consider maps of rings:

$$R \xrightarrow{\alpha} S \xrightarrow{\beta} T .$$

The map α gives S an R -algebra structure. On the other hand T has two different algebra structures: we may think of T as an S -algebra via the map β or think of T as an R -algebra via $\beta \circ \alpha$.

1. If S is algebra-finite over R and $J \subseteq S$ is an ideal, prove that S/J is algebra-finite over R .
2. If S is algebra-finite over R and T is algebra-finite over S , prove that T is algebra-finite over R .
3. Give an example where α and β are injective and T is algebra-finite over R , but S is not algebra-finite over R . Hint: Let $R = \mathbb{K}$, and $T = \mathbb{K}[x, y]$, what happens if $S = \mathbb{K}[x, xy, xy^2, xy^3, \dots]$?
4. If S is algebra-finite over R , and $R \rightarrow T$ is a ring map, prove that $T \otimes_R S$ is algebra-finite over T . What are algebra generators?
5. If $U \subset R$ is multiplicatively closed and S is algebra-finite over R , prove that $U^{-1}S$ is algebra-finite over $U^{-1}R$.

Solution.

1. Let $S = R[s_1, \dots, s_n]$. Then the quotient map $\pi : S \rightarrow S/J$ is a surjective homomorphism with $\text{im}(\pi) = R[\bar{s}_1, \dots, \bar{s}_n]$, where $\pi(s_i) = \bar{s}_i$. Hence S/J is algebra-finite over R .
2. Let $S = R[s_1, \dots, s_n]$, and $T = S[t_1, \dots, t_m]$. Then every element of T is a polynomial in $t_1, \dots, t_m$ with coefficients in S , and each coefficient is a polynomial in $s_1, \dots, s_n$ with coefficients in R . Hence every element of T is a polynomial's expression in $\beta(s_1), \dots, \beta(s_n), t_1, \dots, t_m$ with coefficients in R , so

$$T = R[\beta(s_1), \dots, \beta(s_n), t_1, \dots, t_m]$$

which is algebra finite over R .

3. We let $R = \mathbb{K}$, $T = \mathbb{K}[x, y]$ and $S = R[x, xy, xy^2, xy^3, \dots]$.

4. Let $S = R[s_1, \dots, s_n]$. Every element of $T \otimes_R S$ is a finite sum of simple tensors $t \otimes s$. Since $s \in S$, we can write $s = \sum_i r_i s_1^{i_1} \dots s_n^{i_n}$ for $r_i \in R$. Then,

$$\begin{aligned}
 t \otimes s &= t \otimes \sum_i r_i s_1^{i_1} \dots s_n^{i_n} \\
 &= t \sum_i r_i \otimes s_1^{i_1} \dots s_n^{i_n} \\
 &= \sum_i tr_i \otimes s_1^{i_1} \dots s_n^{i_n} \\
 &= \sum_i tr_i (1 \otimes s_1)^{i_1} \dots (1 \otimes s_n)^{i_n}
 \end{aligned}$$

Which is a polynomial in $(1 \otimes s_1), \dots, (1 \otimes s_n)$ with coefficients in T and thus, these are the algebraic generators of $T \otimes_R S$ and $T \otimes_R S = T[1 \otimes s_1, \dots, 1 \otimes s_n]$.

5. Let $S = R[s_1, \dots, s_n]$. Note that the map $\eta : S \rightarrow U^{-1}S$ is a R -algebra homomorphism, and hence a $U^{-1}R$ algebra homomorphism (by extending scalars). Every element of $U^{-1}S$ is of the form $\frac{s}{u} = \frac{1}{u} \cdot \frac{s}{1}$ where $\frac{1}{u} \in U^{-1}R$.

Since every element $s \in S$ is of the form $s = \sum_i r_i s_1^{i_1} \dots s_n^{i_n}$ we get

$$\frac{s}{u} = \frac{1}{u} \sum_i \frac{r_i}{1} \left(\frac{s_1}{1}\right)^{i_1} \dots \left(\frac{s_n}{1}\right)^{i_n}$$

which is a polynomial in $\frac{s_1}{1}, \dots, \frac{s_n}{1}$. So, $U^{-1}S = U^{-1}R[s_1, \dots, s_n]$ which is algebra-finite over R .

Problem 7. Permanence of Module-Finiteness.

The goal of this exercise is to explore how being module-finite plays with many natural algebraic operations: quotienting, tensoring, localizing, and composing the underlying morphisms giving the algebra structure. Consider maps of rings:

$$R \xrightarrow{\alpha} S \xrightarrow{\beta} T .$$

The map α gives S an R -module structure. On the other hand T has two different module structures: we may think of T as an S -module via the map β or think of T as an R -module via $\beta \circ \alpha$.

1. If S is module-finite over R and $J \subseteq S$ is an ideal, prove that S/J is module-finite over R .
2. If S is module-finite over R and T is module-finite over S , prove that T is module-finite over R .
3. If S is module-finite over R , and $R \rightarrow T$ is a ring map, prove that $T \otimes_R S$ is module-finite over T .
4. If $U \subset R$ is multiplicatively closed and S is module-finite over R , prove that $U^{-1}S$ is module-finite over $U^{-1}R$.

Solution.

1. Let $S = R[m_1, \dots, m_n]$. Then, every element in $s \in S$ can be written as $s = \sum_i r_i m_i$ for $r_i \in R$, $m_i \in S$. Then, every element of S/J can be written as $\bar{s} = \sum_i \bar{r}_i \bar{m}_i = \sum_i r_i \bar{m}_i$. Thus S/J is spanned by the $\bar{m}_i$ as a R -module and hence is module-finite.
2. Let $T = S[n_1, \dots, n_m]$ and $S = R[m_1, \dots, m_n]$. Then every element in T can be written as $t = \sum_i s_i n_i$ with $s_i \in S$, and every s_i can be written as $\sum_j r_j m_j$. Then every element in T can be written as

$$t = \sum_i \left(\sum_j r_{ij} m_j \right) n_i = \sum_{i,j} r_{ij} (m_j n_i)$$

Thus T is spanned by the $m_j n_i$ as a R -module and hence is module-finite.

3. Let $S = R[m_1, \dots, m_n]$. Every element of $T \otimes_R S$ is a finite sum of simple tensors

$t \otimes s$. Since $s \in S$, we can write $s = \sum_i r_i m_i$ for $r_i \in R$. Then,

$$\begin{aligned}
 t \otimes s &= t \otimes \sum_i r_i m_i \\
 &= t \sum_i r_i \otimes \sum_i m_i \\
 &= \sum_i t r_i \otimes m_i \\
 &= \sum_i t r_i (1 \otimes m_i)
 \end{aligned}$$

Thus, $T \otimes_R S$ is spanned by the $(1 \otimes m_i)$ and hence is module-finite.

4. Let $S = R[m_1, \dots, m_n]$. Every element of $U^{-1}S$ is of the form,

$$\frac{s}{u} = \frac{1}{u} \cdot \frac{s}{1}$$

with $\frac{1}{u} \in U^{-1}R$ and since every $s \in S$ is of the form $s = \sum_i r_i m_i$ and then we have,

$$\frac{s}{1} = \sum_i \frac{r_i}{1} \cdot \frac{m_i}{1}$$

Thus, $U^{-1}S$ is spanned by $\frac{m_i}{1}$ as a $U^{-1}R$ -module.

Problem 8. Comparing the Two Notions in Examples.

Decide whether each algebra is algebra-finite and whether it is module-finite over the indicated base ring. Justify your answers.

- | | |
|--|---|
| 1. $\mathbb{K}[x]$ over $\mathbb{K}$. | 4. $\mathbb{Z}[1/2]$ over $\mathbb{Z}$. |
| 2. $\mathbb{K}[x]/\langle x^2 \rangle$ over $\mathbb{K}$. | 5. $\mathbb{K}[t]$ over $\mathbb{K}[t^2]$. |
| 3. $\mathbb{Z}[i]$ over $\mathbb{Z}$. | 6. $\mathbb{K}[t, t^{-1}]$ over $\mathbb{K}[t]$. |

Solution.

- | | |
|--------------------------------------|--------------------------------------|
| 1. Algebra-finite not module-finite. | 4. Algebra-finite not module-finite. |
| 2. Algebra-finite and module-finite. | 5. Algebra-finite and module-finite. |
| 3. Algebra-finite and module-finite. | 6. Algebra-finite not module-finite. |

Let S be an R -algebra and let $s \in S$. We say that s is *integral over R* if there exists a monic polynomial with coefficients in R that s satisfies. In other words, s is integral over R if there are elements $a_1, \dots, a_n \in R$ such that

$$s^n + a_1 s^{n-1} + a_2 s^{n-2} + \cdots + a_{n-1} s + a_n = 0$$

in S , where each a_i acts through the structure map $R \rightarrow S$. The word *monic* is essential. For example, every rational number satisfies some nonzero polynomial over $\mathbb{Z}$, but only the integers satisfy monic polynomial equations over $\mathbb{Z}$. The monic condition is exactly what allows us to solve for high powers of s in terms of lower powers.

Problem 9. First Examples of Integral Elements.

Let S be an R -algebra.

1. Show that every element in the image of $R \rightarrow S$ is integral over R .
2. Show that every nilpotent element of S is integral over R .
3. Let $S = R[x]/\langle x^2 - r \rangle$ for some $r \in R$. Show that the image of x in S is integral over R .
4. Show that $i \in \mathbb{Z}[i]$ is integral over $\mathbb{Z}$.
5. Show that $t \in \mathbb{K}[t]$ is integral over $\mathbb{K}[t^2]$.
6. Show that $\frac{1}{2}(\sqrt{-3} + 1) \in \mathbb{Q}(\sqrt{-3})$ is integral over $\mathbb{Z}[\sqrt{-3}]$.
7. Show directly that $1/2 \in \mathbb{Q}$ is not integral over $\mathbb{Z}$. Hint: Suppose $(1/2)^n + a_1(1/2)^{n-1} + \cdots + a_n = 0$ and clear denominators.

Solution.

1. Every element $s \in \text{im}(\alpha)$ in the image of $\alpha : R \rightarrow S$ is of the form $\alpha(r) = s \implies s - \alpha(r) = 0$.

2. Assume $s^n = 0$. Then we have,

$$s^n + 0s^{n-1} + \cdots + 0 = 0$$

Thus, every nilpotent element of S is integral over R .

3. We have,

$$\bar{x}^2 + 0\bar{x} - r = 0$$

Thus, the image of x in S is integral over R .

4. We have,

$$i^2 + 0i + 1 = 0$$

Thus $i \in \mathbb{Z}[i]$ is integral over $\mathbb{Z}$.

5. We have,

$$(t)^2 + 0(t) - t^2 = 0$$

Thus $t \in \mathbb{K}[t]$ is integral over $\mathbb{K}[t^2]$.

6. We have,

$$\begin{aligned}
 \left(\frac{1}{2}(\sqrt{-3}+1)\right)^2 - \frac{1}{2}(\sqrt{-3}+1) + 1 &= \frac{1}{4}(-2+2\sqrt{-3}) - \frac{1}{2}(\sqrt{-3}+1) + 1 \\
 &= \frac{1}{2}(\sqrt{-3}-1) - \frac{1}{2}(\sqrt{-3}+1) + 1 \\
 &= \frac{-2}{2} + 1 = 0
 \end{aligned}$$

Thus $\frac{1}{2}(\sqrt{-3}+1) \in \mathbb{Q}(\sqrt{-3})$ is integral over $\mathbb{Z}[\sqrt{-3}]$.

7. Suppose for the sake of contradiction that $\frac{1}{2}$ was integral over $\mathbb{Z}$. Then for some n we would have,

$$\left(\frac{1}{2}\right)^n + a_{n-1} \left(\frac{1}{2}\right)^{n-1} + \cdots + a_1 \left(\frac{1}{2}\right) + a_0 = 0$$

Then,

$$\begin{aligned}
 0 &= 2^n \left(\left(\frac{1}{2}\right)^n + a_{n-1} \left(\frac{1}{2}\right)^{n-1} + \cdots + a_1 \left(\frac{1}{2}\right) + a_0 \right) \\
 &= 1 + 2a_{n-1} + \cdots + 2^{n-1}a_1 + 2^n a_0 \\
 \implies 1 &= 2(a_{n-1} + \cdots + 2^{n-2}a_1 + 2^{n-1}a_0)
 \end{aligned}$$

But this is a contradiction. Hence $\frac{1}{2}$ is not integral over $\mathbb{Z}$.

Problem 10. Integral Elements Give Finite Modules.

Let S be an R -algebra and let $s \in S$.

1. Suppose that the element $s \in S$ is integral over R , and

$$s^n + a_1 s^{n-1} + \cdots + a_n = 0$$

with $a_i \in R$. Show that every power s^m with $m \geq n$ is an R -linear combination of $1, s, \dots, s^{n-1}$.

2. Deduce that $R[s]$ is generated as an R -module by $1, s, \dots, s^{n-1}$.
3. Find a finite set of R -module generators for $S = \mathbb{K}[t]$ over $R = \mathbb{K}[t^2]$
4. Find a finite set of R -module generators for $S = \mathbb{Z}[i]$ over $R = \mathbb{Z}$.

Solution.

1. We will prove this by induction.

Base Case: $m = n$.

Then we have,

$$s^n = -(a_1 s^{n-1} + \cdots + a_n)$$

which is a R -linear combination of $1, s, \dots, s^{n-1}$.

Inductive Hypothesis: Assume that s^{m-1} can be written as a R -linear combination of $1, s, \dots, s^{n-1}$. Let $s^{m-1} = \sum_{i=0}^{n-1} r_i s^i$.

Inductive Step: Then,

$$\begin{aligned} s^m &= s \cdot s^{m-1} = s \cdot \sum_{i=0}^{n-1} r_i s^i \\ &= \sum_{i=1}^{n-1} r_{i-1} s^i + r_{n-1} s^n \\ &= \sum_{i=1}^{n-1} r_{i-1} s^i + r_{n-1} \left(-(a_1 s^{n-1} + \cdots + a_n) \right) \end{aligned}$$

Thus every power s^m with $m \geq n$ is a R -linear combination of $1, s, \dots, s^{n-1}$.

2. An element of $R[s]$ are finite sums of the form $\sum_i r_i s^i$ with $r_i \in R$. Since s^i with $i \geq n$ can be written as a R -linear combination of $1, s, \dots, s^{n-1}$, $R[s]$ is spanned by $1, s, \dots, s^{n-1}$ and hence is generated as a R -module by $1, s, \dots, s^{n-1}$.

3. $\{1, t\}$

4. $\{1, i\}$

Problem 11. A Fundamental Criterion.

Let S be an R -algebra and let $s \in S$. Prove that the following are equivalent.

1. s is integral over R .
2. $R[s]$ is module-finite over R .
3. There exists a finitely generated R -submodule $M \subseteq S$ such that $1 \in M$ and $sM \subseteq M$.

Solution.

(1 $\implies$ 2)

We have this from problem 10.

(2 $\implies$ 3)

TODO:

(3 $\implies$ 1)

Problem 12. Integral Elements Form a Subring.

Let $R \rightarrow S$ be an R -algebra. Let $\overline{R}^S$ denote the set of elements of S that are integral over R . When R is an integral domain and S is a field we call $\overline{R}^S$ the *integral closure of R in S* . The *integral closure* or *normalization* of a domain R is the integral closure of R in its field of fractions $S = \text{Frac}(R)$. We denote the integral closure of R by $\overline{R}$.

1. If $x \in S$ is integral over R , prove that $R[x]$ is module-finite over R .
2. If $x, y \in S$ are integral over R , prove that $R[x, y]$ is module-finite over R . Hint: First show $R[x]$ is finite over R , then show $R[x, y]$ is finite over $R[x]$.
3. Let $s \in R[x, y]$. Show that multiplication by s defines an R -linear map

$$\mu_s : R[x, y] \longrightarrow R[x, y], \quad f \longmapsto sf.$$

Use Cayley–Hamilton, or the determinant trick, to prove that s is integral over R .

4. Conclude that $x + y$, xy , and $-x$ are integral over R .
5. Conclude that $\overline{R}^S$ is a subring of S containing the image of R .

Solution.

1. We know this from Problem 10.
2. We know that if $x \in S$ is integral over R then, $R[x]$ is module finite over R . We also know that $y \in S$ is integral over R . That is there exist $a_i \in R$ such that,

$$y^n + a_1 y^{n-1} + \cdots + a_n = 0$$

Note that since $a_i \in R$ they are in $R[x]$ and this equation also holds in $R[x]$. That is, y is integral over $R[x]$, and $R[x][y]$ is module-finite over $R[x]$ and thus module-finite over R (Problem 7, part 2).

3. By Cayley-Hamilton we have the characteristic polynomial,

$$p(\mu_s) = \mu_s^n + p_1 \mu_s^{n-1} + \cdots + p_{n-1} \mu_s + p_n = 0$$

with $p_k \in R$. Applying μ_s to $1 \in R[x, y]$ gives us

$$p(s \cdot 1) = s^n + p_1 s^{n-1} + \cdots + p_{n-1} s + p_n = 0$$

which is a polynomial in s with coefficients in R and thus s is integral over R .

4. Then $x + y, xy, -x \in R[x, y]$, and multiplication by each of these is an R -linear map on the finitely generated R -module $R[x, y]$, so Cayley-Hamilton gives us a monic polynomial over R and thus $x + y, xy, -x$ are integral over R .
5. Thus we conclude that for $\alpha : R \rightarrow S$, every $\alpha(r)$ is integral over R and thus $\text{im}(R) \subseteq S$. Moreover, part 4 shows that $\overline{R}^S$ is closed under addition, multiplication and inverses and thus is a subring of S .

Problem 13. Computations with Integrality.

Let $\mathbb{K}$ be a field.

1. Prove that the elements of $\mathbb{Q}$ integral over $\mathbb{Z}$ are exactly the integers. Hint: Write a rational number in lowest terms.
2. Show that $\mathbb{K}[t]$ is module-finite over $\mathbb{K}[t^2, t^3]$. Find explicit module generators.
3. Show that $\mathbb{K}[x, y]/\langle y^2 - x^3 \rangle$ is module-finite over $\mathbb{K}[x]$. Find explicit module generators.
4. Let $S = \mathbb{K}[x, x^{-1}]$, viewed as a $\mathbb{K}[x]$ -algebra. Show that x^{-1} is not integral over $\mathbb{K}[x]$.
5. Deduce that $\mathbb{K}[x, x^{-1}]$ is algebra-finite but not module-finite over $\mathbb{K}[x]$.

Solution.

1. Assume for contradiction that some rational number $\frac{p}{q} \in \mathbb{Q}$, $q \neq 0$ is integral over $\mathbb{Z}$. Without loss of generality we assume $\gcd(p, q) = 1$. Then, there exists a polynomial with coefficients in $\mathbb{Z}$ as follows,

$$\left(\frac{p}{q}\right)^n + a_1 \left(\frac{p}{q}\right)^{n-1} + \cdots + a_n = 0$$

Then,

$$\begin{aligned} 0 &= q^n \left(\left(\frac{p}{q}\right)^n + a_1 \left(\frac{p}{q}\right)^{n-1} + \cdots + a_n \right) \\ &= p^n + a_1 q p^{n-1} + \cdots + a_n q^n \\ \implies p^n &= -q(a_1 p^{n-1} + \cdots + a_n q^{n-1}) \end{aligned}$$

That is $q \mid p^n$, but because $\gcd(p, q) = 1$ we have $\gcd(p^n, q) = 1$, so $q \mid 1 \implies q = \pm 1$. Thus, $\frac{p}{q} \in \mathbb{Z}$. Thus, the only elements of $\mathbb{Q}$ integral over $\mathbb{Z}$ are exactly the integers.

2. Since $\mathbb{K}[t]$ is module-finite over $\mathbb{K}[t^2]$ it is also module-finite over $\mathbb{K}[t^2, t^3]$. The module generators are $\{1, t\}$.
3. Since $\bar{y}^2 = \bar{x}^3$, every element of $\mathbb{K}[x, y]/\langle y^2 - x^3 \rangle$ can be written as $p(x) + q(x)y$ for $p, q \in \mathbb{K}[x]$, so $\{1, \bar{y}\}$ generates it as a $\mathbb{K}[x]$ module.

4. Assume x^{-1} is integral over $\mathbb{K}[x]$. Then, there exists a polynomial with coefficients in $\mathbb{K}[x]$ as follows,

$$\left(\frac{1}{x}\right)^n + a_1 \left(\frac{1}{x}\right)^{n-1} + \cdots + a_n = 0$$

Then,

$$\begin{aligned} 0 &= x^n \left(\left(\frac{1}{x}\right)^n + a_1 \left(\frac{1}{x}\right)^{n-1} + \cdots + a_n \right) \\ &= 1 + a_1 x + \cdots + a_n x^n \\ \implies 1 &= -x(a_1 + \cdots + a_n x^{n-1}) \end{aligned}$$

Hence a contradiction. Thus, x^{-1} is not integral over $\mathbb{K}[x]$.

5. Thus $\mathbb{K}[x, x^{-1}]$ is algebra-finite but not module-finite over $\mathbb{K}[x]$.

Problem 14. UFDs Are Integrally Closed.

Let R be a UFD with fraction field $S = \text{Frac}(R)$.

1. Explain how we can think of S as an R -algebra?
2. Let $\alpha \in S$. Explain why we can write $\alpha = \frac{a}{b}$ with $a, b \in R$, $b \neq 0$, and a and b having no common irreducible factors.
3. Give an example of a domain R that is *not* a UFD for which part (2) is false.
4. Suppose $\alpha = a/b \in S$ is integral over R and

$$\left(\frac{a}{b}\right)^n + r_1 \left(\frac{a}{b}\right)^{n-1} + \cdots + r_{n-1} \left(\frac{a}{b}\right) + r_n = 0$$

for $r_1, \dots, r_n \in R$. By clearing denominators show that b divides a^n .

5. Show that if a and b have no common irreducible factor and $b \mid a^n$, then b is a unit. Hint: Use unique factorization.
6. Conclude that $\alpha = a/b \in R$. Therefore every element of K that is integral over R already lies in R .
7. Conclude that every UFD is integrally closed.
8. Show that $R = \mathbb{K}[t^2, t^3]$ is not integrally closed by finding an explicitly integral element in $S = \text{Frac}(R) = \mathbb{K}(t)$ not contained in R .
9. Show that $\mathbb{Z}[\sqrt{-3}]$ is not integrally closed. What does this mean about it being a UFD?

Solution.

1. We can think of S as an R -algebra with the natural inclusion

$$\iota : R \hookrightarrow S, \quad r \mapsto \frac{r}{1}$$

which is a ring homomorphism.

2. Since $S = \text{Frac}(R)$, every element $\alpha \in S$ is by definition of the form a/b with $a \in R$ and $b \in R \setminus \{0\}$. Since R is a UFD we can cancel common irreducible factors of a and b and get a representation a/b with $\gcd(a, b) = 1$.
3. Let $R = \mathbb{Z}[\sqrt{-5}]$. Then, $\frac{6}{1} = \frac{2 \cdot 3}{1} = \frac{(1+\sqrt{5})(1-\sqrt{5})}{1}$ which fails unique factorization and the argument in part 2 breaks down.

4. We have,

$$\begin{aligned}
0 &= \left(\frac{a}{b}\right)^n + r_1 \left(\frac{a}{b}\right)^{n-1} + \cdots + r_{n-1} \left(\frac{a}{b}\right) + r_n \\
0 &= b^n \left(\left(\frac{a}{b}\right)^n + r_1 \left(\frac{a}{b}\right)^{n-1} + \cdots + r_n \right) \\
0 &= a^n + r_1 b a^{n-1} + \cdots + r_n b^n \\
\implies a^n &= -b(r_1 a^{n-1} + \cdots + r_n b^{n-1}) \\
\implies b &| a^n
\end{aligned}$$

5. If $\gcd(a, b) = 1$ then $\gcd(a^n, b) = 1$, then $b \mid a^n$ implies that b is a unit.

6. Let $bb' = 1$ then $\alpha = ab'/bb' = ab/1 \in R$. Thus, all integral elements of S over R are already in R .

7. Thus, $\overline{R}^S = R$ and R is integrally closed. Since R was an arbitrary UFD this holds for all UFDs.

8. $t \in S$ is integral over R with,

$$1(t^2) - 0(t) - t^2 = 0$$

where $t^2, 0, 1 \in R$. Thus R is not integrally closed.

9. $\mathbb{Z}[\sqrt{3}]$ is a UFD and by part 7 must be integrally closed.

An R -algebra S is called an *integral extension* of R if every element of S is integral over R . Said differently, the structure map $R \rightarrow S$ is integral if all elements of the target satisfy monic polynomial equations over the source. Integral extensions are the precise setting where algebra generation and module generation interact especially well. The exercises below prove the fundamental comparison theorem:

$$\boxed{\text{module-finite}} \iff \boxed{\text{algebra-finite and integral}}.$$

More explicitly, the exercises that follow will guide us through the proof of the following theorem.

Theorem 0.17. *Let $R \rightarrow S$ be an R -algebra. The R -algebra S is module-finite over R if and only if it S is algebra-finite as an R -algebra.*

Note you will often, especially in algebraic geometry, see this expressed as finite is equivalent to finite type and integral. However, again we try to avoid this language for now.

Problem 15. Module-Finite Extensions Are Integral.

Let S be a module-finite R -algebra.

1. Fix $s \in S$. Show that multiplication by s defines an R -linear map

$$\mu_s : S \longrightarrow S \quad \text{given by} \quad x \longmapsto sx.$$

2. Apply the Cayley-Hamilton theorem (explain why you can apply it) to μ_s to deduce that s is integral over R .
3. Conclude that every module-finite R -algebra is integral over R .

Solution.

1. Let $S = R[m_1, \dots, m_n]$. Then from S 's ring axioms we have,

- $\mu_s(x + y) = s(x + y) = sx + sy = \mu_s(x) + \mu_s(y)$
- $\mu_s(rx) = s(rx) = r(sx) = r\mu_s(x)$

Thus multiplication by s defines a R -linear map.

2. Since S is module-finite over R and μ_s is a R -linear map we can apply Cayley Hamilton with $I = R$. Then we get the characteristic polynomial,

$$p(\mu_s) = \mu_s^n + p_1\mu_s^{n-1} + \dots + p_{n-1}\mu_s + p_n = 0$$

with $p_k \in R$. Applying μ_s to $1 \in R$ gives us

$$p(s \cdot 1) = s^n + p_1s^{n-1} + \dots + p_{n-1}s + p_n = 0$$

which is a polynomial in s with coefficients in R and thus s is integral over R .

3. Since s was an arbitrary element in S , this shows that all module-finite R -algebras are integral over R .

Problem 16. Finite Type Integral Extensions Are Module-Finite.

Suppose $S = R[s_1, \dots, s_n]$ and each s_i is integral over R .

1. Prove that $R[s_1]$ is module-finite over R .
2. Prove that $R[s_1, s_2]$ is module-finite over $R[s_1]$, and hence module-finite over R .
3. Continue by induction to prove that S is module-finite over R .
4. Deduce that if S is algebra-finite and integral over R , then S is module-finite over R .

Solution.

1. We know this from Problem 10.
2. We know that if $s_1 \in S$ is integral over R then, $R[s_1]$ is module finite over R . We also know that $s_2 \in S$ is integral over R . That is there exist $a_i \in R$ such that,

$$s_2^n + a_1 s_2^{n-1} + \dots + a_n = 0$$

Note that since $a_i \in R$ they are in $R[s_1]$ and this equation also holds in $R[s_1]$. That is, s_2 is integral over $R[s_1]$, and $R[s_1][s_2]$ is module-finite over $R[s_1]$ and thus module-finite over R (Problem 7, part 2).

3. Assume that $R[s_1, s_2, \dots, s_{n-1}]$ is module-finite over $R[s_1, s_2, \dots, s_{n-2}]$ and hence module-finite over R .

Now consider, $R[s_1, s_2, \dots, s_{n-1}][s_n]$. We know that $s_n \in S$ is integral over R . That is there exist $a_i \in R$ such that,

$$s_n^k + a_1 s_n^{k-1} + \dots + a_k = 0$$

Note that since $a_i \in R$ they are in $R[s_1, s_2, \dots, s_{n-1}]$ and this equation also holds in $R[s_1, s_2, \dots, s_{n-1}]$. That is, s_n is integral over $R[s_1, s_2, \dots, s_{n-1}]$, and $R[s_1, s_2, \dots, s_{n-1}][s_n]$ is module-finite over $R[s_1, s_2, \dots, s_{n-1}]$ and thus module-finite over R (Problem 7, part 2).

4. Since these were arbitrary rings, if S is algebra-finite and integral over R , then S is module-finite over R .

Problem 17. The Comparison Theorem.

Let S be an R -algebra. Prove that the following are equivalent.

1. S is module-finite over R .
2. S is algebra-finite over R and integral over R .

Then revisit Problem 9 and explain how the theorem classifies each example.

Solution.

We showed the direction $(2 \implies 1)$ in the previous question. So we only need to show $(1 \implies 2)$. Assume S is module-finite over R . From Problem 4 part 3 we know that S is algebra-finite over R and from Problem 15 part 3 we know that S is integral over R . Thus $1 \implies 2$.

1. $\mathbb{Z}[i]$ is algebra-finite and integral over $\mathbb{Z}$ and hence is module-finite.
2. $\mathbb{K}[t]$ is algebra-finite and integral over $\mathbb{K}[t^2]$ and hence is module-finite.
3. $\mathbb{Q}(\sqrt{3})$ is not integral over $\mathbb{Z}[\sqrt{3}]$ and hence is not module-finite.
4. $\mathbb{Q}$ is not integral over $\mathbb{Z}$ and hence is not module-finite.

Problem 18. Permanence of Integral Extensions.

Let $R \rightarrow S \rightarrow T$ be maps of rings.

1. Prove that every quotient R/I is integral over R .
2. Suppose S is integral over R and T is integral over S . Prove that T is integral over R . Hint: If $t \in T$ satisfies a monic equation over S , first adjoin to R the finitely many coefficients appearing in that equation.
3. If S is integral over R and $J \subseteq S$ is an ideal, prove that S/J is integral over $R/(J \cap R)$.
4. If S is integral over R and $U \subset R$ is multiplicatively closed, prove that $U^{-1}S$ is integral over $U^{-1}R$.
5. More generally, if S is integral over R and $R \rightarrow R'$ is any ring map, prove that $R' \otimes_R S$ is integral over R' .

Solution.

1. We know that R is a R -module over itself. Then by Problem 7 part 1, R/I is module-finite and thus is integral.
2. Since T is integral over S , for all $t \in T$ we have,

$$t^n + s_1 t^{n-1} + \cdots + s_n = 0$$

where $s_1, \dots, s_n \in S$. But S is integral over R so we have $R[s_1, \dots, s_n]$ a module-finite extension over R . Moreover, t is integral over $R[s_1, \dots, s_n]$ and $R[s_1, \dots, s_n, t] = R[s_1, \dots, s_n][t]$ is module-finite over $R[s_1, \dots, s_n]$ and thus over R . Then by Problem 15, t is integral over R .

3. Since S is integral over R for any $s \in S$ we have,

$$s^n + r_1 s^{n-1} + \cdots + r_n = 0$$

Then for $\bar{s} \in S/J$ we have,

$$\bar{s}^n + \bar{r}_1 \bar{s}^{n-1} + \cdots + \bar{r}_n = 0$$

But $\bar{r}_i = r_i + J \in S = r_i + (J \cap R) \in R$. Hence the coefficients $\bar{r}_i$ lie in $R/\mathfrak{p}$ and $\bar{s}$ is integral over $R/(J \cap R)$.

4. An element in $U^{-1}S$ is of the form $\frac{s}{u} = \frac{1}{u} \cdot \frac{s}{1}$. Since $\frac{1}{u} \in U^{-1}R$ it is integral over $U^{-1}R$. So, we only need to show that $\frac{s}{1}$ is integral over $U^{-1}R$. We know that s is integral over R and there exists some minimal monic polynomial of the form,

$$s^n + r_1 s^{n-1} + \cdots + r_n = 0$$

Then in $U^{-1}S$ by injectivity of η we have,

$$\left(\frac{s}{1}\right)^n + \left(\frac{r_1}{1}\right) \left(\frac{s}{1}\right)^{n-1} + \cdots + \left(\frac{r_n}{1}\right) = \left(\frac{0}{1}\right)$$

with $\frac{r_i}{1} \in U^{-1}R$.

5. We know that $R' \otimes S$ is generated by the tensors $1 \otimes s$. Since s is integral over R we have,

$$s^n + r_1 s^{n-1} + \cdots + r_n = 0$$

Applying the map $S \rightarrow R' \otimes_R S, s \mapsto 1 \otimes s$ gives us

$$(1 \otimes s)^n + r_1 (1 \otimes s)^{n-1} + \cdots + r_n = 0$$

Hence $1 \otimes s$ is integral over R' and since every element of $R' \otimes_R S$ is a finite sum of products of these elements, the integrality is preserved.

Problem 19. A First Consequence for Maximal Ideals.

Let $A \subseteq B$ be an integral extension of domains.

1. Suppose B is a field. Prove that A is a field. Hint: If $0 \neq a \in A$, then $a^{-1} \in B$ is integral over A ; write a monic equation for a^{-1} and multiply by a suitable power of a .
2. Let $R \rightarrow S$ be an integral ring map and let $\mathfrak{q} \in \text{Spec}(S)$ with contraction $\mathfrak{p} = \mathfrak{q} \cap R$. Show that $S/\mathfrak{q}$ is integral over $R/\mathfrak{p}$.
3. Deduce that if $\mathfrak{q}$ is maximal, then $\mathfrak{p}$ is maximal.
4. Prove the converse: if $\mathfrak{p}$ is maximal, then $\mathfrak{q}$ is maximal. Hint: $S/\mathfrak{q}$ is an integral domain integral over the field $R/\mathfrak{p}$.
5. Conclude that under an integral ring map, a prime ideal upstairs is maximal if and only if its contraction downstairs is maximal.

Solution.

1. If $a \neq 0 \in A$ then $a^{-1} \in B$ is integral over A with

$$\left(\frac{1}{a}\right)^n + r_1 \left(\frac{1}{a}\right)^{n-1} + \cdots + r_n = 0$$

And,

$$\begin{aligned} 0 &= a^n \left(\left(\frac{1}{a}\right)^n + r_1 \left(\frac{1}{a}\right)^{n-1} + \cdots + r_n \right) \\ 0 &= 1 + r_1 a + \cdots + r_n a^n \\ \implies 1 &= a(r_1 + \cdots + r_n a^{n-1}) \end{aligned}$$

That is a has an inverse in A . Since a was an arbitrary element this implies that every non-zero element in A has an inverse and A is a field.

2. Since S is integral over R for any $s \in S$ we have,

$$s^n + r_1 s^{n-1} + \cdots + r_n = 0$$

Then for $\bar{s} \in S/\mathfrak{q}$ we have,

$$\bar{s}^n + \bar{r}_1 \bar{s}^{n-1} + \cdots + \bar{r}_n = 0$$

But $\bar{r}_i = r_i + \mathfrak{q} \in S = r_i + \mathfrak{q} \cap R \in R = r_i + \mathfrak{p}$. Hence the coefficients $\bar{r}_i$ lie in $R/\mathfrak{p}$ and $\bar{s}$ is integral over $R/\mathfrak{p}$.

3. If $\mathfrak{q}$ is maximal, then $S/\mathfrak{q}$ is a field and then by part 1 R/fp must be a field and hence $\mathfrak{p}$ must be maximal.
4. If $\mathfrak{p}$ is maximal, then $R/\mathfrak{p}$ is a field, and by part 2 $S/\mathfrak{q}$ is integral over $R/\mathfrak{p}$. Then for any $\bar{s} \neq 0 \in S/\mathfrak{q}$ we have the minimal monic polynomial,

$$\bar{s}^n + \bar{r}_1 \bar{s}^{n-1} + \cdots + \bar{r}_n = 0$$

And since $R/\mathfrak{p}$ is a field we have,

$$\begin{aligned} \bar{r}_n &= -\bar{s}(\bar{s}^{n-1} + \bar{r}_1 \bar{s}^{n-2} + \cdots + \bar{r}_{n-1}) \\ 1 &= \frac{\bar{r}_n}{\bar{r}_n} = -\bar{s} \left(\frac{1}{\bar{r}_n} \bar{s}^{n-1} + \frac{\bar{r}_1}{\bar{r}_n} \bar{s}^{n-2} + \cdots + \frac{\bar{r}_{n-1}}{\bar{r}_n} \right) \end{aligned}$$

Thus $\bar{s}$ has an inverse in $S/\mathfrak{q}$, and hence $S/\mathfrak{q}$ is a field and $\mathfrak{q}$ is maximal.

5. Thus under an integral ring map $\mathfrak{q}$ is maximal if and only if $\mathfrak{p}$ is maximal.

Problem 20. More Examples of Integral Extensions.

Decide whether each map is integral. If it is integral and algebra-finite, give module generators.

1. $\mathbb{Z} \rightarrow \mathbb{Z}[i]$.
2. $\mathbb{K}[t^2] \rightarrow \mathbb{K}[t]$.
3. $\mathbb{K}[t^2, t^3] \rightarrow \mathbb{K}[t]$.
4. $\mathbb{K}[x] \rightarrow \mathbb{K}[x, x^{-1}]$.
5. $\mathbb{K}[x] \rightarrow \mathbb{K}[x, y] / \langle y^2 - x^3 \rangle$.
6. $\mathbb{K}[x^2, xy, y^2] \rightarrow \mathbb{K}[x, y]$. Hint: Show that x and y are integral over the source.

Solution.

1. The map is integral and algebra-finite with $\{1, i\}$ as module generators.
2. The map is integral and algebra-finite with $\{1, t\}$ as module generators.
3. The map is integral and algebra-finite with $\{1, t\}$ as module generators.
4. The map is not integral.
5. The map is module-finite with generators $\{1, y\}$ and thus is integral.
6. The map is integral and algebra-finite with $\{1, x, y\}$ as module generators.

We now introduce a numerical invariant of a ring. A *chain of prime ideals of length d* in a ring R is a strictly increasing chain

$$\mathfrak{p}_0 \subsetneq \mathfrak{p}_1 \subsetneq \cdots \subsetneq \mathfrak{p}_d$$

of prime ideals. Thus the length counts the number of strict inclusions, not the number of prime ideals appearing in the chain. For a nonzero ring R , the *Krull dimension* of R is

$$\dim(R) := \sup \{d \in \mathbb{Z}_{\geq 0} \mid \text{there exists a chain } \mathfrak{p}_0 \subsetneq \cdots \subsetneq \mathfrak{p}_d \text{ in } \operatorname{Spec}(R)\}.$$

If there are chains of arbitrarily large finite length, we write $\dim(R) = \infty$. The zero ring has no prime ideals; different authors use different conventions for its dimension, so we will avoid using the zero ring in basic dimension computations unless a convention is specified. We will eventually prove the following theorem, which will turn out to be key to computing the dimension of most rings.

Theorem 0.18. *If $\mathbb{K}$ is a field then $\dim \mathbb{K}[x_1, x_2, \dots, x_n] = n$.*

Proving that the dimension of the polynomial ring $\mathbb{K}[x_1, x_2, \dots, x_n]$ is at least n is relatively easy, all one must do is find an explicit chain of prime ideals of length n . You will do this in the exercises below. Proving the equality in the theorem requires us to prove the other inequality, i.e. we must show there is no chain of prime ideals of length more than n . This turns out to be surprisingly difficult. For the remainder of the worksheet you may assume this theorem unless it trivializes an exercise.

You will also prove that if R is any ring then $\dim R[x] \geq \dim R + 1$, again by exhibiting an explicit chain of prime ideals. Intuition tells us that this should be an equality, however, this turns out to be false unless we put additional hypotheses on R , namely that R be Noetherian. This highlights a somewhat tedious fact about dimension theory in commutative algebra: unless one assumes some form of finiteness condition (Noetherian, finitely generated $\mathbb{K}$ -algebra, integral, etc.) there are many pathologies.

Using the correspondence between prime ideals and irreducible closed subsets of $\operatorname{Spec}(R)$, the intuition is that longer chains of prime ideals correspond to longer chains of irreducible closed subsets in $\operatorname{Spec}(R)$, ordered in the opposite direction.

Problem 21. Unpacking the Definition.

Let R be a nonzero ring.

1. Show that $\dim(R) = 0$ if and only if every prime ideal of R is maximal.
2. Show that every field has Krull dimension 0.
3. Show that if R is an integral domain that is not a field, then $\dim(R) \geq 1$.
4. Let $I \subseteq R$ be an ideal. Use the correspondence between prime ideals of R/I and prime ideals of R containing I to describe $\dim(R/I)$ in terms of chains of prime ideals in R .
5. Prove that $\dim(R) = \dim(R_{\text{red}})$, where $R_{\text{red}} = R/\text{Nil}(R)$.

Solution.

1. ($\Rightarrow$)

Assume $\dim(R) = 0$. Then the longest chain of strict inclusions in any chain of prime ideals in R is zero. Thus every prime ideal must be maximal (otherwise there would be a maximal ideal containing it).

($\Leftarrow$)

Assume every prime ideal of R is maximal. Then the number of strict inclusions in any chain of prime ideals is zero, and thus $\dim(R) = 0$.

2. Since the only prime ideal in a field is $\langle 0 \rangle$, it is maximal and thus the field has Krull dimension 0.
3. If R is an integral domain $\langle 0 \rangle$ is a prime ideal and if R is not a field then $\langle 0 \rangle$ is not maximal so there must exist at least one maximal ideal containing it (since R is not the zero-ring). Hence, $\dim(R) \geq 1$.
4. $\dim(R/I) = \text{length of longest chain of prime ideals in } R \text{ containing } I$.
5. We know that $\text{Nil}(R) \subseteq \bigcap_i \mathfrak{p}_i$ where $\mathfrak{p}_i$ is a prime ideal in R . Then every chain of prime ideals in R contains $\text{Nil}(R)$ and by part 4,

$$\dim(R) = \dim(R_{\text{red}})$$

Problem 22. Dimension Zero Examples.
Compute the Krull dimension of each ring.

- | | |
|--|---|
| 1. $\mathbb{K}$. | 4. $\mathbb{Z}/12\mathbb{Z}$. |
| 2. $\mathbb{K} \times \mathbb{K}$. | 5. $\mathbb{K}[x]/\langle x^n \rangle$ for $n \geq 1$. |
| 3. $\mathbb{K}[\epsilon]/\langle \epsilon^2 \rangle$. | 6. $\mathbb{K}[x, y]/\langle x, y \rangle$. |

Solution.

- | | |
|--|--|
| 1. $\dim(\mathbb{K}) = 0$. | 4. $\dim(\mathbb{Z}/12\mathbb{Z}) = 0$. |
| 2. $\dim(\mathbb{K} \times \mathbb{K}) = 1$. | 5. $\dim(\mathbb{K}[x]/\langle x^n \rangle) = 0$ |
| 3. $\dim(\mathbb{K}[\epsilon]/\langle \epsilon^2 \rangle) = 0$. | 6. $\dim(\mathbb{K}[x, y]/\langle x, y \rangle) = 0$. |

Problem 23. Dimension One Examples.

1. Prove that $\dim(\mathbb{Z}) = 1$.
2. More generally, prove that every principal ideal domain that is not a field has Krull dimension 1.
3. Prove that $\dim(\mathbb{K}[x]) = 1$.
4. Compute $\dim(\mathbb{K}[x]/\langle f \rangle)$ for a nonzero nonunit polynomial $f \in \mathbb{K}[x]$.
5. Compute $\dim(\mathbb{Z}[1/n])$ for an integer $n \geq 2$.

Solution.

1. We know that every non-zero ideal in $\mathbb{Z}$ is maximal (and the zero ideal is not maximal), so $\dim(\mathbb{Z}) = 1$.
2. We know from Math 71 that every non-zero prime ideal is maximal in a PID (and the zero ideal is not maximal), so every PID that is not a field has Krull dimension 1.
3. $\mathbb{K}[x]$ is a PID, so by part 2 $\dim(\mathbb{K}[x]) = 1$.
4. By Problem 21, part 4 we have that $\dim(\mathbb{K}[x]/\langle f \rangle) = \text{length of longest chain of prime ideals in } \mathbb{K}[x] \text{ containing } \langle f \rangle$. But since f is non-zero and $\mathbb{K}[x]$ is a PID the longest chain containing $\langle f \rangle$ has length 1. Thus, $\dim(\mathbb{K}[x]/\langle f \rangle) = 1$.
5. Let $\mathbb{Z}[1/n] = U^{-1}\mathbb{Z}$, $U = \{1, n, n^2, \dots\}$. Then prime ideals in $\mathbb{Z}[1/n]$ are prime ideals $\mathfrak{p}$ such that $\mathfrak{p} \cap U = \emptyset$. So $\langle 0 \rangle$ and any prime p such that $p \nmid n$ are prime ideals in $\mathbb{Z}[1/n]$. Thus, $\dim(\mathbb{Z}[1/n]) = 1$.

Problem 24. Building Longer Chains.

Let $\mathbb{K}$ be a field.

1. Prove that

$$\langle 0 \rangle \subsetneq \langle x_1 \rangle \subsetneq \langle x_1, x_2 \rangle \subsetneq \cdots \subsetneq \langle x_1, \dots, x_n \rangle$$

is a chain of prime ideals in $\mathbb{K}[x_1, \dots, x_n]$.

2. Deduce that $\dim(\mathbb{K}[x_1, \dots, x_n]) \geq n$.
3. Let R be any nonzero ring. Show that $\dim(R[x]) \geq \dim(R) + 1$ whenever $\dim(R)$ is finite. Hint: Extend a chain in R to $R[x]$, then add the ideal generated by the final prime together with x .

Solution.

1. First note that $x_i \notin \langle x_1, \dots, x_{i-1} \rangle$ since $\langle x_1, \dots, x_{i-1} \rangle$ is of the form $\sum_j k_j x_1^{j_1} \cdots x_{i-1}^{j_{i-1}}$ and x_i cannot be expressed as a combination of $x_1, \dots, x_{i-1}$. Hence the containments are strict.

Moreover the ideals are prime since for any ideal $\langle x_1, \dots, x_{i-1} \rangle$ the quotient ring $\mathbb{K}[x_1, \dots, x_n] / \langle x_1, \dots, x_{i-1} \rangle = \mathbb{K}[x_i, \dots, x_n]$ which is an integral domain.

Hence,

$$\langle 0 \rangle \subsetneq \langle x_1 \rangle \subsetneq \langle x_1, x_2 \rangle \subsetneq \cdots \subsetneq \langle x_1, \dots, x_n \rangle$$

is a chain of prime ideals in $\mathbb{K}[x_1, \dots, x_n]$.

2. The chain in part 1 has length n so $\dim(\mathbb{K}[x_1, \dots, x_n]) \geq n$.
3. Suppose $\dim(R) = n$. Then there exists a chain of length n in R ,

$$\mathfrak{p}_0 \subsetneq \mathfrak{p}_1 \subsetneq \cdots \subsetneq \mathfrak{p}_n$$

We extend this to a chain in $R[x]$ by

$$\mathfrak{p}_0 R[x] \subsetneq \mathfrak{p}_1 R[x] \subsetneq \cdots \subsetneq \mathfrak{p}_n R[x] \subsetneq \mathfrak{p}_n R[x] + \langle x \rangle.$$

Now note that Each $\mathfrak{p}_i R[x]$ is prime since the quotient $R[x] / \mathfrak{p}_i R[x] \cong (R / \mathfrak{p}_i)[x]$ is an integral domain since $R / \mathfrak{p}_i$ is. And $\mathfrak{p}_n R[x] + \langle x \rangle$ is prime since the quotient

$$R[x] / (\mathfrak{p}_n R[x] + \langle x \rangle) \cong R / \mathfrak{p}_n,$$

which is an integral domain.

Moreover the inclusions are strict. Suppose not, then if $\mathfrak{p}_{i-1} R[x] = \mathfrak{p}_i R[x]$ then contracting to R gives $\mathfrak{p}_{i-1} = \mathfrak{p}_i$, a contradiction. And the last inclusion is strict since $x \in \mathfrak{p}_n R[x] + \langle x \rangle$ but $x \notin \mathfrak{p}_n R[x]$ since elements of $\mathfrak{p}_n R[x]$ have all coefficients in $\mathfrak{p}_n$, and $1 \notin \mathfrak{p}_n$. Hence we have a chain of length $n + 1$ in $R[x]$, so $\dim(R[x]) \geq \dim(R) + 1$.

Problem 25. Dimension and Quotients.

Let $\mathbb{K}$ be a field.

1. Compute $\dim(\mathbb{K}[x, y] / \langle xy \rangle)$. Hint: Every prime ideal containing xy contains x or y .
2. Compute $\dim(\mathbb{K}[x, y] / \langle x^2, xy \rangle)$. Hint: First pass to the reduced ring.
3. Compute $\dim(\mathbb{K}[x, y] / \langle y - x^2 \rangle)$.
4. Find a chain of prime ideals of length 2 in $\mathbb{K}[x, y, z] / \langle xy \rangle$.
5. Assuming $\dim(\mathbb{K}[x_1, \dots, x_n]) = n$, compute the dimension of $\mathbb{K}[x, y, z] / \langle z - y^2 + x^3 \rangle$.

Solution.

1. We know that $\dim(\mathbb{K}[x, y] / \langle xy \rangle) =$ Longest chain containing $\langle xy \rangle$ in $\mathbb{K}[x, y]$.
Every chain containing $\langle xy \rangle$ lies either above $\langle x \rangle$ or above $\langle y \rangle$, and by symmetry both give chains of length 2. So, $\dim(\mathbb{K}[x, y] / \langle xy \rangle) = 2$.

2. Let $R = \mathbb{K}[x, y] / \langle x^2, xy \rangle$. Then we have,

$$\dim(R) = \dim(R_{\text{red}}) = \dim(R / \langle \bar{x} \rangle) \cong \dim(\mathbb{K}[y]) = 1$$

3. We have,

$$\mathbb{K}[x, y] / \langle y - x^2 \rangle \cong \mathbb{K}[x, x^2] = \mathbb{K}[x]$$

Then,

$$\dim(\mathbb{K}[x]) = 1$$

4. A chain of prime ideals of length 2 is,

$$\langle \bar{x} \rangle \subsetneq \langle \bar{x}, \bar{z} \rangle \subsetneq \langle \bar{x}, \bar{y}, \bar{z} \rangle$$

5. We have,

$$\mathbb{K}[x, y, z] / \langle z - y^2 + x^3 \rangle \cong \mathbb{K}[x, y, y^2 - x^3] = \mathbb{K}[x, y]$$

Then,

$$\dim(\mathbb{K}[x, y]) = 2$$

Problem 26. Local Dimension.

Let $\mathfrak{p} \in \text{Spec}(R)$.

1. Recall that prime ideals of $R_{\mathfrak{p}}$ correspond to prime ideals $\mathfrak{q} \subseteq R$ with $\mathfrak{q} \subseteq \mathfrak{p}$. Use this to describe $\dim(R_{\mathfrak{p}})$ in terms of chains of prime ideals in R ending inside $\mathfrak{p}$.
2. Compute $\dim(\mathbb{K}[x]_{\langle x \rangle})$.
3. Compute $\dim(\mathbb{Z}_{\langle p \rangle})$ for a prime number p .
4. Find a chain showing that $\dim(\mathbb{K}[x, y]_{\langle x, y \rangle}) \geq 2$.
5. Assuming $\dim(\mathbb{K}[x, y]) = 2$, prove that $\dim(\mathbb{K}[x, y]_{\langle x, y \rangle}) = 2$.

Solution.

1. Since that prime ideals of $R_{\mathfrak{p}}$ correspond to prime ideals $\mathfrak{q} \subseteq R$ with $\mathfrak{q} \subseteq \mathfrak{p}$, $\dim(R_{\mathfrak{p}}) =$ Longest chain of prime ideals in R with $\mathfrak{p}$ as the maximal ideal.
2. $\dim(\mathbb{K}[x]_{\langle x \rangle}) = 1$.
3. $\dim(\mathbb{Z}_{\langle p \rangle}) = 1$.
4. $\langle 0 \rangle \subsetneq \langle x \rangle \subsetneq \langle x, y \rangle$. So $\dim(\mathbb{K}[x, y]_{\langle x, y \rangle}) \geq 2$.
5. If $\dim(\mathbb{K}[x, y]) = 2$ then the longest chain of prime ideals in $\mathbb{K}[x, y]$ is of length 2 so $\dim(\mathbb{K}[x, y]_{\langle x, y \rangle}) \leq 2$. Thus,

$$\dim(\mathbb{K}[x, y]_{\langle x, y \rangle}) = 2$$

Worksheet 7.2: Integral Extension Pt. II

Throughout this course, “ring” means *commutative* ring with unity and all ring homomorphisms preserve unity. In the previous worksheet we introduced integral elements and integral extensions. We saw that integrality is the mechanism by which monic polynomial equations turn algebraic generation into module generation. The goal of this worksheet is to see what integrality does to prime ideals and to use this to make dimension computable for finitely generated algebras over a field.

Let $\phi : R \rightarrow S$ be a ring map. Recall that ϕ induces a continuous map

$$\mathrm{Spec}(S) \xrightarrow{\phi^\#} \mathrm{Spec}(R), \quad \mathfrak{q} \longmapsto \phi^{-1}(\mathfrak{q}).$$

The central question of the first half of this worksheet is: *how much of $\mathrm{Spec}(R)$ is seen by $\mathrm{Spec}(S)$, and how do chains of prime ideals compare?* For integral extensions the answer is surprisingly rigid. Prime ideals downstairs have primes upstairs, chains can be lifted upward, and no two comparable upstairs primes can lie over the same downstairs prime. These facts are called *Lying Over*, *Going Up*, and *Incomparability*. A complementary statement, *Going Down*, is more delicate: it is true for flat maps, and for integral extensions of domains when the base is integrally closed.

In the second half of the worksheet we use these theorems to prove and apply *Noether Normalization*. The slogan is that every finitely generated algebra over a field is finite over a polynomial ring. Thus, after a finite map, an affine algebra behaves like affine space of some dimension. This is one of the main reasons polynomial rings control dimension theory.

Problem 1. A First Consequence for Maximal Ideals.

Let $\phi : R \rightarrow S$ be an injective integral ring map.

1. Suppose S is a field and R is a domain. Prove that R is a field. Hint: If $r \neq 0 \in R$, then $\phi(r)^{-1} \in S$ is integral over R ; write a monic equation for $\phi(r)^{-1}$ and multiply by a suitable power of $\phi(r)$.
2. Suppose S is a domain and R is a field. Prove that S is a field.
3. Let $\phi : R \rightarrow S$ be an integral ring map and let $\mathfrak{q} \in \text{Spec}(S)$ with contraction $\mathfrak{p} = \phi^{-1}(\mathfrak{q})$. Show that $S/\mathfrak{q}$ is integral over $R/\mathfrak{p}$.
4. Deduce that if $\mathfrak{q}$ is maximal, then $\mathfrak{p}$ is maximal.
5. Prove the converse: If $\mathfrak{p}$ is maximal, then $\mathfrak{q}$ is maximal. Hint: $S/\mathfrak{q}$ is an integral domain integral over the field $R/\mathfrak{p}$.
6. Conclude that under an integral ring map, a prime ideal upstairs is maximal if and only if its contraction downstairs is maximal.

Solution.

1. If $r \neq 0 \in R$ then $\phi(r)^{-1} \in S$ is integral over R with

$$(\phi(r)^{-1})^n + r_1(\phi(r)^{-1})^{n-1} + \cdots + r_n = 0$$

And,

$$\begin{aligned} 0 &= \phi(r)^n \left((\phi(r)^{-1})^n + r_1(\phi(r)^{-1})^{n-1} + \cdots + r_n \right) \\ 0 &= 1 + r_1\phi(r) + \cdots + r_n\phi(r)^n \\ \implies 1 &= \phi(r)(r_1 + \cdots + r_n\phi(r)^{n-1}) \end{aligned}$$

That is $\phi(r)$ has an inverse in $\phi(R)$. Since ϕ is injective this implies that r has an inverse in R . Since $\phi(r)$ was an arbitrary element this implies that every non-zero element in R has an inverse and R is a field.

2. We know that $s \neq 0 \in S$ is integral over R . That is, we have the minimal monic polynomial,

$$s^n + r_1s^{n-1} + \cdots + r_n = 0$$

Since R is a field we have,

$$\begin{aligned}
0 &= s^n + r_1 s^{n-1} + \cdots + r_n \\
r_n &= -(s^n + r_1 s^{n-1} + \cdots + r_{n-1} s) \\
-r_n &= s(s^{n-1} + r_1 s^{n-2} + \cdots + r_{n-1}) \\
1 &= \frac{-r_n}{-r_n} = s \left(\frac{1}{r_n} s^{n-1} + \frac{r_1}{r_n} s^{n-2} + \cdots + \frac{r_{n-1}}{r_n} \right)
\end{aligned}$$

That is s has an integral inverse in S . Since s was an arbitrary non-zero element this shows that S is a field.

3. Since S is integral over R for any $s \in S$ we have,

$$s^n + r_1 s^{n-1} + \cdots + r_n = 0$$

Then for $\bar{s} \in S/\mathfrak{q}$ we have,

$$\bar{s}^n + \bar{r}_1 \bar{s}^{n-1} + \cdots + \bar{r}_n = 0$$

But $\bar{r}_i = r_i + \mathfrak{q} \in S = r_i + \mathfrak{q} \cap R \in R = r_i + \mathfrak{p}$. Hence the coefficients $\bar{r}_i$ lie in $R/\mathfrak{p}$ and $\bar{s}$ is integral over $R/\mathfrak{p}$.

4. If $\mathfrak{q}$ is maximal, then $S/\mathfrak{q}$ is a field and then by part 1 $R/\mathfrak{f}\mathfrak{p}$ must be a field and hence $\mathfrak{p}$ must be maximal.
5. If $\mathfrak{p}$ is maximal, then $R/\mathfrak{p}$ is a field, and by part 2 $S/\mathfrak{q}$ is integral over $R/\mathfrak{p}$. Then for any $\bar{s} \neq 0 \in S/\mathfrak{q}$ we have the minimal monic polynomial,

$$\bar{s}^n + \bar{r}_1 \bar{s}^{n-1} + \cdots + \bar{r}_n = 0$$

And since $R/\mathfrak{p}$ is a field we have,

$$\begin{aligned}
\bar{r}_n &= -\bar{s}(\bar{s}^{n-1} + \bar{r}_1 \bar{s}^{n-2} + \cdots + \bar{r}_{n-1}) \\
1 &= \frac{\bar{r}_n}{\bar{r}_n} = -\bar{s} \left(\frac{1}{\bar{r}_n} \bar{s}^{n-1} + \frac{\bar{r}_1}{\bar{r}_n} \bar{s}^{n-2} + \cdots + \frac{\bar{r}_{n-1}}{\bar{r}_n} \right)
\end{aligned}$$

Thus $\bar{s}$ has an inverse in $S/\mathfrak{q}$, and hence $S/\mathfrak{q}$ is a field and $\mathfrak{q}$ is maximal.

6. Thus under an integral ring map $\mathfrak{q}$ is maximal if and only if $\mathfrak{p}$ is maximal.

Problem 2. Prime Ideals Near a Point.

Let $\phi : R \rightarrow S$ be a ring map and let $\mathfrak{p} \in \text{Spec}(R)$. Write $U = R \setminus \mathfrak{p}$ and $S_{\mathfrak{p}} = U^{-1}S$.

1. Prove that if $\mathfrak{q} \in \text{Spec}(S)$, then $\phi^{-1}(\mathfrak{q})$ is a prime ideal of R .
2. Prove that prime ideals of $S_{\mathfrak{p}}$ are in bijection with prime ideals $\mathfrak{q} \in \text{Spec}(S)$ such that $\mathfrak{q} \cap \phi(U) = \emptyset$. Equivalently, these are the primes $\mathfrak{q}$ such that $\phi^{-1}(\mathfrak{q}) \subseteq \mathfrak{p}$.
3. Suppose that $R \subseteq S$ is an injective ring map. Show that $S_{\mathfrak{p}} \neq 0$. Hint: If $1 = 0$ in $S_{\mathfrak{p}}$, then some element of $R \setminus \mathfrak{p}$ becomes zero in S .
4. Suppose that S is integral over R . Prove that $S_{\mathfrak{p}}$ is integral over $R_{\mathfrak{p}}$.
5. Let $\mathfrak{Q} \in \text{Spec}(S_{\mathfrak{p}})$ and let $\mathfrak{q} \subset S$ be the corresponding prime ideal. Show that if $\phi_{\mathfrak{p}}^{-1}(\mathfrak{Q}) = \mathfrak{p} R_{\mathfrak{p}}$, then $\phi^{-1}(\mathfrak{q}) = \mathfrak{p}$.

Solution.

1. If $\mathfrak{q} \in \text{Spec}(S)$ then $\mathfrak{q}$ is a prime ideal and we know that pre-images of prime ideals are prime, hence $\phi^{-1}(\mathfrak{q})$ is a prime ideal of R .
2. We have already proved this.
3. Assume for contradiction that $S_{\mathfrak{p}} = 0$. Then $\frac{0}{1} \sim \frac{1}{1} \implies \exists t \in \phi(R \setminus \mathfrak{p})$ such that $t(1 - 0) = 0 \implies t = 0 \implies t \in \ker(\phi)$ but ϕ is injective so t must be 0 and $0 \notin R \setminus \mathfrak{p}$. A contradiction. Hence, $S_{\mathfrak{p}} \neq 0$.
4. In Worksheet 7.1 Problem 17 that if S is integral over R and $U \subset R$ is multiplicatively closed, then $U^{-1}S$ is integral over $U^{-1}R$. Since $S_{\mathfrak{p}} = U^{-1}S$ with $U = R \setminus \mathfrak{p}$, we have $S_{\mathfrak{p}}$ is integral over $R_{\mathfrak{p}}$.
5. We have,

$$\begin{array}{ccc} R & \xrightarrow{\phi} & S \\ \downarrow & & \downarrow \\ R_{\mathfrak{p}} & \xrightarrow{\phi_{\mathfrak{p}}} & S_{\mathfrak{p}} \end{array}$$

Where

$$\phi_{\mathfrak{p}}^{-1}(\mathfrak{Q}) = \mathfrak{p} R_{\mathfrak{p}} = \mathfrak{Q} \cap R_{\mathfrak{p}}$$

And

$$\phi^{-1}(\mathfrak{q}) = \phi^{-1}(\mathfrak{Q} \cap S) \cap R = \phi_{\mathfrak{p}}^{-1}(\mathfrak{Q}) \cap R = \mathfrak{p} R_{\mathfrak{p}} \cap R = \mathfrak{p}$$

Problem 3. Lying Over.

Let $R \subseteq S$ be an integral extension of rings. The goal of this exercise is to prove the *Lying Over Theorem*: for every $\mathfrak{p} \in \text{Spec}(R)$, there exists $\mathfrak{q} \in \text{Spec}(S)$ such that $\mathfrak{q} \cap R = \mathfrak{p}$.

1. Fix $\mathfrak{p} \in \text{Spec}(R)$. Explain why $S_{\mathfrak{p}}$ is a nonzero integral extension of $R_{\mathfrak{p}}$.
2. Choose a maximal ideal $\mathfrak{N} \subset S_{\mathfrak{p}}$. Use the theorem from the previous worksheet saying that maximal ideals contract to maximal ideals under integral extensions to prove

$$\mathfrak{N} \cap R_{\mathfrak{p}} = \mathfrak{p} R_{\mathfrak{p}}.$$

3. Pull $\mathfrak{N}$ back to a prime ideal $\mathfrak{q} \subset S$ and prove that $\mathfrak{q} \cap R = \mathfrak{p}$.
4. State the correct version of Lying Over for a not-necessarily-injective integral map $\phi : R \rightarrow S$. Hint: Which prime ideals of R can possibly be contractions of prime ideals of S ?
5. Translate Lying Over into a statement about the image of the map $\phi^{\#} : \text{Spec}(S) \rightarrow \text{Spec}(R)$.

Solution.

1. In Worksheet 7.1 Problem 17 that if S is integral over R and $U \subset R$ is multiplicatively closed, then $U^{-1}S$ is integral over $U^{-1}R$. Since $S_{\mathfrak{p}} = U^{-1}S$ with $U = R \setminus \mathfrak{p}$, we have $S_{\mathfrak{p}}$ is integral over $R_{\mathfrak{p}}$.
2. Since maximal ideals contract to maximal ideals under integral extensions we know that $\phi_{\mathfrak{p}}^{-1}(\mathfrak{N})$ is a maximal ideal in $R_{\mathfrak{p}}$. Since there is only one maximal ideal in $R_{\mathfrak{p}}$, namely $\mathfrak{p} R_{\mathfrak{p}}$, we must have

$$\mathfrak{N} \cap R_{\mathfrak{p}} = \phi_{\mathfrak{p}}^{-1}(\mathfrak{N}) = \mathfrak{p} R_{\mathfrak{p}}$$

3. From Problem 2 part 5, we have that if $\mathfrak{q} \subset S$ corresponds to $\mathfrak{N}$ then

$$\phi^{-1}(\mathfrak{q}) = \mathfrak{q} \cap R = \mathfrak{p}$$

4. From the first isomorphism theorem we have $R/\ker(\phi) \cong \phi(R) \subseteq S$. Since ϕ is integral, prime ideals in S correspond to prime ideals in $R/\ker(\phi)$ which also correspond to prime ideals in R containing $\ker(\phi)$. So the Lying Over Theorem for not injective integral mps says that for every $\mathfrak{p} \in \text{Spec}(R)$ such that $\ker(\phi) \subseteq \mathfrak{p}$, there exists $\mathfrak{q} \in \text{Spec}(S)$ such that $\mathfrak{q} \cap R = \mathfrak{p}$.
5. The image of the map $\phi^{\#}$ is surjective onto $\text{Spec}(R/\ker(\phi))$, or the $\text{im}(\phi^{\#})$ is precisely the set of prime ideals in R containing the kernel of the map ϕ .

Problem 4. First Examples of Lying Over.

Let $\mathbb{K}$ be a field.

1. Show that $\mathbb{K}[t]$ is integral over $\mathbb{K}[t^2]$. For a prime ideal $\mathfrak{q} \subset \mathbb{K}[t]$, compute $\mathfrak{q} \cap \mathbb{K}[t^2]$ when $\mathfrak{q} = \langle t - a \rangle$.
2. Assume $\mathbb{K}$ is algebraically closed and $\text{char}(\mathbb{K}) \neq 2$. For $u = t^2$, describe the primes of $\mathbb{K}[t]$ lying over the maximal ideal $\langle u - a \rangle \subset \mathbb{K}[u]$ when $a = 0$ and when $a \neq 0$.
3. Show that $\mathbb{K}[t]$ is integral over $\mathbb{K}[t^2, t^3]$. Identify the prime of $\mathbb{K}[t]$ lying over the cusp point $\langle t^2, t^3 \rangle \subset \mathbb{K}[t^2, t^3]$.
4. Show that $\mathbb{Z}[i]$ is integral over $\mathbb{Z}$. Compute the primes of $\mathbb{Z}[i]$ lying over $\langle 2 \rangle$, $\langle 3 \rangle$, and $\langle 5 \rangle$ by factoring $x^2 + 1$ modulo 2, 3, and 5.
5. Let $R \subseteq S$ be integral and let $I \subset R$ be an ideal. Prove that the induced map

$$\{\mathfrak{q} \in \text{Spec}(S) \mid \mathfrak{q} \supseteq IS\} \longrightarrow \{\mathfrak{p} \in \text{Spec}(R) \mid \mathfrak{p} \supseteq I\}$$

is surjective.

Solution.

1. $\mathbb{K}[t]$ is module-finite over $\mathbb{K}[t^2]$ and hence is integral over $\mathbb{K}[t^2]$.

$$\mathfrak{q} \cap \mathbb{K}[t^2] = \{p(t^2) \in \mathbb{K}[t^2] \mid p(a^2) = 0\} = \langle t^2 - a^2 \rangle$$

2. Prime ideals that lie over the maximal ideal $\mathfrak{p} = \langle t^2 - a \rangle \subseteq \mathbb{K}[t^2]$ are maximal ideals $\mathfrak{q}$ in $\mathbb{K}[t]$ such that $\mathfrak{q} \cap \mathbb{K}[t^2] = \mathfrak{p}$. Maximal ideals in $\mathbb{K}[t]$ are of the form $\langle t - b \rangle$, and by part 1 we know that $\langle t - b \rangle$ lies over $\mathfrak{p}$ iff $b^2 = a$.

If $a = 0$,

$$\langle t - b \rangle \cap \mathbb{K}[t^2] = \langle t^2 \rangle \implies b = 0$$

So there is a single prime ideal $\langle t \rangle$ lying over $\langle t^2 \rangle$.

If $a \neq 0$, then $b = \pm\sqrt{a}$ since $\mathbb{K}$ is algebraically closed. Hence, two prime ideals lie over $\mathfrak{p}$, $\langle t - \sqrt{a} \rangle$ and $\langle t + \sqrt{a} \rangle$.

3. $\mathbb{K}[t]$ is module-finite over $\mathbb{K}[t^2, t^3]$ and hence is integral over $\mathbb{K}[t^2, t^3]$. Primes of $\mathbb{K}[t]$ are of the form $\langle t - b \rangle$ for $b \in \mathbb{K}$ and $\langle 0 \rangle$. If $\langle t - b \rangle$ lies over $\langle t^2, t^3 \rangle$ then

$$t^2, t^3 \in \langle t - b \rangle \implies b^2 = b^3 = 0 \implies b = 0$$

Thus, $\langle t \rangle$ lies over $\langle t^2, t^3 \rangle$.

4. $\mathbb{Z}[i]$ is module-finite over $\mathbb{Z}$ and hence is integral over $\mathbb{Z}$. We know $\mathbb{Z}[i] \cong \mathbb{Z}[x]/\langle x^2 + 1 \rangle$, and primes of $\mathbb{Z}[i]$ containing $\langle p \rangle$ correspond to prime ideals of $\mathbb{Z}[i]/\langle p \rangle$. Then,

$$\mathbb{Z}[i]/\langle p \rangle \cong \mathbb{F}_p[i] \cong \mathbb{F}_p[x]/\langle x^2 + 1 \rangle$$

so primes correspond to irreducible factors of $\langle x^2 + 1 \rangle$ in $\mathbb{F}_p[x]$ since they generate the ideals containing it. Therefore,

- $x^2 + 1 = (x + 1)^2 \pmod{2}$ so $\langle 1 + i \rangle$ lies over $\langle 2 \rangle$.
- $x^2 + 1$ is irreducible modulo 3 since $3 \pmod{4} \equiv 3$ and thus -1 is not a square, so $\langle 3 \rangle$ lies over $\langle 3 \rangle$.
- $x^2 + 1 = (x + 2)(x - 2) \pmod{5}$ so $\langle i + 2 \rangle$ and $\langle i - 2 \rangle$ lie over $\langle 5 \rangle$.

5. Let $\mathfrak{p} \in \text{Spec}(R)$ such that $I \subseteq \mathfrak{p}$. Then by laying over we know that there exists $\mathfrak{q}$ in $\text{Spec}(S)$ such that $\mathfrak{q} \cap R = \mathfrak{p}$. Since $\mathfrak{q}$ is an ideal and we have $I \subseteq \mathfrak{p} \subseteq \mathfrak{q}$ as sets, $IS \subseteq \mathfrak{q}$. Hence, proving surjectivity.

The next theorem explains how integral extensions interact with inclusions of prime ideals. Geometrically, if $\mathfrak{p}_0 \subseteq \mathfrak{p}_1$ in $\text{Spec}(R)$, then $\mathfrak{p}_1$ is a specialization of $\mathfrak{p}_0$. The theorem says that specializations can be lifted through an integral extension once the starting point upstairs has been chosen.

Theorem 0.19 (Going Up). *Let $R \subseteq S$ be an integral extension. Suppose*

$$\mathfrak{p}_0 \subseteq \mathfrak{p}_1$$

are prime ideals of R , and suppose $\mathfrak{q}_0 \in \text{Spec}(S)$ satisfies $\mathfrak{q}_0 \cap R = \mathfrak{p}_0$. Then there exists $\mathfrak{q}_1 \in \text{Spec}(S)$ such that

$$\mathfrak{q}_0 \subseteq \mathfrak{q}_1 \quad \text{and} \quad \mathfrak{q}_1 \cap R = \mathfrak{p}_1.$$

More generally, every chain of prime ideals downstairs can be lifted to a chain of prime ideals upstairs, after choosing a prime upstairs over the first prime in the chain.

Problem 5. Proof of Going Up.

Let $R \subseteq S$ be integral, let $\mathfrak{p}_0 \subseteq \mathfrak{p}_1$ be prime ideals of R , and let $\mathfrak{q}_0 \in \text{Spec}(S)$ lie over $\mathfrak{p}_0$.

1. Prove that $S/\mathfrak{q}_0$ is integral over $R/\mathfrak{p}_0$.
2. Show that $\mathfrak{p}_1/\mathfrak{p}_0$ is a prime ideal of $R/\mathfrak{p}_0$.
3. Apply Lying Over to the integral extension $R/\mathfrak{p}_0 \subseteq S/\mathfrak{q}_0$ to produce a prime ideal $\mathfrak{q}_1/\mathfrak{q}_0$ of $S/\mathfrak{q}_0$ lying over $\mathfrak{p}_1/\mathfrak{p}_0$.
4. Check that the corresponding prime ideal $\mathfrak{q}_1 \subset S$ satisfies $\mathfrak{q}_0 \subseteq \mathfrak{q}_1$ and $\mathfrak{q}_1 \cap R = \mathfrak{p}_1$.
5. Prove the chain version of Going Up by induction: if

$$\mathfrak{p}_0 \subsetneq \mathfrak{p}_1 \subsetneq \cdots \subsetneq \mathfrak{p}_n$$

is a chain in R and $\mathfrak{q}_0$ lies over $\mathfrak{p}_0$, then there is a chain

$$\mathfrak{q}_0 \subsetneq \mathfrak{q}_1 \subsetneq \cdots \subsetneq \mathfrak{q}_n$$

in S with $\mathfrak{q}_i \cap R = \mathfrak{p}_i$ for all i .

Solution.

1. This follows from Problem 1 part 5.
2. We know that prime ideals of $R/\mathfrak{p}_0$ correspond to prime ideals of R containing $\mathfrak{p}_0$. Since $\mathfrak{p}_0 \subseteq \mathfrak{p}_1$, $\overline{\mathfrak{p}_1} = \mathfrak{p}_1/\mathfrak{p}_0$ is a prime ideal of $R/\mathfrak{p}_0$.
3. We know that the map $R/\mathfrak{p}_0 \rightarrow S/\mathfrak{q}_0$ is an integral map. By laying over, we know there exists $\overline{\mathfrak{q}_1} \subseteq S/\mathfrak{q}_0$ that lies over $\overline{\mathfrak{p}_1}$. Since prime ideals of $S/\mathfrak{q}_0$ correspond to prime ideals of S containing $\mathfrak{q}_0$, $\overline{\mathfrak{q}_1} = \mathfrak{q}_1/\mathfrak{q}_0$.
4. Thus, $\mathfrak{q}_0 \subseteq \mathfrak{q}_1$. We know $\mathfrak{q}_1/\mathfrak{q}_0 \cap R/\mathfrak{p}_0 = \mathfrak{p}_1/\mathfrak{p}_0$. $\mathfrak{q}_1 \cap R$ is the pre-image of the quotient map $\pi : R \rightarrow R/\mathfrak{p}_0$, then,

$$\pi^{-1}((\mathfrak{q}_1/\mathfrak{q}_0) \cap (R/\mathfrak{p}_0)) = \pi^{-1}(\mathfrak{p}_1/\mathfrak{p}_0) = \mathfrak{p}_1$$

5. Assume for $m < n$ that if

$$\mathfrak{p}_0 \subsetneq \mathfrak{p}_1 \subsetneq \cdots \subsetneq \mathfrak{p}_m$$

is a chain in R and $\mathfrak{q}_0$ lies over $\mathfrak{p}_0$, then there is a chain

$$\mathfrak{q}_0 \subsetneq \mathfrak{q}_1 \subsetneq \cdots \subsetneq \mathfrak{q}_m$$

in S with $q_i \cap R = p_i$ for all i .

Then if $p_m \subsetneq p_n$, then by parts 3 and 4, we know there exists $q_n \subseteq S$ such that $q_m \subset q_n$ and $q_n \cap R = p_n$. Note that the inclusion $q_m \subsetneq q_n$ is strict since the contractions correspond to p_m and p_n , and the inclusion $p_m \subsetneq p_n$ is strict.

Thus for any n , if

$$p_0 \subsetneq p_1 \subsetneq \cdots \subsetneq p_n$$

is a chain in R and q_0 lies over p_0 , then there is a corresponding chain

$$q_0 \subsetneq q_1 \subsetneq \cdots \subsetneq q_n$$

in S with $q_i \cap R = p_i$ for all i .

Problem 6. Incomparability.

Let $R \subseteq S$ be an integral extension. The goal of this exercise is to prove: if $\mathfrak{q} \subseteq \mathfrak{q}'$ are prime ideals of S and $\mathfrak{q} \cap R = \mathfrak{q}' \cap R$, then $\mathfrak{q} = \mathfrak{q}'$.

1. Set $\mathfrak{p} = \mathfrak{q} \cap R = \mathfrak{q}' \cap R$. Explain why $R/\mathfrak{p} \subseteq S/\mathfrak{q}$ is an integral extension of domains.
2. Let $\mathfrak{P} = \mathfrak{q}'/\mathfrak{q} \subset S/\mathfrak{q}$. Show that $\mathfrak{P} \cap (R/\mathfrak{p}) = \langle 0 \rangle$.
3. Suppose $0 \neq b \in \mathfrak{P}$. Choose a monic equation of minimal degree

$$b^n + a_1 b^{n-1} + \cdots + a_{n-1} b + a_n = 0$$

with $a_i \in R/\mathfrak{p}$. Prove that $a_n = 0$.

4. Use the fact that $S/\mathfrak{q}$ is a domain to get a monic equation for b of smaller degree, a contradiction.
5. Conclude Incomparability.

Solution.

1. This follows from Problem 1 part 5.

2. If $r + \mathfrak{q}, r' + \mathfrak{q} \in \mathfrak{q}'/\mathfrak{q}$ then $r - r' \in \mathfrak{q} \subseteq \mathfrak{q}'$, that is,

$$\mathfrak{P} \cap (R/\mathfrak{p}) = \{r + \mathfrak{p} \mid r + \mathfrak{q} \in \mathfrak{q}'/\mathfrak{q}\} = \{r + \mathfrak{p} \mid r \in \mathfrak{q}' \cap R\} = \{r + \mathfrak{p} \mid r \in \mathfrak{p}\} = \langle 0 \rangle$$

3. Since $b \in \mathfrak{P}$ we have $b^n + a_1 b^{n-1} + \cdots + a_{n-1} b \in \mathfrak{P}$ which implies that $-a_n \in \mathfrak{P}$. But since $a_n \in R/\mathfrak{p} \implies a_n \in \mathfrak{P} \cap (R/\mathfrak{p}) = \langle 0 \rangle \implies a_n = 0$.
4. We have,

$$b(b^{n-1} + a_1 b^{n-2} + \cdots + a_{n-1}) = 0$$

Since $S/\mathfrak{q}$ is a domain and $b \neq 0$ we must have

$$b^{n-1} + a_1 b^{n-2} + \cdots + a_{n-1} = 0$$

Thus giving us a monic equation for b of smaller degree, which is a contradiction since we assumed minimality. Thus $b = 0$.

5. Part 4 shows us that $\mathfrak{P}$ has no non-zero elements and thus $\mathfrak{q}'/\mathfrak{q} = \langle 0 \rangle \implies \mathfrak{q}' = \mathfrak{q}$.

Problem 7. Dimension is Preserved by Integral Extensions.

Let $R \subseteq S$ be an integral extension of nonzero rings.

1. Let

$$\mathfrak{q}_0 \subsetneq \mathfrak{q}_1 \subsetneq \cdots \subsetneq \mathfrak{q}_n$$

be a chain of prime ideals in S . Use Incomparability to show that

$$\mathfrak{q}_0 \cap R \subsetneq \mathfrak{q}_1 \cap R \subsetneq \cdots \subsetneq \mathfrak{q}_n \cap R$$

is a chain of prime ideals in R . Deduce that $\dim(S) \leq \dim(R)$.

2. Let

$$\mathfrak{p}_0 \subsetneq \mathfrak{p}_1 \subsetneq \cdots \subsetneq \mathfrak{p}_n$$

be a chain of prime ideals in R . Use Lying Over and Going Up to lift it to a chain of prime ideals in S . Deduce that $\dim(R) \leq \dim(S)$.

3. Conclude that $\dim(R) = \dim(S)$.

4. Use this theorem to compute the Krull dimension of each ring below.

(a) $\mathbb{K}[t^2, t^3]$.

(c) $\mathbb{K}[x^2, xy, y^2]$.

(b) $\mathbb{K}[x, y] / \langle y^2 - x^3 \rangle$.

(d) $\mathbb{Z}[i]$.

Solution.

1. Since the preimage of prime ideals is prime we know that each $\mathfrak{q}_i \cap R$ is prime, and by Incomparability all containments are strict. Then if

$$\mathfrak{q}_0 \subsetneq \mathfrak{q}_1 \subsetneq \cdots \subsetneq \mathfrak{q}_n$$

is the longest chain of prime ideals in S , we get a chain of prime ideals in R that has the same length

$$\mathfrak{q}_0 \cap R \subsetneq \mathfrak{q}_1 \cap R \subsetneq \cdots \subsetneq \mathfrak{q}_n \cap R$$

Hence $\dim(S) \leq \dim(R)$.

2. If

$$\mathfrak{p}_0 \subsetneq \mathfrak{p}_1 \subsetneq \cdots \subsetneq \mathfrak{p}_n$$

is a chain of prime ideals in R . Then by laying over we know that there exist prime ideals

$$\mathfrak{q}_0, \mathfrak{q}_1, \dots, \mathfrak{q}_n$$

lying over each corresponding ideal in R . Moreover from going over we know that for each pair $\mathfrak{p}_i \subseteq \mathfrak{p}_{i+1}$ such that there exists a $\mathfrak{q}_i$ that lies over $\mathfrak{p}_i$, there exists an prime ideal $\mathfrak{q}_{i+1} \subseteq S$ that lies over $\mathfrak{p}_{i+1}$ and contains $\mathfrak{q}_i$. Thus we get a chain of prime ideals in S ,

$$\mathfrak{q}_0 \subsetneq \mathfrak{q}_1 \subsetneq \cdots \subsetneq \mathfrak{q}_n$$

Since the chains are of equal length we have $\dim(S) \geq \dim(R)$.

3. Thus, $\dim(S) = \dim(R)$.

4. (a) Since $\mathbb{K}[t]$ is integral over $\mathbb{K}[t^2, t^3]$,

$$\dim(\mathbb{K}[t^2, t^3]) = \dim(\mathbb{K}[t]) = 1$$

(b) Since $\mathbb{K}[x, y] / \langle y^2 - x^3 \rangle$ is module-finite (and thus integral over) $\mathbb{K}[x]$,

$$\dim(\mathbb{K}[x, y] / \langle y^2 - x^3 \rangle) = \dim(\mathbb{K}[x]) = 1$$

(c) Since $\mathbb{K}[x, y]$ is integral over $\mathbb{K}[x^2, xy, y^2]$,

$$\dim(\mathbb{K}[x, y]) = \dim(\mathbb{K}[x^2, xy, y^2]) = 2$$

(d) Since $\mathbb{Z}[i]$ is integral over $\mathbb{Z}$,

$$\dim(\mathbb{Z}[i]) = \dim(\mathbb{Z}) = 1$$

Problem 8. Closed Sets Under Integral Maps.

Let $R \subseteq S$ be integral and let $\phi^\# : \text{Spec}(S) \rightarrow \text{Spec}(R)$ be the induced map.

1. Let $J \subseteq S$ be an ideal. Show that

$$\phi^\#(\mathbb{V}(J)) = \mathbb{V}(J \cap R).$$

Hint: One inclusion is direct. For the other, apply Lying Over to $R/(J \cap R) \subseteq S/J$ after checking the kernel carefully.

2. Deduce that $\phi^\#$ is a closed map.
3. If $R \subseteq S$ is finite and κ is algebraically closed, explain why the induced map on affine algebraic sets should be thought of as a finite-to-one map. Make this precise by showing that for each $\mathfrak{p} \in \text{Spec}(R)$, the fibre ring $\kappa(\mathfrak{p}) \otimes_R S$ is finite-dimensional over $\kappa(\mathfrak{p})$.

Solution.

1. $\phi^\#(\mathbb{V}(J)) \subseteq \mathbb{V}(J \cap R)$

Let $\mathfrak{p} \in \phi^\#(\mathbb{V}(J))$. Then there exists some $\mathfrak{q} \in \mathbb{V}(J)$ such that $\phi^\#(\mathfrak{q}) = \mathfrak{q} \cap R = \mathfrak{p}$. Since $J \subseteq \mathfrak{q}$ we must have $J \cap R \subseteq \mathfrak{q} \cap R = \mathfrak{p}$.

$$\phi^\#(\mathbb{V}(J)) \supseteq \mathbb{V}(J \cap R)$$

Let $\mathfrak{p} \in \mathbb{V}(J \cap R)$. Then by going up and laying over we have,

$$\begin{array}{ccc} J & \subseteq & \mathfrak{q} \\ \downarrow & & \downarrow \\ J \cap R & \subseteq & \mathfrak{p} \end{array}$$

Thus there exists an $\mathfrak{q} \subset S$ such that $\phi^\#(\mathfrak{q}) = \mathfrak{p}$.

2. Thus, $\phi^\#$ maps closed sets to closed sets and hence is a closed map.
3. If $R \subseteq S$ is integral and finite we have $S = R[s_1, \dots, s_n]$. And the following diagram commutes,

$$\begin{array}{ccc} R & \longrightarrow & \kappa(\mathfrak{p}) \\ \downarrow & & \downarrow \\ S & \longrightarrow & \kappa(\mathfrak{p}) \otimes_R S \end{array}$$

Note that every element of $\kappa(\mathfrak{p}) \otimes_R S$ is a finite sum of simple tensors $k \otimes s$. Since $s \in S$, we can write $s = \sum_i r_i s_i$ for $r_i \in R$. Then,

$$\begin{aligned} k \otimes s &= k \otimes \sum_i r_i s_i \\ &= k \sum_i r_i \otimes \sum_i s_i \\ &= \sum_i k r_i \otimes s_i \\ &= \sum_i k r_i (1 \otimes s_i) \end{aligned}$$

Hence $\kappa(\mathfrak{p}) \otimes_R S$ is spanned by simple tensors $(1 \otimes s_1), \dots, (1 \otimes s_n)$ and thus is finite dimensional over $\kappa(\mathfrak{p})$.

We now turn to *Going Down*. For a ring map $R \rightarrow S$, Going Down means that chains can be lifted in the opposite direction: if $\mathfrak{p}_0 \subseteq \mathfrak{p}_1$ in R and a prime $\mathfrak{q}_1$ upstairs has already been chosen over $\mathfrak{p}_1$, then there should exist a prime $\mathfrak{q}_0 \subseteq \mathfrak{q}_1$ lying over $\mathfrak{p}_0$.

Going Down is not automatic for integral extensions. There are two especially important settings where it holds. The first is flatness, which is a module-theoretic condition. The second is the classical integral theorem: an integral extension of domains satisfies Going Down when the base is integrally closed. We will not prove these now, but will explore their consequences.

Theorem 0.20 (Flat Going Down). *Let $\phi : R \rightarrow S$ be a flat ring map. Suppose $\mathfrak{p}_0 \subseteq \mathfrak{p}_1$ are prime ideals of R and $\mathfrak{q}_1 \in \text{Spec}(S)$ satisfies $\phi^{-1}(\mathfrak{q}_1) = \mathfrak{p}_1$. Then there exists $\mathfrak{q}_0 \in \text{Spec}(S)$ such that*

$$\mathfrak{q}_0 \subseteq \mathfrak{q}_1 \quad \text{and} \quad \phi^{-1}(\mathfrak{q}_0) = \mathfrak{p}_0.$$

Theorem 0.21 (Going Down for Normal Domains). *Let $A \subseteq B$ be an integral extension of domains. If A is integrally closed in its fraction field, then $A \subseteq B$ satisfies Going Down.*

Problem 9. Unpacking Going Down.

Let $\phi : R \rightarrow S$ be a ring map.

1. Write the definition of Going Up for ϕ using a diagram of prime ideals.
2. Write the definition of Going Down for ϕ using a diagram of prime ideals.
3. Explain why Going Up is about lifting specializations and Going Down is about lifting generalizations.
4. Prove directly that a localization map $R \rightarrow U^{-1}R$ satisfies Going Down. Hint: Use the correspondence between primes of $U^{-1}R$ and primes of R disjoint from U .
5. Give an example of a flat map that is not integral and an integral map that is not flat.

Solution.

1. Going up says, given $\mathfrak{q}_0$ over $\mathfrak{p}_0$ and $\mathfrak{p}_0 \subseteq \mathfrak{p}_1$ we have

$$\begin{array}{ccc} \mathfrak{q}_0 & \overset{\subseteq}{\dashrightarrow} & \mathfrak{q}_1 \\ \cap R \downarrow & & \downarrow \cap R \\ \mathfrak{p}_0 & \xrightarrow{\subseteq} & \mathfrak{p}_1 \end{array}$$

2. Going down says, given $\mathfrak{q}_1$ over $\mathfrak{p}_1$ and $\mathfrak{p}_0 \subseteq \mathfrak{p}_1$ we have

$$\begin{array}{ccc} \mathfrak{q}_0 & \overset{\subseteq}{\dashleftarrow} & \mathfrak{q}_1 \\ \cap R \downarrow & & \downarrow \cap R \\ \mathfrak{p}_0 & \xrightarrow{\subseteq} & \mathfrak{p}_1 \end{array}$$

3. In $\text{Spec}(S)$ going up enables us to go from $\mathfrak{q}_0$ (over $\mathfrak{p}_0$) to $\mathfrak{q}_0 \subseteq \mathfrak{q}_1$, so $\mathfrak{q}_1 \in \mathbb{V}(\mathfrak{q}_0)$ and going down lets us find $\mathfrak{q}_0$ such that $\mathfrak{q}_1 \in \mathbb{V}(\mathfrak{q}_0)$.
4. In worksheet 6.1 we showed that a localized ring is a flat module and thus localization satisfies the flat going down theorem.
5. $\mathbb{K} \rightarrow \mathbb{K}[x]$ is flat but not integral and $\mathbb{Z} \rightarrow \mathbb{Z}/p\mathbb{Z}$ is integral but not flat.

Problem 10. Consequences of Going Down.

1. Let $R \rightarrow S$ be faithfully flat. Prove that $\text{Spec}(S) \rightarrow \text{Spec}(R)$ is surjective. Hint: For $\mathfrak{p} \in \text{Spec}(R)$, show that $\kappa(\mathfrak{p}) \otimes_R S \neq 0$.
2. Deduce that if $R \rightarrow S$ is faithfully flat, then every finite chain of primes in R can be lifted to a chain of primes in S . Conclude that $\dim(S) \geq \dim(R)$.
3. Apply this to $R \rightarrow R[x]$ to recover the inequality $\dim(R[x]) \geq \dim(R) + 1$ for nonzero R . Hint: if $\mathfrak{p}_0 \subsetneq \cdots \subsetneq \mathfrak{p}_n$ is a chain in R , use the explicit chain $\mathfrak{p}_0 R[x] \subsetneq \cdots \subsetneq \mathfrak{p}_n R[x] \subsetneq \mathfrak{p}_n R[x] + \langle x \rangle$.
4. Let $A \subseteq B$ be an integral extension of domains with A integrally closed. Let $\mathfrak{q} \in \text{Spec}(B)$ and $\mathfrak{p} = \mathfrak{q} \cap A$. Use Going Up, Going Down for Normal Domains, and Incomparability to prove

$$\text{ht}(\mathfrak{q}) = \text{ht}(\mathfrak{p}).$$

5. Specialize the previous part to a finite extension $\mathbb{K}[y_1, \dots, y_d] \subseteq B$ where B is a domain. What does it say about heights of primes in B ?

Solution.

1. Since $R \rightarrow S$ is faithfully flat we know that $\kappa(\mathfrak{p}) \otimes_R S = 0 \iff \kappa(\mathfrak{p}) = 0$, as $\kappa(\mathfrak{p}) \neq 0$ for any prime ideal we know $\kappa(\mathfrak{p}) \otimes_R S \neq 0$. That is $\kappa(\mathfrak{p}) \otimes_R S$ has at least one prime ideal, let this ideal be $\bar{\mathfrak{q}}$. Since pre-images of prime ideals are prime, $\bar{\mathfrak{q}}$ gives us a prime ideal $\mathfrak{q} \in S$.

$$\begin{array}{ccc} S & \longrightarrow & \kappa(\mathfrak{p}) \otimes_R S \\ \uparrow & & \uparrow \\ R & \longrightarrow & \kappa(\mathfrak{p}). \end{array}$$

Since the diagram commutes, the two maps $R \rightarrow S \rightarrow \kappa(\mathfrak{p}) \otimes_R S$ and $R \rightarrow \kappa(\mathfrak{p}) \rightarrow \kappa(\mathfrak{p}) \otimes_R S$ must agree. Pulling back $\bar{\mathfrak{q}}$ along the bottom-right path gives us $(0) \in \text{Spec}(\kappa(\mathfrak{p}))$, and (0) contracts to $\mathfrak{p} \in \text{Spec}(R)$. Thus $\mathfrak{q}$ lies over $\mathfrak{p}$. Hence every $\mathfrak{p} \in \text{Spec}(R)$ is in the image and thus the map $\text{Spec}(S) \rightarrow \text{Spec}(R)$ is surjective.

2. Since $\text{Spec}(S) \rightarrow \text{Spec}(R)$ is surjective, given a longest prime chain in R ,

$$\mathfrak{p}_0 \subsetneq \mathfrak{p}_1 \subsetneq \cdots \subsetneq \mathfrak{p}_n$$

we can find a lift of $\mathfrak{p}_n, \mathfrak{q}_n \in \text{Spec}(S)$. Then by going down, we can inductively find the corresponding prime chain of the same length in S ,

$$\mathfrak{q}_0 \subsetneq \mathfrak{q}_1 \subsetneq \cdots \subsetneq \mathfrak{q}_n$$

so $\dim(S) \geq \dim(R)$.

3. Let

$$\mathfrak{p}_0 \subsetneq \mathfrak{p}_1 \subsetneq \cdots \subsetneq \mathfrak{p}_n$$

be any chain in R and define the ideal lying over $\mathfrak{p}_n$ as $\mathfrak{q}_n = \mathfrak{p}_n R[x]$ since $\mathfrak{q} \cap R = \mathfrak{p}$ and

$$R[x] / \mathfrak{q}_n = R / \mathfrak{p}_n R[x] = (R / \mathfrak{p}_n)[x]$$

which is a domain. Then by part (2) we can inductively find a chain such that

$$\mathfrak{q}_0 \subsetneq \mathfrak{q}_1 \subsetneq \cdots \subsetneq \mathfrak{q}_n = \mathfrak{p}_n R[x]$$

Note that $\mathfrak{p}_n R[x] \subsetneq \mathfrak{p}_n R[x] + \langle x \rangle$ since $x \notin \mathfrak{p}_n R[x]$ because every element of $\mathfrak{p}_n R[x]$ has its coefficients in $\mathfrak{p}_n$ but x has leading coefficient 1.

So we have a chain of length $n + 1$ in S

$$\mathfrak{q}_0 \subsetneq \mathfrak{q}_1 \subsetneq \cdots \subsetneq \mathfrak{q}_n = \mathfrak{p}_n R[x] \subsetneq \mathfrak{p}_n R[x] + \langle x \rangle$$

so $\dim(S) \geq \dim(R) + 1$.

4. Going down for normal domains gives us that $\text{ht}(\mathfrak{q}) \geq \text{ht}(\mathfrak{p})$. To see that $\text{ht}(\mathfrak{q}) \leq \text{ht}(\mathfrak{p})$ we consider a longest chain that ends at $\mathfrak{q}$ in B . Restricting it to A gives us a chain that ends at $\mathfrak{p}$. Moreover we know all inclusions in the chain are strict by Incomparability. Hence, $\text{ht}(\mathfrak{q}) = \text{ht}(\mathfrak{p})$.
5. Since $\mathbb{K}[y_1, \dots, y_d]$ is a polynomial ring over a field it is a UFD and hence is integrally closed and the extension $\mathbb{K}[y_1, \dots, y_d] \subseteq B$ is integral. Moreover note that any prime in $\mathbb{K}[y_1, \dots, y_d]$ has height at most d . Then by part 4 for any prime ideal $\mathfrak{q} \in B$ we have,

$$\text{ht}(\mathfrak{q}) = \text{ht}(\mathfrak{q} \cap \mathbb{K}[y_1, \dots, y_d]) \leq d$$

We now move from arbitrary integral extensions to finitely generated algebras over a field. The next theorem is one of the main structural results in commutative algebra. It says that a finitely generated algebra over a field is finite over a polynomial subring.

Theorem 0.22 (Noether Normalization). *Let $\mathbb{K}$ be a field and let A be an algebra-finite $\mathbb{K}$ -algebra. Then there exist algebraically independent elements $y_1, \dots, y_d \in A$ such that A is module-finite over the polynomial subring*

$$\mathbb{K}[y_1, \dots, y_d] \subseteq A.$$

If A is a domain, then $d = \text{tr.deg}_{\mathbb{K}} \text{Frac}(A)$.

The proof is an induction on the number of algebra generators. The key step is a change of variables trick: given a nonzero polynomial relation, we choose new variables so that one old variable satisfies a monic equation over the others.

Problem 11. The Monic Change of Variables Trick.

Let $0 \neq f \in \mathbb{K}[x_1, \dots, x_n]$. The goal of this exercise is to make a change of variables so that f becomes monic in one variable. Write

$$f = \sum_{\alpha} c_{\alpha} x_1^{\alpha_1} \cdots x_n^{\alpha_n},$$

where $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{Z}_{\geq 0}^n$, only finitely many c_{α} are nonzero, and $c_{\alpha} \in \mathbb{K}$. Choose an integer N larger than every exponent appearing in every monomial of f ; that is, $N > \alpha_i$ for every exponent vector α appearing in f and every i .

1. Define

$$w(\alpha) = \alpha_n + \alpha_1 N + \alpha_2 N^2 + \cdots + \alpha_{n-1} N^{n-1}.$$

Explain why this is the base- N expansion of an integer whose “digits” are $\alpha_n, \alpha_1, \alpha_2, \dots, \alpha_{n-1}$

2. Since all these digits are strictly smaller than N , prove that if $w(\alpha) = w(\beta)$ then $\alpha = \beta$. Thus the weights $w(\alpha)$ are distinct for distinct monomials of f .
3. Introduce new variables

$$y_i = x_i - x_n^{N^i} \quad \text{for } 1 \leq i \leq n-1.$$

Equivalently, $x_i = y_i + x_n^{N^i}$. Substitute these expressions into a monomial

$$x_1^{\alpha_1} \cdots x_{n-1}^{\alpha_{n-1}} x_n^{\alpha_n}.$$

Show that the unique term obtained by choosing $x_n^{N^i}$ from every factor $(y_i + x_n^{N^i})^{\alpha_i}$ is $x_n^{w(\alpha)}$. All other terms contain at least one y_i and have strictly smaller degree in x_n .

4. Since the weights $w(\alpha)$ are distinct, there is a unique exponent vector $\alpha^{\max}$ appearing in f for which $w(\alpha)$ is largest. Show that, after the substitution, the highest power of x_n appearing in f is

$$c_{\alpha^{\max}} x_n^{w(\alpha^{\max})}.$$

5. Conclude that, after multiplying f by the nonzero scalar $c_{\alpha^{\max}}^{-1}$ we may regard f as a monic polynomial in x_n with coefficients in

$$\mathbb{K}[y_1, \dots, y_{n-1}].$$

6. Let $A = \mathbb{K}[x_1, \dots, x_n] / \langle f \rangle$. Prove that the image of x_n in A satisfies a monic polynomial with coefficients in the subalgebra generated by the images of

$$y_1, \dots, y_{n-1}.$$

Therefore the image of x_n is integral over this subalgebra.

7. Show A is generated as an algebra over $\mathbb{K}[y_1, \dots, y_{n-1}]$ by the single integral element x_n . Deduce that A is module-finite over $\mathbb{K}[y_1, \dots, y_{n-1}]$.

Solution.

1. Since $\alpha \in \mathbb{Z}$, and all $\alpha_i < N$ we can write a base- N expansion of the integer $(\alpha_n \alpha_1 \alpha_2 \cdots \alpha_{n-1})_N$ in this manner.
2. Assume $w(\alpha) = w(\beta)$. Then $(\alpha_n \alpha_1 \alpha_2 \cdots \alpha_{n-1})_N = (\beta_n \beta_1 \beta_2 \cdots \beta_{n-1})_N \implies \alpha = \beta$.
3. We have $\prod_i^n x_i^{\alpha_i} = x_n^{\alpha_n} \prod_i^{n-1} (y_i + x_n^{N^i})^{\alpha_i}$. Picking just $x_n^{N^i}$ from every factor gives us,

$$x_n^{\alpha_n} \prod_i^{n-1} (x_n^{N^i})^{\alpha_i} = x_n^{\alpha_n} \prod_i^{n-1} x_n^{\alpha_i N^i} = x_n^{\alpha_n + \alpha_1 N^1 + \cdots + \alpha_{n-1} N^{n-1}} = x_n^{w(\alpha)}$$

By binomial expansion we know that all other terms contain at least one y_i and hence have strictly smaller degree in x_n .

4. Thus after substitution, the leading term contributed by each monomial $c_\alpha x_1^{\alpha_1} \cdots x_n^{\alpha_n}$ in f is $c_\alpha x_n^{w(\alpha)}$. From part 2 we know that distinct α 's have distinct $w(\alpha)$'s, and thus no two monomials of f contribute leading terms of the same order. Therefore, there exists some $\alpha^{\max}$ whose weight $w(\alpha^{\max})$ is the largest, and thus the highest power of x_n in f is $c_{\alpha^{\max}} x_n^{w(\alpha^{\max})}$.
5. Since $c_\alpha \in \mathbb{K}$, we can multiply the term in the previous part by $c_{\alpha^{\max}}^{-1}$ to ensure the coefficient is one, thus making the polynomial f monic in x_n with coefficients in $\mathbb{K}[y_1, \dots, y_{n-1}]$.
6. $x_n \in \mathbb{K}[x_1, \dots, x_n]$ satisfies the monic polynomial

$$c_{\alpha^{\max}}^{-1} \cdot f = x_n^{w(\alpha^{\max})} + g_{d-1}(y_1, \dots, y_{n-1}) x_n^{w(\alpha^{\max})-1} + \cdots + g_0(y_1, \dots, y_{n-1}) = 0$$

with coefficients in $\mathbb{K}[y_1, \dots, y_{n-1}]$. In A we get,

$$\overline{x_n} = \overline{c_{\alpha^{\max}}^{-1} \cdot f} = \overline{x_n^{w(\alpha^{\max})}} + \overline{g_{d-1} x_n^{w(\alpha^{\max})-1}} + \cdots + \overline{g_0} = 0$$

where each $\overline{g_i}$ is the image of $g_i(y_1, \dots, y_{n-1})$ in A . Hence $\overline{x_n}$ is integral over the subalgebra generated by the image of $y_1, \dots, y_{n-1}$ in $\mathbb{K}[x_1, \dots, x_n] / \langle f \rangle$.

7. Note that there is an isomorphism $\mathbb{K}[y_1, \dots, y_{n-1}][x_n] \cong \mathbb{K}[x_1, \dots, x_n]$ given by the maps,

$$\phi : x_i \mapsto y_i + x_n^{N_i}, \quad x_n \mapsto x_n$$

$$\psi : y_i \mapsto x_i - x_n^{N_i}, \quad x_n \mapsto x_n$$

Now we show that the map $\pi : \mathbb{K}[y_1, \dots, y_{n-1}] \rightarrow A$ is injective. Assume for contradiction $p(y_1, \dots, y_{n-1}) \in \ker(\pi)$. That is, $p(y_1, \dots, y_{n-1}) \in \langle f \rangle$ or $p(y_1, \dots, y_{n-1}) = g \cdot f$ for some $g \in \mathbb{K}[x_1, \dots, x_n]$. But $x_n - \deg(p) = 0$ and $x_n - \deg(g \cdot f) \geq w(\alpha^{\max})$. But this is a contradiction. Hence π must be injective.

From the isomorphism, we know that every element of A can be represented by $h \in \mathbb{K}[x_1, \dots, x_n] = \mathbb{K}[y_1, \dots, y_{n-1}][x_n]$. Since f is monic we get,

$$h = f \cdot q + r, \quad x_n - \deg(r) \leq w(\alpha^{\max})$$

Since $f \equiv 0$ in A , we get $\bar{h} = \bar{r}$ where $x_n - \deg(r) \leq w(\alpha^{\max})$ with coefficients in $\mathbb{K}[y_1, \dots, y_{n-1}]$. Thus A is spanned by $1, x_n, \dots, x_n^{w(\alpha^{\max})-1}$ and hence is module-finite.

Problem 12. Proof of Noether Normalization.

Let $A = \mathbb{K}[x_1, \dots, x_n]/I$ be an algebra-finite $\mathbb{K}$ -algebra.

1. If $I = \langle 0 \rangle$, prove the theorem.
2. If $I \neq \langle 0 \rangle$, choose $0 \neq f \in I$. Use the previous exercise to make a change of variables so that A is module-finite over the image of a polynomial ring in $n - 1$ variables.
3. Let A_1 denote this image subalgebra. Explain why A_1 is algebra-finite over $\mathbb{K}$ using at most $n - 1$ algebra generators.
4. Apply induction to A_1 to find a polynomial subring $\mathbb{K}[y_1, \dots, y_d] \subseteq A_1$ such that A_1 is module-finite over $\mathbb{K}[y_1, \dots, y_d]$.
5. Prove that A is module-finite over $\mathbb{K}[y_1, \dots, y_d]$.
6. Explain why the elements $y_1, \dots, y_d$ are algebraically independent over $\mathbb{K}$.

Solution.

1. If $I = \langle 0 \rangle$, then the x_i are algebraically independent and $A = \mathbb{K}[x_1, \dots, x_n]$ is itself a polynomial ring and hence is trivially module-finite over itself.
2. Assume $I \neq \langle 0 \rangle$. Now let f be a non-zero polynomial in I . By Problem 11 we know that $\mathbb{K}[x_1, \dots, x_n]/\langle f \rangle$ is module-finite over $\mathbb{K}[y_1, \dots, y_{n-1}]$ for some algebraically independent $y_1, \dots, y_{n-1}$.
Since $f \in I$ we have $\langle f \rangle \subseteq I$ and we have the quotient map $\mathbb{K}[x_1, \dots, x_n]/\langle f \rangle \rightarrow A$, and hence A is module-finite over the image of $\mathbb{K}[y_1, \dots, y_{n-1}]$ in A .
3. Since A_1 is the image of a polynomial ring in $n - 1$ variables, it has at most $n - 1$ algebra-generators over $\mathbb{K}$.
4. By part 3, A_1 has at most $n - 1$ generators, so by the induction hypothesis there exists some $d \leq n - 1$ and algebraically independent elements $y_1, \dots, y_d$ such that A_1 is module finite over $\mathbb{K}[y_1, \dots, y_d]$.
5. Since A is module-finite over A_1 and A_1 is module-finite over $\mathbb{K}[y_1, \dots, y_d]$, A is module-finite over $\mathbb{K}[y_1, \dots, y_d]$ (Worksheet 7.1, Problem 7).
6. $\mathbb{K}[y_1, \dots, y_d]$ is a polynomial ring and thus the y_i are algebraically independent over $\mathbb{K}$.

Problem 13. Finding Normalizations by Hand.

For each algebra below, find a polynomial subring over which the algebra is module-finite. When possible, give explicit module generators.

- | | |
|--|--|
| 1. $\mathbb{K}[t^2, t^3]$. | 4. $\mathbb{K}[x, y, z] / \langle z^2 - xy \rangle$. |
| 2. $\mathbb{K}[x, y] / \langle y^2 - x^3 \rangle$. | 5. $\mathbb{K}[x^2, xy, y^2] \subseteq \mathbb{K}[x, y]$. |
| 3. $\mathbb{K}[x, y] / \langle xy \rangle$. Hint: Try $x + y$. | 6. $\mathbb{K}[s^2, s^3, t]$. |

Solution.

1. $\mathbb{K}[t^2, t^3]$ is module-finite over $\mathbb{K}[t^3 - t^2]$ with generators $\{1, t^2, t^4\}$.
2. $\mathbb{K}[x, y] / \langle y^2 - x^3 \rangle$ is module-finite over $\mathbb{K}[x]$ with generators $\{1, y\}$.
3. $\mathbb{K}[x, y] / \langle xy \rangle$ is module-finite over $\mathbb{K}[x + y]$ with generators $\{1, x, y\}$.
4. $\mathbb{K}[x, y, z] / \langle z^2 - xy \rangle$ is module-finite over $\mathbb{K}[x + y, z]$ with generators $\{1, x\}$.
5. $\mathbb{K}[x^2, xy, y^2] \subseteq \mathbb{K}[x, y]$ is module-finite over $\mathbb{K}[x^2, y^2]$ with generators $\{1, xy\}$.
6. $\mathbb{K}[s^2, s^3, t]$ is module-finite over $\mathbb{K}[s^3 - s^2, t]$ with generators $\{1, s^2, s^4\}$.

Problem 14. Zariski's Lemma from Noether Normalization.

Let A be an algebra-finite $\mathbb{K}$ -algebra that is also a field.

1. By Noether Normalization, choose $\mathbb{K}[y_1, \dots, y_d] \subseteq A$ such that A is module-finite over $\mathbb{K}[y_1, \dots, y_d]$. Explain why this extension is integral.
2. Use the earlier fact that if B is a field integral over a domain A_0 , then A_0 is a field, to prove that $\mathbb{K}[y_1, \dots, y_d]$ is a field.
3. Prove that $d = 0$.
4. Conclude that A is a finite-dimensional vector space over $\mathbb{K}$. Equivalently, every field that is algebra-finite over $\mathbb{K}$ is algebraic and finite over $\mathbb{K}$.
5. Deduce the weak Nullstellensatz: if $\mathbb{K}$ is algebraically closed and $\mathfrak{m} \subset \mathbb{K}[x_1, \dots, x_n]$ is maximal, then

$$\mathfrak{m} = \langle x_1 - a_1, \dots, x_n - a_n \rangle$$

for some $a_1, \dots, a_n \in \mathbb{K}$.

Solution.

1. Module-finiteness implies integrality hence A is integral over $\mathbb{K}[y_1, \dots, y_d]$.
2. We have an integral extension $\mathbb{K}[y_1, \dots, y_d] \rightarrow A$, and we know that $\mathbb{K}[y_1, \dots, y_d]$ is a polynomial ring over a field and hence a UFD. Then by Problem 1.2, since A is a field, $\mathbb{K}[y_1, \dots, y_d]$ is also a field.
3. Assume $d \neq 0$. Then $y_1 \in \mathbb{K}[y_1, \dots, y_d]$ doesn't have an inverse. A contradiction. Hence d must be 0.
4. Thus A is module-finite over the field $\mathbb{K}$, i.e. is a finite-dimensional vector space over $\mathbb{K}$, i.e. every element of A is algebraic over $\mathbb{K}$. So A is an algebra-finite over $\mathbb{K}$.
5. Let $\mathfrak{m} \subset \mathbb{K}[x_1, \dots, x_n]$ be maximal. Then $\mathbb{K}[x_1, \dots, x_n]/\mathfrak{m}$ is a field. By Noether Normalization $\mathbb{K}[x_1, \dots, x_n]/\mathfrak{m}$ is algebra-finite over $\mathbb{K}$ and hence a finite algebraic extension of $\mathbb{K}$. Since $\mathbb{K}$ is algebraically closed it has no finite proper extensions and hence $\mathbb{K}[x_1, \dots, x_n]/\mathfrak{m} \cong \mathbb{K}$. Then $x_i - a_i \in \mathfrak{m}$ and hence $\langle x_1 - a_1, \dots, x_n - a_n \rangle = \mathfrak{m}$.

We finish by extracting consequences for dimension. Noether Normalization reduces the dimension theory of affine algebras to the dimension theory of polynomial rings, while Lying Over, Going Up, and Incomparability tell us that finite integral extensions preserve lengths of prime chains.

Theorem 0.23 (Dimension of Affine Domains). *Let A be an algebra-finite $\mathbb{K}$ -algebra that is a domain. Then*

$$\dim(A) = \text{tr.deg}_{\mathbb{K}} \text{Frac}(A).$$

Equivalently, if $\mathbb{K}[y_1, \dots, y_d] \subseteq A$ is a Noether normalization, then $\dim(A) = d$.

Problem 15. Dimension of Affine Domains.

Let A be an algebra-finite $\mathbb{K}$ -algebra that is a domain.

1. Let $\mathbb{K}[y_1, \dots, y_d] \subseteq A$ be a Noether normalization. Use dimension preservation for integral extensions to prove $\dim(A) = d$.
2. Show that $\text{Frac}(A)$ is algebraic over $\mathbb{K}(y_1, \dots, y_d)$. Hint: First prove that every element of A is algebraic over $\mathbb{K}(y_1, \dots, y_d)$.
3. Deduce that $\text{tr.deg}_{\mathbb{K}} \text{Frac}(A) = d$.
4. Conversely, explain why d is determined by A and not by the choice of Noether normalization.

Solution.

1. A is integral over $\mathbb{K}[y_1, \dots, y_d]$. We know that $\dim(\mathbb{K}[y_1, \dots, y_d]) = d$ and by laying over and going up $\dim(A) \geq d$. Let

$$\mathfrak{q}_0 \subsetneq \mathfrak{q}_1 \subsetneq \cdots \subsetneq \mathfrak{q}_n$$

be a prime ideal chain in A . Then by Incomparability we get the chain

$$\mathfrak{q}_0 \cap \mathbb{K}[y_1, \dots, y_d] \subsetneq \mathfrak{q}_1 \cap \mathbb{K}[y_1, \dots, y_d] \subsetneq \cdots \subsetneq \mathfrak{q}_n \cap \mathbb{K}[y_1, \dots, y_d]$$

Hence every length in A corresponds to a chain of equal length in $\mathbb{K}[y_1, \dots, y_d]$ and $\dim(A) = d$.

2. Since A is integral over $\mathbb{K}[y_1, \dots, y_d]$ every element of A satisfies some monic polynomial with coefficients in $\mathbb{K}[y_1, \dots, y_d]$, i.e. every element in A is algebraic over $\mathbb{K}[y_1, \dots, y_d]$. Since $\mathbb{K}[y_1, \dots, y_d] \subseteq \mathbb{K}(y_1, \dots, y_d)$, every element of A is algebraic over it.

Now note that elements of $\text{Frac}(A)$ are of the form $\frac{a}{b}$ where $a, b \in A$. Since both a, b are algebraic over $\mathbb{K}(y_1, \dots, y_d)$ and algebraic extensions are closed under field operations $\frac{a}{b}$ is also algebraic over $\mathbb{K}(y_1, \dots, y_d)$.

3. We know that $\text{tr.deg}_{\mathbb{K}} \text{Frac}(A) \geq d$ since $y_1, \dots, y_d$ are algebraically independent over $\mathbb{K}$. By part 2, since $\text{Frac}(A)$ is algebraic over $\mathbb{K}(y_1, \dots, y_d)$, $\text{tr.deg}_{\mathbb{K}} \text{Frac}(A) \leq d$ since every element except $y_1, \dots, y_d$ are algebraic over $\mathbb{K}(y_1, \dots, y_d)$. Thus, $\text{tr.deg}_{\mathbb{K}} \text{Frac}(A) = d$.
4. Because $\text{tr.deg}_{\mathbb{K}} \text{Frac}(A) = d$, which depends only on A and $\mathbb{K}$, not on any choices made in the Noether normalization.

Problem 16. Computations Using Noether Normalization.

Compute the Krull dimension of each ring. In each case, try to exhibit a finite extension from a polynomial subring or reduce to irreducible components.

- | | |
|---|--|
| 1. $\mathbb{K}[t^2, t^3]$. | 5. $\mathbb{K}[x, y, z] / \langle xz, yz \rangle$. |
| 2. $\mathbb{K}[s^2, s^3, t]$. | 6. $\mathbb{K}[x_1, \dots, x_n] / \langle x_1, \dots, x_r \rangle$. |
| 3. $\mathbb{K}[x, y, z] / \langle z^2 - xy \rangle$. | 7. $\mathbb{Z}[x] / \langle x^2 + 1 \rangle$. |
| 4. $\mathbb{K}[x, y, z] / \langle xy, z \rangle$. | 8. $\mathbb{K}[x^2, xy, y^2]$. |

Solution.

1. Since $\mathbb{K}[t^2, t^3]$ is module-finite over $\mathbb{K}[t^3 - t^2]$ we have $\dim(\mathbb{K}[t^2, t^3]) = \dim(\mathbb{K}[t^3 - t^2]) = 1$.
2. Since $\mathbb{K}[s^2, s^3, t]$ is module-finite over $\mathbb{K}[s^3 - s^2, t]$ we have $\dim(\mathbb{K}[s^2, s^3, t]) = \dim(\mathbb{K}[s^3 - s^2, t]) = 2$.
3. Since $\mathbb{K}[x, y, z] / \langle z^2 - xy \rangle$ is module-finite over $\mathbb{K}[x + y, z]$ we have $\dim(\mathbb{K}[x, y, z] / \langle z^2 - xy \rangle) = \dim(\mathbb{K}[x + y, z]) = 2$.
4. Since $\mathbb{K}[x, y, z] / \langle xy, z \rangle$ is module-finite over $\mathbb{K}[x + y]$ we have $\dim(\mathbb{K}[x, y, z] / \langle xy, z \rangle) = \dim(\mathbb{K}[x + y]) = 1$.
5. $\mathbb{K}[x, y, z] / \langle xz, yz \rangle = \mathbb{K}[x, y, z] / \langle z \rangle \cap \langle x, y \rangle$. Then,

$$\mathbb{K}[x, y, z] / \langle z \rangle \cong \mathbb{K}[x, y] \implies \dim(\mathbb{K}[x, y, z] / \langle xz, yz \rangle) = 2$$

or

$$\mathbb{K}[x, y, z] / \langle x, y \rangle \cong \mathbb{K}[z] \implies \dim(\mathbb{K}[x, y, z] / \langle xz, yz \rangle) = 1$$

Since the Krull dimension is the sup, $\dim(\mathbb{K}[x, y, z] / \langle xz, yz \rangle) = 2$.

6. Since $\mathbb{K}[x_1, \dots, x_n] / \langle x_1, \dots, x_r \rangle$ is module-finite over $\mathbb{K}[x_{r+1}, \dots, x_n]$ we have $\dim(\mathbb{K}[x_1, \dots, x_n] / \langle x_1, \dots, x_r \rangle) = \dim(\mathbb{K}[x_{r+1}, \dots, x_n]) = n - r$.
7. Since $\mathbb{Z}[x] / \langle x^2 + 1 \rangle$ is module-finite over $\mathbb{Z}$ we have $\dim(\mathbb{Z}[x] / \langle x^2 + 1 \rangle) = \dim(\mathbb{Z}) = 1$.
8. Since $\mathbb{K}[x^2, xy, y^2]$ is module-finite over $\mathbb{K}[x^2, y^2]$ we have $\dim(\mathbb{K}[x^2, xy, y^2]) = \dim(\mathbb{K}[x^2, y^2]) = 2$.

Problem 17. Finite Maps to Affine Space.

Let A be an algebra-finite $\mathbb{K}$ -algebra and let $P = \mathbb{K}[y_1, \dots, y_d] \subseteq A$ be a Noether normalization.

1. Show that the induced map $\text{Spec}(A) \longrightarrow \text{Spec}(P) \cong \mathbb{A}_{\mathbb{K}}^d$ is surjective.
2. For $\mathfrak{p} \in \text{Spec}(P)$, prove that the fibre ring $\kappa(\mathfrak{p}) \otimes_P A$ is finite-dimensional over $\kappa(\mathfrak{p})$.
3. Assume $\mathbb{K}$ is algebraically closed and $\mathfrak{p} = \langle y_1 - a_1, \dots, y_d - a_d \rangle$ is a closed point of $\mathbb{A}_{\mathbb{K}}^d$. Show that the closed points in the fibre over $\mathfrak{p}$ correspond to maximal ideals of the finite-dimensional $\mathbb{K}$ -algebra $A/\mathfrak{p}A$.
4. Prove that a finite-dimensional algebra over a field has only finitely many maximal ideals. Hint: Distinct maximal ideals are comaximal, and the Chinese Remainder Theorem gives a surjection onto a product of fields.
5. Explain the geometric slogan: Noether normalization gives a finite-to-one projection from an affine algebraic set to affine space of the same dimension.

Solution.

1. By Laying over every ideal $\mathfrak{p} \in \text{Spec}(P)$ corresponds to an ideal $\mathfrak{q} \in \text{Spec}(A)$ such that $\mathfrak{q} \cap P = \mathfrak{p}$. Hence the map $\text{Spec}(A) \rightarrow \text{Spec}(P)$ is surjective.

2. We proved this in Problem 8.

3. Since $\mathbb{K}$ is algebraically closed, $\mathfrak{p}$ is maximal in P and $\kappa(\mathfrak{p}) = P/\mathfrak{p} \cong \mathbb{K}$. So the fibre ring is

$$\kappa(\mathfrak{p}) \otimes_P A = (P/\mathfrak{p}) \otimes_P A \cong A/\mathfrak{p}A$$

The closed points of the fiber are the maximal ideals of this fiber ring, which are exactly the maximal ideals of A containing $\mathfrak{p}A$ i.e. the maximal ideals of $A/\mathfrak{p}A$.

4. Let $\mathfrak{m}_1, \dots, \mathfrak{m}_r$ be distinct maximal ideals of $A/\mathfrak{p}A$. Since they are distinct maximal ideals, they are pairwise comaximal and by the CRT, there is a surjection

$$A/\mathfrak{p}A \twoheadrightarrow \prod_{i=1}^r A/\mathfrak{m}_i$$

Each $\mathfrak{m}_i$ is a maximal ideal of $A/\mathfrak{p}A$, which is a finite-dimensional $\mathbb{K}$ -algebra and hence algebra-finite over $\mathbb{K}$. By the Weak Nullstellensatz, $A/\mathfrak{m}_i \cong \mathbb{K}$ for each i . Thus we have a surjection

$$A/\mathfrak{p}A \twoheadrightarrow \prod_{i=1}^r \mathbb{K}$$

which gives

$$\dim_{\mathbb{K}}(A/\mathfrak{p}A) \geq \dim_{\mathbb{K}} \prod_{i=1}^r \mathbb{K} = r$$

Since $A/\mathfrak{p}A$ is finite-dimensional over $\mathbb{K}$ by part (2) and $r \leq \dim_{\mathbb{K}}(A/\mathfrak{p}A)$, $A/\mathfrak{p}A$ has only finitely many maximal ideals.

5. Noether normalization gives a surjection $\text{Spec}(A) \twoheadrightarrow A_{\mathbb{K}}^d$ where $d = \dim(A)$. Parts (3) and (4) say that each fiber over a closed point is finite and nonempty. So geometrically, every point of $A_{\mathbb{K}}^d$ has finitely many points of $\text{Spec}(A)$ above it and hence Noether normalization gives a finite-to-one projection from an affine algebraic set to affine space of the same dimension.