



Joint Research Centre

# A Parallelization Algorithm for Adequacy Assessment of the Electrical Grid

Gabor Riccardi

Università di Pavia

16/02/24



We formulate the Stochastic Capacity Expansion Problem as a two-stage stochastic program.

We formulate the Stochastic Capacity Expansion Problem as a two-stage stochastic program.

$$\min_{x} c'x + \mathbb{E}_{\omega} [\mathcal{V}(x, \omega)]$$

$$s.t. \ 0 \le x_{n,g} \le X_{n,g}$$
(CEP)

We formulate the Stochastic Capacity Expansion Problem as a two-stage stochastic program.

$$\min_{x} c'x + \mathbb{E}_{\omega} [\mathcal{V}(x, \omega)]$$

$$s.t. \ 0 \le x_{n,g} \le X_{n,g}$$
(CEP)

lacktriangle The first stage determines the capacity expansion  $x_{n,g}$  for each generator  $g\in\mathcal{G}$ 

We formulate the Stochastic Capacity Expansion Problem as a two-stage stochastic program.

$$\min_{x} c'x + \mathbb{E}_{\omega} [\mathcal{V}(x,\omega)]$$

$$s.t. \ 0 \le x_{n,g} \le X_{n,g}$$
(CEP)

- The first stage determines the capacity expansion  $x_{n,g}$  for each generator  $g \in \mathcal{G}$
- The second stage solves the Economic Dispatch (ED).

We formulate the Stochastic Capacity Expansion Problem as a two-stage stochastic program.

$$\min_{x} c'x + \mathbb{E}_{\omega} [\mathcal{V}(x,\omega)]$$

$$s.t. \ 0 \le x_{n,g} \le X_{n,g}$$
 (CEP)

- lacktriangle The first stage determines the capacity expansion  $x_{n,g}$  for each generator  $g\in\mathcal{G}$
- The second stage solves the Economic Dispatch (ED).

Where  $\mathcal{V}(x,\omega)$  is the solution to (ED) in function of the expanded capacities x and the scenario  $\omega$ .

Gabor Riccardi 16/02/24~~3~/~15

# Economic Dispatch (ED) model Scary Slide

For a fixed scenarios  $\omega = (\mathcal{PV}, \mathcal{W}, \mathcal{D})$  comprising of respectively solar power, wind power and loads.

For a fixed scenarios  $\omega = (\mathcal{PV}, \mathcal{W}, \mathcal{D})$  comprising of respectively solar power, wind power and loads, let  $y_{\omega} = (p_{\omega}, f_{\omega}, ls_{\omega}, s)'$  be the vector containing the power generation, power flows, line shedding and spillage variables.

For a fixed scenarios  $\omega = (\mathcal{PV}, \mathcal{W}, \mathcal{D})$  comprising of respectively solar power, wind power and loads, let  $y_{\omega} = (p_{\omega}, f_{\omega}, ls_{\omega}, s)'$  be the vector containing the power generation, power flows, line shedding and spillage variables.

$$\min_{y} q' y_{\omega} \tag{1}$$

$$s.t. p_{n,g,t,\omega} + bd_{n,t,\omega} + \sum_{l \in \mathcal{L}(n)} f_{n,l,t,\omega} + ls_{n,t,\omega} + \mathcal{PV}_{n,t,\omega} + \mathcal{W}_{n,t,\omega} =$$
(2)

$$=\mathcal{D}_{n,t,\omega}+s_{nt,\omega}+bc_{n,t,\omega}$$

$$v_{n,t,\omega} = v_{n,t-1,\omega} + BCE \cdot bc_{n,t,\omega} - BDE \cdot bd_{n,t,\omega} + A_{n,t,\omega}$$
(3)

$$(v_{n,t,\omega}, bc_{n,t,\omega}, bd_{n,t,\omega}) \le (BV, BC, BD) \tag{4}$$

$$p_{n,g,t,\omega} \le p_{n,g}^{\mathsf{max}} + x_{n,g} \tag{5}$$

$$L_{n,l}^{\min} \le f_{n,l,t,\omega} \le L_{n,l}^{\max} \tag{6}$$

Where  $v_{n,t,w}$  is the power stored at bus n at time t.

• We divide the time horizon into K intervals,  $\{t_0 = 0 := 1, \ldots, t_1\}, \{t_1 + 1, \ldots, t_2\}, \ldots, \{t_{K-1} + 1, \ldots, t_K := T\}$ 

- We divide the time horizon into K intervals,  $\{t_0 = 0 := 1, \ldots, t_1\}, \{t_1 + 1, \ldots, t_2\}, \ldots, \{t_{K-1} + 1, \ldots, t_K := T\}$
- We fix a priori the intermediate storage values  $v_{t_k}$  for  $k=1,\ldots,K$ .

- We divide the time horizon into K intervals,  $\{t_0 = 0 := 1, \ldots, t_1\}, \{t_1 + 1, \ldots, t_2\}, \ldots, \{t_{K-1} + 1, \ldots, t_K := T\}$
- We fix a priori the intermediate storage values  $v_{t_k}$  for k = 1, ..., K.
- We refer to the (ED) problems restricted to each time interval with fixed initial and final storage values as (ED-k) and to its optimal values as and with optimal value  $V_{\mathbf{k}}(\mathbf{x}, \mathbf{v_{t_k}}, \mathbf{v_{t_{k+1},\omega}})$

- We divide the time horizon into K intervals,  $\{t_0 = 0 \coloneqq 1, \dots, t_1\}, \{t_1 + 1, \dots, t_2\}, \dots, \{t_{K-1} + 1, \dots, t_K \coloneqq T\}$
- We fix a priori the intermediate storage values  $v_{t_k}$  for k = 1, ..., K.
- We refer to the (ED) problems restricted to each time interval with fixed initial and final storage values as (ED-k) and to its optimal values as and with optimal value  $V_{\mathbf{k}}(\mathbf{x}, \mathbf{v_{t_k}}, \mathbf{v_{t_{k+1},\omega}})$

#### Observation

$$\mathcal{V}(x,\omega) = \min_{\{v_{t_k}\}_{k=1}^K} \sum_{k=0}^{K-1} \mathcal{V}_k(x, v_{t_k}, v_{t_k+1}, \omega)$$
 (7)

Since each function  $\mathcal{V}_k$  is piecewise linear convex in  $x, v_{t_K}, v_{t_{K+1}}$ , it can be approximated by a collection of supporting hyperplanes  $\{\pi_{i,k}^w(x, v_{t_k}, v_{t_{k+1}})\}$  of each  $\mathcal{V}_k$ .

Since each function  $\mathcal{V}_k$  is piecewise linear convex in  $x, v_{t_K}, v_{t_{K+1}}$ , it can be approximated by a collection of supporting hyperplanes  $\{\pi_{i,k}^w(x, v_{t_k}, v_{t_{k+1}})\}$  of each  $\mathcal{V}_k$ . An approximation of (ED) is given by:

Since each function  $\mathcal{V}_k$  is piecewise linear convex in  $x, v_{t_K}, v_{t_{K+1}}$ , it can be approximated by a collection of supporting hyperplanes  $\{\pi_{i,k}^w(x, v_{t_k}, v_{t_{k+1}})\}$  of each  $\mathcal{V}_k$ . An approximation of (ED) is given by:

$$\hat{\mathcal{V}}(x,\omega) = \min_{\{v_{t_k}\}_{k=1}^K} \sum_{k=0}^K \hat{\mathcal{V}}_k(x, v_{t_k}, v_{t_{k+1}}) =$$

$$= \min_{\{v_{t_k}\}_{k=1}^K} \sum_{k=0}^K \theta_k^{\omega}$$

$$\text{s.t.} \quad \theta_k^{\omega} \ge \pi_{i,k}^{\omega}(x, v_{t_k}, v_{t_{k+1}}) \quad \forall i, k$$
(ISP)

We refer to this problem as the Intermediate Storage Problem (ISP)

Since each function  $\mathcal{V}_k$  is piecewise linear convex in  $x, v_{t_K}, v_{t_{K+1}}$ , it can be approximated by a collection of supporting hyperplanes  $\{\pi_{i,k}^w(x,v_{t_k},v_{t_{k+1}})\}$  of each  $\mathcal{V}_k$ . An approximation of (ED) is given by:

$$\begin{split} \hat{\mathcal{V}}(x,\omega) &= \min_{\left\{v_{t_k}\right\}_{k=1}^K} \sum_{k=0}^K \hat{\mathcal{V}}_k(x,v_{t_k},v_{t_{k+1}}) = \\ &= \min_{\left\{v_{t_k}\right\}_{k=1}^K} \sum_{k=0}^K \theta_k^{\omega} \\ \text{s.t.} \quad \theta_k^{\omega} \geq \pi_{i,k}^{\omega}(x,v_{t_k},v_{t_{k+1}}) \quad \forall i,k \end{split} \tag{ISP}$$

We refer to this problem as the **Intermediate Storage Problem (ISP)** (I know, very original)

# Model description: Relaxed Capacity Expansion(CEP-R)

$$\min_{x} c'x + \mathbb{E}_{\omega} [\mathcal{V}(x, \omega)]$$

$$s.t. \ 0 \le x_{n,g} \le X_{n,g}$$
(CEP)

# Model description: Relaxed Capacity Expansion(CEP-R)

$$\min_{x} c'x + \mathbb{E}_{\omega} [\mathcal{V}(x,\omega)]$$
 $s.t. \ 0 \le x_{n,g} \le X_{n,g}$  (CEP)

$$\min_{x} c'x + \mathbb{E}_{w} \left[ \hat{\mathcal{V}}(x, w) \right]$$
 $s.t. \ 0 \le x_{n,g} \le X_{n,g}$  (CEP-R)


# Model description: Relaxed Capacity Expansion(CEP-R)

$$\min_{x} c'x + \mathbb{E}_{\omega} \left[ \mathcal{V}(x, \omega) \right] \\ s.t. \ 0 \leq x_{n,g} \leq X_{n,g}$$
 (CEP)

$$\min_{x} c'x + \mathbb{E}_{w} \left[ \hat{\mathcal{V}}(x, w) \right]$$
 $s.t. \ 0 \le x_{n,g} \le X_{n,g}$  (CEP-R)

Since calculating  $\hat{\mathcal{V}}$  is straightforward, solving (CEP-R) can be done efficiently with L-shaped or subgradient schemes.

### Algorithm



■ Since  $(CEP - R) \le (CEP)$  if a (CEP - R) optimal solution has the same cost for (CEP) then it's also (CEP)-optimal.

■ Since  $(CEP - R) \le (CEP)$  if a (CEP - R) optimal solution has the same cost for (CEP) then it's also (CEP)-optimal.

**Remark 1:** It is sufficient to prove that after a finite number of steps (i) of the algorithm we have:

$$\hat{\mathcal{V}}(\hat{x}^i, \omega) = \mathcal{V}(\hat{x}^i, \omega) \text{ for all } \omega \in \Omega$$
 (8)

Observation

After a finite number of iterations no new cuts are found for  $V_k$ .

Proof.

(9)

Observation

After a finite number of iterations no new cuts are found for  $V_k$ .

Proof.

 $\#\{p\mid p \text{ is a normal vector of a supporting hyperplane of } \mathcal{V}_k\} \leq$ 

(9)

#### Observation

After a finite number of iterations no new cuts are found for  $V_k$ .

Proof.

```
\#\{p\mid p \text{ is a normal vector of a supporting hyperplane of } \mathcal{V}_k\} \leq \ \#\{\text{dual solutions } p=q'B^{-1} \text{ of (ED-k) for varying } x,v_{t_k},v_{t_{k+1}}\} \leq \ \#\{\text{dual solutions } p=q'B^{-1} \text{ of (ED-k) for varying } x,v_{t_k},v_{t_{k+1}}\} \leq \ \#\{\text{dual solutions } p=q'B^{-1} \text{ of (ED-k) for varying } x,v_{t_k},v_{t_{k+1}}\} \leq \ \#\{\text{dual solutions } p=q'B^{-1} \text{ of (ED-k) for varying } x,v_{t_k},v_{t_{k+1}}\} \leq \ \#\{\text{dual solutions } p=q'B^{-1} \text{ of (ED-k) for varying } x,v_{t_k},v_{t_{k+1}}\} \leq \ \#\{\text{dual solutions } p=q'B^{-1} \text{ of (ED-k) for varying } x,v_{t_k},v_{t_{k+1}}\} \leq \ \#\{\text{dual solutions } p=q'B^{-1} \text{ of (ED-k) for varying } x,v_{t_k},v_{t_k}\} \leq \ \#\{\text{dual solutions } p=q'B^{-1} \text{ of (ED-k) for varying } x,v_{t_k},v_{t_k}\} \leq \ \#\{\text{dual solutions } p=q'B^{-1} \text{ of (ED-k) for varying } x,v_{t_k},v_{t_k}\} \leq \ \#\{\text{dual solutions } p=q'B^{-1} \text{ of (ED-k) for varying } x,v_{t_k},v_{t_k}\} \leq \ \#\{\text{dual solutions } p=q'B^{-1} \text{ of (ED-k) for varying } x,v_{t_k}\} \leq \ \#\{\text{dual solutions } p=q'B^{-1} \text{ of (ED-k) for varying } x,v_{t_k}\} \leq \ \#\{\text{dual solutions } p=q'B^{-1} \text{ of (ED-k) for varying } x,v_{t_k}\} \leq \ \#\{\text{dual solutions } p=q'B^{-1} \text{ of (ED-k) for varying } x,v_{t_k}\} \leq \ \#\{\text{dual solutions } p=q'B^{-1} \text{ of (ED-k) for varying } x,v_{t_k}\} \leq \ \#\{\text{dual solutions } p=q'B^{-1} \text{ of (ED-k) for varying } x,v_{t_k}\} \leq \ \#\{\text{dual solutions } p=q'B^{-1} \text{ of (ED-k) for varying } x,v_{t_k}\} \leq \ \#\{\text{dual solutions } p=q'B^{-1} \text{ of (ED-k) for varying } x,v_{t_k}\} \leq \ \#\{\text{dual solutions } p=q'B^{-1} \text{ of (ED-k) for varying } x,v_{t_k}\} \leq \ \#\{\text{dual solutions } x,v_{t_k
```

Gabor Riccardi 16/02/24 9 / 15

(9)

#### Observation

After a finite number of iterations no new cuts are found for  $V_k$ .

Proof.

```
\#\{p \mid p \text{ is a normal vector of a supporting hyperplane of } \mathcal{V}_k\} \le \\ \#\{\text{dual solutions } p = q'B^{-1} \text{ of (ED-k) for varying } x, v_{t_k}, v_{t_{k+1}}\} \le \\ \#\{\text{basis matrices of (ED-k)}\} < \infty  (9)
```

#### Observation

After a finite number of iterations no new cuts are found for  $V_k$ .

Proof.

$$\#\{p \mid p \text{ is a normal vector of a supporting hyperplane of } \mathcal{V}_k\} \le \\ \#\{\text{dual solutions } p = q'B^{-1} \text{ of (ED-k) for varying } x, v_{t_k}, v_{t_{k+1}}\} \le \\ \#\{\text{basis matrices of (ED-k)}\} < \infty$$
 (9)

After a finite number of steps:

#### Observation

After a finite number of iterations no new cuts are found for  $V_k$ .

Proof.

$$\#\{p \mid p \text{ is a normal vector of a supporting hyperplane of } \mathcal{V}_k\} \le \\ \#\{\text{dual solutions } p = q'B^{-1} \text{ of (ED-k) for varying } x, v_{t_k}, v_{t_{k+1}}\} \le \\ \#\{\text{basis matrices of (ED-k)}\} < \infty$$
 (9)

After a finite number of steps:

• new cut:  $\bar{c}(x, v) = p'(x, v) + b$ 

#### Observation


After a finite number of iterations no new cuts are found for  $V_k$ .


Proof.

$$\#\{p \mid p \text{ is a normal vector of a supporting hyperplane of } \mathcal{V}_k\} \le \\ \#\{\text{dual solutions } p = q'B^{-1} \text{ of (ED-k) for varying } x, v_{t_k}, v_{t_{k+1}}\} \le \\ \#\{\text{basis matrices of (ED-k)}\} < \infty$$
 (9)

After a finite number of steps:

- new cut:  $\bar{c}(x, v) = p'(x, v) + b$
- an old cut:  $\pi(x,v) = p'(x,v) + \bar{b}$





Since both are supporting hyperplanes it follows that  $b=\bar{b}$  (and therefore  $\bar{c}$  is not a new cut).

#### Observation

If after the i-iteration no new cuts are added for some i and k then  $\hat{\mathcal{V}}_k(\hat{x}^i,\hat{v}_k,\hat{v}_{k+1}) = \mathcal{V}_k(\hat{x}^i,\hat{v}_k,\hat{v}_{k+1})$ .

#### Observation

If after the i-iteration no new cuts are added for some i and k then  $\hat{\mathcal{V}}_k(\hat{x}^i,\hat{v}_k,\hat{v}_{k+1}) = \mathcal{V}_k(\hat{x}^i,\hat{v}_k,\hat{v}_{k+1}).$  Where  $\hat{\mathcal{V}}_k(\hat{x}^i,\hat{v}_{t_k},\hat{v}_{t_{k+1}},\omega) = \max_i \pi_{i,k}^{\omega}(\hat{x}^i,\hat{v}_{t_k},\hat{v}_{t_{k+1}}) \text{ be the current supporting hyperplane approximation of } \mathcal{V}_k.$ 

#### Observation

If after the i-iteration no new cuts are added for some i and k then  $\hat{\mathcal{V}}_k(\hat{x}^i,\hat{v}_k,\hat{v}_{k+1}) = \mathcal{V}_k(\hat{x}^i,\hat{v}_k,\hat{v}_{k+1}).$  Where  $\hat{\mathcal{V}}_k(\hat{x}^i,\hat{v}_{t_k},\hat{v}_{t_{k+1}},\omega) = \max_i \pi_{i,k}^{\omega}(\hat{x}^i,\hat{v}_{t_k},\hat{v}_{t_{k+1}}) \text{ be the current supporting hyperplane approximation of } \mathcal{V}_k.$ 

#### Proof.

Let  $\bar{c}_k^{\omega}(x, v_{t_k}) := p'(x - \hat{x}^i, v_{t_k} - \hat{v}_{t_k}) + V_k(\hat{x}^i, \hat{v}_{t_k})$  be the new cut found after the *i*-th iteration.

#### Observation

If after the i-iteration no new cuts are added for some i and k then  $\hat{\mathcal{V}}_k(\hat{x}^i,\hat{v}_k,\hat{v}_{k+1}) = \mathcal{V}_k(\hat{x}^i,\hat{v}_k,\hat{v}_{k+1}).$  Where  $\hat{\mathcal{V}}_k(\hat{x}^i,\hat{v}_{t_k},\hat{v}_{t_{k+1}},\omega) = \max_i \pi_{i,k}^{\omega}(\hat{x}^i,\hat{v}_{t_k},\hat{v}_{t_{k+1}}) \text{ be the current supporting hyperplane approximation of } \mathcal{V}_k.$ 

#### Proof.

Let  $\bar{c}_k^{\omega}(x, v_{t_k}) := p'(x - \hat{x}^i, v_{t_k} - \hat{v}_{t_k}) + V_k(\hat{x}^i, \hat{v}_{t_k})$  be the new cut found after the *i*-th iteration.

Since  $\bar{c}$  is not a new cut we have  $\bar{c}(\hat{x}^i, v_{t_k}) \leq \hat{\mathcal{V}}_k(\hat{x}^i, v_{t_k})$ .

#### Observation

If after the i-iteration no new cuts are added for some i and k then  $\hat{\mathcal{V}}_k(\hat{x}^i,\hat{v}_k,\hat{v}_{k+1}) = \mathcal{V}_k(\hat{x}^i,\hat{v}_k,\hat{v}_{k+1}).$  Where  $\hat{\mathcal{V}}_k(\hat{x}^i,\hat{v}_{t_k},\hat{v}_{t_{k+1}},\omega) = \max_i \pi_{i,k}^{\omega}(\hat{x}^i,\hat{v}_{t_k},\hat{v}_{t_{k+1}}) \text{ be the current supporting hyperplane approximation of } \mathcal{V}_k.$ 

#### Proof.

Let  $\bar{c}_k^{\omega}(x, v_{t_k}) := p'(x - \hat{x}^i, v_{t_k} - \hat{v}_{t_k}) + V_k(\hat{x}^i, \hat{v}_{t_k})$  be the new cut found after the *i*-th iteration.

Since  $\bar{c}$  is not a new cut we have  $\bar{c}(\hat{x}^i, v_{t_k}) \leq \hat{\mathcal{V}}_k(\hat{x}^i, v_{t_k})$ . We have thus

$$\mathcal{V}_k(\hat{x}^i, \hat{v}_{t_k}) \geq \hat{\mathcal{V}}_k(\hat{x}^i, \hat{v}_{t_k}) \geq \bar{c}(\hat{x}^i, \hat{v}_{t_k}) = \mathcal{V}_k(\hat{x}^i, \hat{v}_{t_k})$$

#### Observation

If after the i-iteration no new cuts are added for some i and k then  $\hat{\mathcal{V}}_k(\hat{x}^i,\hat{v}_k,\hat{v}_{k+1}) = \mathcal{V}_k(\hat{x}^i,\hat{v}_k,\hat{v}_{k+1}).$  Where  $\hat{\mathcal{V}}_k(\hat{x}^i,\hat{v}_{t_k},\hat{v}_{t_{k+1}},\omega) = \max_i \pi_{i,k}^{\omega}(\hat{x}^i,\hat{v}_{t_k},\hat{v}_{t_{k+1}}) \text{ be the current supporting hyperplane approximation of } \mathcal{V}_k.$ 

#### Proof.

Let  $\bar{c}_k^{\omega}(x, v_{t_k}) := p'(x - \hat{x}^i, v_{t_k} - \hat{v}_{t_k}) + V_k(\hat{x}^i, \hat{v}_{t_k})$  be the new cut found after the *i*-th iteration.

Since  $\bar{c}$  is not a new cut we have  $\bar{c}(\hat{x}^i, v_{t_k}) \leq \hat{\mathcal{V}}_k(\hat{x}^i, v_{t_k})$ . We have thus

$$\mathcal{V}_k(\hat{x}^i, \hat{v}_{t_k}) \geq \hat{\mathcal{V}}_k(\hat{x}^i, \hat{v}_{t_k}) \geq \bar{c}(\hat{x}^i, \hat{v}_{t_k}) = \mathcal{V}_k(\hat{x}^i, \hat{v}_{t_k})$$

which concludes the proof.

In conclusion, we have  $\hat{\mathcal{V}}_k(\hat{x}^i, v_{t_k}, v_{t_{k+1}}, \omega) = \mathcal{V}_k(\hat{x}^i, v_{t_k}, v_{t_{k+1}}, \omega)$  for all  $\omega, k$ .

In conclusion, we have  $\hat{\mathcal{V}}_k(\hat{x}^i, v_{t_k}, v_{t_{k+1}}, \omega) = \mathcal{V}_k(\hat{x}^i, v_{t_k}, v_{t_{k+1}}, \omega)$  for all  $\omega, k$ . Thus  $\hat{\mathcal{V}}(\hat{x}^i, \omega) = \mathcal{V}(\hat{x}^i, \omega)$ .

In conclusion, we have 
$$\hat{\mathcal{V}}_k(\hat{x}^i, v_{t_k}, v_{t_{k+1}}, \omega) = \mathcal{V}_k(\hat{x}^i, v_{t_k}, v_{t_{k+1}}, \omega)$$
 for all  $\omega, k$ . Thus  $\hat{\mathcal{V}}(\hat{x}^i, \omega) = \mathcal{V}(\hat{x}^i, \omega)$ .

#### Proposition

The algorithm converges after a finite number of iterations and  $\hat{x}^i$  is an optimal solution for (CEP).

# Future implementation in Pypsa

 We are currently implementing this and other stochastic methods within the Pypsa [BHS18] framework using the Linopy [Hof23] modeling package in Python.

# Future implementation in Pypsa

- We are currently implementing this and other stochastic methods within the Pypsa [BHS18] framework using the Linopy [Hof23] modeling package in Python.
- We expect improved convergence speed respect to the L-shaped method, especially when leveraging parallel processing capabilities.

 The specific structure of the intertemporal constraints makes it possible do develop tailored optimization algorithms for (CEP).

- The specific structure of the intertemporal constraints makes it possible do develop tailored optimization algorithms for (CEP).
- The presented method can be also applied to other linear models such as DC-OPF.

- The specific structure of the intertemporal constraints makes it possible do develop tailored optimization algorithms for (CEP).
- The presented method can be also applied to other linear models such as DC-OPF.
- Care must be taken for intra-temporal constraints such as ramp up or ramp down constraints.

- The specific structure of the intertemporal constraints makes it possible do develop tailored optimization algorithms for (CEP).
- The presented method can be also applied to other linear models such as DC-OPF.
- Care must be taken for intra-temporal constraints such as ramp up or ramp down constraints.
- Develop heuristics for the choice of the number of time intervals K.

- The specific structure of the intertemporal constraints makes it possible do develop tailored optimization algorithms for (CEP).
- The presented method can be also applied to other linear models such as DC-OPF.
- Care must be taken for intra-temporal constraints such as ramp up or ramp down constraints.
- Develop heuristics for the choice of the number of time intervals K.
- Examine generalizations to multistage stochastic programs.

- The specific structure of the intertemporal constraints makes it possible do develop tailored optimization algorithms for (CEP).
- The presented method can be also applied to other linear models such as DC-OPF.
- Care must be taken for intra-temporal constraints such as ramp up or ramp down constraints.
- Develop heuristics for the choice of the number of time intervals K.
- Examine generalizations to multistage stochastic programs.

Thank you for your attention.

gabor.riccardi01@universitadipavia.it
https://www.compopt.it

#### Some references:

- [Ávi+23] Daniel Ávila, Anthony Papavasiliou, Mauricio Junca, and Lazaros Exizidis. "Applying High-Performance Computing to the European Resource Adequacy Assessment". In: *IEEE Transactions on Power Systems* (2023), pp. 1–13. DOI: 10.1109/TPWRS.2023.3304717.
- [Bie+20] Daniel Bienstock, Mauro Escobar, Claudio Gentile, and Leo Liberti. "Mathematical Programming formulations for the Alternating Current Optimal Power Flow problem". In: 4OR 18.3 (July 2020), pp. 249–292. DOI: 10.1007/s10288-020-00455-w.
- [BM14] Daniel Bienstock and Gonzalo Munoz. "On linear relaxations of OPF problems". In: (Nov. 2014).
- [BHS18] T. Brown, J. Hörsch, and D. Schlachtberger. "PyPSA: Python for Power System Analysis". In: Journal of Open Research Software 6.4 (1 2018). DOI: 10.5334/jors.188. eprint: 1707.09913. URL: https://doi.org/10.5334/jors.188.
- [Hof23] Fabian Hofmann. "Linopy: Linear optimization with n-dimensional labeled variables". In: Journal of Open Source Software 8.84 (2023), p. 4823. DOI: 10.21105/joss.04823. URL: https://doi.org/10.21105/joss.04823.

# Adequacy Assessment of the Electrical Grid

- Measuring the ability of the electric power system to react to adverse uncertain condition has become increasingly importan.
- Member States wishing to introduce capacity mechanisms can do so if an adequacy concern is identified in the ERAA study, a pan-European adequacy assessment for up to 10 years ahead.
- Due to the scale of the ERAA study, ERAA 2022 considered a reduced stochastic problem with three scenarios.
- In [Ávi+23], Daniel A'vila introduced a decomposition algorithm based on subgradient approximations was introduced

Optimal Power Flow (OPF)

[Bie+20]

- AC OPF: exact physical model
- Security-Constrained OPF (SCOPF) Includes contingencies to guarantee system security under failures.
- DC OPF and other linearized models

[BM14]

- other relaxations.
- Unit Commitment Determines on/off status of power units, ignoring grid constraints.
- Economic Dispatch (ED) Minimizes generation cost, ignoring grid constraints.

**Capacity expansion problem:** Based on Economic Dispatch models with added flow balance at bus nodes and various scenarios.