

A Parallelization Algorithm for Adequacy Assessment of the Electrical Grid




Gabor Riccardi

Università di Pavia

16/02/24



Joint Research Centre



8th AIROYoung WORKSHOP

February 14-16, 2024

UNIVERSITY OF CALABRIA

*"Emerging technologies for decision support
systems and innovative optimization paradigms"*

Stochastic Capacity Expansion Problem (CEP)

We formulate the Stochastic Capacity Expansion Problem as a two-stage stochastic program.

Stochastic Capacity Expansion Problem (CEP)

We formulate the Stochastic Capacity Expansion Problem as a two-stage stochastic program.

$$\begin{aligned} \min_x \quad & c'x + \mathbb{E}_\omega [\mathcal{V}(x, \omega)] \\ \text{s.t.} \quad & 0 \leq x_{n,g} \leq X_{n,g} \end{aligned} \tag{CEP}$$

Stochastic Capacity Expansion Problem (CEP)

We formulate the Stochastic Capacity Expansion Problem as a two-stage stochastic program.

$$\begin{aligned} \min_x \quad & c'x + \mathbb{E}_\omega [\mathcal{V}(x, \omega)] \\ \text{s.t.} \quad & 0 \leq x_{n,g} \leq X_{n,g} \end{aligned} \tag{CEP}$$

- The first stage determines the capacity expansion $x_{n,g}$ for each generator $g \in \mathcal{G}$

Stochastic Capacity Expansion Problem (CEP)

We formulate the Stochastic Capacity Expansion Problem as a two-stage stochastic program.

$$\begin{aligned} \min_x \quad & c'x + \mathbb{E}_\omega [\mathcal{V}(x, \omega)] \\ \text{s.t.} \quad & 0 \leq x_{n,g} \leq X_{n,g} \end{aligned} \tag{CEP}$$

- The first stage determines the capacity expansion $x_{n,g}$ for each generator $g \in \mathcal{G}$
- The second stage solves the Economic Dispatch (ED).

Stochastic Capacity Expansion Problem (CEP)

We formulate the Stochastic Capacity Expansion Problem as a two-stage stochastic program.

$$\begin{aligned} \min_x \quad & c'x + \mathbb{E}_\omega [\mathcal{V}(x, \omega)] \\ \text{s.t.} \quad & 0 \leq x_{n,g} \leq X_{n,g} \end{aligned} \tag{CEP}$$

- The first stage determines the capacity expansion $x_{n,g}$ for each generator $g \in \mathcal{G}$
- The second stage solves the Economic Dispatch (ED).

Where $\mathcal{V}(x, \omega)$ is the solution to (ED) in function of the expanded capacities x and the scenario ω .

Economic Dispatch (ED) model

Economic Dispatch (ED) model Scary Slide

Economic Dispatch (ED) model

For a fixed scenarios $\omega = (\mathcal{PV}, \mathcal{W}, \mathcal{D})$ comprising of respectively solar power, wind power and loads,

Economic Dispatch (ED) model

For a fixed scenarios $\omega = (\mathcal{PV}, \mathcal{W}, \mathcal{D})$ comprising of respectively solar power, wind power and loads, let $y_\omega = (p_\omega, f_\omega, ls_\omega)'$ be the vector containing the power generation, power flows and line shedding variables.

Economic Dispatch (ED) model

For a fixed scenarios $\omega = (\mathcal{PV}, \mathcal{W}, \mathcal{D})$ comprising of respectively solar power, wind power and loads, let $y_\omega = (p_\omega, f_\omega, ls_\omega)'$ be the vector containing the power generation, power flows and line shedding variables.

$$\min_y q' y_\omega \quad (1)$$

$$s.t. \quad p_{n,g,t,\omega} + bd_{n,t,\omega} + \sum_{l \in \mathcal{L}(n)} f_{n,l,t,\omega} + ls_{n,t,\omega} + \mathcal{PV}_{n,t,\omega} + \mathcal{W}_{n,t,\omega} = \quad (2)$$

$$= \mathcal{D}_{n,t,\omega} + ps_{nt,\omega} + bc_{n,t,\omega}$$

$$v_{n,t,\omega} = v_{n,t-1,\omega} + BCE \cdot bc_{n,t,\omega} - BDE \cdot bd_{n,t,\omega} + A_{n,t,\omega} \quad (3)$$

$$(v_{n,t,\omega}, bc_{n,t,\omega}, bd_{n,t,\omega}) \leq (BV, BC, BD) \quad (4)$$

$$p_{n,g,t,\omega} \leq p_{n,g}^{\max} + x_{n,g} \quad (5)$$

$$L_{n,l}^{\min} \leq f_{n,l,t,\omega} \leq L_{n,l}^{\max} \quad (6)$$

Where $v_{n,t,\omega}$ is the power stored at bus n at time t .

Model relaxation description: Intermediate Economic Dispatches (ED-k)

- We divide the time horizon into K intervals, $\{t_0 = 0 := 1, \dots, t_1\}, \{t_1 + 1, \dots, t_2\}, \dots, \{t_{K-1} + 1, \dots, t_K := T\}$

Model relaxation description: Intermediate Economic Dispatches (ED-k)

- We divide the time horizon into K intervals, $\{t_0 = 0 := 1, \dots, t_1\}, \{t_1 + 1, \dots, t_2\}, \dots, \{t_{K-1} + 1, \dots, t_K := T\}$
- We fix a priori the intermediate storage values v_{t_k} for $k = 1, \dots, K$.

Model relaxation description: Intermediate Economic Dispatches (ED-k)

- We divide the time horizon into K intervals, $\{t_0 = 0 := 1, \dots, t_1\}, \{t_1 + 1, \dots, t_2\}, \dots, \{t_{K-1} + 1, \dots, t_K := T\}$
- We fix a priori the intermediate storage values v_{t_k} for $k = 1, \dots, K$.
- We refer to the (ED) problems restricted to each time interval with fixed initial and final storage values as **(ED-k)** and to its optimal values as and with optimal value $\mathcal{V}_k(\mathbf{x}, \mathbf{v}_{t_k}, \mathbf{v}_{t_{k+1}}, \omega)$

Model relaxation description: Intermediate Economic Dispatches (ED-k)

- We divide the time horizon into K intervals, $\{t_0 = 0 := 1, \dots, t_1\}, \{t_1 + 1, \dots, t_2\}, \dots, \{t_{K-1} + 1, \dots, t_K := T\}$
- We fix a priori the intermediate storage values v_{t_k} for $k = 1, \dots, K$.
- We refer to the (ED) problems restricted to each time interval with fixed initial and final storage values as **(ED-k)** and to its optimal values as and with optimal value $\mathcal{V}_k(\mathbf{x}, \mathbf{v}_{t_k}, \mathbf{v}_{t_{k+1}}, \omega)$

Observation

$$\mathcal{V}(x, \omega) = \min_{\{v_{t_k}\}_{k=1}^K} \sum_{k=0}^{K-1} \mathcal{V}_k(x, v_{t_k}, v_{t_{k+1}}, \omega) \quad (7)$$

Model relaxation description: Lower Approximation of (ED)

Since each function \mathcal{V}_k is piecewise linear convex in $x, v_{t_K}, v_{t_{K+1}}$, it can be approximated by a collection of supporting hyperplanes $\{\pi_{i,k}^w(x, v_{t_k}, v_{t_{k+1}})\}$ of each \mathcal{V}_k .

Model relaxation description: Lower Approximation of (ED)

Since each function \mathcal{V}_k is piecewise linear convex in $x, v_{t_K}, v_{t_{K+1}}$, it can be approximated by a collection of supporting hyperplanes $\{\pi_{i,k}^w(x, v_{t_k}, v_{t_{k+1}})\}$ of each \mathcal{V}_k .

An approximation of (ED) is given by:

Model relaxation description: Lower Approximation of (ED)

Since each function \mathcal{V}_k is piecewise linear convex in $x, v_{t_k}, v_{t_{k+1}}$, it can be approximated by a collection of supporting hyperplanes $\{\pi_{i,k}^w(x, v_{t_k}, v_{t_{k+1}})\}$ of each \mathcal{V}_k .

An approximation of (ED) is given by:

$$\begin{aligned}\hat{\mathcal{V}}(x, \omega) &= \min_{\{v_{t_k}\}_{k=1}^K} \sum_{k=0}^K \hat{\mathcal{V}}_k(x, v_{t_k}, v_{t_{k+1}}) = \\ &= \min_{\{v_{t_k}\}_{k=1}^K} \sum_{k=0}^K \theta_k^\omega \\ &\quad \text{s.t.} \quad \theta_k^\omega \geq \pi_{i,k}^\omega(x, v_{t_k}, v_{t_{k+1}}) \quad \forall i, k\end{aligned}\tag{ISP}$$

We refer to this problem as the **Intermediate Storage Problem (ISP)**

Model relaxation description: Lower Approximation of (ED)

Since each function \mathcal{V}_k is piecewise linear convex in $x, v_{t_K}, v_{t_{K+1}}$, it can be approximated by a collection of supporting hyperplanes $\{\pi_{i,k}^w(x, v_{t_k}, v_{t_{k+1}})\}$ of each \mathcal{V}_k .

An approximation of (ED) is given by:

$$\begin{aligned}\hat{\mathcal{V}}(x, \omega) &= \min_{\{v_{t_k}\}_{k=1}^K} \sum_{k=0}^K \hat{\mathcal{V}}_k(x, v_{t_k}, v_{t_{k+1}}) = \\ &= \min_{\{v_{t_k}\}_{k=1}^K} \sum_{k=0}^K \theta_k^\omega \\ &\text{s.t. } \theta_k^\omega \geq \pi_{i,k}^\omega(x, v_{t_k}, v_{t_{k+1}}) \quad \forall i, k\end{aligned}\tag{ISP}$$

We refer to this problem as the **Intermediate Storage Problem (ISP)**
(I know, very original)

Model description: Relaxed Capacity Expansion(CEP-R)

$$\begin{aligned} \min_x \quad & c'x + \mathbb{E}_\omega [\mathcal{V}(x, \omega)] \\ \text{s.t.} \quad & 0 \leq x_{n,g} \leq X_{n,g} \end{aligned} \quad (\text{CEP})$$

Model description: Relaxed Capacity Expansion(CEP-R)

$$\begin{aligned} \min_x \quad & c'x + \mathbb{E}_\omega [\mathcal{V}(x, \omega)] \\ \text{s.t.} \quad & 0 \leq x_{n,g} \leq X_{n,g} \end{aligned} \tag{CEP}$$

$$\begin{aligned} \min_x \quad & c'x + \mathbb{E}_w [\hat{\mathcal{V}}(x, w)] \\ \text{s.t.} \quad & 0 \leq x_{n,g} \leq X_{n,g} \end{aligned} \tag{CEP-R}$$


Model description: Relaxed Capacity Expansion(CEP-R)

$$\begin{aligned} \min_x \quad & c'x + \mathbb{E}_\omega [\mathcal{V}(x, \omega)] \\ \text{s.t.} \quad & 0 \leq x_{n,g} \leq X_{n,g} \end{aligned} \tag{CEP}$$

$$\begin{aligned} \min_x \quad & c'x + \mathbb{E}_w [\hat{\mathcal{V}}(x, w)] \\ \text{s.t.} \quad & 0 \leq x_{n,g} \leq X_{n,g} \end{aligned} \tag{CEP-R}$$

Since calculating $\hat{\mathcal{V}}$ is straightforward, solving (CEP-R) can be done efficiently with L-shaped or subgradient schemes.

Algorithm



Convergence results 2

- Since $(CEP - R) \leq (CEP)$ if a $(CEP - R)$ optimal solution has the same cost for (CEP) then it's also (CEP) -optimal.

Convergence results 2

- Since $(CEP - R) \leq (CEP)$ if a $(CEP - R)$ optimal solution has the same cost for (CEP) then it's also (CEP) -optimal.

Remark 1: It is sufficient to prove that after a finite number of steps (i) of the algorithm we have:

$$\hat{\mathcal{V}}(\hat{x}^i, \omega) = \mathcal{V}(\hat{x}^i, \omega) \text{ for all } \omega \in \Omega \quad (8)$$

Convergence results 3

Observation

After a finite number of iterations no new cuts are found for \mathcal{V}_k .

Proof.

(9)

Convergence results 3

Observation

After a finite number of iterations no new cuts are found for \mathcal{V}_k .

Proof.

$$\#\{p \mid p \text{ is a normal vector of a supporting hyperplane of } \mathcal{V}_k\} \leq$$

(9)

Convergence results 3

Observation

After a finite number of iterations no new cuts are found for \mathcal{V}_k .

Proof.

$$\begin{aligned} \#\{p \mid p \text{ is a normal vector of a supporting hyperplane of } \mathcal{V}_k\} &\leq \\ \#\{\text{dual solutions } p = q' B^{-1} \text{ of (ED-k) for varying } x, v_{t_k}, v_{t_{k+1}}\} &\leq \end{aligned} \tag{9}$$

Convergence results 3

Observation

After a finite number of iterations no new cuts are found for \mathcal{V}_k .

Proof.

$$\begin{aligned} \#\{p \mid p \text{ is a normal vector of a supporting hyperplane of } \mathcal{V}_k\} &\leq \\ \#\{\text{dual solutions } p = q'B^{-1} \text{ of (ED-k) for varying } x, v_{t_k}, v_{t_{k+1}}\} &\leq \\ \#\{\text{basis matrices of (ED-k)}\} &< \infty \end{aligned} \tag{9}$$

Convergence results 3

Observation

After a finite number of iterations no new cuts are found for \mathcal{V}_k .

Proof.

$$\begin{aligned} \#\{p \mid p \text{ is a normal vector of a supporting hyperplane of } \mathcal{V}_k\} &\leq \\ \#\{\text{dual solutions } p = q'B^{-1} \text{ of (ED-k) for varying } x, v_{t_k}, v_{t_{k+1}}\} &\leq \\ \#\{\text{basis matrices of (ED-k)}\} &< \infty \end{aligned} \tag{9}$$

After a finite number of steps:

Convergence results 3

Observation

After a finite number of iterations no new cuts are found for \mathcal{V}_k .

Proof.

$$\begin{aligned} \#\{p \mid p \text{ is a normal vector of a supporting hyperplane of } \mathcal{V}_k\} &\leq \\ \#\{\text{dual solutions } p = q'B^{-1} \text{ of (ED-k) for varying } x, v_{t_k}, v_{t_{k+1}}\} &\leq \\ \#\{\text{basis matrices of (ED-k)}\} &< \infty \end{aligned} \tag{9}$$

After a finite number of steps:

- new cut: $\bar{c}(x, v) = p'(x, v) + b$

Convergence results 3

Observation

After a finite number of iterations no new cuts are found for \mathcal{V}_k .

Proof.


$$\begin{aligned} \#\{p \mid p \text{ is a normal vector of a supporting hyperplane of } \mathcal{V}_k\} &\leq \\ \#\{\text{dual solutions } p = q'B^{-1} \text{ of (ED-k) for varying } x, v_{t_k}, v_{t_{k+1}}\} &\leq \\ \#\{\text{basis matrices of (ED-k)}\} &< \infty \end{aligned} \tag{9}$$

After a finite number of steps:


- new cut: $\bar{c}(x, v) = p'(x, v) + b$
- an old cut: $\pi(x, v) = p'(x, v) + \bar{b}$



Convergence results 3



Convergence results 3



Since both are supporting hyperplanes it follows that $b = \bar{b}$
(and therefore \bar{c} is not a new cut).

Convergence results 4

Observation

If after the i -iteration no new cuts are added for some i and k then

$$\hat{\mathcal{V}}_k(\hat{x}^i, \hat{v}_k, \hat{v}_{k+1}) = \mathcal{V}_k(\hat{x}^i, \hat{v}_k, \hat{v}_{k+1}).$$

Convergence results 4

Observation

If after the i -iteration no new cuts are added for some i and k then

$$\hat{\mathcal{V}}_k(\hat{x}^i, \hat{v}_k, \hat{v}_{k+1}) = \mathcal{V}_k(\hat{x}^i, \hat{v}_k, \hat{v}_{k+1}).$$

Where $\hat{\mathcal{V}}_k(\hat{x}^i, \hat{v}_{t_k}, \hat{v}_{t_{k+1}}, \omega) = \max_i \pi_{i,k}^\omega(\hat{x}^i, \hat{v}_{t_k}, \hat{v}_{t_{k+1}})$ be the current supporting hyperplane approximation of \mathcal{V}_k .

Convergence results 4

Observation

If after the i -iteration no new cuts are added for some i and k then

$$\hat{\mathcal{V}}_k(\hat{x}^i, \hat{v}_k, \hat{v}_{k+1}) = \mathcal{V}_k(\hat{x}^i, \hat{v}_k, \hat{v}_{k+1}).$$

Where $\hat{\mathcal{V}}_k(\hat{x}^i, \hat{v}_{t_k}, \hat{v}_{t_{k+1}}, \omega) = \max_i \pi_{i,k}^\omega(\hat{x}^i, \hat{v}_{t_k}, \hat{v}_{t_{k+1}})$ be the current supporting hyperplane approximation of \mathcal{V}_k .

Proof.

Let $\bar{c}_k^\omega(x, v_{t_k}) := p'(x - \hat{x}^i, v_{t_k} - \hat{v}_{t_k}) + V_k(\hat{x}^i, \hat{v}_{t_k})$ be the new cut found after the i -th iteration.

Convergence results 4

Observation

If after the i -iteration no new cuts are added for some i and k then

$$\hat{\mathcal{V}}_k(\hat{x}^i, \hat{v}_k, \hat{v}_{k+1}) = \mathcal{V}_k(\hat{x}^i, \hat{v}_k, \hat{v}_{k+1}).$$

Where $\hat{\mathcal{V}}_k(\hat{x}^i, \hat{v}_{t_k}, \hat{v}_{t_{k+1}}, \omega) = \max_i \pi_{i,k}^\omega(\hat{x}^i, \hat{v}_{t_k}, \hat{v}_{t_{k+1}})$ be the current supporting hyperplane approximation of \mathcal{V}_k .

Proof.

Let $\bar{c}_k^\omega(x, v_{t_k}) := p'(x - \hat{x}^i, v_{t_k} - \hat{v}_{t_k}) + V_k(\hat{x}^i, \hat{v}_{t_k})$ be the new cut found after the i -th iteration.

Since \bar{c} is not a new cut we have $\bar{c}(\hat{x}^i, v_{t_k}) \leq \hat{\mathcal{V}}_k(\hat{x}^i, v_{t_k})$.

Convergence results 4

Observation

If after the i -iteration no new cuts are added for some i and k then

$$\hat{\mathcal{V}}_k(\hat{x}^i, \hat{v}_k, \hat{v}_{k+1}) = \mathcal{V}_k(\hat{x}^i, \hat{v}_k, \hat{v}_{k+1}).$$

Where $\hat{\mathcal{V}}_k(\hat{x}^i, \hat{v}_{t_k}, \hat{v}_{t_{k+1}}, \omega) = \max_i \pi_{i,k}^\omega(\hat{x}^i, \hat{v}_{t_k}, \hat{v}_{t_{k+1}})$ be the current supporting hyperplane approximation of \mathcal{V}_k .

Proof.

Let $\bar{c}_k^\omega(x, v_{t_k}) := p'(x - \hat{x}^i, v_{t_k} - \hat{v}_{t_k}) + V_k(\hat{x}^i, \hat{v}_{t_k})$ be the new cut found after the i -th iteration.

Since \bar{c} is not a new cut we have $\bar{c}(\hat{x}^i, v_{t_k}) \leq \hat{\mathcal{V}}_k(\hat{x}^i, v_{t_k})$. We have thus

$$\mathcal{V}_k(\hat{x}^i, \hat{v}_{t_k}) \geq \hat{\mathcal{V}}_k(\hat{x}^i, \hat{v}_{t_k}) \geq \bar{c}(\hat{x}^i, \hat{v}_{t_k}) = \mathcal{V}_k(\hat{x}^i, \hat{v}_{t_k})$$

Convergence results 4

Observation

If after the i -iteration no new cuts are added for some i and k then

$$\hat{\mathcal{V}}_k(\hat{x}^i, \hat{v}_k, \hat{v}_{k+1}) = \mathcal{V}_k(\hat{x}^i, \hat{v}_k, \hat{v}_{k+1}).$$

Where $\hat{\mathcal{V}}_k(\hat{x}^i, \hat{v}_{t_k}, \hat{v}_{t_{k+1}}, \omega) = \max_i \pi_{i,k}^\omega(\hat{x}^i, \hat{v}_{t_k}, \hat{v}_{t_{k+1}})$ be the current supporting hyperplane approximation of \mathcal{V}_k .

Proof.

Let $\bar{c}_k^\omega(x, v_{t_k}) := p'(x - \hat{x}^i, v_{t_k} - \hat{v}_{t_k}) + V_k(\hat{x}^i, \hat{v}_{t_k})$ be the new cut found after the i -th iteration.

Since \bar{c} is not a new cut we have $\bar{c}(\hat{x}^i, v_{t_k}) \leq \hat{\mathcal{V}}_k(\hat{x}^i, v_{t_k})$. We have thus

$$\mathcal{V}_k(\hat{x}^i, \hat{v}_{t_k}) \geq \hat{\mathcal{V}}_k(\hat{x}^i, \hat{v}_{t_k}) \geq \bar{c}(\hat{x}^i, \hat{v}_{t_k}) = \mathcal{V}_k(\hat{x}^i, \hat{v}_{t_k})$$

which concludes the proof. □

Convergence results 5

In conclusion, we have $\hat{\mathcal{V}}_k(\hat{x}^i, v_{t_k}, v_{t_{k+1}}, \omega) = \mathcal{V}_k(\hat{x}^i, v_{t_k}, v_{t_{k+1}}, \omega)$ for all ω, k .

Convergence results 5

In conclusion, we have $\hat{\mathcal{V}}_k(\hat{x}^i, v_{t_k}, v_{t_{k+1}}, \omega) = \mathcal{V}_k(\hat{x}^i, v_{t_k}, v_{t_{k+1}}, \omega)$ for all ω, k .
Thus $\hat{\mathcal{V}}(\hat{x}^i, \omega) = \mathcal{V}(\hat{x}^i, \omega)$.

Convergence results 5

In conclusion, we have $\hat{\mathcal{V}}_k(\hat{x}^i, v_{t_k}, v_{t_{k+1}}, \omega) = \mathcal{V}_k(\hat{x}^i, v_{t_k}, v_{t_{k+1}}, \omega)$ for all ω, k .
Thus $\hat{\mathcal{V}}(\hat{x}^i, \omega) = \mathcal{V}(\hat{x}^i, \omega)$.

Proposition

The algorithm converges after a finite number of iterations and \hat{x}^i is an optimal solution for (CEP).

Future implementation in Pypsa

- We are currently implementing this and other stochastic methods within the Pypsa [BHS18] framework using the Linopy [Hof23] modeling package in Python.

Future implementation in Pypsa

- We are currently implementing this and other stochastic methods within the Pypsa [BHS18] framework using the Linopy [Hof23] modeling package in Python.
- We expect improved convergence speed respect to the L-shaped method, especially when leveraging parallel processing capabilities.

Conclusions and future perspectives.

- The specific structure of the intertemporal constraints makes it possible to develop tailored optimization algorithms for (CEP).

Conclusions and future perspectives.

- The specific structure of the intertemporal constraints makes it possible to develop tailored optimization algorithms for (CEP).
- The presented method can be also applied to other linear models such as DC-OPF.

Conclusions and future perspectives.

- The specific structure of the intertemporal constraints makes it possible to develop tailored optimization algorithms for (CEP).
- The presented method can be also applied to other linear models such as DC-OPF.
- Care must be taken for intra-temporal constraints such as ramp up or ramp down constraints.

Conclusions and future perspectives.

- The specific structure of the intertemporal constraints makes it possible to develop tailored optimization algorithms for (CEP).
- The presented method can be also applied to other linear models such as DC-OPF.
- Care must be taken for intra-temporal constraints such as ramp up or ramp down constraints.
- Develop heuristics for the choice of the number of time intervals K .

Conclusions and future perspectives.

- The specific structure of the intertemporal constraints makes it possible to develop tailored optimization algorithms for (CEP).
- The presented method can be also applied to other linear models such as DC-OPF.
- Care must be taken for intra-temporal constraints such as ramp up or ramp down constraints.
- Develop heuristics for the choice of the number of time intervals K .
- Examine generalizations to multistage stochastic programs.

Conclusions and future perspectives.

- The specific structure of the intertemporal constraints makes it possible to develop tailored optimization algorithms for (CEP).
- The presented method can be also applied to other linear models such as DC-OPF.
- Care must be taken for intra-temporal constraints such as ramp up or ramp down constraints.
- Develop heuristics for the choice of the number of time intervals K .
- Examine generalizations to multistage stochastic programs.

Thank you for your attention.

`gabor.riccardi01@universitadipavia.it`

`https://www.compopt.it`

Some references:

- [Ávi+23] Daniel Ávila, Anthony Papavasiliou, Mauricio Junca, and Lazaros Exizidis. “Applying High-Performance Computing to the European Resource Adequacy Assessment”. In: *IEEE Transactions on Power Systems* (2023), pp. 1–13. DOI: [10.1109/TPWRS.2023.3304717](https://doi.org/10.1109/TPWRS.2023.3304717).
- [Bie+20] Daniel Bienstock, Mauro Escobar, Claudio Gentile, and Leo Liberti. “Mathematical Programming formulations for the Alternating Current Optimal Power Flow problem”. In: *4OR* 18.3 (July 2020), pp. 249–292. DOI: [10.1007/s10288-020-00455-w](https://doi.org/10.1007/s10288-020-00455-w).
- [BM14] Daniel Bienstock and Gonzalo Munoz. “On linear relaxations of OPF problems”. In: (Nov. 2014).
- [BHS18] T. Brown, J. Hörsch, and D. Schlachtberger. “PyPSA: Python for Power System Analysis”. In: *Journal of Open Research Software* 6.4 (1 2018). DOI: [10.5334/jors.188](https://doi.org/10.5334/jors.188). eprint: 1707.09913. URL: <https://doi.org/10.5334/jors.188>.
- [Hof23] Fabian Hofmann. “Linopy: Linear optimization with n-dimensional labeled variables”. In: *Journal of Open Source Software* 8.84 (2023), p. 4823. DOI: [10.21105/joss.04823](https://doi.org/10.21105/joss.04823). URL: <https://doi.org/10.21105/joss.04823>.

Adequacy Assessment of the Electrical Grid

- Measuring the ability of the electric power system to react to adverse uncertain condition has become increasingly important.
- Member States wishing to introduce capacity mechanisms can do so if an adequacy concern is identified in the ERAA study, a pan-European adequacy assessment for up to 10 years ahead.
- Due to the scale of the ERAA study, ERAA 2022 considered a reduced stochastic problem with three scenarios.
- In [Ávi+23], Daniel A'vila introduced a decomposition algorithm based on subgradient approximations was introduced

Power Grid Optimization

Stochasticity —
Time / Exactness ↑

- Optimal Power Flow (OPF) [Bie+20]
 - AC OPF: exact physical model
 - Security-Constrained OPF (SCOPF) – Includes contingencies to guarantee system security under failures.
 - DC OPF and other linearized models [BM14]
 - other relaxations.
- Unit Commitment – Determines on/off status of power units, ignoring grid constraints.
- Economic Dispatch (ED) – Minimizes generation cost, ignoring grid constraints.

Capacity expansion problem: Based on Economic Dispatch models with added flow balance at bus nodes and various scenarios.