



# *Hypergeometric Motives*

Fernando Rodriguez Villegas

The Abdus Salam International Centre for Theoretical Physics

May 2015

## *Collaborators*

H. Cohen, Ph. Candelas, S. Baig, F. Beukers, X. de la Ossa,  
A. Mellit, P. Molin, D. Roberts, M. Vashenko, M. Watkins

## *L-functions*



$$\Lambda(s) = N^{s/2} L_{\infty}(s) \prod_p L_p(p^{-s})^{-1}$$

## *L-functions*



$$\Lambda(s) = N^{s/2} L_{\infty}(s) \prod_p L_p(p^{-s})^{-1}$$

- ▶ Conductor:  $N$ , a positive integer.

# *L-functions*



$$\Lambda(s) = N^{s/2} L_{\infty}(s) \prod_p L_p(p^{-s})^{-1}$$

- ▶ Conductor:  $N$ , a positive integer.
- ▶ Euler factors:  $L_p(T) \in \mathbb{Z}[T]$  polynomial generically of degree  $d$ .

# *L-functions*



$$\Lambda(s) = N^{s/2} L_{\infty}(s) \prod_p L_p(p^{-s})^{-1}$$

- ▶ Conductor:  $N$ , a positive integer.
- ▶ Euler factors:  $L_p(T) \in \mathbb{Z}[T]$  polynomial generically of degree  $d$ .
- ▶ Weight  $w$

$$L_p(T) = \prod_{i=1}^d (1 - \xi_i T), \quad |\xi_i| = p^{w/2}, \quad p \nmid N.$$

# *L-functions*



$$\Lambda(s) = N^{s/2} L_{\infty}(s) \prod_p L_p(p^{-s})^{-1}$$

- ▶ Conductor:  $N$ , a positive integer.
- ▶ Euler factors:  $L_p(T) \in \mathbb{Z}[T]$  polynomial generically of degree  $d$ .
- ▶ Weight  $w$

$$L_p(T) = \prod_{i=1}^d (1 - \xi_i T), \quad |\xi_i| = p^{w/2}, \quad p \nmid N.$$

- ▶  $L_{\infty}(s)$  product of gamma factors



# *L-functions*



$$\Lambda(s) = N^{s/2} L_{\infty}(s) \prod_p L_p(p^{-s})^{-1}$$

- ▶ Conductor:  $N$ , a positive integer.
- ▶ Euler factors:  $L_p(T) \in \mathbb{Z}[T]$  polynomial generically of degree  $d$ .
- ▶ Weight  $w$

$$L_p(T) = \prod_{i=1}^d (1 - \xi_i T), \quad |\xi_i| = p^{w/2}, \quad p \nmid N.$$

- ▶  $L_{\infty}(s)$  product of gamma factors
- ▶ Expect functional equation

$$\Lambda(w + 1 - s) = \epsilon \Lambda(s), \quad \epsilon = \pm 1$$

## Computing $\Lambda$

- ▶ Inverse Mellin transform

$$L_{\infty}(s) = \int_0^{\infty} k_{\infty}(t) t^s \frac{dt}{t}$$

## Computing $\Lambda$

- ▶ Inverse Mellin transform

$$L_{\infty}(s) = \int_0^{\infty} k_{\infty}(t) t^s \frac{dt}{t}$$



$$\sum_{n \geq 1} \frac{a_n}{n^s} := \prod_p L_p(p^{-s})^{-1}.$$

converges in some half-plane.

## Computing $\Lambda$

- ▶ Inverse Mellin transform

$$L_{\infty}(s) = \int_0^{\infty} k_{\infty}(t) t^s \frac{dt}{t}$$



$$\sum_{n \geq 1} \frac{a_n}{n^s} := \prod_p L_p(p^{-s})^{-1}.$$

converges in some half-plane.



$$\Lambda(s) = \int_0^{\infty} \phi(t) t^s \frac{dt}{t}, \quad \phi(t) := \sum_{n \geq 1} a_n k_{\infty} \left( \frac{nt}{\sqrt{N}} \right).$$

## Computing $\Lambda$

- ▶ Inverse Mellin transform

$$L_{\infty}(s) = \int_0^{\infty} k_{\infty}(t) t^s \frac{dt}{t}$$



$$\sum_{n \geq 1} \frac{a_n}{n^s} := \prod_p L_p(p^{-s})^{-1}.$$

converges in some half-plane.



$$\Lambda(s) = \int_0^{\infty} \phi(t) t^s \frac{dt}{t}, \quad \phi(t) := \sum_{n \geq 1} a_n k_{\infty} \left( \frac{nt}{\sqrt{N}} \right).$$

- ▶ Functional equation

$$\phi(t^{-1}) = \epsilon t^{w+1} \phi(t)$$

## *Computing $\Lambda$ (cont'd)*

- ▶  $k_\infty$  is of rapid decay  $\implies$  can compute  $\phi$  if we know

## Computing $\Lambda$ (cont'd)

- ▶  $k_\infty$  is of rapid decay  $\implies$  can compute  $\phi$  if we know



$$k_\infty, \quad N, \quad \epsilon, \quad a_n, \quad n = 1, 2, \dots, \approx \sqrt{N}$$

## Computing $\Lambda$ (cont'd)

- ▶  $k_\infty$  is of rapid decay  $\implies$  can compute  $\phi$  if we know



$$k_\infty, \quad N, \quad \epsilon, \quad a_n, \quad n = 1, 2, \dots, \approx \sqrt{N}$$

- ▶ Functional equation for  $\phi \implies$  can compute  $\Lambda(s)$  (Riemann).



## Computing $\Lambda$ (cont'd)

- ▶  $k_\infty$  is of rapid decay  $\implies$  can compute  $\phi$  if we know



$$k_\infty, \quad N, \quad \epsilon, \quad a_n, \quad n = 1, 2, \dots, \approx \sqrt{N}$$

- ▶ Functional equation for  $\phi \implies$  can compute  $\Lambda(s)$  (Riemann).

- ▶ Typically calculation breaks up into:

$$L_p(T), \quad p \notin S, \quad L_p(T), \quad p \in S, \quad L_\infty(s), \quad N, \quad \epsilon$$

$S$  finite set of primes.

## *Computing $\Lambda$ (cont'd)*

- ▶ In practice: only know some of the data.

## *Computing $\Lambda$ (cont'd)*

- ▶ In practice: only know some of the data.
- ▶ Test functional equation for  $\phi$ : evaluating both sides at  $t \approx 1$ .

## *Computing $\Lambda$ (cont'd)*

- ▶ In practice: only know some of the data.
- ▶ Test functional equation for  $\phi$ : evaluating both sides at  $t \approx 1$ .
- ▶ Use test to find missing data.

## Computing $\Lambda$ (cont'd)

- ▶ In practice: only know some of the data.
- ▶ Test functional equation for  $\phi$ : evaluating both sides at  $t \approx 1$ .
- ▶ Use test to find missing data.
- ▶ Robust test.

## Computing $\Lambda$ (cont'd)

- ▶ In practice: only know some of the data.
- ▶ Test functional equation for  $\phi$ : evaluating both sides at  $t \approx 1$ .
- ▶ Use test to find missing data.
- ▶ Robust test.
- ▶ Bootstrap.

## Computing $\Lambda$ (cont'd)

- ▶ In practice: only know some of the data.
- ▶ Test functional equation for  $\phi$ : evaluating both sides at  $t \approx 1$ .
- ▶ Use test to find missing data.
- ▶ Robust test.
- ▶ Bootstrap.
- ▶ HGM implementation in MAGMA (M. Watkins)

## *Source of $L$ -functions*

- ▶ Automorphic Forms.



## *Source of $L$ -functions*

- ▶ Automorphic Forms.
- ▶ Cohomology of algebraic varieties.

## *Source of L-functions*

- ▶ Automorphic Forms.
- ▶ Cohomology of algebraic varieties.
- ▶ Typically appear as piece of a bigger object

## *Source of L-functions*

- ▶ Automorphic Forms.
- ▶ Cohomology of algebraic varieties.
- ▶ Typically appear as piece of a bigger object
- ▶ Subspace  $V \subseteq H$

$$L_p(T) = \det(1 - \text{Frob}_p|_V T), \quad p \nmid N$$

# *Automorphic Forms*

- ▶ Usual modular forms:

$$d = \dim V = 2, \qquad \dim H \approx cN$$

# *Automorphic Forms*

- ▶ Usual modular forms:

$$d = \dim V = 2, \qquad \dim H \approx cN$$

- ▶ Siegel modular forms  $\mathrm{Sp}_4$ : can compute for  $N \approx 500$

# *Automorphic Forms*

- ▶ Usual modular forms:

$$d = \dim V = 2, \qquad \dim H \approx cN$$

- ▶ Siegel modular forms  $\mathrm{Sp}_4$ : can compute for  $N \approx 500$
- ▶  $\mathrm{SL}_4$  happy if can compute  $L_2(T)$ .

# *Algebraic Varieties*

- ▶ Quintic threefold

$$X : F(x_1, \dots, x_5) = 0$$

# *Algebraic Varieties*

- ▶ Quintic threefold

$$X : F(x_1, \dots, x_5) = 0$$



$$H := H^3(X, \mathbb{Q}), \quad \dim H = 204.$$



# *Algebraic Varieties*

- ▶ Quintic threefold

$$X : F(x_1, \dots, x_5) = 0$$



$$H := H^3(X, \mathbb{Q}), \quad \dim H = 204.$$

- ▶ Dwork pencil

$$X_\psi : x_1^5 + \dots x_5^5 - 5\psi x_1 \dots x_5 = 0$$

# *Algebraic Varieties*

- ▶ Quintic threefold

$$X : F(x_1, \dots, x_5) = 0$$



$$H := H^3(X, \mathbb{Q}), \quad \dim H = 204.$$

- ▶ Dwork pencil

$$X_\psi : x_1^5 + \dots x_5^5 - 5\psi x_1 \dots x_5 = 0$$



$$A \subseteq \text{Aut}(X_\psi), \quad x_i \mapsto \zeta_i x_i, \quad \zeta_1^5 = \dots = \zeta_5^5 = \zeta_1 \dots \zeta_5 = 1$$

# *Algebraic Varieties*

- ▶ Quintic threefold

$$X : F(x_1, \dots, x_5) = 0$$



$$H := H^3(X, \mathbb{Q}), \quad \dim H = 204.$$

- ▶ Dwork pencil

$$X_\psi : x_1^5 + \dots x_5^5 - 5\psi x_1 \dots x_5 = 0$$



$$A \subseteq \text{Aut}(X_\psi), \quad x_i \mapsto \zeta_i x_i, \quad \zeta_1^5 = \dots = \zeta_5^5 = \zeta_1 \dots \zeta_5 = 1$$



$$V := H^A, \quad \dim V = 4$$

## *Hodge numbers*

- ▶ Refinement of rank  $d$  that determines  $L_{\infty}(s)$ .

# *Hodge numbers*

- ▶ Refinement of rank  $d$  that determines  $L_\infty(s)$ .



$$h^{p,q} \in \mathbb{Z}_{\geq 0}, \quad p + q = w$$

# Hodge numbers

- ▶ Refinement of rank  $d$  that determines  $L_\infty(s)$ .



$$h^{p,q} \in \mathbb{Z}_{\geq 0}, \quad p + q = w$$



$$h^{p,q} = h^{q,p}, \quad \sum_{p,q} h^{p,q} = d$$

# Hodge numbers

- ▶ Refinement of rank  $d$  that determines  $L_\infty(s)$ .



$$h^{p,q} \in \mathbb{Z}_{\geq 0}, \quad p + q = w$$



$$h^{p,q} = h^{q,p}, \quad \sum_{p,q} h^{p,q} = d$$

- ▶ Hodge vector (up to Tate twists)

$$\mathbf{h} := (h^{w,0}, h^{w-1,1}, \dots, h^{0,w})$$

# Hodge numbers

- ▶ Refinement of rank  $d$  that determines  $L_\infty(s)$ .



$$h^{p,q} \in \mathbb{Z}_{\geq 0}, \quad p + q = w$$



$$h^{p,q} = h^{q,p}, \quad \sum_{p,q} h^{p,q} = d$$

- ▶ Hodge vector (up to Tate twists)

$$\mathbf{h} := (h^{w,0}, h^{w-1,1}, \dots, h^{0,w})$$



$$h^{p,p} = h_+^{p,p} + h_-^{p,p}$$



# *Hypergeometric Motives*

- ▶  $q_0, q_\infty \in \mathbb{Z}[T]$ , coprime, same degree  $d$ , roots are roots of unity.

# *Hypergeometric Motives*

- ▶  $q_0, q_\infty \in \mathbb{Z}[T]$ , coprime, same degree  $d$ , roots are roots of unity.
- ▶ Get associated family of motives  $\mathcal{H}(t)$  with  $t \in \mathbb{P}^1 \setminus \{0, 1, \infty\}$ .

# *Hypergeometric Motives*

- ▶  $q_0, q_\infty \in \mathbb{Z}[T]$ , coprime, same degree  $d$ , roots are roots of unity.
- ▶ Get associated family of motives  $\mathcal{H}(t)$  with  $t \in \mathbb{P}^1 \setminus \{0, 1, \infty\}$ .
- ▶  $\mathcal{H}(t)$  has rank  $d$  and a computable weight  $w$  in terms of  $q_0, q_\infty$ .

# Hypergeometric Motives

- ▶  $q_0, q_\infty \in \mathbb{Z}[T]$ , coprime, same degree  $d$ , roots are roots of unity.
- ▶ Get associated family of motives  $\mathcal{H}(t)$  with  $t \in \mathbb{P}^1 \setminus \{0, 1, \infty\}$ .
- ▶  $\mathcal{H}(t)$  has rank  $d$  and a computable weight  $w$  in terms of  $q_0, q_\infty$ .
- ▶ Can also compute the Hodge numbers:  $\implies L_\infty(s)$

# Hypergeometric Motives

- ▶  $q_0, q_\infty \in \mathbb{Z}[T]$ , coprime, same degree  $d$ , roots are roots of unity.
- ▶ Get associated family of motives  $\mathcal{H}(t)$  with  $t \in \mathbb{P}^1 \setminus \{0, 1, \infty\}$ .
- ▶  $\mathcal{H}(t)$  has rank  $d$  and a computable weight  $w$  in terms of  $q_0, q_\infty$ .
- ▶ Can also compute the Hodge numbers:  $\implies L_\infty(s)$
- ▶ For fixed  $t \in \mathbb{Q}$ : formula for  $L_p(T)$  for  $p \notin S$  (Katz's hypergeometric trace)

## Examples

- ▶ Belyi polynomials  $c := a + b$

$$\mathbb{Q}[x]/(B(a, b; t)), \quad B(a, b; t) := x^a(1-x)^b - \frac{a^a b^b}{c^c} t$$

## Examples

- ▶ Belyi polynomials  $c := a + b$

$$\mathbb{Q}[x]/(B(a, b; t)), \quad B(a, b; t) := x^a(1-x)^b - \frac{a^a b^b}{c^c} t$$



$$\frac{q_\infty}{q_0} = \frac{T^c - 1}{(T^a - 1)(T^b - 1)}$$

## Examples

- ▶ Belyi polynomials  $c := a + b$

$$\mathbb{Q}[x]/(B(a, b; t)), \quad B(a, b; t) := x^a(1-x)^b - \frac{a^a b^b}{c^c} t$$



$$\frac{q_\infty}{q_0} = \frac{T^c - 1}{(T^a - 1)(T^b - 1)}$$

- ▶ Legendre family of elliptic curves:  $H^1(E_t)$

$$E_t : y^2 = x(x-1)(x-t)$$



## Examples

- ▶ Belyi polynomials  $c := a + b$

$$\mathbb{Q}[x]/(B(a, b; t)), \quad B(a, b; t) := x^a(1-x)^b - \frac{a^a b^b}{c^c} t$$



$$\frac{q_\infty}{q_0} = \frac{T^c - 1}{(T^a - 1)(T^b - 1)}$$

- ▶ Legendre family of elliptic curves:  $H^1(E_t)$

$$E_t : y^2 = x(x-1)(x-t)$$



$$\frac{q_\infty}{q_0} = \frac{(T+1)^2}{(T-1)^2}$$

## Examples

- ▶ Belyi polynomials  $c := a + b$

$$\mathbb{Q}[x]/(B(a, b; t)), \quad B(a, b; t) := x^a(1-x)^b - \frac{a^a b^b}{c^c} t$$



$$\frac{q_\infty}{q_0} = \frac{T^c - 1}{(T^a - 1)(T^b - 1)}$$

- ▶ Legendre family of elliptic curves:  $H^1(E_t)$

$$E_t : y^2 = x(x-1)(x-t)$$



$$\frac{q_\infty}{q_0} = \frac{(T+1)^2}{(T-1)^2}$$

- ▶ Dwork pencil piece:  $V$

$$\frac{q_\infty}{q_0} = \frac{T^5 - 1}{(T - 1)^5}$$

## *Hodge vectors*

- ▶ By Griffiths transversality  $\mathbf{h}$  is a symmetric composition of  $d$ .

## *Hodge vectors*

- ▶ By Griffiths transversality  $\mathbf{h}$  is a symmetric composition of  $d$ .
- ▶ Total number:  $2^{\lfloor d/2 \rfloor}$

## *Hodge vectors*

► By Griffiths transversality  $\mathbf{h}$  is a symmetric composition of  $d$ .

► Total number:  $2^{\lfloor d/2 \rfloor}$

►

$(2), \quad (1, 1)$

$(3), \quad (1, 1, 1)$

$(4), \quad (2, 2), \quad (1, 2, 1), \quad (1, 1, 1, 1)$

$(5), \quad (2, 1, 2), \quad (1, 3, 1), \quad (1, 1, 1, 1, 1)$

## *Rank 24*

- ▶ Rank  $d = 24$ . Number of possible Hodge vectors: 4096.

## *Rank 24*

- ▶ Rank  $d = 24$ . Number of possible Hodge vectors: 4096.
- ▶ Total number of family of motives: 464, 247, 183

## Rank 24

- ▶ Rank  $d = 24$ . Number of possible Hodge vectors: 4096.
- ▶ Total number of family of motives: 464, 247, 183

	<b>h</b>	<b>#</b>
	[9, 1, 1, 2, 1, 1, 9]	0
	[7, 1, 1, 1, 1, 2, 1, 1, 1, 1, 7]	0
	[1, 6, 1, 1, 1, 1, 2, 1, 1, 1, 1, 6, 1]	0
▶	[4, 1, 3, 1, 1, 1, 2, 1, 1, 1, 3, 1, 4]	0
	[5, 1, 2, 1, 1, 1, 2, 1, 1, 1, 2, 1, 5]	0
	[6, 1, 1, 1, 1, 1, 2, 1, 1, 1, 1, 1, 6]	0
	[4, 1, 1, 2, 1, 1, 1, 2, 1, 1, 1, 2, 1, 1, 4]	0
	[4, 1, 2, 1, 1, 1, 1, 2, 1, 1, 1, 1, 2, 1, 4]	0




# Rank 24

<b>h</b>	<b>#</b>
[6, 2, 1, 1, 1, 2, 1, 1, 1, 2, 6]	2
[8, 1, 1, 1, 2, 1, 1, 1, 8]	4
[1, 22, 1]	4
[8, 1, 1, 4, 1, 1, 8]	6
[6, 1, 2, 1, 1, 2, 1, 1, 2, 1, 6]	8
[6, 1, 3, 1, 2, 1, 3, 1, 6]	8
[10, 1, 2, 1, 10]	10
⋮	⋮
[1, 3, 4, 4, 4, 4, 3, 1]	6082776
[2, 5, 5, 5, 5, 2]	6850823
[1, 3, 8, 8, 3, 1]	6868016
[1, 5, 6, 6, 5, 1]	7637828
[1, 2, 4, 5, 5, 4, 2, 1]	7982874
[2, 4, 6, 6, 4, 2]	9504072
[1, 4, 7, 7, 4, 1]	9905208

# Densities

Graph of logarithmic densities, rank  $d = 24$



## *Useful features*

- ▶  $\mathcal{H}(t)$ : family over  $\mathbb{P}^1 \setminus \{0, 1, \infty\}$ .

## *Useful features*

- ▶  $\mathcal{H}(t)$ : family over  $\mathbb{P}^1 \setminus \{0, 1, \infty\}$ .
- ▶ Study of  $L_p(T)$  for tame primes  $p$  from behaviour at cusps.

## *Useful features*

- ▶  $\mathcal{H}(t)$ : family over  $\mathbb{P}^1 \setminus \{0, 1, \infty\}$ .
- ▶ Study of  $L_p(T)$  for tame primes  $p$  from behaviour at cusps.
- ▶ Cusps  $0, \infty$  with full Jordan blocks  $\implies$  Hecke characters.

## *Useful features*

- ▶  $\mathcal{H}(t)$ : family over  $\mathbb{P}^1 \setminus \{0, 1, \infty\}$ .
- ▶ Study of  $L_p(T)$  for tame primes  $p$  from behaviour at cusps.
- ▶ Cusps  $0, \infty$  with full Jordan blocks  $\implies$  Hecke characters.
- ▶ Families  $\mathcal{H}(t)$  share many properties.

## *Useful features*

- ▶  $\mathcal{H}(t)$ : family over  $\mathbb{P}^1 \setminus \{0, 1, \infty\}$ .
- ▶ Study of  $L_p(T)$  for tame primes  $p$  from behaviour at cusps.
- ▶ Cusps  $0, \infty$  with full Jordan blocks  $\implies$  Hecke characters.
- ▶ Families  $\mathcal{H}(t)$  share many properties.
- ▶ Study simple cases to uncover them.

## *Hypergeometric series*

- ▶ Hypergeometric series  $|t| < 1$

$$u(t) = {}_dF_{d-1} \left[ \begin{matrix} \alpha_1 & \cdots & \alpha_d \\ \beta_1 & \cdots & \beta_{d-1} \end{matrix} \mid t \right] := \sum_{n \geq 0} \frac{(\alpha_1)_n \cdots (\alpha_d)_n}{(\beta_1)_n \cdots (\beta_{d-1})_n} \frac{t^n}{n!},$$



## Hypergeometric series

- ▶ Hypergeometric series  $|t| < 1$

$$u(t) = {}_dF_{d-1} \left[ \begin{matrix} \alpha_1 & \cdots & \alpha_d \\ \beta_1 & \cdots & \beta_{d-1} \end{matrix} \mid t \right] := \sum_{n \geq 0} \frac{(\alpha_1)_n \cdots (\alpha_d)_n}{(\beta_1)_n \cdots (\beta_{d-1})_n} \frac{t^n}{n!},$$



$$(\alpha)_n := \alpha(\alpha + 1) \cdots (\alpha + n - 1)$$

is the Pochhammer symbol.

# Hypergeometric series

- ▶ Hypergeometric series  $|t| < 1$

$$u(t) = {}_dF_{d-1} \left[ \begin{matrix} \alpha_1 & \cdots & \alpha_d \\ \beta_1 & \cdots & \beta_{d-1} \end{matrix} \mid t \right] := \sum_{n \geq 0} \frac{(\alpha_1)_n \cdots (\alpha_d)_n}{(\beta_1)_n \cdots (\beta_{d-1})_n} \frac{t^n}{n!},$$



$$(\alpha)_n := \alpha(\alpha + 1) \cdots (\alpha + n - 1)$$

is the Pochhammer symbol.

- ▶ Satisfies linear differential equation of order  $d$  with regular singularities at  $t = 0, 1, \infty$ .

# Hypergeometric series

- ▶ Hypergeometric series  $|t| < 1$

$$u(t) = {}_dF_{d-1} \left[ \begin{matrix} \alpha_1 & \cdots & \alpha_d \\ \beta_1 & \cdots & \beta_{d-1} \end{matrix} \mid t \right] := \sum_{n \geq 0} \frac{(\alpha_1)_n \cdots (\alpha_d)_n}{(\beta_1)_n \cdots (\beta_{d-1})_n} \frac{t^n}{n!},$$



$$(\alpha)_n := \alpha(\alpha + 1) \cdots (\alpha + n - 1)$$

is the Pochhammer symbol.

- ▶ Satisfies linear differential equation of order  $d$  with regular singularities at  $t = 0, 1, \infty$ .
- ▶ Gives rise to a monodromy representation

$$\rho : \pi_1(\mathbb{P}^1 \setminus \{0, 1, \infty\}) \rightarrow \mathrm{GL}(V)$$

# Hypergeometric series

- ▶ Hypergeometric series  $|t| < 1$

$$u(t) = {}_dF_{d-1} \left[ \begin{matrix} \alpha_1 & \cdots & \alpha_d \\ \beta_1 & \cdots & \beta_{d-1} \end{matrix} \mid t \right] := \sum_{n \geq 0} \frac{(\alpha_1)_n \cdots (\alpha_d)_n}{(\beta_1)_n \cdots (\beta_{d-1})_n} \frac{t^n}{n!},$$



$$(\alpha)_n := \alpha(\alpha + 1) \cdots (\alpha + n - 1)$$

is the Pochhammer symbol.

- ▶ Satisfies linear differential equation of order  $d$  with regular singularities at  $t = 0, 1, \infty$ .
- ▶ Gives rise to a monodromy representation

$$\rho : \pi_1(\mathbb{P}^1 \setminus \{0, 1, \infty\}) \rightarrow \mathrm{GL}(V)$$

- ▶  $V :=$  space of local solutions of the DE at  $z = t \in \mathbb{P}^1 \setminus \{0, 1, \infty\}$ .

## *Hypergeometric series (cont'd)*

- ▶ This local system is rigid.

## *Hypergeometric series (cont'd)*

- ▶ This local system is rigid.
- ▶ The local monodromies  $h_0, h_1, h_\infty$  satisfy

$$h_0 h_1 h_\infty = \text{id}_V$$

## *Hypergeometric series (cont'd)*

- ▶ This local system is rigid.
- ▶ The local monodromies  $h_0, h_1, h_\infty$  satisfy

$$h_0 h_1 h_\infty = \text{id}_V$$

- ▶  $h_1$  is a pseudo-reflection

## Hypergeometric series (cont'd)

- ▶ This local system is rigid.
- ▶ The local monodromies  $h_0, h_1, h_\infty$  satisfy

$$h_0 h_1 h_\infty = \text{id}_V$$

- ▶  $h_1$  is a pseudo-reflection
- ▶ Characteristic polynomials

$$q_\infty(T) := \prod_{j=1}^d (T - e^{2\pi i \alpha_j}), \quad q_0(T) := \prod_{j=1}^d (T - e^{2\pi i (1 - \beta_j)})$$

and  $h_\infty, h_0$  have full Jordan blocks.



## Hypergeometric series (cont'd)

- ▶ This local system is rigid.
- ▶ The local monodromies  $h_0, h_1, h_\infty$  satisfy

$$h_0 h_1 h_\infty = \text{id}_V$$

- ▶  $h_1$  is a pseudo-reflection
- ▶ Characteristic polynomials

$$q_\infty(T) := \prod_{j=1}^d (T - e^{2\pi i \alpha_j}), \quad q_0(T) := \prod_{j=1}^d (T - e^{2\pi i (1 - \beta_j)})$$

and  $h_\infty, h_0$  have full Jordan blocks.

- ▶  $\rho$  is irreducible if  $q_\infty$  and  $q_0$  are coprime.

## Hypergeometric trace



$$H_p(\alpha, \beta \mid t) := \frac{1}{1-p} \sum_{\varrho} (-p)^{\eta_f(\varrho)} \frac{(\alpha_1)_{\infty, \varrho} \cdots (\alpha_r)_{\infty, \varrho}}{(\beta_1)_{0, \varrho} \cdots (\beta_s)_{0, \varrho}} \omega(t)^{(p-1)\varrho},$$

## Hypergeometric trace



$$H_p(\alpha, \beta \mid t) := \frac{1}{1-p} \sum_{\varrho} (-p)^{\eta_f(\varrho)} \frac{(\alpha_1)_{\infty, \varrho} \cdots (\alpha_r)_{\infty, \varrho}}{(\beta_1)_{0, \varrho} \cdots (\beta_s)_{0, \varrho}} \omega(t)^{(p-1)\varrho},$$

- For  $x \in \mathbb{Q}/\mathbb{Z}$  with denominator coprime to  $p$

$$(x)_{\nu, \varrho} := \frac{\Gamma_p(\{x - \varrho\}_{\nu})}{\Gamma_p(\{x\}_{\nu})}, \quad \nu = 0, \infty, \quad \varrho \in \frac{1}{p-1} \mathbb{Z}/\mathbb{Z},$$

## Hypergeometric trace



$$H_p(\alpha, \beta \mid t) := \frac{1}{1-p} \sum_{\varrho} (-p)^{\eta_f(\varrho)} \frac{(\alpha_1)_{\infty, \varrho} \cdots (\alpha_r)_{\infty, \varrho}}{(\beta_1)_{0, \varrho} \cdots (\beta_s)_{0, \varrho}} \omega(t)^{(p-1)\varrho},$$

- ▶ For  $x \in \mathbb{Q}/\mathbb{Z}$  with denominator coprime to  $p$

$$(x)_{\nu, \varrho} := \frac{\Gamma_p(\{x - \varrho\}_{\nu})}{\Gamma_p(\{x\}_{\nu})}, \quad \nu = 0, \infty, \quad \varrho \in \frac{1}{p-1} \mathbb{Z}/\mathbb{Z},$$

- ▶  $\Gamma_p$ : Morita's  $p$ -adic gamma function

## Hypergeometric trace



$$H_p(\alpha, \beta \mid t) := \frac{1}{1-p} \sum_{\varrho} (-p)^{\eta_f(\varrho)} \frac{(\alpha_1)_{\infty, \varrho} \cdots (\alpha_r)_{\infty, \varrho}}{(\beta_1)_{0, \varrho} \cdots (\beta_s)_{0, \varrho}} \omega(t)^{(p-1)\varrho},$$

- ▶ For  $x \in \mathbb{Q}/\mathbb{Z}$  with denominator coprime to  $p$

$$(x)_{\nu, \varrho} := \frac{\Gamma_p(\{x - \varrho\}_{\nu})}{\Gamma_p(\{x\}_{\nu})}, \quad \nu = 0, \infty, \quad \varrho \in \frac{1}{p-1} \mathbb{Z}/\mathbb{Z},$$

- ▶  $\Gamma_p$ : Morita's  $p$ -adic gamma function



$$\{x\}_{\infty} := \{x\}, \quad \{x\}_0 := 1 - \{-x\},$$

## *Chebyshev example*

- ▶ Interlacing roots

$$q_{\infty} = \Phi_{30},$$

$$q_0 = \Phi_1 \Phi_2 \Phi_3 \Phi_5,$$

## *Chebyshev example*

- ▶ Interlacing roots

$$q_{\infty} = \Phi_{30}, \quad q_0 = \Phi_1 \Phi_2 \Phi_3 \Phi_5,$$



$$\begin{aligned} \alpha &= 1/30, 7/30, 11/30, 13/30, 17/30, 19/30, 23/30, 29/30 \\ \beta &= 1, 1/2, 1/3, 2/3, 1/5, 2/5, 3/5, 4/5 \end{aligned}$$

## *Chebyshev example*

- ▶ Interlacing roots

$$q_{\infty} = \Phi_{30}, \quad q_0 = \Phi_1 \Phi_2 \Phi_3 \Phi_5,$$



$$\begin{aligned} \alpha &= 1/30, 7/30, 11/30, 13/30, 17/30, 19/30, 23/30, 29/30 \\ \beta &= 1, 1/2, 1/3, 2/3, 1/5, 2/5, 3/5, 4/5 \end{aligned}$$



$$\frac{q_{\infty}}{q_0} = \frac{(T^{30} - 1)(T - 1)}{(T^{15} - 1)(T^{10} - 1)(T^6 - 1)}.$$



## Chebyshev example

- ▶ Interlacing roots

$$q_{\infty} = \Phi_{30}, \quad q_0 = \Phi_1 \Phi_2 \Phi_3 \Phi_5,$$



$$\begin{aligned} \alpha &= 1/30, 7/30, 11/30, 13/30, 17/30, 19/30, 23/30, 29/30 \\ \beta &= 1, 1/2, 1/3, 2/3, 1/5, 2/5, 3/5, 4/5 \end{aligned}$$



$$\frac{q_{\infty}}{q_0} = \frac{(T^{30} - 1)(T - 1)}{(T^{15} - 1)(T^{10} - 1)(T^6 - 1)}.$$



$$u(t) := \sum_{n \geq 0} \frac{(30n)!n!}{(15n)!(10n)!(6n)!} \left( \frac{t}{M} \right)^n, \quad M := \frac{30^{30}}{15^{15} \cdot 10^{10} \cdot 6^6}.$$

## Chebyshev example

- ▶ Interlacing roots

$$q_{\infty} = \Phi_{30}, \quad q_0 = \Phi_1 \Phi_2 \Phi_3 \Phi_5,$$



$$\begin{aligned} \alpha &= 1/30, 7/30, 11/30, 13/30, 17/30, 19/30, 23/30, 29/30 \\ \beta &= 1, 1/2, 1/3, 2/3, 1/5, 2/5, 3/5, 4/5 \end{aligned}$$



$$\frac{q_{\infty}}{q_0} = \frac{(T^{30} - 1)(T - 1)}{(T^{15} - 1)(T^{10} - 1)(T^6 - 1)}.$$



$$u(t) := \sum_{n \geq 0} \frac{(30n)!n!}{(15n)!(10n)!(6n)!} \left( \frac{t}{M} \right)^n, \quad M := \frac{30^{30}}{15^{15} \cdot 10^{10} \cdot 6^6}.$$



$$\frac{(30n)!n!}{(15n)!(10n)!(6n)!} = 1, 77636318760, 53837289804317953893960, \dots$$

are integral for every  $n$ .

## *Chebyshev example (cont'd)*

- ▶ Monodromy group is finite.

## *Chebyshev example (cont'd)*

- ▶ Monodromy group is finite.
- ▶ Series  $u(t)$ : Taylor expansion of an algebraic function of  $t$ .

## *Chebyshev example (cont'd)*

- ▶ Monodromy group is finite.
- ▶ Series  $u(t)$ : Taylor expansion of an algebraic function of  $t$ .
- ▶ Degree over  $\bar{\mathbb{Q}}(t)$ : 483,840.

## *Chebyshev example (cont'd)*

- ▶ Monodromy group is finite.
- ▶ Series  $u(t)$ : Taylor expansion of an algebraic function of  $t$ .
- ▶ Degree over  $\bar{\mathbb{Q}}(t)$ : 483,840.
- ▶  $\mathcal{H}(t)$ : Artin with  $\text{Gal} \leq W(E_8)$  (generically =).