

Planar Keller Maps after the Counterexample: Machine-Certified Newton-Polygon Exclusions, the Staircase Transport, and the Recalibrated Frontier

Felipe Santibañez-Leal

CAOS Research program, github.com/fsantibanezleal/CAOS_RESEARCH

2026-07-22 (working manuscript, v0.01; split from the foundational companion)

Abstract

The July 2026 counterexample settled the Jacobian conjecture in dimensions $N \geq 3$ and left $JC(2)$ as the open case. This manuscript records an exact-arithmetic program on the planar case. The Keller condition $J(P, Q) = 1$ is linear in Q for fixed P ; the resulting machine (eliminate, parametrize the consistency variety, complete, invert) closes the columns $\min \deg \leq 2$ uniformly (one closed inverse for the whole quasi-triangular class), forces perfect-power tops at bidegrees $(2, n)$ and $(3, n)$, and inverts every consistent stratum it has met. Four unconditional exclusion theorems follow from a weight-class calculus: a two-term $P = x + ax^u y^v$ is never a Keller component at any partner degree; neither is any lower-weight perturbation of it; no Keller component carries a Newton top edge anchored at the linear vertex x with interior direction; and if x is a vertex of the Newton polygon then $P = x + f(y)$ exactly (the vertex dichotomy). The residual open core is the transport of the Keller constant along the staircase of Newton vertices swallowing x : we show the completion window is block-triangular under the edge grading, exhibit the classwise transport chain as an executable elimination whose obstruction always localizes at the constant's class, and prove a window form of the corresponding fifth exclusion with cleared all-parameter certificates (monomial pairings such as $-13860 a^7$). A source-audited recalibration places the program honestly: our composite-gcd certificates replicate literature-covered territory (the verified floor is $\gcd \leq 8$, primes, $2p$, then Heitmann's $B \geq 16$ and the Guccione–Guccione–Valqui dichotomy $B = 16$ or $B > 20$), and the live frontier is the $B = 16$ normal form together with the single surviving degree pair $(72, 108)$ below 125. A new multiplication formula for x^m -anchored edges shows the $B = 16$ leading form cannot produce the Keller constant on its top edge for $m \geq 2$: that case, too, is a staircase transport. All computations are exact and reproducible; every claim is labeled machine-verified [MV], derived [D], or conjectural [C].

1 Introduction

Let $P, Q \in \mathbb{C}[x, y]$ with $J(P, Q) := P_x Q_y - P_y Q_x \in \mathbb{C}^\times$ (a *Keller pair*; each member is a *Keller component*). The two-variable Jacobian conjecture $JC(2)$ asserts every Keller pair is an automorphism pair. Following the July 2026 counterexample in dimension 3 [1], $JC(2)$ is the surviving case, and the classical planar theory (Newton polygons, approximate roots, gcd coverage) is its context. This manuscript is the planar half of a two-manuscript record: the foundational companion documents the three-dimensional aftermath (validation, the seed family, the geometry of non-properness, two-dimensional equivariant rigidity); this one documents the

planar program. The split preserves history: every statement cites its experiment record (EXP-*nnn*) in the repository, where hypotheses are declared before runs and refuted attempts are kept.

Labels: [MV] machine-verified exact identity; [D] derived analytically and certified on instances; [C] conjecture. Machine-verification is exact sympy arithmetic over $\mathbb{Q}$ or $\mathbb{Q}(i)$; no floats anywhere.

Two structural inputs from the companion are used freely. First, every $\mathbb{G}_m$ -equivariant Keller map of $\mathbb{C}^2$ is linear (equivariant rigidity, EXP-010), so nothing planar is equivariant beyond the affine group. Second, at every weight ray the leading pair of a planar Keller map is Jacobian-dependent (the tropical bridge, EXP-013), which is the entry point of all edge analyses below.

2 The planar machine

The Keller condition is bilinear in the coefficient vectors of P and Q , hence *linear in Q given P* . In the affine gauge (identity linear parts) this yields a machine per degree pair: eliminate Q (the consistency ideal in P 's coefficients), parametrize, complete linearly, and invert or fiber-test (EXP-017, EXP-019, EXP-020).

Theorem 2.1 ([MV]; uniform closure at $\min \deg \leq 2$ (EXP-021/022)). *Write $\ell = \alpha x + \beta y$, $\beta \neq 0$. In shear coordinates the Keller condition for $P = x + f(\ell)$ (any polynomial f) is a linear first-order PDE with characteristic invariant P ; its complete polynomial solution space is $Q = \ell/\beta + H(P)$, H arbitrary, and the pair inverts by one closed formula. Consequently every planar Keller map with $\min(\deg P, \deg Q) \leq 2$ is invertible, and the whole quasi-triangular class $P \sim x + f(\ell)$ is invertible at every degree.*

Theorem 2.2 ([MV]; forced tops at $(2, n)$ and $(3, n)$ (EXP-019/020/022/039)). *The $(2, 3)$ consistency ideal is the single equation $(4A_0A_2 - A_1^2)^2 = 0$ (P 's top form is a perfect square); the $(2, 4)$ elimination returns the same ideal. At $(3, 4)$, the 6×5 linear map $Q_4 \mapsto J(P_3, Q_4)$ is rank-deficient exactly when P_3 is a perfect cube: all 5×5 minors vanish on the cube parametrization, every minor lies in the Hessian-covariant ideal, and completeness follows from GL_2 -equivariance ($J(f \circ A, g \circ A) = \det(A) J(f, g) \circ A$, verified identically) plus full rank at the two non-cube orbit representatives x^2y and $xy(x+y)$. On the cube stratum, sampled completions are Keller and invert with explicit polynomial inverses (4/4 by lex Gröbner extraction). At $(3, 3)$ the consistency ideal forces the quasi-triangular alignment at the radical level.*

The degree columns close by classical coverage: $\gcd(2, n) \leq 2$ and $\gcd(3, n) \in \{1, 3\}$ are inside the verified gcd landscape of Section 5, so the machine's contribution there is the *constructive* loop (explicit consistency varieties and inverses), stated as replication. The classical degree descent is an algorithm on this machine: subtract cP^k while the tops are power-proportional, finish with Theorem 2.1; it explicitly inverted a deterministic library of composed Keller maps, and rotate-descent (rotation of linear-base tops plus de Jonquieres subtraction, EXP-034) linearizes that entire library in 0 to 6 steps. Its exact limit: mixed-corner tops stay mixed under rotation; those are the classical hard shapes, and they are where the exclusion theorems below take over.

3 Four unconditional exclusions

Throughout, P has linear part x after affine gauge. The weight-class calculus: under $w = (v, 1 - u)$, the operator $J(P, \cdot)$ shifts weight by a constant, so the Keller equation decouples into weight classes; the technique goes back to Dixmier and Joseph [11] and is standard in the polygon literature [8]; the statements below appear to be new in the stated generality (Section 6).

Theorem 3.1 ([MV/D]; the weight-class theorem (EXP-029)). *For $u \geq 2$, $v \geq 1$, $a \neq 0$, the polynomial $P = x + ax^uy^v$ is not a Keller component: no Q , of any degree, satisfies $J(P, Q) \in \mathbb{C}^\times$.*

Proof shape. P is $(v, 1 - u)$ -homogeneous; the constant is reachable only from the class of y , spanned by the ray $g_s = x^{ks}y^{ms+1}$ ($k = (u - 1)/d$, $m = v/d$, $d = \gcd(u - 1, v)$); there the operator is banded, $J(P, g_s) = (ms + 1)e_s + aB_s e_{s+d}$ with B_s a positive integer, and the chain recursion contradicts the last row of every truncation. $\square$

Theorem 3.2 ([MV/D]; lower-weight perturbations never rescue (EXP-030/031)). *With u, v, a as above and R any polynomial whose monomials have degree ≥ 2 and $(v, 1 - u)$ -weight strictly below v : $P = x + ax^uy^v + R$ is not a Keller component.*

Theorem 3.3 ([MV/D]; every x -anchored edge falls (EXP-032)). *Let $z = x^k y^m$, $k, m \geq 1$, and $E = x\varphi(z)$, $\deg \varphi = g \geq 1$. On the y -class the edge operator is multiplication: $J(E, g_s) = [(ms + 1)\varphi + kz\varphi']z^s$, and the top coefficient $(mD + kg + 1)\varphi_g$ of the class ODE kills every polynomial candidate. Hence E is never the leading form of a Keller component, and $P = E + R$ (any strictly lower-weight tail) is never a Keller component.*

Theorem 3.4 ([MV/D]; the vertex dichotomy (EXP-033)). *If the monomial x is a vertex of the Newton polygon $N(P)$ (convex hull of the support with the origin) of a Keller component with linear part x , then $P = x + f(y)$.*

Lemma 3.5 (Annihilation (EXP-036)). *For any polynomial P , $J(m, P^k) = -kL(P^{k-1}m)$ with $L = J(P, \cdot)$ (Leibniz). Any covector annihilating the image of L on a space containing the preimage therefore kills the source; the certificate covectors used throughout are left-null on the completion matrix, whose columns are exactly $L(m)$, so the tail steps of Theorems 3.2–3.4 are unconditional. The finite-window subtlety (spurious pairings for out-of-window preimages) reproduces the recorded artifact values exactly.*

The dichotomy's other side is real: $x + (x + y)^2$ and $x + (x + 2y)^3$ are components in which x is *swallowed* inside the polygon (equivalently: a pure power x^d , $d \geq 2$, appears in the support). Every non-triangular component is of that kind. The residual open core is therefore precise: components whose polygon swallows x with a mixed staircase.

4 The corner is diagonal; the staircase transport

Proposition 4.1 ([MV]; the diagonal corner (EXP-035)). *For $B = x^i y^j$, $i, j \geq 1$: $J(B^p, x^a y^b) = p(ib - ja)x^{ip+a-1}y^{jp+b-1}$, with kernel exactly the powers of B ; no monomial gives $J(B^p, \cdot)$ a constant term. The mixed corner is clean; the obstruction lives on the staircase of Newton vertices joining the swallowed linear vertex to the corner.*

Theorem 4.2 ([MV]; block triangularity and the transport chain (EXP-037)). *Fix the x -edge grading $w = (v, 1 - u)$ and let s_0 be the minimal w -class of P . In the completion window (unknown Q of bounded degree), every nonzero matrix entry sits at a class offset $s - \sigma$ for a P -class s ($\sigma = v + 1 - u$), and the offset $s_0 - \sigma$ is minimal: ordering Q -classes by w -class, the system is block triangular. Classwise elimination (solve the diagonal block, inject its kernel parameters, emit its cokernel obstructions) is therefore well-defined; it reproduces the monolithic consistency verdict on every tested sample, including a consistent positive control. On the pure above-weight staircase family the diagonal blocks are the banded operators of Theorem 3.1, and on all 8 grid samples the first inconsistent transport equation sits at the constant's class.*

Theorem 4.3 ([MV]; the fifth exclusion, window form (EXP-042)). *Let $P = x + ax^uy^v + bx^d$ ($u \geq 2, v, d \geq 1$; the minimal shape swallowing x). For every grid point $(u, v, d) \in \{2, 3\} \times \{1, 2\} \times \{2, 3\}$ and window $N = 7$, a cleared certificate covector with polynomial entries satisfies $c^T M = 0$ identically and pairs with the right-hand side to a monomial:*

$$-13860a^7, \quad -32a^4, \quad -630a^4, \quad -2730a^7, \quad -1155a^3, \quad -165a^4, \quad -4620a^7b, \quad -81b^4,$$

so the completion window is empty for all parameter values with $a \neq 0$ (the b -factors vanish only where Theorem 3.1 applies anyway). At $(u, v, d) = (2, 1, 2)$ the pairing at window N is $-c_N a^N$ with $c_N \in \{420, 1260, 13860, 180180, \dots\}$ for $N = 5..9$: a nonzero monomial in a alone at every certified window. The annihilation mechanism of Lemma 3.5 transfers verbatim (it is P -agnostic), and in-window sources pair to zero exactly as the window criterion demands. A three-parameter tail certificate $(-27720a^7c_3^4)$ extends the family.

Conjecture 4.4 ([C]; the fifth exclusion, all degrees). *The constant-class transport equation is a nonzero monomial multiple of a at every window: no above-weight perturbation of $x + ax^uy^v$ is ever a Keller component. The missing step is the closed form of the transport chain, the staircase analog of the banded formula behind Theorem 3.1.*

Two instrument-level results complete the toolkit. The *matched-pair law* (EXP-038): for cornered pairs $P = \alpha B^p + P_{\text{sub}}, Q = \beta B^q + Q_{\text{sub}}$, the second class of $J(P, Q)$ is exactly $\alpha p(ib - ja)q_{a,b} - \beta q(id - jc)p_{c,d}$ over outputs matched by $(a, b) = (c, d) + (q - p)(i, j)$; imposing the pair shape does not change the obstruction depth on any tested sample, but it halves the window unknowns: an efficiency tool, not a second obstruction source. And the *x^m -anchored edge operator* (EXP-043): for $E = x^m \varphi(z), z = x^k y^l$,

$$J\left(x^m \varphi(z), x^\alpha y^\beta\right) = x^{m+\alpha-1} y^{\beta-1} [\beta m \varphi(z) + (\beta k - \alpha l) z \varphi'(z)],$$

verified identically in symbolic edge coefficients; Theorem 3.3's operator is the case $m = 1$.

5 The frontier, recalibrated from primary sources

An audited literature pass (2026-07-22; dossier in the repository with per-claim sources and explicit UNVERIFIED flags) fixes the honest coordinates of the program.

Verified gcd coverage. For a counterexample pair, $B := \gcd(\deg P, \deg Q)$ satisfies: $B \geq 9$ is forced by the classical interval results ($B = 1$: Magnus [2]; $B \leq 2$: Nakai–Baba; $B \leq 8$ or prime: Appelgate–Onishi [3] and Nagata [4]); $B \neq p$ [7]; $B \geq 16$ (Heitmann [6]); $B \neq 2p$ and $B = 16$ or $B > 20$ [8]. Zoladek's claimed $B > 33$ has a documented gap [8] and is not used.

Consequently our machine certificates at $\gcd = 2, 9, 12, 18$ (EXP-024..028), stated in earlier revisions as frontier contact, are *replications* of literature-covered territory; we keep them as independent verifications and relabel them accordingly.

Verified degree floor. Moh’s classical analysis discards $\max \deg \leq 100$ [5], with complete published detail only for the largest case (64); the four exceptional cases correspond to six configurations in the modern corner terminology, which the algorithmic audit of Guccione–Guccione–Horruitiner–Valqui enumerates and discards [9, 10]. The current floor is: any counterexample has $\max \deg \geq 125$, or $(\deg P, \deg Q) \in \{(72, 108), (108, 72)\}$ [10]; that single pair was left open explicitly for compute reasons.

The live targets. (i) The $B = 16$ case comes with a forced direction set and leading forms $l_{4,-1}(P) = \lambda_P R_0^m$, $R_0 = x(xy^4 - \lambda_0)^3$, and a reduced normal form [8]. We verify [MV] that R_0 is $(4, -1)$ -homogeneous with collinear x -anchored support, and, by the x^m -anchored operator above, that for $m \geq 2$ *no monomial reaches the Keller constant from the R_0^m edge*: the $B = 16$ problem is a staircase transport problem, the exact object of Section 4. The pure $m = 1$ shape has x as a Newton vertex and is excluded outright by Theorem 3.4 (window replication empty); swallowed variants land in Theorem 4.3’s territory (sampled windows empty). (ii) For the $(72, 108)$ system, the classwise decomposition is block-cheap: exact histograms give 5992 window unknowns in 535 classes with largest diagonal block 22 (and 2142 unknowns, largest block 13, for the $(48, 64)$ validation case): the attack cost is coupling bookkeeping, not linear algebra, and the transport instrument automates exactly that.

6 Positioning and novelty

Per the adversarial novelty pass (same dossier): no prior statement of Theorems 3.1, 3.2, or of the companion’s equivariant rigidity theorem was found (for the latter, the C^* -equivariant analog *fails* on pseudo-planes [14], and the finite-group analog is Miyanishi [13], so the $\mathbb{C}^2$ statement has content). Theorem 3.3 was not found as stated, but its operator identity is within the standard bracket calculus [11, 8]: the statement appears new, the method is classical. Theorem 3.4’s counterexample half is essentially contained in [8] (Prop. 4.1) with van den Essen’s subrectangular normalization [12]; we claim only the sharp dichotomy form. All proofs are elementary once seen; a folklore risk remains and is recorded, with the primary-source audit trail (including its open verification items) persisted in the repository.

7 Program

Priorities, from the standing routes evaluation: (1) close Conjecture 4.4 by the transport chain’s closed form (the cleared-certificate and annihilation ingredients are already in place); (2) run the transport sweep on the $(48, 64)$ validation case and then the open $B = 16$ territory and $(72, 108)$; (3) the properness reformulation ($JC(2)$ holds iff every planar Keller map has empty Jelonek set; exact certificates shipped, EXP-014) as a cross-check instrument. The reproducibility contract: every [MV] claim has a persisted script and output; hypotheses are declared before runs; refuted predictions (three among the experiments cited here) stay in the record.

References

- [1] L. Alpöge, announcement of the counterexample, X (@__alpoge__), 2026-07-19/20; map produced with Claude Fable 5 (Anthropic); question posed by Akhil.
- [2] A. Magnus, On polynomial solutions of a differential equation, *Math. Scand.* **3** (1955), 255–260. (The $\gcd = 1$ theorem; Magnus’ formula is in *Proc. Amer. Math. Soc.* **5** (1954), 256–266.)
- [3] H. Appelgate, H. Onishi, The Jacobian conjecture in two variables, *J. Pure Appl. Algebra* **37** (1985), 215–227.
- [4] M. Nagata, Some remarks on the two-dimensional Jacobian conjecture, *Chinese J. Math.* **17** (1989), 1–7.
- [5] T.-T. Moh, On the Jacobian conjecture and the configurations of roots, *J. Reine Angew. Math.* **340** (1983), 140–212.
- [6] R. Heitmann, On the Jacobian conjecture, *J. Pure Appl. Algebra* **64** (1990), 35–72.
- [7] S. S. Abhyankar, Some thoughts on the Jacobian conjecture, Parts II–III, *J. Algebra* **319** (2008), 1154–1248; **320** (2008), 2720–2826.
- [8] J. A. Guccione, J. J. Guccione, C. Valqui, On the shape of possible counterexamples to the Jacobian conjecture, *J. Algebra* **471** (2017), 13–74; arXiv:1401.1784.
- [9] J. A. Guccione, J. J. Guccione, R. Horruitiner, C. Valqui, Some algorithms related to the Jacobian Conjecture, arXiv:1708.07936.
- [10] J. A. Guccione, J. J. Guccione, R. Horruitiner, C. Valqui, Increasing the degree of a possible counterexample to the Jacobian Conjecture from 100 to 108, arXiv:2204.14178.
- [11] A. Joseph, The Weyl algebra: semisimple and nilpotent elements, *Amer. J. Math.* **97** (1975), 597–615.
- [12] A. van den Essen, *Polynomial Automorphisms and the Jacobian Conjecture*, Progress in Mathematics 190, Birkhäuser, 2000.
- [13] M. Miyanishi, Equivariant Jacobian Conjecture in dimension two, *Transform. Groups* **28** (2023), 951–971.
- [14] A. Dubouloz, K. Palka, The Jacobian Conjecture fails for pseudo-planes, *Adv. Math.* **339** (2018), 248–284.