

Curvilinear Geometry and Primary Structure of Conductor Fiber Cones in a Huneke–Wiegand Family

Felipe Santibañez-Leal 

CAOS Research programme, github.com/fsantibanezleal/CAOS_RESEARCH

2026-08-18

Version 0.01

PREPRINT · not peer reviewed · CAOS Research programme

Companion paper in the Huneke–Wiegand extension record · version 0.01 · 2026-08-18

Version DOI: [10.5281/zenodo.21997378](https://doi.org/10.5281/zenodo.21997378) · licensed CC BY 4.0

Concept DOI: [10.5281/zenodo.21997377](https://doi.org/10.5281/zenodo.21997377)

Sources and machine records: github.com/fsantibanezleal/CAOS_RESEARCH

Provenance. Son Pham discovered the first public counterexample addressed by the wider programme, and Professor Craig Huneke independently verified it. This companion paper concerns a separate CAOS infinite family and the projective geometry of its conductor fiber cones.

Abstract

For every integer $p \geq 4$, a previously constructed symmetric numerical semigroup ring R_p and rigid two-generated monomial ideal determine a conductor ideal T_p whose special fiber $\mathcal{C}_p = \mathcal{F}(T_p)$ is one-dimensional Cohen–Macaulay, generated in degree one by $10p$ monomials, and has multiplicity $24p$. We prove that this fiber cone admits the explicit standard-graded parametrization

$$\mathcal{C}_p \cong \mathbb{k}[xy^a : a \in G_p] \subseteq \mathbb{k}[x, y]/(y^{24p}), \quad \deg(xy^a) = 1,$$

where G_p is the exact degree-one offset set. If $P_p = \mathbb{k}[X_a : a \in G_p]$ and J_p is the defining ideal, then

$$\sqrt{J_p} = L_p = (X_a : a > 0), \quad J_p \text{ is } L_p\text{-primary.}$$

Thus the complete primary decomposition has one component. The nilradical of $\mathcal{C}_p$ has sharp nilpotency index $24p$: its $(24p - 1)$ st power is nonzero, while its $24p$ th power vanishes. Dehomogenization at X_0 gives

$$\mathcal{C}_p/(X_0 - 1) \cong \mathbb{k}[y]/(y^{24p}).$$

Consequently $\text{Proj}(\mathcal{C}_p)$ is a saturated length- $24p$ curvilinear fat point with one-dimensional Zariski tangent space. It is locally Gorenstein, although its homogeneous coordinate ring has Cohen–Macaulay type $10p + 1$ and is neither level nor Gorenstein. We also compute its affine Kähler differential module, including the exact characteristic split. The proof is deductive from frozen offset-basis and Cohen–Macaulay input theorems. An exact campaign for every $p = 4, \dots, 300$ and an independently encoded all-row audit validate the implementation and adversarial controls but are not used to extrapolate the infinite statement.

1 Introduction

Fiber cones encode the number and relations of generators of powers of an ideal. Even for monomial ideals, their reducedness and projective geometry can vary sharply. Explicit defining

ideals are known for several convex, concave, symmetric, and low-generator classes [4, 5]; in particular, nonradical monomial fiber cones occur naturally. Curvilinear zero-dimensional schemes provide a complementary geometric language: their local maximal ideals are generated by one element, and their differential modules give effective signatures of this property [6].

This paper brings those two viewpoints together for the conductor fiber cones attached to an explicit infinite family in the Huneke–Wiegand extension programme. The family was constructed in [3] after Son Pham’s discovery of a first public counterexample [2]; the discovery priority and the present contribution are distinct. The same earlier work determines the conductor fiber cone, its complete degreewise offset bases, its Cohen–Macaulay type, its defining ideal, and exact edges of its graded resolution. Here we use that information from a different angle. The presence of offsets 0 and 1, combined with a hard truncation at $24p$, makes the entire projective scheme one-variable after dehomogenization.

The main result is stronger than a radical calculation. It gives a concrete truncated monomial model, a complete one-component primary decomposition, the exact nilpotency exponent, and the scheme-theoretic affine chart. It also separates two notions that are easy to conflate: the projective scheme is locally Gorenstein, while its given homogeneous embedding is not arithmetically Gorenstein.

2 The conductor family and frozen input

Fix $p \geq 4$ and put $s = 6p$ and $q = 4s = 24p$. Define residue sets in $[0, s - 1]$ by

$$\begin{aligned} A_p &= [0, p] \cup [3p, 4p - 2], \\ B_p &= ([p + 1, 3p - 1] \setminus \{2p - 1\}) \cup \{4p\} \cup [5p - 1, 6p - 1], \\ D_p &= [0, 2p] \cup [3p, 5p - 2]. \end{aligned}$$

The numerical semigroup Γ_p contains

$$\{0\}, \quad 4s + A_p, \quad [5s, 6s - 1], \quad 6s + B_p, \quad 8s + D_p, \quad [9s, 13s - 2], \quad [13s, \infty),$$

and no other nonnegative integers. The earlier family theorem proves that Γ_p is symmetric, with multiplicity $24p$ and Frobenius number $78p - 1$, and that $I_p = (t^{24p}, t^{30p})$ is a nonprincipal rigid ideal in the localization of $\mathbb{k}[t^{\Gamma_p}]$ at its positive-degree maximal ideal [3].

Let $E_p = \text{End}_{R_p}(I_p)$ and let $T_p = R_p : E_p$ be the conductor. The object studied here is its special fiber

$$\mathcal{C}_p = \mathcal{F}(T_p) = \bigoplus_{n \geq 0} T_p^n / \mathfrak{m}_p T_p^n.$$

We recall the exact input in the form needed below.

Proposition 2.1 (Frozen offset-basis input). *There is a presentation*

$$P_p = \mathbb{k}[X_a : a \in G_p] \longrightarrow \mathcal{C}_p, \quad X_a \longmapsto [t^{q+a}],$$

where $|G_p| = 10p$ and

$$\begin{aligned} G_p &= \{0\} \cup [1, p] \cup [3p, 4p - 2] \cup [6p, 8p - 2] \cup [8p, 10p - 2] \cup \{10p\} \\ &\quad \cup [11p - 1, 12p - 1] \cup [13p + 1, 14p - 2] \cup [14p, 15p - 1] \cup \{16p\} \cup [17p - 1, 18p - 1]. \end{aligned}$$

If J_p is the kernel, a degree- n monomial is zero precisely when its total offset is absent from E_n , and two nonzero monomials have the same class precisely when their total offsets agree. The cumulative bases are

$$\begin{aligned} E_1 &= G_p, \\ E_2 &= [0, 2p] \cup [3p, 5p - 2] \cup [6p, q - 1], \\ E_3 &= [0, q - 1] \setminus \{6p - 1\}, \\ E_n &= [0, q - 1] \quad (n \geq 4). \end{aligned}$$

Moreover, $\mathcal{C}_p$ is a one-dimensional Cohen–Macaulay standard graded algebra, free of rank q over $\mathbb{k}[X_0]$, and has Cohen–Macaulay type $10p + 1$. Its Artinian reduction by X_0 has socle in two degrees, so $\mathcal{C}_p$ is neither level nor Gorenstein.

Remark 2.2. Proposition 2.1 packages the previously proved conductor, fiber-cone, presentation, and offset-congruence theorems. The present paper does not infer it from the finite campaign below. Its exact all-parameter component step retains the solver/encoding boundary disclosed in [3].

3 The truncated monomial model

Assign standard degree one to each monomial xy^a and define

$$\mathcal{A}_p = \mathbb{k}[xy^a : a \in G_p] \subseteq \mathbb{k}[x, y]/(y^q).$$

Theorem 3.1 (Truncated parametrization). *For every $p \geq 4$, the homomorphism*

$$P_p \longrightarrow \mathcal{A}_p, \quad X_a \longmapsto xy^a,$$

induces a standard-graded isomorphism $\mathcal{C}_p \cong \mathcal{A}_p$.

Proof. A degree- n monomial with total offset b maps to $x^n y^b$. This is zero exactly when $b \geq q$, and distinct surviving offsets give linearly independent monomials. By Proposition 2.1, the corresponding class in $(\mathcal{C}_p)_n$ is zero precisely when the offset is absent from E_n , and otherwise depends only on the offset. The achieved surviving offsets are exactly E_n in both quotients. Hence the two kernels agree in every graded degree. $\square$

Corollary 3.2 (Exact dehomogenization). *There is an isomorphism*

$$\mathcal{C}_p/(X_0 - 1) \cong \mathbb{k}[y]/(y^q), \quad X_a \longmapsto y^a.$$

Proof. The offsets 0 and 1 belong to G_p . For every positive $a \in G_p$, the two degree- a monomials $X_0^{a-1}X_a$ and X_1^a have the same surviving offset $a < q$, so

$$X_0^{a-1}X_a = X_1^a.$$

After setting $X_0 = 1$, every generator is a power of X_1 , and $X_1^q = 0$. This gives a surjection from $\mathbb{k}[y]/(y^q)$. Freeness of rank q over $\mathbb{k}[X_0]$ shows that the target also has vector-space dimension q , hence the surjection is an isomorphism. $\square$

4 Primary decomposition and sharp nilpotence

Let

$$L_p = (X_a : a > 0) \subseteq P_p, \quad N_p = L_p/J_p \subseteq \mathcal{C}_p.$$

Theorem 4.1 (One primary component). *For every $p \geq 4$,*

$$\sqrt{J_p} = L_p, \quad J_p \text{ is } L_p\text{-primary.}$$

Thus J_p itself is the complete primary decomposition. The nilradical N_p has exact nilpotency index $q = 24p$:

$$N_p^{q-1} \neq 0, \quad N_p^q = 0.$$

Proof. The quotient $\mathcal{C}_p/N_p$ is $\mathbb{k}[X_0]$, so it is reduced. Every product of q generators with positive offset has total offset at least q and vanishes by Theorem 3.1. Hence N_p is nilpotent and equals the nilradical. It is the unique minimal prime. The element X_1^{q-1} maps to $x^{q-1}y^{q-1}$ and is nonzero, proving the sharp exponent.

The unique-minimal-prime statement alone would not prove primaryness. Here the Cohen–Macaulay hypothesis is essential: a one-dimensional Cohen–Macaulay ring has no embedded associated primes. Therefore $\text{Ass}(\mathcal{C}_p) = \{N_p\}$, so the zero ideal is N_p -primary. Taking its inverse image in P_p proves the assertion about J_p . $\square$

Remark 4.2 (Why Cohen–Macaulayness is load-bearing). The ring $\mathbb{k}[u, v]/(u^2, uv)$ has a unique minimal prime but also an embedded associated prime; its zero ideal is not primary. This is an explicit control against replacing the associated-prime argument by an invalid radical-only inference.

5 Curvilinear geometry and Gorenstein separation

Theorem 5.1 (The projective fat point). *The ideal J_p is saturated for the irrelevant ideal. The projective scheme $\text{Proj}(\mathcal{C}_p)$ is supported at $[1 : 0 : \cdots : 0]$ and is a length- q curvilinear fat point. Its Zariski tangent space has dimension one. The scheme is locally Gorenstein, while the homogeneous coordinate ring $\mathcal{C}_p$ is not level and not Gorenstein.*

Proof. Because $\mathcal{C}_p$ is one-dimensional Cohen–Macaulay, its zeroth local cohomology for the irrelevant ideal vanishes. The identity $J_p^{\text{sat}}/J_p = H_{P_{p,+}}^0(\mathcal{C}_p)$ gives saturation. Theorem 4.1 shows that the support is the single coordinate point and hence lies entirely in $D_+(X_0)$. Corollary 3.2 identifies its affine coordinate ring with $\mathbb{k}[y]/(y^q)$. This has length q and principal maximal ideal (y) , so the scheme is curvilinear and its tangent space $(y)/(y^2)$ is one-dimensional. Its socle is generated by y^{q-1} , proving local Gorensteinness. Proposition 2.1 gives type $10p+1$ and nonlevelness for the homogeneous coordinate ring, proving the arithmetic contrast. $\square$

Remark 5.2. This example makes the embedding distinction concrete:

$$\text{locally Gorenstein} \not\Rightarrow \text{arithmetically Gorenstein.}$$

The local scheme is always the one-variable truncation, while the standard grading retains the nonlevel offset-layer structure.

6 Kähler differentials

The curvilinear chart has a short differential fingerprint which also records the ground-field characteristic.

Proposition 6.1 (Differential signature). *Let $B_p = \mathcal{C}_p/(X_0 - 1)$. Then*

$$\Omega_{B_p/\mathbb{k}}^1 \cong (B_p/(qy^{q-1})) dy, \quad \bigwedge^m \Omega_{B_p/\mathbb{k}}^1 = 0 \quad (m \geq 2).$$

Consequently,

$$\dim_{\mathbb{k}} \Omega_{B_p/\mathbb{k}}^1 = \begin{cases} q-1, & \text{char}(\mathbb{k}) \nmid q, \\ q, & \text{char}(\mathbb{k}) \mid q. \end{cases}$$

The cotangent space at the closed point has dimension one in every characteristic.

Proof. Apply the conormal sequence to $B_p = \mathbb{k}[y]/(y^q)$. Differentiating the defining equation gives $qy^{q-1}dy = 0$, which proves the presentation and the characteristic split. The differential module is cyclic, so every exterior power of degree at least two vanishes. Tensoring with $B_p/(y)$ leaves the single generator dy . $\square$

This agrees with the general differential characterization of weakly curvilinear schemes in [6], while the exact truncation and characteristic formula here come from the conductor-family parametrization.

7 Exact verification record

The symbolic proofs above use the frozen results in Proposition 2.1. Computation is used to validate their translation into the new parametrization and to exercise negative controls. The campaign computes truncated sumsets directly from G_p for every $p = 4, \dots, 300$. The audit is separately encoded: it starts from the five disjoint Artinian layers, rebuilds cumulative bases, rehashes all 297 rows, and performs detailed reconstructions at $p = 4, 5, 17, 73, 151, 300$.

record	coverage	SHA-256 aggregate
campaign	all $p = 4, \dots, 300$	f3373f4f58287fd3...f9d6b39b6
independent audit	all rows; six detailed rebuilds	84c00be8ff64002...f5b6c8f143e

The controls delete offset 1, perturb the truncation and nilpotency exponents, corrupt the degree-three hole, omit a radical coordinate, conflate local and arithmetic Gorensteinness, erase the differential characteristic split, and remove the no-embedded-primes premise. Every mutation is rejected. Exact scripts, artifacts, premise hashes, proof, and verdict are archived as `EXP-025-curvilinear-primary-structure` in the public repository.

The finite sweep does not prove the theorem for all p . The infinite conclusion is supplied by Theorems 3.1–5.1 and Proposition 6.1. A defect in a frozen input theorem would propagate even if both implementations agreed.

8 Scope and further questions

The result concerns the conductor fiber cones of the explicit family recalled in Section 2. It does not assert that arbitrary monomial fiber cones are primary, curvilinear, Cohen–Macaulay, or locally Gorenstein. It does not strengthen the underlying Huneke–Wiegand counterexample theorem for arbitrary modules or arbitrary one-dimensional Gorenstein domains.

The primary decomposition is now complete, but two algebraic questions remain open for this family:

1. Is there a uniform explicit Gröbner basis refining the known minimal set of $50p^2 - 17p$ quadrics and one cubic?
2. Can the unresolved interior of the graded Betti table be expressed by closed formulas, or does it exhibit parameter-dependent combinatorial phases?

The truncated parametrization may help with both questions by replacing value-semigroup products with the combinatorics of monomials below one nilpotent boundary.

Acknowledgments

The author thanks Son Pham for making the first counterexample and its verification materials public, and Craig Huneke for the independent verification recorded with that release. Neither is an author of this paper, and this acknowledgment does not imply endorsement of the present theorem. All authorship and accountability for this manuscript remain with the named human author.

References

- [1] C. Huneke and R. Wiegand, *Tensor products of modules and the rigidity of Tor*, Math. Ann. 299 (1994), 449–476. doi:10.1007/BF01459793.
- [2] S. Pham, *Huneke–Wiegand candidate verification*, public software and certificate repository (2026). <https://github.com/sonpham-org/huneke-wiegand-candidate-verification>.
- [3] F. Santibañez-Leal, *Frobenius minimality, minimum-layer uniqueness, and an infinite family of Huneke–Wiegand counterexamples*, CAOS Research preprint, version 0.13 (2026). doi:10.5281/zenodo.21995498.
- [4] J. Herzog, A. A. Qureshi, and M. Mohammadi Saem, *The fiber cone of a monomial ideal in two variables*, arXiv:1711.08775 (2017). <https://arxiv.org/abs/1711.08775>.
- [5] J. Herzog and G. Zhu, *On the fiber cone of monomial ideals*, arXiv:1904.04988 (2019). <https://arxiv.org/abs/1904.04988>.
- [6] M. Kreuzer, T. N. K. Linh, and L. N. Long, *Differential theory of zero-dimensional schemes*, arXiv:2302.11903 (2023). <https://arxiv.org/abs/2302.11903>.