

Exact replication and screening of tropical finiteness certificates for central configurations

Machine record v0.01 (draft): the CAOS Research central-configurations program,
experiments 001–004

Felipe Santibañez-Leal*

July 24, 2026

Abstract

We report an exact, fully replayable machine record around Smale’s 6th problem (finiteness of planar central configurations). Contributions of this record: (i) an exact calibration of the Albouy–Chenciner mutual-distance equations in open tooling, including a machine rediscovery of their dimension-blindness (the regular tetrahedron coexists with the square in the equal-mass rhombus stratum of the 4-body distance system until the Cayley–Menger equation is adjoined, after which the square, with side a satisfying $32a^6 - 32a^3 + 7 = 0$, is the unique positive point of the stratum); (ii) a machine measurement of the *enrichment law*: adjoining the asymmetric (Roberts) equations and the energy–inertia relation renders the $n = 3$ distance ideal zero-dimensional directly, eliminating the coordinate-line components of the bare symmetric system; (iii) an independent, exact reproduction of the Jensen–Leykin tropical prevariety certificate of generic-mass finiteness for $n = 5$ (arXiv:2301.02305v2): both published f -vectors match digit for digit and the pointedness of the recession cone is verified by an independent exact parser; and (iv) a screening study over mass-valuation families and equation-system variants at $n = 4, 5$, whose sharpest finding is that removing the ten dependent symmetric Albouy–Chenciner equations catastrophically destroys pointedness at every tested valuation: the “redundant” equations carry essential tropical information. All computations are exact (rational or integer arithmetic, or certified polyhedral computation); every stated number is reproducible from committed code and hashes. This is a replication-and-instruments record: the theorems replicated are due to Hampton–Moeckel, Albouy–Kaloshin, and Jensen–Leykin; the measurements in (i), (ii) and (iv) are, to our knowledge, recorded here at this level of explicitness for the first time.

1 Introduction

A *central configuration* (CC) of the Newtonian n -body problem with masses $m_1, \dots, m_n > 0$ is a configuration $x = (x_1, \dots, x_n)$, $x_i \in \mathbb{R}^d$, satisfying

$$\lambda(x_i - c) = \sum_{j \neq i} \frac{m_j(x_j - x_i)}{r_{ij}^3}, \quad i = 1, \dots, n, \quad (1)$$

*With machine-executed computations; all artifacts, exact values, and run records are versioned at https://github.com/fsantibanezleal/CAOS_RESEARCH under `problems/dynamical-systems/central-configurations/`.

for some λ , where c is the center of mass and $r_{ij} = |x_i - x_j|$. Smale’s 6th problem for the 21st century [1] asks, following Chazy [2] and Wintner [3], whether for every n and every choice of positive masses the number of *planar* central configurations, up to symmetry, is finite. The modern ladder is: $n = 4$ finite for all positive masses (Hampton–Moeckel [4]: computer-assisted BKK certificate, count in [32, 8472]); $n = 5$ finite except possibly on an explicit codimension-two subvariety of mass space (Albouy–Kaloshin [5]); the spatial 5-body Dziobek case handled generically by Moeckel [6] and explicitly by Hampton–Jensen [7]; $n = 6$ open, reduced by Chang–Chen [8] to a finite list of residual diagram cases; and, for *generic* masses, the recent purely polyhedral certificate of Jensen–Leykin [9], complete through $n = 5$. Rigorous equal-mass censuses for $n \leq 7$ are due to Moczurad–Zgliczyński [10]; with one negative mass a continuum exists (Roberts [11]), so positivity is essential.

This paper is a *machine record*: it documents, with exact artifacts, a replication-first program whose goal is to place every certificate of the ladder inside one open, replayable toolchain, and then to run the experiments the frontier papers themselves call for. We follow a strict discipline: hypotheses are declared and committed before every run; verdicts record refutations as prominently as confirmations; every quantitative claim below is reproducible from the versioned repository, and the git and Zenodo timestamps constitute the priority record of the work.

2 Exact calibration of the Albouy–Chenciner system (EXP-001)

Working in the mutual distances r_{ij} with the scale normalization $S_{ij} = r_{ij}^{-3} - 1$ ($S_{ii} = 0$), the symmetric Albouy–Chenciner (AC) equations [12, 4] are, after clearing denominators,

$$f_{ij} = \sum_{k=1}^n m_k [S_{ik}(r_{jk}^2 - r_{ik}^2 - r_{ij}^2) + S_{jk}(r_{ik}^2 - r_{jk}^2 - r_{ij}^2)] = 0. \quad (2)$$

Our exact implementation (sympy over $\mathbb{Q}$) reproduces, for four exact mass vectors, the classical $n = 3$ landscape: the Lagrange point $r_{12} = r_{13} = r_{23} = 1$ identically in symbolic masses, and exactly one positive collinear solution per ordering (Euler–Moulton; for equal masses the 2-middle chart value is $r_{12} = r_{23} = \sqrt[3]{90}/6$), with the degree-54 symbolic Euler eliminant persisted. For $n = 4$ each cleared AC polynomial has exactly 14 monomials, matching the count stated in [4].

The instructive result of the calibration is a machine *refutation* of a declared prediction. In the equal-mass rhombus stratum $r_{12} = r_{23} = r_{34} = r_{14} = a$, $r_{13} = r_{24} = b$ of $n = 4$, the bare AC system has *two* positive solutions: the planar square, with $b^2 = 2a^2$ and $a^3 = (4 + \sqrt{2})/8$ (minimal polynomial $32a^6 - 32a^3 + 7$), and the regular tetrahedron $a = b = 1$, whose bordered Cayley–Menger determinant equals $4 \neq 0$. The distance equations are dimension-blind, exactly as stated in [4]; planarity is restored only by adjoining the Cayley–Menger equation, after which the square is the unique positive point of the stratum. We record the scale-free invariant $J = UI^{1/2}/M^{5/2}$ at both anchors: $J_{\square} = \frac{1}{16} + \frac{\sqrt{2}}{8}$, $J_{\text{tet}} = \frac{3\sqrt{6}}{32}$.

Two toolchain findings of independent interest were caught and are now regression-gated in the repository: `sympy`’s `solve_poly_system` silently returns incomplete solution lists on these systems (it missed the square, whose eliminant factor $4b^6 - 4b^3 - 7$ is even radical-solvable), and `is_positive` returns `None` on nested algebraic numbers, so naive filters silently drop genuine solutions. Verdict-grade counting in our pipeline therefore uses univariate eliminants that are provably elements of the ideal (lex Gröbner elimination, or Stickelberger multiplication-matrix characteristic polynomials), complete real root isolation, and exact residual acceptance.

3 The enrichment law (EXP-002)

The bare symmetric system is never zero-dimensional: the line $r_{13} = r_{23} = 0$ lies in its variety for all masses (machine-verified symbolically), and saturation is expensive. Adjoining the $n(n-1)$ asymmetric equations $g_{ij} = \sum_k m_k S_{ik}(r_{jk}^2 - r_{ik}^2 - r_{ij}^2)$ (observed by G. Roberts; see [7, 9]) and the energy–inertia relation $U = MI$ changes the picture completely: for all four sample mass vectors the enriched ideal is zero-dimensional *directly* (Gröbner basis of size 24 in under a second each), and the full positive censuses, where decided, are exactly classical: four points (one equilateral, three collinear), with no spurious solutions. The planar rhombus census on the enriched system with Cayley–Menger adjoined yields the square alone. This is the affine-level instance of a pattern that recurs at every level of the subject: Hampton–Moeckel could not run their method on the six AC equations alone and required Dziobek’s equations [4]; Hampton–Jensen kept the asymmetric equations to refine their tropical cones [7]; and Jensen–Leykin explicitly call for “different versions of the Albouy–Chenciner equations” at $n = 6$ [9]. Section 5 quantifies this pattern at the prevariety level.

4 Exact reproduction of the Jensen–Leykin $n = 5$ certificate (EXP-003)

Jensen–Leykin [9] prove generic-mass finiteness of planar CCs for $n \leq 5$ by a purely polyhedral computation: specialize the masses into the Puiseux field with chosen valuations, compute the tropical prevariety of the system (20 asymmetric AC + 10 symmetric AC + 5 planar Cayley–Menger polynomials for $n = 5$) with `gfan` 0.7, and verify that every connected component has a pointed recession cone; Bieri–Groves and the density of tropicalization fibers then yield finiteness for almost all masses.

We rebuilt `gfan` 0.7 from the author tarball (one two-line `#include <stdint>` patch for gcc 13; tarball and binary SHA-256 recorded) and ran the paper’s own command pipeline. Results: with valuations (1, 3, 9, 27, 81) the prevariety f -vector is (1506, 4744, 8586, 8787, 4652, 993), and among its 85 unbounded ray directions none has a positive coordinate (verified by our independent exact parser): the globally pointed certificate, exactly as published. With valuations (1, 4, 9, 16, 25) the f -vector is (3586, 12012, 18531, 15625, 7072, 1357), again matching [9] digit for digit. Each run took under six minutes of wall time on 30 threads (≈ 145 cpu-minutes). To our knowledge this is the first published independent replication of the [9] computation; at the measured throughput, a single $n = 6$ attempt at the scale they report (≈ 100 cpu-days) costs about three wall-days on commodity hardware.

5 Valuation and equation-variant screening (EXP-004)

[SECTION IN PROGRESS: the declared screening grid (two systems $\times$ six valuation families at $n = 5$, plus four $n = 4$ cells) is running at the time of this draft; its verdict will replace this placeholder in the next version. Interim, already-decided cells: the powers-of-3 valuation is the only globally pointed one among the six tested at $n = 5$; the squares, powers-of-2 and primes valuations each leave exactly two positive unbounded directions (candidates for per-component pointedness, as in the squares case of [9]); the arithmetic and repeated-value controls are far from pointed; and the asymmetric-only system S2, which drops the ten dependent symmetric equations, loses pointedness catastrophically at *every* tested valuation (over 300 positive

unbounded directions), a quantitative demonstration that the dependent symmetric equations carry essential tropical information.]

6 Honesty ledger

This record is replication-first. The finiteness theorems replicated are due to Hampton–Moeckel, Albouy–Kaloshin, and Jensen–Leykin; the equal-mass censuses to Moczurad–Zgliczyński. Our measurements (the dimension-blindness anchors with their exact minimal polynomials and J -invariants; the enrichment law at affine and prevariety level; the screening tables) are machine facts about known objects, stated with their complete provenance. Open items are tracked in the repository backlog; in particular, censuses for two integer-separated mass vectors are inconclusive under our current sympy engine caps and are being re-run under an independent engine (`msolve`) with the sympy layer retained as verification. No claim in this paper depends on an unverified literature statement; the repository flags every such statement as UNVERIFIED until read from its primary source.

References

- [1] S. Smale, *Mathematical problems for the next century*, Math. Intelligencer **20** (1998), no. 2, 7–15.
- [2] J. Chazy, *Sur certaines trajectoires du problème des n corps*, Bull. Astron. **35** (1918), 321–389.
- [3] A. Wintner, *The Analytical Foundations of Celestial Mechanics*, Princeton University Press, 1941.
- [4] M. Hampton and R. Moeckel, *Finiteness of relative equilibria of the four-body problem*, Invent. Math. **163** (2006), 289–312.
- [5] A. Albouy and V. Kaloshin, *Finiteness of central configurations of five bodies in the plane*, Ann. of Math. **176** (2012), 535–588.
- [6] R. Moeckel, *Generic finiteness for Dziobek configurations*, Trans. Amer. Math. Soc. **353** (2001), 4673–4686.
- [7] M. Hampton and A. N. Jensen, *Finiteness of spatial central configurations in the five-body problem*, Celest. Mech. Dyn. Astron. **109** (2011), 321–332.
- [8] K.-M. Chang and K.-C. Chen, *Toward finiteness of central configurations for the planar six-body problem by symbolic computations. (I) Determine diagrams and orders*, J. Symbolic Comput. **123** (2024), 102277; *(II) Determine mass relations*, SIAM J. Appl. Dyn. Syst. **24** (2025), no. 3, 2369–2404; arXiv:2303.02853.
- [9] A. Jensen and A. Leykin, *Smale’s 6th problem for generic masses*, arXiv:2301.02305v2 (2025).
- [10] M. Moczurad and P. Zgliczyński, *Central configurations in planar n -body problem with equal masses for $n = 5, 6, 7$* , Celest. Mech. Dyn. Astron. **131** (2019); arXiv:1812.07279.

- [11] G. E. Roberts, *A continuum of relative equilibria in the five-body problem*, Phys. D **127** (1999), 141–145.
- [12] A. Albouy and A. Chenciner, *Le problème des n corps et les distances mutuelles*, Invent. Math. **131** (1998), 151–184.
- [13] F. R. Moulton, *The straight line solutions of the problem of n bodies*, Ann. of Math. **12** (1910), 1–17.