

Frobenius Minimality, Minimum-Layer Uniqueness, and an Infinite Family of Huneke–Wiegand Counterexamples

Felipe Santibañez-Leal 

CAOS Research program, github.com/fsantibanezleal/CAOS_RESEARCH

2026-08-30

Version 0.22

PREPRINT · not peer reviewed · CAOS Research programme

First paper of the Huneke–Wiegand extension record · version 0.22 · 2026-08-30

Concept DOI [10.5281/zenodo.21763582](https://doi.org/10.5281/zenodo.21763582) (resolves to latest) · version DOI reserved after
prepublication QA

Licensed CC BY 4.0 · sources and machine records:

github.com/fsantibanezleal/CAOS_RESEARCH

Provenance. Son Pham discovered the counterexample; Professor Craig Huneke independently verified it. This paper contributes certified Frobenius minimality, minimum-layer uniqueness, a separate explicit infinite family, and its uniform endomorphism, type, trace, conductor, stability, reduction, tangent-cone, fiber-cone, defining-ideal, syzygy, and characteristic-dependence anatomy.

Abstract

Son Pham publicly identified a counterexample to the Huneke–Wiegand conjecture in the class of two-generated monomial ideals over symmetric numerical semigroup rings, subsequently verified by Professor Craig Huneke. We preserve that discovery priority and answer two different questions: how small can such an example be, and does the same semigroup/ideal class contain an explicit infinite family? We prove by proof-carrying exact computation that the least Frobenius number is 181, attained by Pham’s semigroup at ideal shift 14. First, an implementation of the complete Blanco–Rosales tree exhausts all 48,954 symmetric semigroups with Frobenius number below 69 and all 1,503,391 associated gap cases. An independent Boolean route checks 1,156 fixed-pair UNSAT certificates for the same range. We then introduce a one-hot shift-selector CNF that asks one existence question per Frobenius number. Its DRAT certificates exclude every odd value from 69 through 179, while the formula at 181 is satisfiable and decodes exactly to the public semigroup and shift. A separately generated fixed-pair formula returns the same model. An independent audit rehashes all 228 files in the minimality search, totaling 760,081,739 bytes, rechecks both SAT models, and reproduces the strict-order manifest with no mismatch. Complete theorem-tree enumeration at $F = 69, 71, 73, 75$ provides an object-level cross-check on another 79,790 semigroups and 2,933,163 gap cases. A second projected enumeration proves that shift 14 is the only feasible shift at $F = 181$ and that the public membership vector is the only semigroup there. Both terminal DRAT proofs are freshly rechecked in an independent 12-file audit. Thus Pham’s pair is the unique normalized pair attaining the Frobenius minimum in this class. Finally, for every integer $p \geq 4$ we construct a symmetric numerical semigroup Γ_p of multiplicity $24p$, Frobenius number $78p - 1$, conductor $78p$, and embedding dimension $11p$, for which $I_p = (t^{24p}, t^{30p})$ is nonprincipal and rigid. This family theorem is deductive: seven exact interval-sum identities prove closure, symmetry, and $D_s = E_s + E_s$; a finite sweep through $p = 300$ and an

independently implemented audit are supporting checks. For every family member we also determine the value semigroup of the endomorphism ring exactly. It has multiplicity $24p$, Frobenius number $54p - 1$, conductor $54p$, genus $38p - 1$, and embedding dimension $12p$. It is nonsymmetric, and the endomorphism ring is strictly larger than the original ring. Recent endomorphism-center criteria then show uniformly that the ideal remains rigid over its endomorphism ring but is not reflexive there, with the adjacent Ext and Tor obstruction groups nonzero. We further prove that this endomorphism semigroup has Cohen–Macaulay type and reduced type $10p$, hence maximal reduced type, and that its completed semigroup ring is not almost Gorenstein. Finally, the traces of the ideal and its endomorphism ring both equal the conductor of the finite birational extension. We determine that common value ideal exactly and prove the identity

$$\ell_{R_p}(R_p/(R_p : E_p)) = \ell_{R_p}(E_p/R_p) = p + 1.$$

The equality of the two lengths is a general consequence of one-dimensional Gorenstein local duality, while the exact common ideal and its value $p + 1$ are the family-specific data. We further prove that this conductor is nonstable and compute its complete one-step stability defect:

$$\ell_{R_p}(T_p^2/t^{4s}T_p) = 14p.$$

The full reduction sequence is also exact: $t^{4s}R_p$ is a minimal reduction of T_p with reduction number four, successive Sally-quotient lengths $23p - 1, 14p, 2p, 1, 0$, and Hilbert–Samuel coefficients $e_0(T_p) = 24p$ and $e_1(T_p) = 39p$. The associated graded ring of T_p nevertheless has depth zero: its complete Valabrega–Valla module is concentrated in one degree, where it has length p . We compute the exact Hilbert series and show that its numerator has only positive coefficients despite this failure of Cohen–Macaulayness. We then determine the complete zeroth local cohomology: it is k^p , concentrated in degree zero and annihilated by the full homogeneous maximal ideal. Hence the tangent cone is Buchsbaum but not Cohen–Macaulay, with unbounded Buchsbaum invariant p ; its quotient by the finite-length torsion is Cohen–Macaulay with an exact Hilbert series. Over the polynomial Noether normalization induced by the minimal reduction, we determine the entire graded module: the torsion consists of p exponent-one cyclic summands, while the free quotient has rank $24p$ and explicit shifts. This yields the complete minimal resolution, projective dimension one, regularity four, top-local-cohomology a -invariant three, and the exact parameter-section identity $\ell(G_p/x_p G_p) = 25p = e_0(T_p) + I(G_p)$. Finally, the exact identity $T_p^2 = \mathfrak{m}_p T_p$ identifies the special fiber canonically with $G_p/H^0(G_p)$. It is Cohen–Macaulay of type $10p + 1$, but its Artinian socle lies in degrees two and four, so it is neither level nor Gorenstein. We then determine its defining ideal: in the polynomial ring on its $10p$ minimal degree-one generators, it has exactly $50p^2 - 17p$ minimal quadrics and the single additional cubic $X_0^2 X_{3p} - X_p^3$, with no other minimal equation. Thus its relation type is three and it is not Koszul. The all-parameter component calculation is an exact Presburger verification, supported by a separately encoded graph audit; its solver trust boundary is stated explicitly. From the same exact h-vector, Artinian socle, and first Betti row, we then determine the homological edges over the full presentation ring. Its projective dimension is $10p - 1$, its regularity is four,

$$\beta_{2,3} = \frac{2p(500p^2 - 330p + 31)}{3},$$

the complete last row has ranks $10p$ in shift $c + 2$ and one in shift $c + 4$ for $c = 10p - 1$, and $\beta_{c-1, c+3} = 8p$. The canonical module has $10p$ minimal generators in degree -1 and one in degree -3 . We then determine the first previously unknown interior strand. With the offset grading $\deg X_a = (1, a)$,

$$\beta_{2,4} = 8p, \quad \beta_{3,4} = \frac{p(5p - 1)(500p^2 - 440p + 47)}{2},$$

and the $8p$ second syzygies occur with multiplicity one exactly at offsets $\{3p + a : a \in \mathcal{E}_{p,1}, a \geq 6p\}$. Relative squarefree-divisor chains and an integral unit matching give the upper bound;

an exact quadratic colon and minimal mapping cone give the matching lower bound. We then complete the second Betti row. The only remaining possible entries are

$$\beta_{2,5} = p(2p - 3), \quad \beta_{2,6} = 0.$$

The degree-five support is the union of three explicit intervals, with pair-sum multiplicities on the outer blocks and a plateau profile on the middle block. An integral offset matching gives a unit Smith form, so the result again holds over every field. This completes one homological row, not the remaining Betti table. We next determine the degree-five third syzygies. If $H_p = \{a \in \mathcal{E}_{p,1} : a \geq 6p\}$, then

$$\beta_{3,(5,b)} = \#\{\{a, c\} \subset H_p : a < c, a + c = b - 3p\}, \quad \beta_{3,5} = 4p(8p - 1).$$

The support is $[15p + 1, 39p - 3] \setminus \{33p - 1\}$. An integral normal form leaves a primitive free basis indexed by unordered pairs of the $8p$ high cubic-colon variables. Combining this with the complete second row and the Hilbert numerator gives

$$\beta_{4,5} = \frac{2p(5p - 1)(10p - 3)(100p^2 - 110p + 13)}{3},$$

completing internal degree five over every field. Finally, we identify the complete cubic-colon quotient with the canonical idealization of the p -th Veronese rational normal curve ring. Its multigraded Hilbert numerator and an integral relative normal form give

$$\beta_{3,6} = \frac{8p(7p^2 - 12p + 2)}{3}$$

over every field, with support $[3p + 4, 29p - 5] \setminus ([6p - 3, 6p + 1] \cup [9p - 3, 9p])$. In total degree seven, a zero-vertex Boolean matching leaves only triangles at the single residual hole $6p - 1$. Adjoining a missing vertex from $\{1, 2, 3, 4\}$ gives each one a same-offset tetrahedron whose reduced boundary has that triangle as its unique critical face with coefficient ± 1 . Hence $\beta_{3,7} = 0$ integrally, and the complete third row over every field is

$$\begin{aligned} \beta_{3,4} &= \frac{p(5p - 1)(500p^2 - 440p + 47)}{2}, & \beta_{3,5} &= 4p(8p - 1), \\ \beta_{3,6} &= \frac{8p(7p^2 - 12p + 2)}{3}, & \beta_{3,7} &= 0. \end{aligned}$$

with no other shifts. Its total rank is $\beta_3 = p(7500p^3 - 7988p^2 + 2025p - 133)/6$. The same canonical idealization determines much more about the cubic-colon quotient $D_p = P_p/(\mathfrak{q}_p : f_p)$. Put $c = 2p - 2$, $m = 8p$, and $\lambda_{c,a} = c \binom{c}{a} - \binom{c}{a+1} - \binom{c}{a-1}$. Its complete graded Betti polynomial is

$$B_{D_p}^{P_p}(x, z) = (1 + xz)^m \left(1 + \sum_{a=1}^{c-1} \lambda_{c,a} x^a z^{a+1} + x^c z^{c+2} \right).$$

Hence every free-module rank and shift is known over every field, with projective dimension $10p - 2$ and regularity two. A colon-intersection pullback then proves that the quadratic quotient $A_p = P_p/\mathfrak{q}_p$ has depth one and regularity two. The grading gap makes the cubic mapping cone minimal, so

$$B_{C_p}^{P_p}(x, z) = B_{A_p}^{P_p}(x, z) + xz^3 B_{D_p}^{P_p}(x, z).$$

Consequently the complete regularity-three and regularity-four strands of the special fiber are explicit binomial convolutions of the displayed colon polynomial, over every field. This removes every comparison-rank ambiguity from the cubic cone. Finally, put $\tau_p = 8p - 1 + p(p + 1)/2$. The Artinian reduction of the high-variable kernel is a two-layer incidence module. Its first

missing offset gives a unique primitive cokernel cell, and a separate unit pivot proves that the connecting map from the colon quotient cannot hit it. Thus, over every field,

$$\beta_{p,(p+2,\tau_p)}(K_p) = \beta_{p,(p+2,\tau_p)}(A_p) = \beta_{p,(p+2,\tau_p)}(C_p) = 1.$$

We then classify every primitive zero row of the incidence cokernel. If $R_b = \{g : b - g \in H_p\}$, the coordinate $e_F \otimes v_b$ is a zero row exactly when $R_b \subseteq F$, giving a closed free-summand count in every homological degree. The block $b = 10p + t$, $2 \leq t \leq p - 2$, supplies consecutive kernel classes in degrees $p + 1, \dots, 2p - 3$. The first connecting cell does not repeat the unit-pivot mechanism, but its complete target quotient exposes characteristic dependence:

$$\beta_{5,(7,87)}(A_4) = \beta_{5,(7,87)}(C_4) = \begin{cases} 4, & \mathbb{k} = \mathbf{F}_2, \\ 3, & \mathbb{k} = \mathbf{F}_3. \end{cases}$$

The corresponding integral kernel cokernel is $\mathbb{Z}^4 \oplus \mathbb{Z}/2\mathbb{Z}$. This determines exact lower-strand points and a characteristic-dependent multigraded cell, not the two complete lower strands, the differential matrices, or the full special-fiber resolution. The results remain confined to two-generated monomial ideals over symmetric numerical semigroup rings and do not classify arbitrary Gorenstein domains and modules.

1 Introduction

Let R be a one-dimensional Gorenstein local domain and let M be a nonzero finitely generated non-free module. The Huneke–Wiegand conjecture predicted nontrivial torsion in $M \otimes_R \operatorname{Hom}_R(M, R)$ [1]. For a numerical semigroup $\Gamma \subseteq \mathbb{N}$, the semigroup ring $\mathbb{k}[t^\Gamma]$ is Gorenstein exactly when Γ is symmetric. This specialization converts part of the module problem into finite additive combinatorics. García-Sánchez and Leamer proved broad positive classes and reported a NumericalSgps check for every symmetric semigroup with Frobenius number below 69 [2].

In 2026, Son Pham published a symmetric numerical semigroup and a two-generated monomial ideal for which the predicted torsion is absent [5]. Professor Craig Huneke supplied an independent verification archived with the public candidate record. That example has Frobenius number 181. The discovery settles the original conjecture negatively, but does not by itself say whether 181 is small, minimal, or typical.

This paper reports computer-assisted classification theorems and a deductive family theorem. Claims are labeled [MV] when they are verified by exact machine computation and independently checked artifacts, and [D] when derived in the text. No conjectural claim is needed for the main theorem.

Contributions.

1. We give a self-contained finite-window reduction and a direct proof that the selector CNF is equisatisfiable with existence of a rigid two-generated monomial ideal.
2. We reproduce the published $F < 69$ boundary by two independent complete routes: the Blanco–Rosales tree and checked DRAT proofs.
3. We certify UNSAT for every odd $F = 69, 71, \dots, 179$, then recover the exact public model at $(F, s) = (181, 14)$.
4. We independently rehash and inspect the complete search artifact set, and cross-check the first four new odd Frobenius values by explicit theorem-tree enumeration.

5. We classify the complete minimum layer by projected enumeration: the only feasible shift at $F = 181$ is 14, and the public semigroup is the only membership vector at that shift.
6. We identify two structural boundaries: the public semigroup lies outside the generalized arithmetic-sequence positive family, and the visible fixed-offset generator blocks do not extend to a naive parametric family.
7. We use the failed fixed-width ansatz to derive and prove an explicit infinite family for every integer $p \geq 4$, with a separate exact implementation and independent audit.
8. We compute the endomorphism overring of every family member and prove that the nonreflexive Ext/Tor escape identified in the public seed persists uniformly.
9. We compute the complete pseudo-Frobenius set of every endomorphism semigroup, proving unbounded maximal reduced type and a uniform non-almost-Gorenstein boundary.
10. We compute both trace ideals and the conductor exactly, obtaining a common ideal whose colength equals the birational extension length, namely $p + 1$.
11. We separate that general duality balance from the family-specific calculation, prove that the common conductor is nonstable, and compute its exact stability defect $14p$.
12. We determine every later power of the conductor, proving reduction number four, the full Sally-quotient profile, and Hilbert–Samuel coefficients $(e_0, e_1) = (24p, 39p)$.
13. We prove that the conductor tangent cone has depth zero, determine its unique Valabrega–Valla defect of length p , and compute its coefficientwise-positive Hilbert numerator exactly.
14. We identify that defect with the complete zeroth local cohomology, prove that the full homogeneous maximal ideal annihilates it, and obtain a Buchsbaum non-Cohen–Macaulay family with unbounded Buchsbaum invariant p and an exact Cohen–Macaulay quotient series.
15. We determine the complete graded module over the minimal-reduction Noether normalization, its minimal resolution and Betti numbers, regularity four, a -invariant three, and the exact parameter-section identity $\ell(G_p/x_p G_p) = 25p = e_0(T_p) + I(G_p)$.
16. We prove $T_p^2 = \mathfrak{m}_p T_p$, identify the conductor special fiber naturally with $G_p/H^0(G_p)$, and compute its type $10p + 1$ and nonlevel Artinian socle.
17. We determine the complete defining ideal of that special fiber: $50p^2 - 17p$ minimal quadrics and one primitive cubic, giving relation type three and a non-Koszul family.
18. We determine exact homological edges over its full presentation ring: projective dimension and regularity, the linear first-syzygy count, the complete last row, one penultimate extremal entry, and the canonical-module generator degrees.
19. We determine the first interior Betti strand over every field: $\beta_{2,4} = 8p$ with complete multiplicity-free offset support, and the resulting exact formula for $\beta_{3,4}$.
20. We complete the second Betti row over every field: $\beta_{2,5} = p(2p-3)$ with complete three-block offset multiplicities and $\beta_{2,6} = 0$.

21. We determine the complete degree-five third-syzygy profile from the high cubic colon, prove $\beta_{3,5} = 4p(8p - 1)$ over every field, and complete the internal-degree-five diagonal.
22. We identify the complete cubic-colon quotient as a canonical idealization and determine the complete degree-six third-syzygy profile, including $\beta_{3,6} = 8p(7p^2 - 12p + 2)/3$ over every field.
23. We prove the integral vanishing $\beta_{3,7} = 0$ by a zero-vertex matching and a signed unit filler block, thereby determining the complete third homological row over every field.
24. We determine the complete graded Betti polynomial of the cubic-colon quotient over every field, including every free-module rank and shift.
25. We prove that the quadratic quotient has depth one and regularity two, so the full cubic mapping cone is minimal; this determines the two highest regularity strands of the special fiber.
26. We reduce the high-variable kernel to a two-layer incidence module and prove the first exact surviving regularity-two class in the kernel, quadratic quotient, and special fiber over every field.
27. We classify all primitive zero rows of that incidence cokernel, obtain consecutive kernel classes in homological degrees $p + 1, \dots, 2p - 3$, and prove that the first complete target quotient has characteristic-dependent multigraded Betti number: four over $\mathbf{F}_2$ and three over $\mathbf{F}_3$ for both the quadratic quotient and the special fiber.

2 The finite semigroup problem

Definition 2.1. A *numerical semigroup* is a cofinite additive submonoid $\Gamma \subseteq \mathbb{N}$. Its Frobenius number is $F(\Gamma) = \max(\mathbb{N} \setminus \Gamma)$. It is *symmetric* when

$$n \in \Gamma \iff F(\Gamma) - n \notin \Gamma \quad (0 \leq n \leq F(\Gamma)).$$

Fix a symmetric semigroup Γ with Frobenius number F and a positive gap $s \notin \Gamma$. Normalize the corresponding two-generated relative ideal to $(0, s)$. Define

$$\begin{aligned} E_s &= \{n \in \mathbb{N} : n, n + s \in \Gamma\}, \\ D_s &= \{n \in \mathbb{N} : n, n + s, n + 2s \in \Gamma\}. \end{aligned}$$

The García-Sánchez-Leamer reduction identifies the counterexample condition with

$$D_s = E_s + E_s. \tag{1}$$

Equivalently, every length-two arithmetic sequence of step s in Γ decomposes as a sum of two length-one sequences [2, 4].

Lemma 2.2 (Automatic inclusion, [D]). *For every numerical semigroup Γ and every positive integer s , $E_s + E_s \subseteq D_s$.*

Proof. If $a, b \in E_s$, then $a, a + s, b, b + s \in \Gamma$. Closure gives $a + b \in \Gamma$, $a + b + s = a + (b + s) \in \Gamma$, and $a + b + 2s = (a + s) + (b + s) \in \Gamma$. Hence $a + b \in D_s$. $\square$

Lemma 2.3 (Finite window, [D]). *Equation (1) holds if and only if $D_s \cap [0, 2F + 1] \subseteq E_s + E_s$.*

Proof. Only the reverse inclusion needs checking by Lemma 2.2. Every integer at least $F + 1$ belongs to Γ , hence to E_s and D_s . Let $m = \min E_s$; then $m \leq F + 1$. For every $n \geq m + F + 1$, both m and $n - m$ lie in E_s , so $n \in E_s + E_s$. Since $m + F + 1 \leq 2F + 2$, every $n \geq 2F + 2$ is automatic. The displayed finite window is therefore complete. $\square$

Lemma 2.4 (Parity, [D]). *The Frobenius number of a symmetric numerical semigroup is odd.*

Proof. If F were even, symmetry at $n = F/2$ would require $F/2$ to belong to Γ exactly when it does not belong to Γ . $\square$

3 The public model and independent calibration

Pham’s public semigroup is

$$\Gamma = \langle 56, 57, 58, 63, 64, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, \\ 87, 89, 90, 93, 95, 96, 97 \rangle,$$

with the ideal generated by t^{56} and t^{70} , hence normalized shift $s = 14$ [5]. An independent Singular/4ti2 calculation and a separate standard-library finite route verify $F = 181$, conductor 182, genus 91, symmetry, and equality (1) [MV]. The Singular route uses a 322-generator toric standard basis; both reduced ideal differences vanish. The hypersurface control $\langle 4, 5 \rangle$ rejects equality with explicit residues [MV].

The endomorphism overring of the ideal has value semigroup

$$v(\text{End}_R(I)) = \Gamma \cup \{101, 107, 181\},$$

with Frobenius number 125, genus 88, and type 24 [MV]. This explains why positive criteria that require a Gorenstein endomorphism ring do not apply, but it does not imply minimality.

4 Complete reproduction below 69

For fixed odd F , irreducible numerical semigroups are exactly the symmetric ones. Blanco and Rosales construct all of them as a rooted tree [3]. Starting from

$$C(F) = \{0, (F + 1)/2, (F + 1)/2 + 1, \dots\} \setminus \{F\},$$

a child replaces a suitable minimal generator $x > F/2$ by $F - x$. Their theorem supplies exact child conditions, completeness, and a unique parent.

Our direct implementation validates every node for closure, symmetry, genus, minimal generators, and its theorem parent. It then checks every gap by Lemma 2.3 and retains a first element of $D_s \setminus (E_s + E_s)$ whenever equality fails. The full range $F = 1, 3, \dots, 67$ contains 48,954 semigroups and 1,503,391 gap cases; none is rigid [MV]. At $F = 11$ the implementation returns the six nodes listed by Blanco–Rosales, while an invalid mutation is rejected by closure at $(1, 1)$.

An independent route generates one fixed-pair DIMACS formula for each (F, s) , including shifts that are forced not to be gaps. CaDiCaL emits a DRAT proof for every UNSAT result, and DRAT-trim checks each proof against the exact CNF [20, 21]. All 1,156 formulas are UNSAT and independently verified [MV]. A committed auditor rehashes 4,624 external files totaling 211,671,241 bytes with zero mismatch.

5 One selector formula per Frobenius value

The fixed-pair route duplicates the same semigroup constraints once per shift. We instead introduce Boolean membership variables h_n for $0 \leq n \leq F$ and one-hot variables q_s for $1 \leq s \leq F$.

5.1 Semigroup and selector clauses

The CNF contains units h_0 and $\neg h_F$, the symmetry clauses

$$(h_n \vee h_{F-n}) \wedge (\neg h_n \vee \neg h_{F-n}),$$

and closure clauses $\neg h_a \vee \neg h_b \vee h_{a+b}$ for $a + b \leq F$. The selectors satisfy one at-least-one clause, all pairwise at-most-one clauses, and $q_s \Rightarrow \neg h_s$. Thus every model chooses exactly one gap.

For $0 \leq n \leq 2F + 1$, guarded Tseitin clauses impose, under the selected q_s ,

$$\begin{aligned} e_n &\longleftrightarrow h_n \wedge h_{n+s}, \\ d_n &\longleftrightarrow h_n \wedge h_{n+s} \wedge h_{n+2s}, \end{aligned}$$

where h_j is the constant true for $j > F$. For each unordered decomposition $n = a + b$, a variable $p_{n,a,b}$ is equivalent to $e_a \wedge e_b$. Finally,

$$d_n \Rightarrow \bigvee_{a+b=n} p_{n,a,b}.$$

The automatic inclusion from Lemma 2.2 need not be duplicated in the formula.

Proposition 5.1 (Selector equisatisfiability, [D]). *For positive odd F , the selector CNF is satisfiable if and only if there exist a symmetric numerical semigroup Γ with Frobenius number F and a gap s satisfying $D_s = E_s + E_s$.*

Proof. The semigroup clauses decode exactly a symmetric semigroup of Frobenius number F , and the one-hot clauses decode a unique gap. Because exactly one guard is true, all e_n and d_n equal the definitions for that gap. The p equivalences and final disjunctions impose $D_s \subseteq E_s + E_s$ through $2F + 1$. Lemmas 2.2 and 2.3 give global equality. Conversely, from a rigid pair assign h and q from the pair, assign e, d by their definitions, and assign each p by its conjunction. Every clause is then true. $\square$

The selector encoding passes two adversarial calibrations [MV]. Every odd $F \leq 67$ is UNSAT with an accepted proof, while $F = 181$ is SAT. The latter formula has 34,397 variables and 363,912 clauses, selects $s = 14$, and decodes exactly to the public membership vector. A newly generated fixed-pair formula at $(181, 14)$ returns the same vector.

6 The certified search and minimality theorem

We scan odd F in strict increasing order. Table 1 summarizes the closed run.

Theorem 6.1 (Frobenius minimality, [MV+D]). *Among symmetric numerical semigroups admitting a nonprincipal two-generated monomial ideal that is a counterexample to the Huneke–Wiegand prediction, the least Frobenius number is 181. It is attained by Pham’s public example at shift 14.*

Table 1: Selector-CNF minimality search.

Quantity	Value
Odd Frobenius values requested	57 (69 through 181)
UNSAT with accepted DRAT proof	56 (69 through 179)
Validated SAT models	1 (181, shift 14)
UNKNOWN or timeout	0
Wall time	4,303.86 seconds
CNF bytes	169,745,433
DRAT proof bytes	586,785,257
Slowest solve	$F = 175$, 218.83 seconds

Proof. By Lemma 2.4, only odd Frobenius values occur. Complete tree enumeration and the independent fixed-pair proof suite exclude every odd $F \leq 67$. Proposition 5.1 and the accepted selector DRAT proofs exclude every odd F from 69 through 179. The validated selector and fixed-pair models at (181, 14) provide existence. These cases exhaust all smaller positive Frobenius values. $\square$

The search audit independently rehashes all 228 expected CNF, proof, solver-log, and checker-log files, totaling 760,081,739 bytes. It also reconstructs both SAT membership masks, validates closure, symmetry, and rigidity, checks strict odd ordering, and recomputes aggregate SHA-256

0f580de2707a00fdd52e1b3c04e7767b97ce7b0a826593b119e9a49ae04da743.

No file is missing, no hash differs, and no status marker or semantic check fails [MV].

7 Complete classification at the minimum

The selector search proves existence at $F = 181$ but, by itself, does not classify every model at that value. We close this gap in two projected layers. First, after discovering a valid selector model at shift s , we add the unit clause $\neg q_s$. Second, for each feasible shift we enumerate the fixed-shift formula and block each complete membership projection.

Lemma 7.1 (Projected blocker, [D]). *Let $x_1, \dots, x_k$ be distinct Boolean variables with complete assignment a . The clause*

$$B(a) = \bigvee_{a_i=1} \neg x_i \vee \bigvee_{a_i=0} x_i$$

is false exactly on assignment a . Its truth is independent of every variable outside the projection.

Proof. Every displayed literal is false under a . Any other complete assignment differs at some coordinate j , where the corresponding displayed literal is true. Variables absent from the clause cannot affect either statement. $\square$

For semigroup enumeration, $B(a)$ contains one signed literal for every $h_0, \dots, h_{181}$. Consequently, it excludes one mathematical membership vector and every auxiliary extension of that vector, while preserving every different semigroup. Blocking only the positive coordinates would not have this property.

Theorem 7.2 (Minimum-layer uniqueness, [MV+D]). *Up to translation and interchange of the two monomial generators, Pham’s pair is the unique nonprincipal two-generated monomial-ideal counterexample over a symmetric numerical semigroup ring with Frobenius number 181.*

Proof. The unblocked selector formula returns the exact public pair at shift 14. After adding $\neg q_{14}$, the formula is UNSAT and its DRAT proof is accepted, so no other shift is feasible. The fixed (181, 14) formula returns the exact public membership vector. After adding its complete 182-literal blocker from Lemma 7.1, that formula is UNSAT and its DRAT proof is accepted. Thus no second semigroup exists at the only feasible shift. Translation and interchange reduce any two monomial generators to one positive normalized shift, which proves the statement. $\square$

Table 2: Certificate summary for the complete $F = 181$ layer.

Quantity	Value
Feasible shifts	1, namely {14}
Normalized rigid pairs	1
Distinct membership vectors	1
Support terminal proof bytes	45,867,741
Fixed-class terminal proof bytes	1,608,691
Audited external files	12
Audited external bytes	63,609,504
Fresh proof-recheck time	232.88 seconds

The independent auditor reconstructs all four CNFs byte-for-byte, decodes both SAT logs, reruns closure, symmetry, minimal-generator and full-window/tail rigidity checks, and freshly checks both terminal proofs [MV]. It accounts for every external file and reproduces manifest SHA-256

02d1265b75e886bc6ec693e60b367b661476170c755e38a220342396564ef5d3.

The compact classification aggregate is

b37d9410fa07c00f22ccc0f83796644db3b44f0dd654c06933aa5d22cb1ed788.

8 Independent object-level checks beyond the old frontier

The theorem-tree route was continued beyond 67 without consulting SAT models. Table 3 lists the first four new odd values.

This is not needed to trust a DRAT derivation, but it attacks the custom encoding at the object level and supplies explicit failed-rigidity witnesses for every tested gap. The routes agree.

9 From a failed ansatz to an infinite family

Landeros et al. prove the positive result for semigroups generated by a generalized arithmetic sequence $\langle a, ah + d, \dots, ah + kd \rangle$ [4]. The public semigroup cannot lie in this class [D]. Its multiplicity forces $a = 56$. Since $57 \in \Gamma$ and no sum of two positive semigroup elements is below

Table 3: Complete Blanco–Rosales tree cross-checks beyond the published boundary.

F	Symmetric semigroups	Gap cases	Rigid cases	Seconds
69	11,276	394,660	0	137.00
71	19,812	713,232	0	235.37
73	25,405	939,985	0	333.21
75	23,297	885,286	0	330.10
Total	79,790	2,933,163	0	1,035.68

112, the presentation forces $h = d = 1$. Since $63 \in \Gamma$, it would then include 59, 60, 61, 62, but all four are gaps.

The seed also displays $56 = 4s$ and $181 = 13s - 1$ at $s = 14$, with generator blocks

$$4s + \{0, 1, 2, 7, 8\}, \quad 5s + [0, s - 1], \quad 6s + \{3, 5, 6, 9, 11, 12, 13\}.$$

A predeclared exact sweep refutes the naive fixed-offset lift [MV]. Only $s = 14$ passes among even $s \leq 100$. At $s = 16$, the generated semigroup has $F = 205$ rather than 207; at $s = 28$ the predicted Frobenius value returns but symmetry fails. This negative result prevents the minimality theorem from being overinterpreted as evidence of an immediate family.

We next retained only the structural scaffold $m = 4s$, $F = 13s - 1$, the full interval $[5s, 6s - 1] \subseteq \Gamma$, and rigidity at shift s . A proof-carrying Boolean search certifies that the scaffold is infeasible at $s = 16, 18$ and returns independently validated models for every even $s = 20, 22, \dots, 40$ [MV]. The eleven non-seed models justified a separate extraction stage, but did not themselves prove recurrence. The first fixed-width interval formula passed three parameters and then failed: at $s = 38$, $9s + 7$ lies in D_s but not in $E_s + E_s$. For every later parameter in that formula, carried sums from the first block end at residue 6, while the next summand block begins at residue at least 8. This uniform obstruction suggests widening the first interval with the parameter.

9.1 The parametric blocks

Fix an integer $p \geq 4$ and put $s = 6p$. All intervals below are integer intervals. Define subsets of $[0, s - 1]$ by

$$\begin{aligned} A_p &= [0, p] \cup [3p, 4p - 2], \\ B_p &= ([p + 1, 3p - 1] \setminus \{2p - 1\}) \cup \{4p\} \cup [5p - 1, 6p - 1], \\ C_p &= [0, 2p] \cup [3p, 5p - 2]. \end{aligned}$$

Define $\Gamma_p \subseteq \mathbb{N}$ by the membership blocks

$$\{0\}, \quad 4s + A_p, \quad [5s, 6s - 1], \quad 6s + B_p, \quad 8s + C_p, \quad [9s, 13s - 2], \quad [13s, \infty), \quad (2)$$

with every unlisted nonnegative integer below $13s$ a gap.

For subsets of $[0, s - 1]$, let $\text{low}(X + Y)$ denote the sums below s , and let $\text{carry}(X + Y)$ denote the sums at least s after subtraction of s .

Lemma 9.1 (Residue identities, [D]). *For every $p \geq 4$,*

$$\begin{aligned} \text{low}(A_p + A_p) &= C_p, & \text{carry}(A_p + A_p) &= [0, 2p - 4], \\ \text{low}(A_p + B_p) &= [p + 1, 6p - 1], & \text{carry}(A_p + B_p) &= [0, 4p - 3], \\ \text{low}(B_p + B_p) &= [2p + 2, 6p - 2], & \text{carry}(B_p + B_p) &= [0, 6p - 2], \\ \text{low}(A_p + C_p) &= [0, 6p - 2]. \end{aligned}$$

Moreover,

$$[0, 2p - 4] \cup [p + 1, 6p - 1] = [0, 6p - 1], \quad (3)$$

$$[0, 4p - 3] \cup [2p + 2, 6p - 2] = [0, 6p - 2]. \quad (4)$$

Proof. Split A_p into its two displayed intervals and split the punctured component of B_p at $2p - 1$. Pairwise interval addition, followed by separating sums below and above $s = 6p$, gives the seven displayed identities. The singleton $4p$ fills the endpoint removed from the middle of B_p . All interval joins are immediate except (3): its two pieces meet exactly when $2p - 3 \geq p + 1$, equivalently $p \geq 4$. For (4), the overlap condition is $4p - 2 \geq 2p + 2$, which also holds for $p \geq 4$. $\square$

Theorem 9.2 (Infinite family, [D]). *For every integer $p \geq 4$, the set Γ_p in (2) is a symmetric numerical semigroup with multiplicity $24p$, Frobenius number $78p - 1$, conductor $78p$, and embedding dimension $11p$. Let*

$$R_p = \mathbb{k}[t^{\Gamma_p}]_{(t^n : n \in \Gamma_p \setminus \{0\})}, \quad I_p = (t^{24p}, t^{30p})R_p.$$

Then R_p is Gorenstein and I_p is nonprincipal and rigid. Hence the pairs (R_p, I_p) form an infinite family of counterexamples in the two-generated monomial-ideal class.

Proof. We first prove the semigroup assertions. The only sums of positive displayed members that can stay below the conductor have starting levels among 4, 5, 6, 8. Lemma 9.1 handles the non-full target blocks: $A_p + A_p$ produces $8s + C_p$ and part of level 9; $A_p + B_p$, $B_p + B_p$, and $A_p + C_p$ land exactly inside the declared high blocks and never produce the Frobenius residue $s - 1$ in level 12. Sums involving the full level 5 block either fill a declared level or enter the conductor tail. Thus Γ_p is closed.

Conversely, the lower three blocks generate every displayed member. The identity $\text{low}(A_p + A_p) = C_p$ generates level 8. The presence of $0 \in A_p$ makes $(4s + A_p) + [5s, 6s - 1]$ fill level 9, and two level-5 elements fill level 10 and residues $0, \dots, s - 2$ of level 11; the residue $s - 1 \in B_p$ fills the remaining point. Three elements from $4s + A_p$ generate residues $0, \dots, 3p$ of level 12, while $\text{low}(B_p + B_p) = [2p + 2, s - 2]$ completes that level through $13s - 2$. Adding $4s$ to $[9s, 13s - 2]$ gives $[13s, 17s - 2]$, and $(4s + 1) + (13s - 2) = 17s - 1$. These are $4s$ consecutive generated integers, so repeated addition of the multiplicity gives the entire tail.

Therefore the multiplicity is $4s = 24p$, the conductor is $13s = 78p$, and the Frobenius number is $13s - 1 = 78p - 1$. Every member of the three lower blocks lies below $7s$, whereas a sum of two positive members is at least $8s$; all are minimal generators. Since $|A_p| = 2p$, $|B_p| = 3p$, and the full block has size $6p$, the embedding dimension is $11p$.

Symmetry is blockwise about $F = 13s - 1$. The residue-reflection complement of A_p is exactly C_p . The full level-5 block reflects to the empty level-7 block. In B_p , exactly one residue from each pair $\{r, s - 1 - r\}$ occurs: the puncture $2p - 1$ pairs with the singleton $4p$, and the remaining displayed intervals pair with their omitted intervals. The gaps below $4s$ reflect to the full range $[9s, 13s - 2]$, and 0 reflects to the gap F . Hence Γ_p is symmetric, so R_p is Gorenstein.

It remains to prove rigidity. Normalize I_p to $J_p = (1, t^s)$ and use E_s, D_s from Section 2. Direct inspection of (2) gives the nontrivial residue layers

$$\begin{aligned} E_4 &= A_p, & E_5 &= B_p, & E_8 &= C_p, \\ E_9 &= E_{10} = [0, s-1], & E_{11} &= E_{12} = [0, s-2], \\ D_8 &= C_p, & D_9 &= [0, s-1], \\ D_{10} &= D_{11} = D_{12} = [0, s-2]. \end{aligned}$$

All lower D layers are empty; in particular D_4 is empty because $A_p \cap B_p = \emptyset$. Lemma 9.1 and (3)–(4) compute the five load-bearing layers of $E_s + E_s$:

$$\begin{aligned} D_8 &= \text{low}(A_p + A_p), & D_9 &= \text{carry}(A_p + A_p) \cup \text{low}(A_p + B_p), \\ D_{10} &= \text{carry}(A_p + B_p) \cup \text{low}(B_p + B_p), & D_{11} &= \text{carry}(B_p + B_p), & D_{12} &= \text{low}(A_p + C_p). \end{aligned}$$

At levels 13 and 14, the zero residue in A_p translates full high E blocks. At levels 15 and 16, those high blocks omit only $s-1$, and the residues $0, 1 \in A_p$ fill both the interval and its final endpoint. Level 17 is filled by $E_4 + E_{13}$; levels 18, 19, and 20 by $E_9 + E_9$, $E_9 + E_{10}$, and $E_{10} + E_{10}$; and level 21 by $E_8 + E_{13}$ because $0 \in C_p$. Every level from 22 onward is filled by $E_9 + E_q$ with $q \geq 13$, where the conductor makes E_q full. Thus $D_s \subseteq E_s + E_s$; Lemma 2.2 gives equality.

Finally, s is a gap. If J_p were principal, write $J_p = xR_p$. Since $1 \in J_p$, there is $r \in R_p$ with $xr = 1$, so $rJ_p = R_p$ and $(r, rt^s) = R_p$. In the local ring, one of these two generators must be a unit. Positive valuation excludes rt^s , so r is a unit and $t^s \in R_p$, a contradiction. The ideal is therefore nonprincipal and rigid. The multiplicities $24p$ are distinct, so the family contains infinitely many distinct rings. $\square$

9.2 Exact support and boundaries

The proof of Theorem 9.2 is independent of a finite sweep. As adversarial support, the implementation evaluated every $p = 2, 3, \dots, 300$ [MV]. Exactly the predicted boundary values $p = 2, 3$ fail, while all 297 values $p \geq 4$ pass the seven identities, closure, generation, symmetry, invariants, and full-window/tail rigidity. The instances $p = 4, 5$ reproduce independently generated Boolean models byte-for-byte. Two corrupted positive controls are rejected. A separate auditor reconstructs all 297 positive membership hashes from the formula and performs fresh semantic checks at $p = 4, 5, 17, 73, 151, 300$. The campaign aggregate is

81d5a8eb6cf2e848807323e3b0bdba58c464779d25cdc788cef027585540dce2.

The independent-audit aggregate is

eb2aaf17650ed99f4e220a43c53bdd8835c82688a37567bb154c30a1ae520ce9.

The family remains outside the generalized-arithmetic-sequence positive class [D]. Membership of $4s+1$ forces unit step, while membership of $4s+3p$ would then force the actual gap $4s+p+1$. Theorem 9.2 does not say that every model in the Boolean scaffold belongs to this family, or that $p=4$ is a global threshold outside the displayed formula.

10 Uniform endomorphism overrings

Normalize I_p to $J_p = (1, t^s)$ and write

$$V_p = v(J_p) = \Gamma_p \cup (s + \Gamma_p), \quad \Lambda_p = v(\text{End}_{R_p}(J_p)).$$

Define

$$Q_p = [p + 1, 2p - 2] \cup \{2p, 4p\}.$$

Theorem 10.1 (Uniform endomorphism anatomy, [D]). *For every integer $p \geq 4$,*

$$\Lambda_p = \Gamma_p \cup (7s + Q_p) \cup \{13s - 1\}.$$

This numerical semigroup has multiplicity $24p$, Frobenius number $54p - 1$, conductor $54p$, genus $38p - 1$, and embedding dimension $12p$. It is nonsymmetric, so the localized semigroup ring $E_p = \text{End}_{R_p}(J_p)$ is not Gorenstein. Moreover,

$$\begin{aligned} \text{Ext}_{E_p}^1(J_p, J_p) &= 0, & J_p &\text{ is not reflexive over } E_p, \\ \text{Ext}_{E_p}^1(J_p, E_p) &\neq 0, & \text{Ext}_{E_p}^2(J_p, J_p) &\neq 0, \\ \text{Tor}_1^{R_p}(J_p, E_p) &\neq 0. \end{aligned}$$

Proof. Let G_k and V_k denote the residue sets at levels ks in Γ_p and V_p , respectively. From (2),

$$V_0 = V_1 = \{0\}, \quad V_2 = V_3 = \emptyset, \quad V_4 = A_p, \quad V_5 = V_6 = [0, s - 1], \quad V_7 = B_p, \quad V_8 = C_p,$$

and $V_k = [0, s - 1]$ for every $k \geq 9$. An exponent $n = ks + r$ stabilizes J_p exactly when both n and $n + s$ belong to V_p . Thus the level- k residue set of Λ_p is $V_k \cap V_{k+1}$.

All adjacent intersections reproduce the corresponding Γ_p block except at level 7 and the old Frobenius point. Directly,

$$B_p \cap C_p = ([p + 1, 2p] \setminus \{2p - 1\}) \cup \{4p\} = Q_p.$$

Every level from 9 onward is full, so the only other added value is $13s - 1$. This proves the formula for Λ_p .

The first positive value remains $4s = 24p$. Level 8 still omits residue $s - 1$, while level 9 and all later levels are full. Hence the Frobenius number is $9s - 1 = 54p - 1$ and the conductor is $9s = 54p$. Counting gaps by levels gives

$$(4s - 1) + (s - |A_p|) + (s - |B_p|) + (s - |Q_p|) + (s - |C_p|) = 38p - 1.$$

A symmetric semigroup with this Frobenius number would have genus $27p$, so Λ_p is nonsymmetric.

The $11p$ generators of Γ_p and the p new values $7s + Q_p$ all lie below twice the multiplicity, hence are minimal. For any $q \in Q_p$, the level-5 block contains $5s + (s - 1 - q)$ and

$$(7s + q) + (5s + (s - 1 - q)) = 13s - 1.$$

Thus the old Frobenius point is generated and there are exactly $12p$ minimal generators.

Finally, R_p is a one-dimensional Gorenstein local domain and J_p is faithful, torsion-free, two-generated, nonprincipal, and rigid by Theorem 9.2. The ring E_p is commutative and is strictly larger than R_p because Q_p is nonempty. Proposition 4.1(2) of Dey and Lyle makes rigidity descend to E_p ; the contrapositives of their Theorems 4.2–4.4 give the displayed nonreflexivity, Ext, and Tor conclusions [6]. $\square$

The symbolic argument is independent of a parameter sweep. As adversarial support, two exact implementations agree for every $p = 4, \dots, 300$ [MV]. A separate auditor rehashes all 297 rows, reconstructs complete semantic windows at $p = 4, 5, 17, 73, 151, 300$, and rejects a missing Q_p value and a missing terminal singleton. The deterministic campaign and audit aggregates are

e21926a689178a6c70b3b6e8319053edd0fd13f164ced9565d3b976e6159c0b0

and

2ed711045ad83a3b47fb3e71d4c75ae9bfa9be1a5dd9a8c4072d5f170510343b.

11 Pseudo-Frobenius anatomy and reduced type

For a numerical semigroup H , write

$$\text{PF}(H) = \{f \notin H : f + h \in H \text{ for every } h \in H \setminus \{0\}\}.$$

Its Cohen–Macaulay type is $|\text{PF}(H)|$. For complements inside $[0, s - 1]$, define

$$\begin{aligned} B_p^c &= [0, p] \cup \{2p - 1\} \cup [3p, 4p - 1] \cup [4p + 1, 5p - 2], \\ Q_p^c &= [0, p] \cup \{2p - 1\} \cup [2p + 1, 4p - 1] \cup [4p + 1, 6p - 1], \\ C_p^c &= [2p + 1, 3p - 1] \cup [5p - 1, 6p - 1]. \end{aligned}$$

Theorem 11.1 (Type anatomy, [D]). *For every integer $p \geq 4$,*

$$\text{PF}(\Lambda_p) = (6s + B_p^c) \cup (7s + Q_p^c) \cup (8s + C_p^c).$$

Consequently the type and reduced type of $\mathbb{k}[[\Lambda_p]]$ are both $10p$. Thus the ring has maximal reduced type. The semigroup is not almost symmetric, and $\mathbb{k}[[\Lambda_p]]$ is not almost Gorenstein.

Proof. The multiplicity of Λ_p is $4s$ and its conductor is $9s$. Every gap in the final multiplicity window $[5s, 9s - 1]$ is therefore pseudo-Frobenius. Since level 5 is full, these gaps are precisely the complements of B_p, Q_p, C_p at levels 6, 7, and 8. Their cardinalities are $3p, 5p, 2p$.

It remains to exclude a pseudo-Frobenius number below $5s$. The full level-5 generator block gives explicit witnesses. For a gap r at level 0 with $r > 0$, add $5s + (s - r)$ to reach the level-6 gap of residue zero. At level 1, use $5s$ when $r = 0$ and $5s + (s - r)$ otherwise. At levels 2 and 3, add $5s + (s - 1 - r)$ to reach residue $s - 1$ in a missing later block.

For a level-4 gap $4s + r$, if $r \notin C_p$ add the generator $4s$. The remaining residues lie in

$$C_p \setminus A_p = [p + 1, 2p] \cup [4p - 1, 5p - 2].$$

In the first interval add $4s + (2p + 1 - r)$; in the second add $4s + (5p - 1 - r)$. Both added residues lie in $[1, p] \subset A_p$, while the resulting level-8 residue is a gap of C_p . Thus no lower gap is pseudo-Frobenius, proving the formula.

Reduced type counts the gaps in the last multiplicity window, so it is also $10p$ [7]. Finally, using $g(\Lambda_p) = 38p - 1$ and $F(\Lambda_p) = 54p - 1$,

$$2g - (F + \text{type}) = 12p - 1 > 0.$$

The almost-symmetric genus identity therefore fails, giving the final claims. $\square$

Two exact implementations confirm the displayed set for all $p = 4, \dots, 300$ [MV]. One checks every gap against the full minimal-generator set; the other extracts maximal elements of the Apéry poset modulo $4s$. The campaign and independent-audit aggregates are

9bed38fb1c786c3740e000dde7ea7d79a7e7c83fa584ff12fc2c4623b5d503ec

and

0315c4c22c41e0d2b8a5abb27f717a4d4a6f7356ef30ca388206d739e0de2c37.

12 Exact trace and conductor ideals

Let $E_p = \text{End}_{R_p}(J_p) = \mathbb{k}[[\Lambda_p]]$ and define the reflected block

$$H_p = (s - 1) - Q_p = \{2p - 1, 4p - 1\} \cup [4p + 1, 5p - 2].$$

Theorem 12.1 (Exact trace and conductor, [D]). *For every integer $p \geq 4$,*

$$\text{tr}_{R_p}(J_p) = R_p : E_p = \text{tr}_{R_p}(E_p) = T_p,$$

where

$$\begin{aligned} v(T_p) &= (4s + A_p) \cup (5s + (A_p \cup B_p)) \cup (6s + B_p) \cup (8s + C_p) \\ &\quad \cup [9s, 13s - 2] \cup [13s, \infty). \end{aligned}$$

Moreover,

$$\ell_{R_p}(R_p/T_p) = \ell_{R_p}(E_p/R_p) = p + 1.$$

Proof. Because $1 \in J_p$,

$$v(R_p : J_p) = W_p = \{n \in \Gamma_p : n + s \in \Gamma_p\}.$$

Intersecting adjacent blocks of (2) gives

$$(W_4, W_5, W_6, W_7, W_8) = (A_p, B_p, \emptyset, \emptyset, C_p),$$

with full blocks at levels 9 and 10, blocks $[0, s - 2]$ at levels 11 and 12, and full blocks from level 13 onward. Since W_p is a Γ_p -ideal,

$$v(\text{tr}_{R_p}(J_p)) = W_p \cup (s + W_p),$$

which is exactly the displayed set $v(T_p)$.

For the conductor $R_p : E_p$, only the values in

$$\Lambda_p \setminus \Gamma_p = (7s + Q_p) \cup \{13s - 1\}$$

add constraints. The terminal singleton excludes $n = 0$ and imposes no condition on positive $n \in \Gamma_p$. Every value at level 4 passes the $7s + Q_p$ translates: a failure in the sole level-12 gap would require $a + q = 2s - 1$, but $a + q \leq 8p - 2 < 12p - 1$. At level 5, residue r fails exactly when $r + q = s - 1$ for some $q \in Q_p$, hence exactly when $r \in H_p$. All values from level 6 onward translate into the conductor tail. Since

$$[0, s - 1] \setminus H_p = A_p \cup B_p,$$

this proves $v(R_p : E_p) = v(T_p)$. The conductor is an E_p -ideal, so $\text{tr}_{R_p}(E_p) = E_p(R_p : E_p) = R_p : E_p$.

Relative to Γ_p , the common ideal omits exactly $\{0\} \cup (5s + H_p)$, giving colength $p + 1$. The endomorphism formula gives exactly $p + 1$ values in $\Lambda_p \setminus \Gamma_p$, proving the balanced identity. $\square$

The equality of the two traces is consistent with the reflexive-trace criterion of Lindo–Maitra–Zhang [8]; over the present Gorenstein base it is not by itself discriminatory. The equality of the two lengths is also general structure, not a new family-specific mechanism. Indeed, applying $\text{Hom}_{R_p}(-, R_p)$ to $0 \rightarrow R_p \rightarrow E_p \rightarrow E_p/R_p \rightarrow 0$ and using local duality gives

$$\ell_{R_p}(R_p/(R_p : E_p)) = \ell_{R_p}(E_p/R_p),$$

as in Herzog–Kumashiro, Proposition 3.1, Claim 1 [9]. The new family-specific data in Theorem 12.1 are the exact common ideal and the evaluation of both lengths as $p + 1$.

The first declared shorthand used the tail $[9s, \infty)$ and was refuted at $13s - 1$ by the $p = 4$ smoke check [MV]. That point is the Frobenius gap of Γ_p , so no R_p -ideal can contain it. The corrected formula above was then checked by two exact routes for all $p = 4, \dots, 300$, with independent reconstruction at six parameters and three rejected corruptions. The campaign and audit aggregates are

77448398a26b958c66818b7ac4aaa4b542bad11cdba5ec7c8f7fe76db37526e2

and

d55ed876d7918cb4c46d8e5f3894a508d172693bde4b4c76cb67c42fcfbf0ac1.

13 Exact conductor-stability defect

Dey proves that for a finite birational extension $R \subseteq S$ of a one-dimensional Gorenstein local ring, the conductor $R : S$ is stable if and only if S is Gorenstein [10, Corollary 3.7]. Theorem 11.1 gives $\text{type}(E_p) = 10p > 1$, so this criterion already shows that $T_p = R_p : E_p$ is nonstable. The following computation quantifies that failure.

Theorem 13.1 (Exact stability defect, [D]). *For every integer $p \geq 4$,*

$$v(T_p^2) = (8s + C_p) \cup [9s, 13s - 2] \cup [13s, \infty).$$

Moreover,

$$\begin{aligned} v(T_p^2) \setminus v(t^{4s}T_p) = & 8s + ([p + 1, 2p] \cup [4p - 1, 5p - 2]) \\ & \cup 9s + (\{2p - 1, 4p - 1\} \cup [4p + 1, 5p - 2]) \\ & \cup 10s + ([0, p] \cup \{2p - 1\} \cup [3p, 4p - 1] \cup [4p + 1, 5p - 2]) \\ & \cup 11s + [0, s - 1] \\ & \cup 12s + ([2p + 1, 3p - 1] \cup [5p - 1, s - 2]) \\ & \cup \{17s - 1\}. \end{aligned}$$

Consequently,

$$\ell_{R_p}(T_p^2/t^{4s}T_p) = 14p.$$

Proof. Put $D = [0, s - 1]$, $D^- = [0, s - 2]$, and $U_p = A_p \cup B_p$. For residue sets $X, Y \subseteq D$, write $L(X, Y)$ for sums below s and $H(X, Y)$ for sums at least s , translated by $-s$. Expanding the

defining sets into integer intervals and using $[a, b] + [c, d] = [a + c, b + d]$ gives

$$\begin{aligned} L(A_p, A_p) &= C_p, \\ H(A_p, A_p) \cup L(A_p, U_p) &= D, \\ H(A_p, U_p) \cup L(A_p, B_p) \cup L(U_p, U_p) &= D, \\ H(A_p, B_p) \cup H(U_p, U_p) \cup L(U_p, B_p) &= D, \\ H(U_p, B_p) \cup L(B_p, B_p) \cup L(A_p, C_p) &= D^-. \end{aligned}$$

These five identities list all contributions, with carries, to levels 8 through 12 of T_p^2 . They give $(8s + C_p) \cup [9s, 13s - 2]$ and exclude the inherited gap $13s - 1$. Multiplication by the least value $4s$ covers $[13s, 17s - 2] \cup [17s, \infty)$, while

$$17s - 1 = (4s + 1) + (13s - 2)$$

fills the intervening endpoint. This proves the square formula.

Shifting Theorem 12.1 by $4s$ and subtracting its blocks from the square gives the displayed defect. The six block cardinalities are $2p, p, 3p, 6p, 2p - 1, 1$, whose sum is $14p$. $\square$

The first EXP-015 prediction incorrectly replaced the two tail intervals by $[9s, \infty)$. The $p = 4$ smoke gate refuted it at $13s - 1$ before a campaign artifact was written [MV]. EXP-016 retained that gap, proved the corrected formula above, and checked two exact routes for all $p = 4, \dots, 300$. Its campaign and independent-audit aggregates are

8c280bf1db017f678896d473461ae245b4fce57d680be94094b0884556d91853

and

ce4ded03761eb44cba766eafb46926c8ca83d0f7f49a26697afed5ed49905653.

14 Conductor reduction number and Hilbert data

For an ideal I , a subideal Q is a *reduction* when $I^{r+1} = QI^r$ for some r ; the least such r is the reduction number with respect to Q [11].

Theorem 14.1 (Exact conductor reduction sequence, [D]). *For every integer $p \geq 4$, put $Q_p = t^{4s}R_p \subset T_p$. Then Q_p is a minimal reduction of T_p with reduction number four. More precisely,*

n	0	1	2	3	$n \geq 4$
$\ell_{R_p}(T_p^{n+1}/Q_p T_p^n)$	$23p - 1$	$14p$	$2p$	1	0.

The Hilbert–Samuel function satisfies

$$\ell_{R_p}(R_p/T_p^n) = 24pn - 39p \quad (n \geq 4),$$

and consequently $e_0(T_p) = 24p$ and $e_1(T_p) = 39p$.

Proof. Theorem 13.1 supplies the square. The residue identity $\text{low}(A_p + C_p) = [0, s - 2]$ then gives

$$v(T_p^3) = [12s, 13s - 2] \cup [13s, \infty).$$

Adding $4s + A_p$ fills $[16s, 17s - 2]$ and the tail from $17s$; because $1 \in A_p$, $(4s + 1) + (13s - 2) = 17s - 1$ fills the intervening endpoint. Hence

$$v(T_p^4) = [16s, \infty), \quad v(T_p^5) = [20s, \infty) = v(Q_p T_p^4).$$

Multiplication proves the same equality for every later power. Subtracting the shifted square and cubic profiles gives

$$\begin{aligned} v(T_p^3) \setminus v(Q_p T_p^2) &= 12s + ([2p + 1, 3p - 1] \cup [5p - 1, 6p - 2]) \cup \{17s - 1\}, \\ v(T_p^4) \setminus v(Q_p T_p^3) &= \{17s - 1\}. \end{aligned}$$

Their cardinalities are $2p$ and 1 . Theorem 13.1 gives $14p$ at the preceding stage. Finally, subtracting $v(Q_p) = 4s + \Gamma_p$ from Theorem 12.1 leaves eight disjoint blocks of sizes

$$2p - 1, \quad 5p, \quad 3p, \quad 2p, \quad 3p, \quad 6p, \quad 2p - 1, \quad 1,$$

whose sum is $23p - 1$. The strict nonzero quotients and terminal equality prove reduction number four. Since T_p is maximal-ideal primary in dimension one, the one-generated reduction is minimal.

The parameter ideal Q_p has colength $4s = 24p$. Multiplication by its nonzerodivisor generator gives

$$\ell(T_p^i / T_p^{i+1}) = 24p - \ell(T_p^{i+1} / Q_p T_p^i).$$

Summing through $i = n - 1$ and using $(23p - 1) + 14p + 2p + 1 = 39p$ proves the Hilbert formula and coefficients. $\square$

Two exact representations confirm every displayed set and length for all $p = 4, \dots, 300$ [MV]. A separate tail-set implementation rehashes all 297 rows, reconstructs six parameters, and rejects deleted-terminal, false-stabilization, altered-interval, and perturbed-length controls. The campaign and audit aggregates are

e9c3c887648f08cf67c614b381f00c8c6520dcd1bb89f8cdece62293bfd06030

and

0f6ed70676ffb8972b8b167ad52c4f9d2851f69c3b1d96f4023e5e3d5825c781.

15 The depth-zero conductor tangent cone

Let

$$G_p = \text{gr}_{T_p}(R_p) = \bigoplus_{n \geq 0} T_p^n / T_p^{n+1}$$

and retain the minimal reduction $Q_p = (x)$, where $x = t^{4s}$. The Valabrega–Valla criterion detects regularity of the initial form x^* through the intersections $Q_p \cap T_p^{n+1}$ [12].

Theorem 15.1 (Exact tangent-cone defect, [D]). *For every integer $p \geq 4$, put*

$$K_p = \{2p - 1, 4p - 1\} \cup [4p + 1, 5p - 2].$$

Then the complete Valabrega–Valla module is concentrated in degree one:

$$v\left(\frac{Q_p \cap T_p^2}{Q_p T_p}\right) = 9s + K_p, \quad \ell_{R_p}\left(\frac{Q_p \cap T_p^2}{Q_p T_p}\right) = p,$$

while

$$Q_p \cap T_p^{n+1} = Q_p T_p^n \quad (n = 0 \text{ and every } n \geq 2).$$

Consequently

$$\dim G_p = 1, \quad \text{depth } G_p = 0,$$

so G_p is not Cohen–Macaulay. Its Hilbert series is

$$H_{G_p}(z) = \frac{(p+1) + (9p-1)z + 12pz^2 + (2p-1)z^3 + z^4}{1-z}.$$

In particular every coefficient of the numerator is positive.

Proof. Write $D = [0, s-1]$ and $U = A_p \cup B_p$. Directly from the residue definitions,

$$D \setminus U = K_p, \quad |K_p| = p.$$

The exact profiles from Theorems 12.1 and 13.1 give

$$\begin{aligned} v(Q_p) &= \{4s\} \cup (8s + A_p) \cup (9s + D) \cup (10s + B_p) \cup (12s + C_p) \cup [13s, \infty), \\ v(Q_p T_p) &= (8s + A_p) \cup (9s + U) \cup (10s + B_p) \cup (12s + C_p) \cup [13s, \infty), \\ v(T_p^2) &= (8s + C_p) \cup ([9s, \infty) \setminus \{13s - 1\}). \end{aligned}$$

Since $A_p \subset C_p$ and $C_p \subset [0, s-2]$, intersection leaves exactly the additional block $9s + (D \setminus U) = 9s + K_p$ beyond $v(Q_p T_p)$. This proves the first formula and its cardinality.

Theorem 14.1 gives

$$v(T_p^3) = [12s, \infty) \setminus \{13s - 1\}, \quad v(T_p^4) = [16s, \infty), \quad v(T_p^5) = [20s, \infty).$$

Intersecting each with $v(Q_p)$ gives respectively $v(Q_p T_p^2)$, $v(Q_p T_p^3)$, and $v(Q_p T_p^4)$. For every $n \geq 4$, stabilization $T_p^{n+1} = Q_p T_p^n \subset Q_p$ proves the intersection equality immediately. The case $n = 0$ is $Q_p \cap T_p = Q_p$.

The ring R_p is a one-dimensional Cohen–Macaulay local domain with infinite residue field, x is regular, and Q_p is a minimal reduction. Hence x^* is a homogeneous parameter of G_p . The

nonzero intersection defect makes x^* a zero divisor by the Valabrega–Valla criterion, so the one-dimensional ring G_p has depth zero.

Finally write

$$a_n = \ell(T_p^{n+1}/Q_p T_p^n).$$

Theorem 14.1 gives

$$(a_0, a_1, a_2, a_3, a_4, \dots) = (23p - 1, 14p, 2p, 1, 0, \dots).$$

Since $\ell(T_p^n/xT_p^n) = \ell(R_p/xR_p) = 24p$, the natural exact sequence gives

$$h_n := \ell(T_p^n/T_p^{n+1}) = 24p - a_n.$$

Thus $(h_0, h_1, h_2, h_3, h_4, \dots) = (p + 1, 10p, 22p, 24p - 1, 24p, \dots)$, and multiplication of $\sum h_n z^n$ by $1 - z$ gives the displayed numerator. $\square$

Two exact implementations verify every intersection, length, and Hilbert coefficient for all $p = 4, \dots, 300$ [MV]. A separately written bounded-bitset audit reconstructs six parameters and rehashes all 297 campaign rows. Deleted-witness, false-later-defect, false-Cohen–Macaulay, and perturbed-numerator controls are rejected. The campaign and audit aggregates are

$$9631c644732f0921be3b3027e18a01110f23dad897fbf2cb14dd3a493eda5971$$

and

$$7c2abcd290bc3461fc5251bc3372e20a7fa25c888e3b2ed635368b4dda0781ff.$$

15.1 Complete graded torsion and Buchsbaumness

Let $\mathcal{M}_p = (\mathfrak{m}_p/T_p) \oplus G_{p,+}$ be the homogeneous maximal ideal of G_p . The dimension-one local-cohomology criterion says that G_p is Buchsbaum precisely when $\mathcal{M}_p H_{\mathcal{M}_p}^0(G_p) = 0$ [16, 17]. The cited semigroup formulas concern maximal-ideal filtrations; here we compute the local cohomology directly for the conductor filtration.

Theorem 15.2 (Buchsbaum tangent cones with unbounded invariant, [D]). *For every integer $p \geq 4$, the complete zeroth local cohomology is concentrated in degree zero:*

$$H_{\mathcal{M}_p}^0(G_p) = \text{span}_{\mathbb{k}}\{t^{5s+h} + T_p : h \in K_p\} \cong \mathbb{k}^p.$$

Moreover

$$\mathcal{M}_p H_{\mathcal{M}_p}^0(G_p) = 0.$$

Consequently G_p is Buchsbaum but not Cohen–Macaulay, and its Buchsbaum invariant is

$$I(G_p) = \ell(H_{\mathcal{M}_p}^0(G_p)) = p.$$

The quotient $G_p/H_{\mathcal{M}_p}^0(G_p)$ is Cohen–Macaulay and has Hilbert series

$$\frac{1 + (10p - 1)z + 12pz^2 + (2p - 1)z^3 + z^4}{1 - z}.$$

Proof. A homogeneous class represented by $a \in T_p^n$ lies in zeroth local cohomology exactly when $aT_p^k \subseteq T_p^{n+k+1}$ for some k . Thus

$$H_{\mathcal{M}_p}^0(G_p)_n = \bigcup_{k \geq 1} \frac{(T_p^{n+k+1} : T_p^k) \cap T_p^n}{T_p^{n+1}}.$$

Here saturation by $G_{p,+}$ and by $\mathcal{M}_p$ agree because the maximal ideal of the Artinian degree-zero ring R_p/T_p is nilpotent.

Theorem 14.1 gives $v(T_p^k) = [4ks, \infty)$ for every $k \geq 4$. Therefore, for a monomial t^v and any $k \geq 4$,

$$t^v T_p^k \subseteq T_p^{n+k+1} \iff v \geq 4(n+1)s.$$

Any containment at a smaller power persists after multiplication to a power at least four, so this stable threshold is the complete colon saturation.

The exact profiles give

$$v(R_p) \setminus v(T_p) = \{0\} \cup (5s + K_p).$$

The degree-zero threshold $4s$ deletes the unit and retains $5s + K_p$. In degrees one, two, and three, the consecutive power profiles agree from the respective thresholds $8s$, $12s$, and $16s$ onward. For every $n \geq 4$, stabilization makes them agree from $4(n+1)s$ onward. Hence the displayed p degree-zero classes are the complete local cohomology.

It remains to test the full homogeneous maximal ideal. If $r = 5s + h$ with $h \in K_p$, every product of t^r with $\mathfrak{m}_p$ or T_p has value at least $9s$. Both T_p and T_p^2 contain all such values except $13s - 1$. Equality $r + a = 13s - 1$ would force $a = 8s - 1 - h$, strictly between levels $7s$ and $8s$, but the exact ring profile has no value there. Thus

$$t^r \mathfrak{m}_p \subseteq T_p, \quad t^r T_p \subseteq T_p^2.$$

The degree-zero and positive parts of $\mathcal{M}_p$ both annihilate every torsion class. The dimension-one criterion proves Buchsbaumness; Theorem 15.1 proves depth zero. The Buchsbaum invariant is therefore the displayed length p .

Quotienting by the full zeroth local cohomology leaves a one-dimensional ring of positive depth, hence a Cohen–Macaulay ring. Since the torsion Hilbert series is the constant p , subtracting it from Theorem 15.1’s Hilbert series gives the asserted numerator. $\square$

Two exact implementations verify the full colon saturation and both maximal-ideal actions for all $p = 4, \dots, 300$ [MV]. An independently written bounded-bitset audit reconstructs six parameters and rehashes all 297 rows. Unit, deleted-torsion, false-positive-degree, degree-zero-action, false-invariant, and perturbed-series controls are rejected. The campaign and audit aggregates are

854d7889d9d7b911b462e4d483e021210ae2873ae0ec0091ec30e8fb29d6dbf7

and

0b01853febc9e9754e28abcd099a7ae3a97f4cc0ab92f3a345ab2ae03cd3c68a.

15.2 The complete module over the Noether normalization

Put $F_p = \mathbb{k}[x_p]$, where $x_p = (t^{4s})^*$ has degree one. The reduction theorem makes $F_p \rightarrow G_p$ a Noether normalization. This viewpoint follows the graded-module framework of Cortadellas Benitez and Zarzuela [13, 15], while the conductor-filtration calculation below is derived directly from Theorems 14.1, 15.1, and 15.2.

Theorem 15.3 (Exact Noether-normalization module, [D]). *For every integer $p \geq 4$, there is a graded F_p -module isomorphism*

$$G_p \cong (F_p/(x_p))^p \oplus F_p \oplus F_p(-1)^{10p-1} \oplus F_p(-2)^{12p} \\ \oplus F_p(-3)^{2p-1} \oplus F_p(-4).$$

Its minimal graded F_p -resolution is

$$0 \longrightarrow F_p(-1)^p \longrightarrow F_p^{p+1} \oplus F_p(-1)^{10p-1} \oplus F_p(-2)^{12p} \oplus F_p(-3)^{2p-1} \oplus F_p(-4) \longrightarrow G_p \longrightarrow 0.$$

Consequently

$$\mathrm{pd}_{F_p}(G_p) = 1, \quad \mathrm{reg}_{F_p}(G_p) = 4, \quad a(G_p) = 3,$$

and

$$\ell(G_p/x_p G_p) = 25p = e_0(T_p) + I(G_p).$$

Proof. Theorem 14.1 gives $T_p^{n+1} = t^{4s} T_p^n$ for $n \geq 4$, so G_p is finite over F_p . Since $G_p/x_p G_p$ has finite length, F_p -torsion equals zeroth local cohomology. Theorem 15.2 therefore gives

$$\mathrm{tors}_{F_p}(G_p) = H_{\mathcal{M}_p}^0(G_p) \cong (F_p/(x_p))^p,$$

with all generators in degree zero. In particular, every torsion elementary divisor has exponent one.

The quotient $C_p = G_p/H_{\mathcal{M}_p}^0(G_p)$ is finite and torsion-free over the principal ideal domain F_p , hence graded free. Its Hilbert series from Theorem 15.2 is

$$H_{C_p}(z) = \frac{1 + (10p-1)z + 12pz^2 + (2p-1)z^3 + z^4}{1-z}.$$

Since $H_{F_p(-d)}(z) = z^d/(1-z)$, the numerator uniquely gives the five displayed free shifts. The quotient is projective, so the exact sequence with the torsion submodule splits after choosing homogeneous lifts of a free basis. This proves the decomposition.

Resolving each $F_p/(x_p)$ by multiplication by x_p gives the displayed minimal resolution. Its only first Betti number is $\beta_{1,1} = p$; its degree-zero through degree-four generator counts are $p+1, 10p-1, 12p, 2p-1, 1$. Therefore the projective dimension is one and the resolution formula for regularity gives four. Finite-length torsion has no first local cohomology, while the largest free shift is four and end $H^1(F_p(-d)) = d-1$; hence $a(G_p) = 3$.

Finally, modulo x_p every cyclic summand contributes one k -dimension. The free rank is

$$1 + (10p-1) + 12p + (2p-1) + 1 = 24p = e_0(T_p),$$

and the torsion contributes $p = I(G_p)$. This proves the parameter-section identity. $\square$

For an independent exact reconstruction, EXP-020 forms the Apéry table of $R_p, T_p, \dots, T_p^5$ modulo $24p$. Each column splits into finite and infinite strings under multiplication by x_p , recovering every cyclic summand. The recursive-power and Hilbert/torsion routes agree for all $p = 4, \dots, 300$ [MV]; a separate closed-form implementation rebuilds six parameters and rehashes all 297 rows. Exponent-two-torsion, deleted-free-summand, perturbed-Betti, false-regularity, and false-section-length controls are rejected. The campaign and audit aggregates are

02cf6f62a71de1a897cd46149e8c89d1c55bf810d28dddc02fc6c5330b9c1aed

and

c439f7e4fbd3cee983f32e5c6a27b347017c165cdf6d9fee54eb8d53ab634eac.

15.3 The conductor special fiber

Let

$$\mathcal{F}_p = \mathcal{F}(T_p) = \bigoplus_{n \geq 0} T_p^n / \mathfrak{m}_p T_p^n$$

be the special fiber. There is a natural graded-algebra surjection $G_p \rightarrow \mathcal{F}_p$ because $T_p^{n+1} \subseteq \mathfrak{m}_p T_p^n$.

Theorem 15.4 (Canonical fiber cone and exact type, [D]). *For every integer $p \geq 4$,*

$$T_p^2 = \mathfrak{m}_p T_p, \quad T_p^{n+1} = \mathfrak{m}_p T_p^n \quad (n \geq 1),$$

and the natural map induces a graded-algebra isomorphism

$$G_p / H_{\mathcal{M}_p}^0(G_p) \cong \mathcal{F}_p.$$

The ring $\mathcal{F}_p$ is one-dimensional Cohen–Macaulay, with

$$\begin{aligned} \mathcal{F}_p \cong & F_p \oplus F_p(-1)^{10p-1} \oplus F_p(-2)^{12p} \\ & \oplus F_p(-3)^{2p-1} \oplus F_p(-4) \end{aligned}$$

as a graded F_p -module. Its multiplicity is $24p$, its reduction number and regularity are four, and $a(\mathcal{F}_p) = 3$. The Artinian reduction $B_p = \mathcal{F}_p / x_p \mathcal{F}_p$ has

$$h(B_p) = (1, 10p - 1, 12p, 2p - 1, 1)$$

and graded socle dimensions

$$(0, 0, 10p, 0, 1).$$

Thus $\text{type}(\mathcal{F}_p) = 10p + 1$, and $\mathcal{F}_p$ is neither level nor Gorenstein.

Proof. Minkowski addition of the exact value blocks in Theorems 12.1 and 13.1 gives

$$v(\mathfrak{m}_p T_p) = (8s + C_p) \cup [9s, 13s - 2] \cup [13s, \infty) = v(T_p^2).$$

Indeed, the lowest block follows from $A_p + A_p = C_p \cup [6p, 8p - 4]$; the latter interval and all remaining block sums lie at level at least $9s$ and fill every ring value there. The Frobenius value $13s - 1$ remains absent. Equality of the monomial value sets proves $T_p^2 = \mathfrak{m}_p T_p$, and multiplication by T_p^{n-1} proves the higher identities.

The degree-zero kernel of $G_p \rightarrow \mathcal{F}_p$ is $\mathfrak{m}_p / T_p$, which is exactly Theorem 15.2's H^0 . In degree $n \geq 1$, the kernel is $\mathfrak{m}_p T_p^n / T_p^{n+1} = 0$. This proves the natural algebra isomorphism. Cohen–Macaulayness and the displayed free module now follow from Theorems 15.2 and 15.3; the Hilbert, multiplicity, reduction, regularity, and a -invariant claims follow immediately.

For the type, reduce modulo the regular element x_p . Direct subtraction of $\mathfrak{m}_p T_p^n + t^{4s} T_p^{n-1}$ from the exact power blocks gives the displayed h -vector. The degree-two nonsocle offsets are precisely

$$[p + 1, 2p] \cup [4p - 1, 5p - 2].$$

They multiply into the two degree-three blocks by degree-one offsets in $[1, p]$. Every other degree-two basis offset kills degree one. No degree-three basis element is socle: if its offset is d , the complementary offset $6p - 1 - d$ belongs to the degree-one basis and their product is the unique degree-four class. That class is socle because degree five vanishes. Hence the only socle dimensions are $10p$ in degree two and one in degree four, proving the final claims. $\square$

EXP-021 reconstructs the square, natural kernel, Artinian basis, and socle for all $p = 4, \dots, 300$ [MV]. The first full attempt passed through $p = 225$ but exceeded its declared budget and is preserved as inconclusive. An exact bitset implementation then completed all 297 parameters, and an independently encoded audit rebuilt six parameters and rehashed every row. The campaign and audit aggregates are

3857877586143a3be5f14852feb12bd9efbfdf7c1cde458f30e8cd689155a95b

and

1779407050b199039d3f6d808a720ea051a81ef11734fd1bccd1a76ec78c0a9c.

15.4 The defining ideal of the special fiber

We now refine Theorem 15.4 from a module decomposition to a minimal algebra presentation. All intervals in this subsection are integer intervals. Let $\mathcal{E}_{p,d}$ be the offset basis of $(\mathcal{F}_p)_d$, after subtracting the base value $4sd$. The exact power calculation gives

$$\begin{aligned} \mathcal{E}_{p,1} = & [0, p] \cup [3p, 4p - 2] \cup [6p, 8p - 2] \cup [8p, 10p - 2] \cup \{10p\} \\ & \cup [11p - 1, 12p - 1] \cup [13p + 1, 14p - 2] \cup [14p, 15p - 1] \cup \{16p\} \cup [17p - 1, 18p - 1], \end{aligned} \quad (5)$$

$$\begin{aligned} \mathcal{E}_{p,2} = & [0, 2p] \cup [3p, 5p - 2] \cup [6p, 24p - 1], \\ \mathcal{E}_{p,3} = & [0, 24p - 1] \setminus \{6p - 1\}, \quad \mathcal{E}_{p,4} = \mathcal{E}_{p,5} = [0, 24p - 1]. \end{aligned} \quad (6)$$

In particular, $|\mathcal{E}_{p,1}| = 10p$. Set

$$P_p = \mathbb{k}[X_a : a \in \mathcal{E}_{p,1}]$$

and let $\varphi_p : P_p \rightarrow \mathcal{F}_p$ send X_a to the class of t^{4s+a} . Write $\mathfrak{a}_p = \ker \varphi_p$.

Theorem 15.5 (One-cubic defining ideal, [MV]). *For every integer $p \geq 4$,*

$$\mathfrak{a}_p = ((\mathfrak{a}_p)_2, f_p), \quad f_p = X_0^2 X_{3p} - X_p^3.$$

Equivalently, the first graded Betti row over P_p is

$$\beta_{1,2}^{P_p}(\mathcal{F}_p) = 50p^2 - 17p, \quad \beta_{1,3}^{P_p}(\mathcal{F}_p) = 1, \quad \beta_{1,j}^{P_p}(\mathcal{F}_p) = 0 \quad (j \geq 4).$$

Hence $\mu(\mathfrak{a}_p) = 50p^2 - 17p + 1$, the relation type is three, and $\mathcal{F}_p$ is not Koszul.

Proof. The semigroup monomials are linearly independent. Consequently, a monomial in the variables X_a is zero in $\mathcal{F}_p$ exactly when its total offset is absent from the corresponding $\mathcal{E}_{p,d}$, and two nonzero monomials have the same image exactly when their total offsets agree. There is no linear equation. Theorem 15.4 gives $\dim_{\mathbb{k}}(\mathcal{F}_p)_2 = 22p$, so

$$\beta_{1,2}^{P_p}(\mathcal{F}_p) = \binom{10p+1}{2} - 22p = 50p^2 - 17p. \quad (7)$$

The offsets $0, p, 3p$ belong to $\mathcal{E}_{p,1}$, and $0 + 0 + 3p = p + p + p$, so $f_p \in \mathfrak{a}_p$. It is not generated by quadrics. Indeed, the only unordered pair in $\mathcal{E}_{p,1}$ with sum $3p$ is $\{0, 3p\}$, the only pair with sum $2p$ is $\{p, p\}$, and the only pair with sum zero is $\{0, 0\}$. Thus neither monomial in f_p admits a nontrivial quadratic replacement, and the two cannot be joined by a chain of quadratic moves.

It remains to prove that this is the only higher equation. Suppose the complete kernel is known through degree $d - 1$, where $d \in \{3, 4, 5\}$. For a fixed total t , form the graph

$$V_{d,t} = \{a \in \mathcal{E}_{p,1} : t - a \in \mathcal{E}_{p,d-1}\}.$$

Two vertices a, c are adjacent when

$$t - a - c \in \mathcal{E}_{p,d-2}.$$

This is exactly the commutative factorization move between X_a times the preceding class of offset $t - a$ and X_c times the class of offset $t - c$. Add a zero vertex when the same factorization makes the alternate degree- $(d - 1)$ remainder invalid. Therefore the degree- d quotient by all lower equations has one basis element for each nonzero graph component. It follows that

$$\beta_{1,d}^{P_p}(\mathcal{F}_p) = \#\{\text{nonzero state components in degree } d\} - |\mathcal{E}_{p,d}|. \quad (8)$$

For degree three, the exact component reduction is

Total range	common hub	path length at most
$[0, 2p]$	0	3
$[2p + 1, 3p - 1]$	$t - 2p$	3
$[3p + 1, 5p - 2]$	0	3
$[5p - 1, 6p - 2]$	$t - (5p - 2)$	3
$[6p, 24p - 1]$	0	3

Every invalid total reaches the zero vertex in at most one move. At the omitted total $t = 3p$, the vertices are exactly $0, p, 3p$; the vertices 0 and $3p$ are joined, while p is isolated. Hence there are exactly two components, represented by the two monomials in f_p . Adjoining f_p joins them.

For degree four, zero is a hub except at $t = 6p - 1$, where p is a hub. To see the only nontrivial reduction, put

$$H = [2p + 1, 3p - 1] \cup [5p - 1, 6p - 2] = [0, 6p - 2] \setminus \mathcal{E}_{p,2}.$$

If $t - a \in \mathcal{E}_{p,2}$, the state is adjacent to zero. If $t - a \in H$, it moves first to p , then either to zero or through $t - 3p$, which belongs to $[1, p - 1]$ or $[3p, 4p - 2]$. The apparent boundary $t = 7p - 1$, where $t - p$ would be the hole in $\mathcal{E}_{p,3}$, cannot occur with both $t - a \in H$ and $a \in \mathcal{E}_{p,1}$. At $t = 6p - 1$, the vertex set is $[1, p] \cup [3p, 4p - 2]$, connected through p . Invalid high totals die directly. Thus the degree-four defect is zero.

For degree five, $\mathcal{E}_{p,4}$ is full and $\mathcal{E}_{p,3}$ misses only $h = 6p - 1$. A state has a direct edge to zero unless $t - a = h$. In the exceptional case, moving through the generator 1, or through 2 when $a = 1$, gives a state with a direct zero edge. Hence the degree-five defect is also zero.

The degree-three table and every boundary assertion in the degree-four and degree-five reductions were encoded as negated Presburger formulas over all integers $p \geq 4$ [MV]. The exact solver closed all 133 terminal formulas as UNSAT, after four affine-cell bisections, with no SAT or unresolved result. A separately encoded total-by-total graph rehashed the complete campaign and rebuilt three spaced parameters. Thus (8) gives one cubic and no new equation in degrees four or five.

Finally, T_p has reduction number four by Theorem 14.1, and $\mathcal{F}_p$ is Cohen–Macaulay by Theorem 15.4. The defining-ideal construction of Abdolmaleki–Kumashiro [14, Theorem 2.8] therefore supplies a complete set of equations through degree five. No higher minimal equation can occur. This proves the Betti row and relation type. A standard graded Koszul algebra has a quadratic defining ideal, while f_p is a necessary minimal cubic, so $\mathcal{F}_p$ is not Koszul. $\square$

The bounded state-graph campaign passes every $p = 4, \dots, 23$ [MV]. Its first requested extension through $p = 24$ completed that row but crossed the declared five-minute budget, and is preserved as inconclusive. The successful campaign, independent audit, and Presburger-query aggregates are respectively

d23792c47a2e07785a27ebc71e99619705f7aa53a38ebe7f66ffa03b0518ce83,
a27b3b13fde197b1f011bf07dc2c321d84ab7c895c9aa02d7c2a073e48f18038,
and
832c8421fe66359b8c246e3465e27de6ea7829215f892ab815e72b1f44787194.

15.5 Homological edges over the presentation ring

The h-vector and socle in Theorem 15.4, together with the complete first Betti row in Theorem 15.5, force more of the minimal resolution than the defining equations alone suggest. Put

$$N = 10p, \quad c = N - 1 = 10p - 1,$$

and write $\beta_{i,j} = \beta_{i,j}^{P_p}(\mathcal{F}_p)$.

Theorem 15.6 (Homological edge data, [D]). *For every integer $p \geq 4$,*

$$\mathrm{pd}_{P_p}(\mathcal{F}_p) = c, \quad \mathrm{reg}_{P_p}(\mathcal{F}_p) = 4,$$

and

$$\sum_{i,j} (-1)^i \beta_{i,j} z^j = (1 - z)^c (1 + (10p - 1)z + 12pz^2 + (2p - 1)z^3 + z^4). \quad (9)$$

The first linear-syzygy count is

$$\beta_{2,3} = \frac{2p(500p^2 - 330p + 31)}{3}. \quad (10)$$

The complete last homological row and a penultimate extremal entry are

$$\beta_{c,c+2} = 10p, \quad \beta_{c,c+4} = 1, \quad \beta_{c,j} = 0 \ (j \notin \{c+2, c+4\}), \quad \beta_{c-1,c+3} = 8p. \quad (11)$$

Finally, the canonical module $\omega_{\mathcal{F}_p}$ has $10p$ minimal generators in degree -1 and one in degree -3 .

Proof. The polynomial ring P_p has depth N , whereas Theorem 15.4 gives $\mathrm{depth}_{P_p}(\mathcal{F}_p) = \dim \mathcal{F}_p = 1$. Auslander–Buchsbaum therefore gives projective dimension $N - 1 = c$. The same theorem gives the h-polynomial

$$h_p(z) = 1 + (10p - 1)z + 12pz^2 + (2p - 1)z^3 + z^4.$$

Since $\mathcal{F}_p$ is Cohen–Macaulay, its regularity is $\deg h_p = 4$. In particular, regularity over the full presentation ring is already determined, rather than a separate open question. Taking Hilbert series in a minimal P_p -resolution gives

$$\frac{h_p(z)}{1 - z} = \frac{\sum_{i,j} (-1)^i \beta_{i,j} z^j}{(1 - z)^N},$$

which proves (9).

There is no linear equation, so minimality gives no shift below $i + 1$ in homological degree i . Consequently the coefficient of z^3 in (9) is $\beta_{2,3} - \beta_{1,3}$. Theorem 15.5 gives $\beta_{1,3} = 1$, and expansion yields

$$\beta_{2,3} = 1 - \binom{c}{3} + c \binom{c}{2} - 12pc + (2p - 1) = \frac{2p(500p^2 - 330p + 31)}{3}.$$

Independently, the degree-three part of the first free module has dimension $N\beta_{1,2} + \beta_{1,3}$ and maps onto $(\mathfrak{a}_p)_3$. Since

$$\dim(P_p)_3 = \binom{N+2}{3}, \quad \dim(\mathcal{F}_p)_3 = 24p - 1,$$

its kernel has dimension

$$N(50p^2 - 17p) + 1 - \left(\binom{N+2}{3} - (24p - 1) \right),$$

which simplifies to the same value.

Now reduce modulo X_0 , whose image x_p is the regular parameter from Theorem 15.4. Because X_0 is regular on both P_p and $\mathcal{F}_p$, tensoring the minimal resolution with $Q_p = P_p/(X_0)$ remains exact and minimal. Hence its Betti numbers equal those of the Artinian reduction $B_p = \mathcal{F}_p/x_p\mathcal{F}_p$ over the polynomial ring Q_p in c variables. Top Koszul homology identifies

$$\mathrm{Tor}_c^{Q_p}(B_p, \mathbb{k})_j \cong \mathrm{Soc}(B_p)_{j-c}.$$

Theorem 15.4 gives socle dimensions $10p$ in degree two, one in degree four, and zero otherwise. This proves the complete last row in (11).

In internal degree $c + 3$, regularity four kills every contribution in homological degree at most $c - 2$, while the last-row calculation kills $\beta_{c,c+3}$. Thus the coefficient of z^{c+3} in (9) is $(-1)^{c-1}\beta_{c-1,c+3}$. Only the last two terms of h_p contribute, giving

$$(-1)^c(2p - 1) + (-1)^{c-1}c = (-1)^{c-1}8p.$$

Therefore $\beta_{c-1,c+3} = 8p$.

Finally,

$$\omega_{\mathcal{F}_p} = \mathrm{Ext}_{P_p}^c(\mathcal{F}_p, P_p(-N)).$$

Minimality lets us read its minimal generators from the dual of the last free module. A summand $P_p(-j)$ contributes in degree $N - j$. The $10p$ last summands with $j = c + 2 = N + 1$ therefore give degree -1 , and the single summand with $j = c + 4 = N + 3$ gives degree -3 . $\square$

EXP-024 checks both derivations of (10), every selected coefficient in (9), and the socle-to-last-row transfer for all $p = 4, \dots, 300$ [MV]. Its independent implementation rebuilds all 297 rows, reconstructs six spaced EXP-021 socle cases, cross-checks all twenty persisted EXP-023 first rows, and rejects seven adversarial corruptions. The campaign and audit aggregates are

baf6200a442be9476cd083fde753bbdd9e623c06aa2528f3a7f138ee825637eb

and

b6035f615f2b2092351b5a42e5a734c72ba4783adf82943ed41b38fe07ef17e2.

The symbolic argument above, not the finite campaign, proves the theorem. Equation (9) is alternating data and does not determine the unresolved interior of the Betti table.

15.6 The first interior Betti strand

We retain the offset in (5) by declaring $\deg X_a = (1, a)$. Write $\beta_{i,(j,b)}$ for the resulting bigraded Betti number and $\beta_{i,j} = \sum_b \beta_{i,(j,b)}$. Let

$$\mathfrak{q}_p = ((\mathfrak{a}_p)_2), \quad \mathfrak{a}_p = (\mathfrak{q}_p, f_p)$$

as in Theorem 15.5.

Theorem 15.7 (First interior Betti strand, [D]). *For every integer $p \geq 4$ and every field $\mathbb{k}$,*

$$\beta_{2,4} = 8p, \quad \beta_{3,4} = \frac{p(5p-1)(500p^2 - 440p + 47)}{2}. \quad (12)$$

Moreover, $\beta_{2,(4,b)} = 1$ exactly for

$$\begin{aligned} \mathcal{S}_p = & [9p, 11p-2] \cup [11p, 13p-2] \cup \{13p\} \cup [14p-1, 15p-1] \\ & \cup [16p+1, 17p-2] \cup [17p, 18p-1] \cup \{19p\} \cup [20p-1, 21p-1], \end{aligned} \quad (13)$$

and it is zero at every other offset. Equivalently,

$$\mathcal{S}_p = \{3p + a : a \in \mathcal{E}_{p,1}, a \geq 6p\}.$$

The corresponding integral relative homology is free of rank one at every supported offset; in particular, the result has no characteristic exception.

Proof. We first identify the exact finite chain complexes. Put

$$H_p = \langle (1, a) : a \in \mathcal{E}_{p,1} \rangle \subseteq \mathbb{N}^2, \quad M_p = (t^{(n,u)} : u \geq 24p) \subseteq \mathbb{Z}[H_p].$$

The offset calculation (6) identifies the special fiber with the base change of $\mathbb{Z}[H_p]/M_p$. For a bidegree $w = (j, b)$, define the squarefree divisor pair

$$\begin{aligned} \Delta_w &= \{F \subseteq \mathcal{E}_{p,1} : w - (|F|, \sum F) \in H_p\}, \\ \Gamma_w &= \{F \in \Delta_w : t^{w-(|F|, \sum F)} \in M_p\}. \end{aligned}$$

The relative squarefree-divisor construction is standard for affine semigroup quotients by monomial ideals [18]; see also [19] for the semigroup-ring Betti framework and its characteristic warning. Here the identification is immediate: the bidegree- w Koszul chain group in homological degree r has one basis vector for every r -subset F satisfying

$$b - \sum F \in \mathcal{E}_{p,j-r},$$

and signed deletion is exactly the relative simplicial boundary. Hence

$$\beta_{i,w} = \dim_{\mathbb{k}} \tilde{H}_{i-1}(\Delta_w, \Gamma_w; \mathbb{k}). \quad (14)$$

We next give an integral upper bound in total degree four. Order $\mathcal{E}_{p,1}$ increasingly. In every offset block, scan the vertices in this order and pair a still-unmatched cell F with $F \cup \{v\}$ whenever both survive; within a scan use increasing dimension and lexicographic order. Every pair is a unit boundary entry, and the pivot order strictly increases along a reversed matched arrow, so this is an acyclic integral matching. Substitution of (5)–(6) leaves the following complete critical-cell list in relative dimensions zero and one. In the middle row write $b = 6p + r$.

Offset	critical 0-cells	critical 1-cells
$b \in \mathcal{S}_p$	$\{0\}$	$\{p, b - 3p\}$
$6p \leq b \leq 7p - 3$	$\{0\}, \{r + 2\}$	$\{p, 3p + r\}$
all other b	immaterial	none

In the middle row, the unique alternating path sends the critical edge to a unit multiple of $\{r + 2\} - \{0\}$, so it contributes no homology. The table follows by scanning the eleven blocks in (5): the only toggle obstructions are the gaps of $\mathcal{E}_{p,2}$, the hole $6p - 1$ of $\mathcal{E}_{p,3}$, and a gap of $\mathcal{E}_{p,1}$. All candidate edges outside the two displayed rows toggle at their first available vertex. Therefore, over every field,

$$\beta_{2,(4,b)} \leq 1 \quad (b \in \mathcal{S}_p), \quad \beta_{2,(4,b)} = 0 \quad (b \notin \mathcal{S}_p). \quad (15)$$

For the matching lower bound, we compute the linear part of the quadratic colon. Define

$$V_p = \{v \in \mathcal{E}_{p,1} : v - p, v + p \in \mathcal{E}_{p,1}\} = [7p, 8p - 2] \cup [8p, 9p - 2] \cup \{9p, 10p\}.$$

For every $a \in \mathcal{E}_{p,1}$ with $a \geq 6p$, the factorizations $(0, 0, 3p, a)$ and (p, p, p, a) are connected by quadratic moves. If $a + p, a + 2p \in \mathcal{E}_{p,1}$, use

$$(0, 0, 3p, a) \sim (0, 0, p, a + 2p) \sim (0, p, p, a + p) \sim (p, p, p, a).$$

For $a = 7p - 1$, use

$$(0, 0, 3p, 7p - 1) \sim (0, 1, 3p, 7p - 2) \sim (0, 1, p, 9p - 2) \sim (1, p, p, 8p - 2) \sim (p, p, p, 7p - 1).$$

In every remaining case write $a = u + v$, with $u \in \mathcal{E}_{p,1}$ and $v \in V_p$, and use

$$(0, 0, 3p, a) \sim (0, 3p, u, v) \sim (p, 3p, u, v - p) \sim (p, p, u, v + p) \sim (p, p, p, a).$$

The block formula (5) verifies that these cases cover all high offsets.

They cover no low offset. For a factorization of total at most $7p - 2$, color a generator odd exactly when it lies in $[3p, 4p - 2]$. Every equality of two pair sums preserves the parity of the number of odd generators: the sole cross-block possibility replaces two middle-block generators by one low and one high generator. If $a \leq p$, the two factorizations of $X_a f_p$ have parities one and zero; if $3p \leq a \leq 4p - 2$, they have parities zero and one. Thus

$$(\mathfrak{q}_p : f_p)_1 = \text{span}_{\mathbb{k}}\{X_a : a \in \mathcal{E}_{p,1}, a \geq 6p\}, \quad (16)$$

which has dimension $8p$.

Apply the mapping cone to

$$0 \longrightarrow P_p/(\mathfrak{q}_p : f_p)(-3) \xrightarrow{f_p} P_p/\mathfrak{q}_p \longrightarrow \mathcal{F}_p \longrightarrow 0.$$

Every linear generator in (16) supplies a homological-degree-two summand shifted by four. No cancellation is possible: its maps to the shift-three free summand by a linear entry and to the quadratic-generator module by degree-two entries. Thus there is one nonzero class at each offset $3p + a$ in (13). Combined with (15), this proves $\beta_{2,4} = 8p$, the exact support, and freeness over $\mathbb{Z}$: the matching is integral and each mapping-cone class is primitive.

Finally, no minimal degree-four equation exists by Theorem 15.5, and minimal shifts exclude homological degree at least four from internal degree four. The coefficient of z^4 in (9) is therefore

$$\beta_{2,4} - \beta_{3,4} = -\frac{p(2500p^3 - 2700p^2 + 675p - 63)}{2}.$$

Substitution of $\beta_{2,4} = 8p$ gives the second formula in (12). $\square$

EXP-027 checks the complete offset chain profile for $p = 4, 5, 6$, with totals 32, 40, 48, and the smallest case agrees over $\text{GF}(2)$ and $\text{GF}(1000003)$ [MV]. The closed formulas and explicit quadratic paths pass every $p = 4, \dots, 300$; six all-parameter counterexample queries are UNSAT. An independently written audit reconstructs the full $p = 4$ chain profile and six spaced formula rows. The campaign, symbolic, and audit aggregates are

9bc93c41d899df4a39b85a452257d114000df84730d44f0661c19b7dd8322b63,

b7e760290880e821fbc2ef86b03279edab1543307fd5c58cbb635d9c4b765db8,

and

f30b94ce86638732a407a3bb5abb4dfde8a1258441df71a36138dd0ae129d454.

The finite campaigns validate the implementation. The relative-chain and matching arguments, together with the colon and mapping cone, prove the theorem.

15.7 The complete second Betti row

The preceding theorem leaves only internal degrees five and six unresolved in homological degree two. We retain the same offset grading and relative complex (14).

Theorem 15.8 (Complete second Betti row, [D]). *For every integer $p \geq 4$ and every field $\mathbb{k}$,*

$$\beta_{2,5} = p(2p - 3), \quad \beta_{2,6} = 0. \quad (17)$$

The degree-five offset support is

$$[3p + 2, 5p - 2] \cup [6p + 1, 8p - 3] \cup [9p, 11p - 4]. \quad (18)$$

For $0 \leq r \leq 2p - 4$, put

$$m_{\text{out}}(r) = \min \left\{ \left\lfloor \frac{r}{2} \right\rfloor + 1, \left\lfloor \frac{2p - 4 - r}{2} \right\rfloor + 1 \right\},$$

$$m_{\text{mid}}(r) = \min\{r + 1, 2p - 3 - r, p - 2\}.$$

Then

$$\begin{aligned} \beta_{2,(5,3p+2+r)} &= m_{\text{out}}(r), \\ \beta_{2,(5,6p+1+r)} &= m_{\text{mid}}(r), \\ \beta_{2,(5,9p+r)} &= m_{\text{out}}(2p - 4 - r). \end{aligned} \quad (19)$$

The integral first homology in total degree five is free abelian, and it vanishes in total degree six. Consequently the complete second row is

$$\beta_{2,3} = \frac{2p(500p^2 - 330p + 31)}{3}, \quad \beta_{2,4} = 8p, \quad \beta_{2,5} = p(2p - 3), \quad \beta_{2,6} = 0,$$

with no other nonzero internal degree.

Proof. The cumulative offset sets needed in total degrees five and six are

$$\begin{aligned}\mathcal{E}_{p,2} &= [0, 2p] \cup [3p, 5p - 2] \cup [6p, 24p - 1], \\ \mathcal{E}_{p,3} &= [0, 24p - 1] \setminus \{6p - 1\}, \\ \mathcal{E}_{p,n} &= [0, 24p - 1] \quad (n \geq 4).\end{aligned}$$

Thus a degree-five vertex, edge, or triangle at offset b occurs precisely when its residual belongs respectively to $\mathcal{E}_{p,4}, \mathcal{E}_{p,3}, \mathcal{E}_{p,2}$. In degree six, the first two residual sets are full intervals and only the triangle condition retains the hole $6p - 1$.

Apply the integral lexicographic matching used in the proof of Theorem 15.7. Substitution of the eleven blocks of $\mathcal{E}_{p,1}$ gives the following complete degree-five normal form. In each row $0 \leq r \leq 2p - 4$.

Offset	critical edge labels	surviving free rank
$3p + 2 + r$	$\{(i, j) : 1 \leq i < j \leq p, i + j = r + 3\}$	$m_{\text{out}}(r)$
$6p + 1 + r$	$\{(i, j) : 1 \leq i < j \leq p, i + j = r + 4\}$ $\cup \{(i, j) : 2 \leq i < j \leq p, i + j = r + 3\}$	$m_{\text{mid}}(r)$
$9p + r$	$\{(i, 3p + j) : 2 \leq i \leq p - 1, 0 \leq j \leq p - 2, i + j = r + 2\}$ $\cup \{(p, 4p - 2) : r = 2p - 4\}$	$m_{\text{out}}(2p - 4 - r)$
all other b	transient labels only	0

For the first row, the absent star triangles are exactly the low-block pairs of sum $r + 3$; every other chord toggles at its least available vertex. For the second row, the hole in $\mathcal{E}_{p,3}$ and the two gaps in $\mathcal{E}_{p,2}$ leave the two adjacent pair-sum diagonals displayed in the table. Their reduced Morse boundary is zero. For the third row, ordering the low-middle critical chords by the low endpoint makes the remaining triangle boundary a unit staircase. Its Smith form has

$$c_C(r) - m_{\text{out}}(2p - 4 - r) \quad \text{unit factors}$$

and $m_{\text{out}}(2p - 4 - r)$ zero factors, where

$$c_C(r) = \begin{cases} 1, & r = 2p - 4, \\ \min\{r + 1, 2p - 4 - r, p - 2\}, & r < 2p - 4. \end{cases}$$

Outside the three rows, each transient edge has a distinct least admissible triangle and hence a unit pivot. All cancellations are over $\mathbb{Z}$.

In total degree six, the edge residual set is full. A chord whose star triangle is deleted at residual $6p - 1$ pairs with the next admissible vertex in its interval block; at a block endpoint the first vertex of the next block supplies the pair. Endpoint substitution leaves no critical edge. Hence the integral first homology is zero in every degree-six offset.

It remains to count the zero factors. The first diagonal consists of unordered pairs in $[1, p]$, and therefore

$$\sum_{r=0}^{2p-4} m_{\text{out}}(r) = \binom{p}{2}.$$

Across all r , the two middle diagonals contain every pair in $[1, p]$ except $(1, 2)$, and every pair in $[2, p]$. Thus

$$\sum_{r=0}^{2p-4} m_{\text{mid}}(r) = \left(\binom{p}{2} - 1 \right) + \binom{p-1}{2} = p(p-2).$$

The reflected third block contributes another $\binom{p}{2}$, proving $\beta_{2,5} = p(2p - 3)$. Unit Smith factors prove integral freeness and therefore characteristic independence. Finally, no linear equations force the minimum second-syzygy shift to be three, while $\text{reg } \mathcal{F}_p = 4$ forces it to be at most six. The known degree-three and degree-four entries and (17) therefore exhaust the row. $\square$

EXP-028 checks all formula identities for $p = 4, \dots, 300$ and complete degree-five offset profiles for $p = 4, 5, 6$, whose totals are 20, 35, 54 [MV]. At $p = 4$, complete degree-five and degree-six profiles agree over $\text{GF}(2)$ and $\text{GF}(1000003)$. An independent SymPy route obtains the same rational ranks and unit Smith factors without importing the canonical rank code. The canonical campaign aggregate is

$$45f08e6a15e321512629fa4b6ab07161ddcc766ddf56e1d9579175f3444ec32f.$$

These calculations test the implementation and endpoints; the integral matching and unit Smith form prove the theorem.

15.8 The colon–Koszul degree-five diagonal

Put

$$H_p = \{a \in \mathcal{E}_{p,1} : a \geq 6p\}. \quad (20)$$

Equation (16) gives $|H_p| = 8p$. The next theorem reads the second Koszul wedges of this exact linear colon inside the total-degree-five relative complex.

Theorem 15.9 (Colon–Koszul degree-five diagonal, [D]). *For every integer $p \geq 4$, every field $\mathbb{k}$, and every offset b ,*

$$\beta_{3,(5,b)} = \#\{\{a, c\} \subset H_p : a < c, a + c = b - 3p\}. \quad (21)$$

Consequently,

$$\beta_{3,5} = \binom{8p}{2} = 4p(8p - 1), \quad (22)$$

and its offset support is

$$[15p + 1, 39p - 3] \setminus \{33p - 1\}. \quad (23)$$

The corresponding integral relative homology is free abelian. The complete internal-degree-five diagonal is therefore

$$\begin{aligned} \beta_{2,5} &= p(2p - 3), \\ \beta_{3,5} &= 4p(8p - 1), \\ \beta_{4,5} &= \frac{2p(5p - 1)(10p - 3)(100p^2 - 110p + 13)}{3}, \end{aligned} \quad (24)$$

with $\beta_{i,5} = 0$ in every other homological degree.

Proof. The eleven-block formula for $\mathcal{E}_{p,1}$ gives the disjoint decomposition

$$\begin{aligned} H_p &= [6p, 8p - 2] \cup [8p, 10p - 2] \cup \{10p\} \cup [11p - 1, 12p - 1] \\ &\quad \cup [13p + 1, 14p - 2] \cup [14p, 15p - 1] \cup \{16p\} \cup [17p - 1, 18p - 1]. \end{aligned} \quad (25)$$

Its block lengths sum to $8p$. The linear strand of $P_p/(\mathfrak{q}_p : f_p)$ is the Koszul strand on these coordinate variables. Its second wedges are indexed by unordered distinct pairs $\{a, c\} \subset H_p$, and

the cubic shift gives bidegree $(5, 3p + a + c)$. We prove that these candidate classes are complete and primitive rather than assuming that the full mapping cone is minimal.

For fixed b , the total-degree-five relative complex from (14) contains a cell F of size s exactly when

$$b - \sum_{a \in F} a \in \mathcal{E}_{p, 5-s}.$$

Apply the same increasing-vertex integral matching as in Theorem 15.8, now through cells of size four. Every matched incidence is a unit. The only failed toggles come from

$$[2p + 1, 3p - 1], \quad [5p - 1, 6p - 1], \quad \{6p - 1\},$$

the gaps of $\mathcal{E}_{p,2}$ and $\mathcal{E}_{p,3}$. Substitution of the eleven blocks of $\mathcal{E}_{p,1}$ gives the following integral normal form. If

$$K_b = \{(p, a, c) : a < c, \ a, c \in H_p, \ a + c = b - 3p\},$$

then, after contractible unit cancellations, the part bearing on second homology is

$$0 \longrightarrow \mathbb{Z}^{T_b} \xrightarrow{(I,0)} \mathbb{Z}^{T_b} \oplus \mathbb{Z}^{K_b} \longrightarrow 0. \quad (26)$$

Here T_b indexes the transient critical triangles. Indeed, a critical triangle not of the form (p, a, c) with $a, c \in H_p$ has a least admissible tetrahedral filler in the same block of (25), or at the first vertex of the next block. Ordering first by the failed toggle and then lexicographically makes this boundary submatrix upper triangular with unit diagonal. For (p, a, c) , the residual is $2p$, the right endpoint of the first component of $\mathcal{E}_{p,2}$; the next toggle crosses one of the three displayed gaps, so no filler remains. The least-toggle order prevents a transient boundary from entering a pair row. These two cases exhaust the block endpoints and give (26). Thus

$$H_2(K_{(5,b)}; \mathbb{Z}) = \mathbb{Z}^{K_b},$$

which proves (21), primitivity, and characteristic independence.

It remains to identify the support. The following pair families already cover every possible sum except one; the letters denote the corresponding nonsingleton blocks of (25).

Pair	Sum interval	Pair	Sum interval
AA	$[12p + 1, 16p - 5]$	AB	$[14p, 18p - 4]$
AD	$[17p - 1, 20p - 3]$	BD	$[19p - 1, 22p - 3]$
BE	$[21p + 1, 24p - 4]$	AK	$[23p - 1, 26p - 3]$
BK	$[25p - 1, 28p - 3]$	DK	$[28p - 2, 30p - 2]$
EK	$[30p, 32p - 3]$	FK	$[31p - 1, 33p - 2]$
GK	$[33p - 1, 34p - 1]$	KK	$[34p - 1, 36p - 3]$

For $p \geq 4$ successive intervals overlap or meet, except between DK and EK , where the sole missing integer is $30p - 1$. Direct substitution in the complementary gaps of (25) shows that no omitted block pair supplies this value. Hence the unshifted support is $[12p + 1, 36p - 3] \setminus \{30p - 1\}$, proving (23). Summing (21) over b counts all unordered pairs and proves (22).

Finally, Theorem 15.8 gives $\beta_{2,5} = p(2p - 3)$. No degree-five minimal equation exists, and the minimum-shift bound excludes homological degree at least five from internal degree five. Taking the coefficient of z^5 in (9) therefore gives

$$[z^5](9) = \beta_{2,5} - \beta_{3,5} + \beta_{4,5}.$$

Substitution and expansion yield the third formula in (24); one of its displayed factors is divisible by three in every residue class of p modulo three. $\square$

EXP-029 checks all pair profiles, endpoints, totals, and Hilbert-coefficient identities for $p = 4, \dots, 300$ [MV]. Complete exact relative H_2 profiles at $p = 4, 5, 6$ have totals 496, 780, 1128, and the complete $p = 4$ profile agrees over $\text{GF}(2)$ and $\text{GF}(1000003)$. An independent rational route reconstructs the left support boundary, and a constant-memory arithmetic/ $\mathbb{Z}_3$ certificate checks the support-chain and coefficient identities through $p = 10000$. The campaign, audit, and symbolic aggregates are, respectively,

7564f15534e8a29f875a367d3a324b95041e8eef836d15deac3e35130e1ad37d,

337854eef5d773c84cdd79c7734e63b295fa0337c5a1852e652559c334949b04,

and

605733497d6fb0ead97bfd25e26daaa66d546c297751960e1c427f29ff69f279.

The first symbolic implementation exceeded its route budget because it materialized every support integer; it is preserved as non-evidence. The optimized endpoint implementation checks the same obligations. These calculations validate the implementation; the integral normal form proves the theorem.

15.9 The cubic-colon idealization and degree-six third strand

The preceding theorem uses only the linear part of the cubic colon. We now identify the complete colon and read its first nonlinear strand inside total degree six. Retain H_p from (20), and put

$$L_p = [0, p] \cup [3p, 4p - 2], \quad A_i = X_i \ (0 \leq i \leq p), \quad B_j = X_{3p+j} \ (0 \leq j \leq p - 2).$$

Let $V_p = \mathbb{k}[s, t]^{(p)}$ be the p -th Veronese ring and let ω_{V_p} carry its natural grading, so $(\omega_{V_p})_d = \mathbb{k}[s, t]_{dp-2}$.

For the multigraded formula, set

$$\begin{aligned} \ell_p(T) &= \sum_{a \in L_p} T^a, & \eta_p(T) &= \sum_{a \in H_p} T^a, \\ e_{p,r}(T) &= \sum_{\substack{F \subset L_p \\ |F|=r}} T^{\sum F}, & u_{p,0}(T) &= 1, \\ u_{p,d}(T) &= \sum_{v=0}^{dp} T^v + \sum_{v=3p}^{(d+3)p-2} T^v & (d \geq 1), \end{aligned}$$

and

$$r_p(T) = \ell_p(T)u_{p,1}(T) - u_{p,2}(T) - e_{p,2}(T), \quad (27)$$

$$s_p(T) = u_{p,3}(T) - \ell_p(T)u_{p,2}(T) + e_{p,2}(T)u_{p,1}(T) - e_{p,3}(T). \quad (28)$$

Theorem 15.10 (Cubic-colon idealization and degree-six third strand, [D]). *For every integer $p \geq 4$ and every field $\mathbb{k}$,*

$$\mathfrak{q}_p : f_p = \mathfrak{q}_p + (X_h : h \in H_p), \quad |H_p| = 8p. \quad (29)$$

Moreover,

$$P_p/(\mathfrak{q}_p : f_p) \cong V_p \rtimes \omega_{V_p}, \quad (30)$$

where the right side is the square-zero canonical idealization under

$$A_i \mapsto s^{p-i}t^i, \quad B_j \mapsto \epsilon s^{p-2-j}t^j, \quad \epsilon^2 = 0.$$

It is a two-dimensional graded Gorenstein algebra with Hilbert series

$$\frac{1 + (2p-2)z + z^2}{(1-z)^2}.$$

The complete total-degree-six third-syzygy profile is

$$\beta_{3,(6,b)} = [T^{b-3p}](s_p(T) + \eta_p(T)r_p(T)). \quad (31)$$

The corresponding integral relative homology is free abelian, its support is

$$[3p+4, 29p-5] \setminus ([6p-3, 6p+1] \cup [9p-3, 9p]), \quad (32)$$

and consequently

$$\beta_{3,6} = \frac{8p(7p^2 - 12p + 2)}{3}. \quad (33)$$

Proof. Kill the variables indexed by H_p , and write $S_p = \mathbb{k}[A_0, \dots, A_p, B_0, \dots, B_{p-2}]$. The image of $\mathfrak{q}_p$ is generated by

$$\begin{aligned} A_i A_j - A_k A_l & \quad (i+j = k+l), \\ A_i B_j - A_k B_l & \quad (i+j = k+l), \\ B_i B_j & \quad \text{for all } i, j. \end{aligned} \quad (34)$$

The first two rows are the equal-offset quadratic fibers. For the third, the offset of $B_i B_j$ lies in $[6p, 8p-4] \subset H_p$, and $B_i B_j - X_0 X_{6p+i+j} \in \mathfrak{q}_p$. Conversely, no AA or AB total has a high-variable factorization or vanishes in degree two, so this list is exact.

Relations (34) give a surjection to $V_p \rtimes \omega_{V_p}$. Every nonzero degree- d monomial has a unique normal form of one of two types: an A -only class indexed by $[0, dp]$, or a class with one B , indexed by $[3p, (d+3)p-2]$. Their images are the monomial bases of $\mathbb{k}[s, t]_{dp}$ and $\epsilon \mathbb{k}[s, t]_{dp-2}$, respectively, and remain independent where the intervals overlap. This proves the idealization presentation after the high variables are killed and gives dimension $2pd$ in every positive degree, hence the displayed Hilbert series.

Equation (16) already places all X_h , $h \in H_p$, in $\mathfrak{q}_p : f_p$. In the idealization, the two monomials of f_p have nonzero components in the ring and canonical-module summands. An annihilator first kills a nonzero element of the Veronese domain and is therefore zero in the ring summand; it then kills a nonzero element on the torsion-free canonical module and is zero there as well. Thus f_p is a nonzerodivisor modulo $\mathfrak{q}_p + (X_h : h \in H_p)$, proving (29) and (30).

The two normal-form intervals give the offset Hilbert profiles $u_{p,d}(T)$. Multiplication by $\prod_{a \in L_p} (1 - zT^a)$ shows that the degree-two numerator coefficient is $-r_p(T)$ and the degree-three coefficient is $s_p(T)$. The ideal in (34) is generated by quadrics and there are no linear equations. Minimum shifts therefore identify these two coefficients with the complete low-variable $\beta_{1,2}$ and $\beta_{2,3}$ profiles. Tensoring with the Koszul complex on the $8p$ killed high variables gives $s_p(T) + \eta_p(T)r_p(T)$ in homological degree two and total degree three.

It remains to prove that the cubic shift survives without cancellation. In the total-degree-six relative complex, a cell F occurs exactly when

$$b - \sum_{a \in F} a \in \mathcal{E}_{p,6-|F|}.$$

The edge residual set is full, the triangle residual set has the single hole $6p-1$, and the tetrahedron residual gaps are $[2p+1, 3p-1]$ and $[5p-1, 6p-1]$. Apply increasing-vertex integral matching through tetrahedra, rooting each equal-sum low quadratic fiber and choosing a lexicographic spanning forest in every low degree-three reduction graph. The only primitive unmatched triangles are the low cycle labels counted by $s_p(T)$, and one high vertex times a rooted low quadratic label counted by $\eta_p(T)r_p(T)$. Every triangle with at least two high vertices receives a least tetrahedral filler in the same high block or at the first vertex of the next block. The apparent endpoint $8p-1$ is not a generator.

Ordering all other triangles first by failed residual gap, then by number of high vertices, then lexicographically makes their tetrahedron boundary block upper triangular with unit diagonal and no entry in a primitive label row. Thus the reduced integral block is

$$0 \longrightarrow \mathbb{Z}^{T_b} \xrightarrow{(I,0)} \mathbb{Z}^{T_b} \oplus \mathbb{Z}^{U_b} \longrightarrow 0,$$

where U_b has generating polynomial $[T^{b-3p}](s_p + \eta_p r_p)$. This proves (31), primitivity, and characteristic independence.

Direct expansion of the interval polynomials gives the support in (32). At $T = 1$, with $c = 2p - 2$,

$$r_p(1) = \frac{c(c+1)}{2} - 1 = p(2p-3), \quad s_p(1) = \frac{c(c-2)(c+2)}{3} = \frac{8p(p-1)(p-2)}{3}.$$

Adding $8pr_p(1)$ gives (33). □

EXP-030 checks all colon and scalar formulas for $p = 4, \dots, 300$ and complete exact degree-six profiles at $p = 4, 5, 6$, with totals (704, 1560, 2912) [MV]. The complete $p = 4$ profile agrees over GF(2) and GF(1000003). An independent idealization-Hilbert implementation matches every coefficient of all three profiles and separately reconstructs selected offsets over $\mathbb{Q}$. Its first attempt incorrectly inserted the forbidden generator $8p-1$; that invalid implementation is preserved as non-evidence, and the corrected route passes. The canonical, audit, and symbolic aggregates are

```
de439ff5cf0784b332fcf811b17217579221afca42510f755963c81ff8beaa4d,
bf5034efc37ec23edbd60d87c1eca36d437a9f9fc1e9d38f59816d8a7d3a7a16,
c519356b98ea0c76ec3d49d5f04e3512f711e601fa6491a8bf28dd337454968c.
```

The calculations validate the implementation; the colon, Hilbert-numerator, and integral normal-form arguments prove the theorem.

15.10 Integral degree-seven contraction and the complete third row

The regularity bound permits one further shift in homological degree three. Its relative complex has a simpler defect than the degree-six strand and admits a uniform integral contraction.

Theorem 15.11 (Complete third homological row, [D]). *For every integer $p \geq 4$, every field $\mathbb{k}$, and every multigraded offset b ,*

$$\beta_{3,(7,b)}(C_p) = 0. \quad (35)$$

Consequently $\beta_{3,7} = 0$, and the complete third homological row is

$$\begin{aligned} \beta_{3,4} &= \frac{p(5p-1)(500p^2-440p+47)}{2}, \\ \beta_{3,5} &= 4p(8p-1), \\ \beta_{3,6} &= \frac{8p(7p^2-12p+2)}{3}, \\ \beta_{3,7} &= 0, \quad \beta_{3,j} = 0 \quad \text{otherwise.} \end{aligned} \quad (36)$$

Its total rank is

$$\beta_3 = \frac{p(7500p^3 - 7988p^2 + 2025p - 133)}{6}. \quad (37)$$

Proof. Fix a total offset b and put $h = 6p - 1$. The relative-complex dictionary used above says that a squarefree cell F occurs in total degree seven exactly when

$$b - \sum_{a \in F} a \in \mathcal{E}_{p,7-|F|}.$$

The three cumulative offset sets governing edges, triangles, and tetrahedra are

$$\mathcal{E}_{p,5} = \mathcal{E}_{p,4} = [0, 24p - 1], \quad \mathcal{E}_{p,3} = [0, 24p - 1] \setminus \{h\}. \quad (38)$$

All cellular incidence coefficients are units over $\mathbb{Z}$.

Use vertex 0 for a partial Boolean matching. Match every edge not containing 0 with the triangle obtained by adjoining 0. Match every triangle F not containing 0 with $F \cup \{0\}$ whenever $b - \sum F \neq h$. Equation (38) makes every pair valid. The matching is acyclic: a reversed matched arrow adds 0, while deletion of any other vertex leaves 0 present and cannot lead to another lower cell of the same matching. The unmatched triangles are therefore precisely

$$0 \notin F, \quad |F| = 3, \quad b - \sum F = h. \quad (39)$$

For such a triangle choose the least $x \in \{1, 2, 3, 4\} \setminus F$ and put $T = F \cup \{x\}$. These four vertices belong to the degree-one offset set for every $p \geq 4$, and

$$b - \sum T = h - x \in [0, 24p - 1] \setminus \{h\},$$

so T is an unmatched tetrahedron in the same offset complex. If $y \in T$, then the residual of its face $T \setminus \{y\}$ is $h - x + y$, which equals h exactly when $y = x$. Thus F is the unique critical triangle face of T , and its direct boundary coefficient is $+1$ or -1 .

There is no additional gradient contribution to another critical triangle. Any other face G of T not containing 0 is paired upward with $G \cup \{0\}$; all its other triangle faces contain 0 and are paired downward with edges. At fixed b , two critical triangles cannot receive the same tetrahedron, since that tetrahedron would then have two faces satisfying (39). Hence the reduced integral boundary from unmatched tetrahedra onto unmatched triangles contains a signed identity block. It is surjective, so the integral second homology vanishes at every offset. Base change to any field proves (35).

The preceding interior-strand, colon–Koszul, and cubic-colon theorems give the entries in degrees four, five, and six. The minimum-shift bound excludes smaller degrees, and regularity four excludes degrees greater than seven. This proves (36). Adding its three nonzero entries and simplifying gives (37). $\square$

EXP-031 checks the residual identities for $p = 4, \dots, 300$, verifies every unit filler for $p = 4, \dots, 12$, and computes complete exact degree-seven profiles at $p = 4, 5$ [MV]. All 374 offsets for $p = 4$ vanish over both $\text{GF}(2)$ and $\text{GF}(1000003)$, and all 470 offsets for $p = 5$ vanish over $\text{GF}(2)$. An independent implementation reverses the filler order and matches all critical counts and offset ranges for $p = 4, \dots, 12$. The first smoke implementation keyed a tetrahedron globally instead of by offset, falsely treating reuse in different multigraded complexes as a collision; it is preserved as invalid non-evidence. The canonical, audit, and symbolic aggregates are

```
d68afb5c54ebb86abbf420c389e1cacf666071cb35f83e5d2b67eccbc354858,
0be4b659126064328b5ef14a40e488a836f874d2eed9b048d4d3f19da971346e,
e4bf2e0ae303e905efc9f985b239d059a5255b02d2ddc1d37abab5cc5cb2fc1f.
```

The computations validate the implementation; the integral matching and signed identity block prove the theorem.

15.11 The complete graded Betti polynomial of the cubic-colon quotient

The canonical idealization in Theorem 15.10 also determines every ordinary graded free-module rank and shift of the colon quotient. In this subsection put

$$c = 2p - 2, \quad m = 8p, \quad D_p = P_p / (\mathfrak{q}_p : f_p),$$

and let $S_p = \mathbb{k}[A_0, \dots, A_p, B_0, \dots, B_{p-2}]$ be the polynomial ring on the $2p$ low variables. Write $E_p = V_p \rtimes \omega_{V_p}$, so $D_p \cong E_p$ as graded algebras, but the two displayed ambient polynomial rings record different Betti tables. For a graded module M , set $B_M(x, z) = \sum_{i,j} \beta_{i,j}(M) x^i z^j$.

Theorem 15.12 (Complete cubic-colon Betti polynomial, [D]). *For every integer $p \geq 4$ and every field $\mathbb{k}$, put*

$$\lambda_{c,a} = c \binom{c}{a} - \binom{c}{a+1} - \binom{c}{a-1} \quad (1 \leq a \leq c-1). \quad (40)$$

Then

$$B_{E_p}^{S_p}(x, z) = 1 + \sum_{a=1}^{c-1} \lambda_{c,a} x^a z^{a+1} + x^c z^{c+2}, \quad (41)$$

$$B_{D_p}^{P_p}(x, z) = (1 + xz)^m B_{E_p}^{S_p}(x, z). \quad (42)$$

Equivalently, with out-of-range binomial coefficients read as zero,

$$\beta_{i,i}^{P_p}(D_p) = \binom{m}{i},$$

$$\beta_{i,i+1}^{P_p}(D_p) = \sum_{a=1}^{c-1} \lambda_{c,a} \binom{m}{i-a}, \quad (43)$$

$$\beta_{i,i+2}^{P_p}(D_p) = \binom{m}{i-c}, \quad (44)$$

and every other entry vanishes. In particular,

$$\mathrm{pd}_{P_p} D_p = 10p - 2, \quad \mathrm{reg}_{P_p} D_p = 2. \quad (45)$$

The total Betti rank is

$$\sum_{i,j} \beta_{i,j}^{P_p}(D_p) = 2^{8p} \left((2p-4)2^{2p-2} + 4 \right). \quad (46)$$

Proof. Theorem 15.10 identifies E_p as a two-dimensional standard graded Gorenstein algebra with Hilbert series

$$H_{E_p}(z) = \frac{1 + cz + z^2}{(1-z)^2}.$$

It has codimension c , no linear equations, regularity two, and a-invariant zero. Minimality therefore excludes $\beta_{i,j}$ for $i > 0$ and $j \leq i$, while regularity excludes $j - i > 2$. Gorenstein self-duality has final shift $(c+2) + 0 = c+2$ and gives

$$\beta_{i,j}^{S_p}(E_p) = \beta_{c-i,c+2-j}^{S_p}(E_p). \quad (47)$$

The row $j - i = 2$ is dual to the diagonal row, which contains only $\beta_{0,0} = 1$. Consequently its only entry is $\beta_{c,c+2} = 1$; every remaining entry lies on the linear strand.

The alternating Betti numerator is

$$(1-z)^{c+2} H_{E_p}(z) = (1 + cz + z^2)(1-z)^c. \quad (48)$$

For $1 \leq a \leq c-1$, its coefficient in degree $a+1$ is

$$(-1)^a \left(c \binom{c}{a} - \binom{c}{a+1} - \binom{c}{a-1} \right).$$

The proved resolution shape leaves only $(-1)^a \beta_{a,a+1}$ in that degree, which proves (40) and (41). The factorization

$$\lambda_{c,a} = \binom{c}{a} \frac{a(c-a)(c+2)}{(a+1)(c-a+1)}$$

proves positivity and makes self-dual symmetry visible. The endpoint-separated binomial sum gives

$$\sum_{a=1}^{c-1} \lambda_{c,a} = (c-2)2^c + 2.$$

Finally, write $P_p = S_p \otimes_{\mathbb{k}} T_p$, where T_p is the polynomial ring in the m high variables. Equation (29) gives

$$D_p = E_p \otimes_{\mathbb{k}} T_p / (\text{all high variables}).$$

Tensor the minimal S_p -resolution of E_p with the Koszul resolution over T_p . The variables are disjoint, so the tensor complex is exact, and every differential entry lies in the homogeneous maximal ideal, so it is minimal. Its Betti polynomial is multiplied by $(1+xz)^m$, which proves (42) and (43). The endpoint, regularity, and evaluation at $x = z = 1$ give (45) and (46). $\square$

EXP-032 checks all 297 parameters $p = 4, \dots, 300$, stores complete tables at $p = 4, 5, 6$, and verifies Hilbert reconstruction, Gorenstein symmetry, positivity, endpoints, total ranks, and four corrupted controls [MV]. An independent implementation reconstructs every rank from

the alternating coefficients of (48) without importing canonical code; a symbolic route checks the closed identities. Two budget-limited attempts are retained as inconclusive. The audit also rejected a wrong post-Koszul Hilbert target and a nonintegral -0.0 serialization before the final artifacts were regenerated. The canonical, audit, and symbolic aggregates are

```
907438b249b98ca9ffef689b7edb9574cdb0044cc3dd4cb52de523129f7d37ee,  
43635c8497dfe57904997326e983c7477e7320809cb2fee661c7933041f47b09,  
f696390447a3ce20397d937aa73baebf23a3c5ae249d4ad1215ff48cb710a2ae.
```

The finite routes validate the implementations. The all-parameter theorem follows from the canonical idealization, Gorenstein duality, the Hilbert numerator, and the minimal Koszul tensor argument. The theorem determines every free-module rank and shift, but not the differential matrices and not the full resolution of C_p .

15.12 The quadratic quotient and the minimal cubic mapping cone

The preceding colon theorem controls the source of the cubic mapping cone. A pullback through the same colon now bounds the target tightly enough to decide all possible cancellations. Retain $c = 2p - 2$ and $m = 8p$, put $N = 10p$, and write

$$A_p = P_p/\mathfrak{q}_p, \quad L_p = \mathfrak{q}_p : f_p, \quad U_p = D_p/f_p D_p.$$

Theorem 15.13 (Minimal cubic mapping cone, [D]). *For every integer $p \geq 4$ and every field $\mathbb{k}$,*

$$\text{depth}_{P_p} A_p = 1, \quad \text{pd}_{P_p} A_p = N - 1, \quad \text{reg}_{P_p} A_p = 2. \quad (49)$$

The mapping cone of

$$0 \longrightarrow D_p(-3) \xrightarrow{f_p} A_p \longrightarrow C_p \longrightarrow 0$$

is minimal. Equivalently,

$$B_{C_p}^{P_p}(x, z) = B_{A_p}^{P_p}(x, z) + xz^3 B_{D_p}^{P_p}(x, z). \quad (50)$$

In particular, with out-of-range binomial coefficients read as zero, the two highest regularity strands of C_p are

$$\beta_{i,i+3}^{P_p}(C_p) = \sum_{a=1}^{c-1} \lambda_{c,a} \binom{m}{i-1-a}, \quad (51)$$

$$\beta_{i,i+4}^{P_p}(C_p) = \binom{m}{i-1-c}. \quad (52)$$

The first of these strands is positive exactly for $2 \leq i \leq N - 2$, while the second is positive exactly for $2p - 1 \leq i \leq N - 1$. Their total ranks are

$$\sum_i \beta_{i,i+3}^{P_p}(C_p) = 2^m((c-2)2^c + 2), \quad \sum_i \beta_{i,i+4}^{P_p}(C_p) = 2^m. \quad (53)$$

Proof. Theorem 15.10 proves that f_p is a nonzerodivisor on $P_p/L_p = D_p$. We first claim

$$\mathfrak{q}_p = (\mathfrak{q}_p, f_p) \cap L_p. \quad (54)$$

Only the reverse inclusion needs proof. If $g = q + af_p$ belongs to the intersection, with $q \in \mathfrak{q}_p$, then $af_p \in L_p$. Regularity of f_p modulo L_p gives $a \in L_p = \mathfrak{q}_p : f_p$, and hence $af_p \in \mathfrak{q}_p$. This proves (54).

The standard intersection sequence is therefore

$$0 \longrightarrow A_p \longrightarrow C_p \oplus D_p \longrightarrow U_p \longrightarrow 0.$$

Projecting this pullback onto D_p gives

$$0 \longrightarrow K_p \longrightarrow A_p \longrightarrow D_p \longrightarrow 0, \quad K_p = \ker(C_p \longrightarrow U_p). \quad (55)$$

Since f_p is D_p -regular of degree three, Theorem 15.12 and the known special-fiber Hilbert series give

$$\begin{aligned} H_{U_p}(z) &= \frac{1 + (2p-1)z + 2pz^2 + (2p-1)z^3 + z^4}{1-z}, \\ H_{K_p}(z) &= H_{C_p}(z) - H_{U_p}(z) = \frac{8pz + 10pz^2}{1-z}. \end{aligned}$$

The linear form X_0 is regular on C_p by the flat degeneration used in the proof of Theorem 15.6; it is therefore regular on the submodule K_p . Thus K_p is one-dimensional Cohen–Macaulay, and

$$H_{K_p/X_0K_p}(z) = 8pz + 10pz^2$$

shows that $\text{reg } K_p = 2$.

The module D_p is two-dimensional Cohen–Macaulay of regularity two. The depth lemma applied to (55) gives $\text{depth } A_p = 1$, while the regularity inequality gives $\text{reg } A_p \leq 2$. Auslander–Buchsbaum yields $\text{pd } A_p = N - 1$. Moreover,

$$H_{A_p}(z) = \frac{1 + (N-2)z + (2p+1)z^2 - Nz^3}{(1-z)^2}. \quad (56)$$

The coefficient of z^{N+1} in $(1-z)^N H_{A_p}(z)$ is $-N$, because N is even. Under the bounds $\text{pd } A_p = N - 1$ and $\text{reg } A_p \leq 2$, only $(-1)^{N-1} \beta_{N-1, N+1}(A_p)$ can contribute there. Hence $\beta_{N-1, N+1}(A_p) = N$, proving equality in the regularity bound and all of (49).

In homological degree i , every source summand in a lifted comparison map from the minimal resolution of $D_p(-3)$ has shift at least $i + 3$, by Theorem 15.12. Every target summand in the minimal resolution of A_p has shift at most $i + 2$, by (49). Every comparison entry therefore has positive degree. The cone admits no scalar cancellation and is minimal, proving (50).

The A_p -summand contributes only regularity at most two. The regularity-three and regularity-four strands consequently come from, respectively, the linear and terminal rows of D_p , shifted by one homological and three internal degrees. Substitution in (43) proves (51) and (52). Positivity of the $\lambda_{c,a}$, their support, and the binomial support give the stated endpoints. Summing the convolution and using $\sum_{a=1}^{c-1} \lambda_{c,a} = (c-2)2^c + 2$ proves (53). $\square$

EXP-033 checks every parameter $p = 4, \dots, 300$, stores complete upper strands at $p = 4, 5, 6$, verifies all Hilbert identities and prior Betti anchors, and rejects six adversarial mutations [MV]. An independent implementation reconstructs all 297 coefficient arrays; a structural audit checks the kernel and the X_0 -action at 25 parameters; and a symbolic route checks the closed identities. Three exact budget stops, at $p = 102, 209, 267$, are preserved as inconclusive non-evidence. The final canonical, independent, and symbolic aggregates are

67bff9217c89f212916220e858ef5168abe2d64cdbc789488e0ce5f49204092a,
 6593291efaf092333bc42972c2f05712a151efb46f3f52ed9d28afd329585a4c,
 58ab24887c79c3c075fdefea1f38ff2e1c1ef539490f7f52359149ed2bb1a4c8.

The all-parameter conclusion is deductive. The campaigns validate its implementations and frozen premises; they do not replace the colon-intersection, depth, regularity, and grading-gap proof.

15.13 The first surviving lower-strand class

The same pullback can determine an individual class in one of the two still-unknown strands. Retain the offset grading $\deg X_g = (1, g)$, and put

$$b_p = 8p - 1, \quad F_p = \{1, \dots, p\}, \quad \tau_p = b_p + \sum_{g=1}^p g = 8p - 1 + \frac{p(p+1)}{2}. \quad (57)$$

Theorem 15.14 (First surviving lower-strand class, [D]). *For every integer $p \geq 4$ and every field $\mathbb{k}$,*

$$\beta_{p, (p+2, \tau_p)}^{P_p}(K_p) = \beta_{p, (p+2, \tau_p)}^{P_p}(A_p) = \beta_{p, (p+2, \tau_p)}^{P_p}(C_p) = 1. \quad (58)$$

Consequently,

$$\beta_{p, p+2}^{P_p}(A_p) \geq 1, \quad \beta_{p, p+2}^{P_p}(C_p) \geq \binom{8p}{p-1} + 1. \quad (59)$$

Proof. The element X_0 is regular on K_p by the preceding proof. It is regular on D_p as well: in the canonical idealization of Theorem 15.10, it acts as s^p on both the Veronese domain and its torsion-free canonical module. Sequence (55) then makes it regular on A_p , and reduction modulo X_0 preserves that exact sequence and all minimal Betti numbers. Write

$$S_p = P_p/(X_0), \quad M_p = K_p/X_0K_p.$$

The offset bases used above give

$$\begin{aligned} (M_p)_1 &= \langle u_h : h \in H_p \rangle, & H_p &= \{g \in \mathcal{E}_{p,1} : g \geq 6p\}, \\ (M_p)_2 &= \langle v_b : b \in B_p \rangle, & B_p &= [6p, 24p-1] \setminus H_p, \end{aligned}$$

with dimensions $8p$ and $10p$, respectively, and $(M_p)_d = 0$ for $d \geq 3$. For $g \in \mathcal{E}_{p,1} \setminus \{0\}$, multiplication is

$$X_g u_h = \begin{cases} v_{g+h}, & g+h \in B_p, \\ 0, & g+h \notin B_p. \end{cases} \quad (60)$$

Indeed, an offset in H_p is the image of $X_0 u_{g+h}$, while an offset outside the stable kernel interval vanishes.

Let V_p have basis e_g , indexed by the positive degree-one offsets. In standard internal degree $i+1$, the Koszul complex of M_p has only the map

$$\delta_i : \bigwedge^i V_p \otimes (M_p)_1 \longrightarrow \bigwedge^{i-1} V_p \otimes (M_p)_2, \quad (61)$$

where

$$\delta_i(e_F \otimes u_h) = \sum_{\substack{g \in F \\ h+g \in B_p}} (-1)^{\text{pos}(g, F)} e_{F \setminus \{g\}} \otimes v_{h+g}.$$

Thus the two Betti strands of K_p are the kernels and cokernels of these integral incidence maps.

The smallest element of B_p is $b_p = 8p - 1$, and the p smallest positive variable offsets are exactly F_p . Hence the codomain of δ_{p+1} in total offset τ_p is one-dimensional, on $e_{F_p} \otimes v_{b_p}$. Its possible incoming variables form

$$R_{b_p} = \{g \in \mathcal{E}_{p,1} \setminus \{0\} : b_p - g \in H_p\}.$$

For $1 \leq g \leq p$, the complement is in $[7p - 1, 8p - 2] \subset H_p$. A variable in the second low block gives a complement at most $5p - 1$, while a high variable gives one at most $2p - 1$. Therefore $R_{b_p} = F_p$. An incoming face would require an element of $R_{b_p} \setminus F_p$, so this integral cokernel is free of rank one.

It remains to exclude the connecting map from D_p . The degree-one basis of D_p/X_0D_p has low offsets

$$L'_p = [1, p] \cup [3p, 4p - 2].$$

A differential contributing to the distinguished coordinate must remove a high exterior variable $h_l = b_p - l$, where $1 \leq l \leq p$, and leaves the low chain $e_{F_p} \otimes X_l$ at low total offset $\sum F_p + l$. This is the only low chain in that multidegree. Indeed, using one exterior variable in the second block raises the exterior sum to at least $3p + 1 + \dots + (p - 1) = \sum F_p + 2p$, and using a coefficient in that block raises it by at least $3p > l$. Without the second block, a size- p exterior set must be F_p , and its coefficient is forced to be l .

The unique chain is not a cycle: removing p from its Koszul boundary gives the unit coordinate

$$(-1)^{p-1} e_{\{1, \dots, p-1\}} \otimes X_p X_l,$$

and $X_p X_l \neq 0$ in the ring summand of D_p/X_0D_p , at offset $p + l \in [p + 1, 2p]$. The high variables annihilate D_p , so its Koszul complex splits by high exterior set. Consequently every source cycle has zero coefficient on all of the selected chains, and the connecting map cannot hit the rank-one class.

The row-two strand of D_p starts at homological degree $c = 2p - 2 > p$, by Theorem 15.12. The long exact Tor sequence therefore gives the exact middle equality in (58). Finally, the multigraded minimal cone could add at this offset only a shifted diagonal wedge of $p - 1$ high variables. Its smallest offset is $3p + \sum_{h=6p}^{7p-2} h$, which exceeds τ_p by $(p - 2)(6p - 1) > 0$. This proves the last equality. Summing offsets and using the shifted diagonal rank $\binom{8p}{p-1}$ proves (59). $\square$

EXP-034 verifies the frozen premises and all 297 parameters $p = 4, \dots, 300$ [MV]. Exact rank probes pass through $p = 8$ over $\mathbf{F}_2$ and $\mathbf{F}_{1000003}$. An independent numerical-semigroup reconstruction matches the two Artinian layers through $p = 25$ and enumerates the rational unit sources through $p = 9$. Eight symbolic interval negations are UNSAT, and 299 separate arithmetic rows pass through $p = 1000$. The canonical, finite-rank, independent, and symbolic aggregates are

```
65ef176dcd9f5bd5467c09e763fdb20c67798de9743443ce5d0e34958c1645ce,
31d70c09d251bb6009b610be05c33a42ccd50e417b84aff2c0db561018e6acc5,
31479abd3c7247fe0ba464ee06e437a595812c3d6055d0de8d0ced25d12794,
b3f461298706a394cc0f1a296557e10f52435f78d2f1039452fb726871b79a4d.
```

The theorem is deductive. These campaigns audit the bases, unit pivots, interval exclusions, and adversarial mutations; they do not replace the displayed all-parameter argument.

15.14 Primitive zero rows and characteristic dependence

For $b \in B_p$, define the representation set

$$R_b = \{g \in \mathcal{E}_{p,1} \setminus \{0\} : b - g \in H_p\}. \quad (62)$$

The EXP-034 cell is the case $R_{8p-1} = \{1, \dots, p\}$. The same invariant classifies a canonical part of every incidence cokernel.

Theorem 15.15 (Zero-row summands and a characteristic-dependent cell, $[\mathbf{D}/\mathbf{MV}]$). *Let $p \geq 4$, and put $n = 10p - 1$. The coordinate $e_F \otimes v_b$ in the codomain of the two-layer incidence map is an integral zero row if and only if $R_b \subseteq F$. Consequently, in homological degree i , these coordinates split off a primitive free summand of rank*

$$z_{p,i} = \sum_{\substack{b \in B_p \\ |R_b| \leq i}} \binom{10p - 1 - |R_b|}{i - |R_b|} \quad (63)$$

from the regularity-two cokernel of K_p .

For every $2 \leq t \leq p - 2$, set

$$b_{p,t} = 10p + t, \quad F_{p,t} = [3p, 4p - 2] \cup \{t\} \cup [t + 2, p]. \quad (64)$$

Then

$$R_{b_{p,t}} = F_{p,t}, \quad |F_{p,t}| = 2p - t - 1, \quad b_{p,t} + \sum F_{p,t} = 4p^2 + 6p - \frac{t(t-1)}{2}. \quad (65)$$

Thus K_p has a primitive zero-coordinate class in every homological degree $p + 1, \dots, 2p - 3$.

At $(p, t) = (4, 2)$, the complete multigraded target satisfies

$$\beta_{5,(7,87)}^{P_4}(K_4) = \begin{cases} 5, & \mathbb{k} = \mathbf{F}_2, \\ 4, & \mathbb{k} = \mathbf{F}_3, \end{cases} \quad (66)$$

$$\beta_{5,(7,87)}^{P_4}(A_4) = \beta_{5,(7,87)}^{P_4}(C_4) = \begin{cases} 4, & \mathbb{k} = \mathbf{F}_2, \\ 3, & \mathbb{k} = \mathbf{F}_3. \end{cases} \quad (67)$$

The integral cokernel in (66) is $\mathbb{Z}^4 \oplus \mathbb{Z}/2\mathbb{Z}$. In particular, the lower multigraded Betti tables of K_4 , A_4 , and C_4 depend on the characteristic.

Proof. By (61), a coefficient can enter the row $e_F \otimes v_b$ only from

$$e_{F \cup \{g\}} \otimes u_{b-g}, \quad g \in R_b \setminus F.$$

If this set is empty, the row is zero. Conversely, any element of it supplies an incidence entry with coefficient 1 or -1 . Hence the zero-row criterion is integral. The image of the incidence map has zero projection onto the coordinate module spanned by all such rows, so that module splits off from the cokernel. For fixed b , the number of size- i supersets of R_b is the binomial coefficient in (63), proving the first claim.

Fix $2 \leq t \leq p - 2$ and $b = 10p + t$. For $1 \leq g < t$, the complement $b - g$ lies strictly between the high singleton $10p$ and the next high block beginning at $11p - 1$. The value $g = t$ gives $b - g = 10p$, while $g = t + 1$ gives the missing offset $10p - 1$. Every $t + 2 \leq g \leq p$ gives $8p \leq b - g \leq 10p - 2$. For $3p \leq g \leq 4p - 2$,

$$6p + t + 2 \leq b - g \leq 7p + t \leq 8p - 2,$$

so the entire second low block represents b . A high variable $g \geq 6p$ gives $b - g < 6p$ and contributes nothing. This proves the set equality in (65). Counting and summing its intervals gives the other two formulas.

It remains to compute the first complete target. Here

$$b = 42, \quad F = \{2, 4, 12, 13, 14\}, \quad i = 5, \quad \tau = 87.$$

The selected coordinate is not protected by the EXP-034 unit-pivot argument. With a fixed high exterior factor e_{40} , an exact ten-term low Koszul cycle has coefficient one on $e_F \otimes X_2$, and its connecting image therefore reaches $e_F \otimes v_{42}$. This refutes coordinatewise survival, but not survival of the full target quotient.

The full target has 79 kernel codomain rows and 119 kernel boundary columns. Let d_D be the boundary on all 710 degree-one source chains, J the connecting chain map, and δ_K the kernel incidence boundary. For

$$M = \begin{pmatrix} d_D & 0 \\ J & \delta_K \end{pmatrix},$$

projection of $\text{im } M$ onto $\text{im } d_D$ has kernel $\text{im } \delta_K + J(\ker d_D)$. Therefore

$$\dim \text{coker}(\text{Tor}(D_4) \rightarrow \text{coker } \delta_K) = 79 + \text{rank } d_D - \text{rank } M. \quad (68)$$

Exact elimination gives

$\mathbb{k}$	$\text{rank } \delta_K$	$\dim \text{coker } \delta_K$	$\text{rank } d_D$	$\text{rank } M$	$\text{rank } J$	$\dim A_4$
$\mathbf{F}_2$	74	5	513	588	1	4
$\mathbf{F}_3$	75	4	513	589	1	3

and the same odd-field row over $\mathbf{F}_5$ and $\mathbf{F}_{1000003}$. Smith reduction of the integral 79-by-119 matrix δ_K has rank 75, free cokernel rank four, and one nonunit invariant factor, equal to 2. This proves (66) and (67) for A_4 .

The minimal cubic cone can add to the displayed C_4 component only from a four-high-variable diagonal at shifted offset $87 - 3p = 75$. Its smallest possible offset is $24 + 25 + 26 + 27 = 102$. That source is absent, so the C_4 values equal the A_4 values. $\square$

EXP-035 checks the zero-row and interval formulas for every $p = 4, \dots, 300$ [MV]. The first failed pivot is retained as an explicit integral cycle. The complete target matrices are rebuilt independently from numerical-semigroup ideal powers, using reversed column and pivot orders; all three basis hashes and every rank agree. Nine symbolic interval negations are UNSAT. The classification aggregate and target, independent, and symbolic artifact hashes are

```
cc98154e60bdc00fe1f503020aa7d5c66b53ff0cc4ce2158f199d03c2a5fda8b,
4072a9fb7844d07763fae1b08e99da3d94d38cf3a40f980316c38f0931091276,
b92e787bc120b5fa12aac1fc4a10792883e699ed7315055958f3916e8d10b60b,
b1bfc105f3e9ace368f181ccf10f367fe1f4d23199e49c14275bd8e9b941569e.
```

The zero-row theorem is deductive for all $p \geq 4$. The characteristic-dependent theorem is the complete exact target at $p = 4$; no extension of its torsion statement to $p > 4$ is claimed.

16 Reproducibility and trust boundary

All compact code, hypotheses, manifests, audits, and verdicts are in the public repository under

[problems/commutative-algebra/huneke-wiegand/](https://github.com/caos-problems/commutative-algebra/huneke-wiegand/).

Heavy CNFs and proofs are retained outside Git with hashes committed in the manifests. The minimum-layer archive adds 12 fully accounted files and a reconstruction auditor. The record also includes the compact EXP-011–035 campaigns, independent reconstruction audits, symbolic proofs, the preserved EXP-013 correction, the EXP-015 refutation, and the EXP-022 quadratic refutation. The proof toolchain is pinned as follows:

Component	Identity
CaDiCaL	reported 1.7.3; Ubuntu package 1.7.4-1
CaDiCaL binary SHA-256	7b73df0a6d9cf3c751a1948300e5baff8e82c4d39bcd88f0c063b5f5cfb8b33e
DRAT-trim commit	2e3b2dc0ecf938addbd779d42877b6ed69d9a985
DRAT-trim binary SHA-256	92f0aa9575ed519d66a99b8b1b3dde6ece4618ae4c202a3a4b200265dda0aa7a
Z3	4.16.0; exact integer Presburger queries for EXP-023
EXP-023 query aggregate	832c8421fe66359b8c246e3465e27de6ea7829215f892ab815e72b1f44787194
EXP-024 campaign/audit	exact integer coefficient and dimension routes; 297 rows plus independent rebuild
EXP-027	relative-chain profiles, six exact Presburger queries, 297 formula rows, and independent reconstruction
EXP-028	integral degree-five/six matching, 297 formula rows, two-field profiles, and independent Smith forms
EXP-029	integral pair-basis normal form, 297 formula rows, exact small profiles, and independent boundary audit
EXP-030	canonical-idealization profile, exact degree-six homology, corrected independent audit, and symbolic support certificate
EXP-031	integral degree-seven contraction, exact small profiles, independent filler audit, and arithmetic certificate
EXP-032	complete colon Betti polynomial, 297 exact rows, independent coefficient audit, and symbolic identities
EXP-033	minimal cubic cone, 297 upper-strand rows, kernel audit, independent reconstruction, and symbolic identities
EXP-034	two-layer incidence class, 297 exact rows, two-field ranks, semi-group reconstruction, and symbolic exclusions
EXP-035	zero-row classification, complete characteristic target, Smith form, semigroup reconstruction, and symbolic exclusions

The trusted base comprises the cited tree theorem, the finite reduction proved here, the small standard-library model checker, and DRAT-trim’s verification of each proof against its exact CNF. The SAT solver need not be trusted for UNSAT. For SAT, the independently extracted model is checked without consulting solver internals.

The EXP-023 trust boundary is different. Z3’s UNSAT leaves do not carry a separately checked proof object. The exact Presburger encoding, the explicit interval reductions in the proof, the separate total-graph implementation, and the finite stress campaign reduce implementation risk, but do not remove reliance on solver soundness. The one-cubic theorem is therefore labeled **[MV]**, not presented as a proof-assistant theorem or a proof-carrying UNSAT result.

EXP-024 introduces no additional solver dependency. Its scripts verify arithmetic identities and the frozen hashes of the EXP-021/023 premises; Theorem 15.6 is a deductive consequence of those premises and standard homological arguments. Any defect in an imported premise would still propagate, and the theorem does not erase EXP-023's disclosed boundary.

EXP-027 uses Z3 only to adversarially negate six interval statements. The proof of Theorem 15.7 does not infer the theorem from those finite or solver results: the relative-chain identification, integral matching, explicit quadratic paths, parity obstruction, and minimal mapping cone are given above. The independent audit does not import the canonical rank implementation. The remaining trust boundary is the correctness of the frozen EXP-021/023/025 premises and the ordinary algebraic arguments displayed in the proof.

EXP-028 reuses the integral relative-chain identification but independently freezes its imported premises. Its finite-field ranks are audits, not the source of characteristic independence. That claim rests on the displayed unit matching and Smith normal form. The arithmetic/Z3 certificate checks endpoints and count identities without replacing the all-parameter interval argument.

EXP-029 likewise uses finite and solver checks only as adversarial validation. The theorem rests on the integral normal form (26), whose unit block proves both completeness and freeness. The arithmetic/Z3 certificate checks the pair-sum support chain and the Hilbert-coefficient identity, but does not certify the relative matching itself. The first over-budget symbolic implementation is retained explicitly as non-evidence.

EXP-030 introduces no solver dependency. Its first independent audit accidentally filled the missing generator offset (8p-1), failed against the canonical profile, and is retained as invalid non-evidence. The corrected audit matches every coefficient and selected rational ranks. The theorem rests on the nonzerodivisor proof of the canonical idealization, the multigraded Hilbert numerator, and the displayed integral unit normal form, not on the finite profiles.

EXP-031 also introduces no solver dependency. Its characteristic-independent vanishing is not extrapolated from the small finite-field profiles: it rests on the all-parameter acyclic matching and the signed identity block in Theorem 15.11. The corrected implementation keys cells by both multigraded offset and tetrahedron; the earlier tuple-only collision is retained as invalid non-evidence.

EXP-032 introduces no solver dependency. The finite campaigns test the coefficient implementation, including corrupted controls, but do not prove the theorem. The proof uses the frozen EXP-030 canonical-idealization premise, standard Gorenstein self-duality, coefficient extraction from the Hilbert numerator, and the minimal Koszul tensor product. The claim is the complete graded Betti polynomial, not explicit differential matrices or the full special-fiber resolution.

EXP-033 has no solver dependency. Its theorem imports the frozen colon, Hilbert-series, and regular-element premises from EXP-023/024/026/030/032. The proof then uses an exact colon intersection, its pullback, the depth and regularity lemmas, the Auslander–Buchsbaum formula, and the strict grading gap in the comparison map. Finite, structural, independent, and symbolic routes audit these steps but are not extrapolated into the theorem. The minimal cone fixes both upper regularity strands; it does not determine the two lower strands of the quadratic quotient.

EXP-034 uses Z3 only to negate eight elementary interval inequalities; its theorem does not rely on solver output or finite extrapolation. The proof rests on regular reduction, the displayed two-layer incidence map, uniqueness of the minimum-offset target, a characteristic-free unit source boundary, and the long exact Tor sequence. The independent route reconstructs the Artinian fiber bases from numerical-semigroup ideal powers rather than importing the canonical offset functions. The result is one exact lower-strand class, not a complete strand.

EXP-035 preserves the first failed connecting pivot rather than hiding it. The all-parameter

zero-row theorem follows directly from the displayed integral incidence formula and interval partition. Characteristic dependence is an exact finite-target statement: the canonical route uses every row and column, the independent route reconstructs the Artinian bases from semigroup ideal powers and reverses the elimination order, and Smith form identifies the single factor two. The result proves one characteristic-dependent multigraded cell at $p = 4$, not a complete lower strand or an all-parameter torsion theorem.

17 Scope, attribution, and open questions

Son Pham retains discovery priority for the counterexample. Professor Craig Huneke’s verification is independent external evidence. The contributions proved here are Frobenius minimality, uniqueness of the normalized pair at that minimum, the separate infinite family in Theorem 9.2, and its uniform endomorphism anatomy in Theorem 10.1, pseudo-Frobenius anatomy in Theorem 11.1, exact trace/conductor formula in Theorem 12.1, stability defect in Theorem 13.1, and conductor reduction/Hilbert data in Theorem 14.1, tangent-cone depth in Theorem 15.1, and complete graded-torsion/Buchsbaum anatomy in Theorem 15.2, and the complete Noether-normalization module in Theorem 15.3, and the canonical fiber cone and type in Theorem 15.4, and its complete defining ideal in Theorem 15.5, and its exact homological edges in Theorem 15.6, its first interior Betti strand in Theorem 15.7, and its complete second Betti row in Theorem 15.8, and the colon–Koszul degree-five diagonal in Theorem 15.9, and the cubic-colon idealization and degree-six strand in Theorem 15.10, and the complete third homological row in Theorem 15.11, and the complete cubic-colon Betti polynomial in Theorem 15.12, and the minimal cubic mapping cone and complete upper special-fiber strands in Theorem 15.13, and the first surviving lower-strand class in Theorem 15.14, and the zero-row and characteristic-dependence theorem in Theorem 15.15, together with their reproducible records.

Theorems 6.1, 7.2, 9.2, 10.1, 11.1, 12.1, 13.1, 14.1, 15.1, 15.2, 15.3, 15.4, 15.5, 15.6, 15.7, 15.8, 15.9, 15.10, 15.11, 15.12, 15.13, 15.14, and 15.15 do not classify all higher Frobenius values, prove a minimum multiplicity or embedding dimension across all counterexamples, or establish corresponding statements for arbitrary Gorenstein domains and modules. The following questions remain open:

1. How does the number of normalized rigid pairs grow above $F = 181$?
2. Is there a global minimum under multiplicity, embedding dimension, or number of minimal generators?
3. Which other Boolean-scaffold models lie on affine or residue-class families, and is the family in Theorem 9.2 maximal inside a natural Kunz face?
4. Which additional hypotheses on the endomorphism ring recover a positive theorem while excluding both the public example and the full family proved here?
5. Can the finite certificates be imported into a smaller formally verified checker?
6. Can the remaining incidence cokernel cells and their connecting maps be classified to determine the two complete lower strands of the quadratic quotient in (55), thereby completing the presentation-ring Betti table and the full minimal resolution of the special fiber?
7. Does the factor-two incidence torsion in Theorem 15.15 extend to infinitely many parameters or multidegrees, and can its supporting subcomplex be classified integrally?

Acknowledgments

The author thanks Son Pham for making the candidate and verification materials public, and Craig Huneke for the independent verification recorded with that release. Neither is an author of this paper, and this acknowledgment does not imply endorsement of the minimality result.

References

- [1] C. Huneke and R. Wiegand, *Tensor products of modules and the rigidity of Tor*, Math. Ann. 299 (1994), 449–476. doi:10.1007/BF01459793.
- [2] P. A. García-Sánchez and M. J. Leamer, *Huneke–Wiegand conjecture for complete intersection numerical semigroup rings*, J. Algebra 391 (2013), 114–124. doi:10.1016/j.jalgebra.2013.06.007.
- [3] V. Blanco and J. C. Rosales, *The tree of irreducible numerical semigroups with fixed Frobenius number*, Forum Math. 25 (2013), 1249–1261. doi:10.1515/form.2011.151.
- [4] M. Landeros, C. O’Neill, R. Pelayo, K. Peña, J. Ren, and B. Wissman, *Families of numerical semigroups and a special case of the Huneke–Wiegand conjecture*, Commun. Algebra 52 (2024), 5196–5206. doi:10.1080/00927872.2024.2362934.
- [5] S. Pham, *Huneke–Wiegand candidate verification*, public software and certificate repository (2026). <https://github.com/sonpham-org/huneke-wiegand-candidate-verification>.
- [6] S. Dey and J. Lyle, *Centers of endomorphism rings and reflexivity*, arXiv:2510.02210v2 (2025). <https://arxiv.org/abs/2510.02210v2>.
- [7] S. Maitra and V. Mukundan, *Extremal behavior of reduced type of one dimensional rings*, arXiv:2306.17069v1 (2023). <https://arxiv.org/abs/2306.17069v1>.
- [8] H. Lindo, S. Maitra, and W. Zhang, *Trace does not preserve reflexivity*, Collect. Math. (2026). <https://doi.org/10.1007/s13348-026-00515-0>.
- [9] J. Herzog and S. Kumashiro, *Upper bound on the colength of the trace of the canonical module in dimension one*, Arch. Math. 119 (2022), 237–246. doi:10.1007/s00013-022-01764-8.
- [10] S. Dey, *Finite birational extension with stable conductor*, arXiv:2212.09087v1 (2022). <https://arxiv.org/abs/2212.09087v1>.
- [11] D. G. Northcott and D. Rees, *Reductions of ideals in local rings*, Proc. Cambridge Philos. Soc. 50 (1954), 145–158. doi:10.1017/S0305004100029194.
- [12] P. Valabrega and G. Valla, *Form rings and regular sequences*, Nagoya Math. J. 72 (1978), 93–101. doi:10.1017/S0027763000018225.
- [13] T. Cortadellas and S. Zarzuela, *On the structure of the fiber cone of ideals with analytic spread one*, J. Algebra 317 (2007), 759–785. doi:10.1016/j.jalgebra.2007.02.044.
- [14] R. Abdolmaleki and S. Kumashiro, *Defining ideals of Cohen–Macaulay fiber cones*, Int. J. Algebra Comput. 34 (2024), no. 7, 1099–1109. doi:10.1142/S0218196724500437.
- [15] T. Cortadellas and S. Zarzuela, *Tangent cones of numerical semigroup rings*, arXiv:0906.0911 (2009).
- [16] M. D’Anna, M. Mezzasalma, and V. Micale, *On the Buchsbaumness of the associated graded ring of a one-dimensional local ring*, Commun. Algebra 37 (2009), 1594–1603. doi:10.1080/00927870802116521.
- [17] M. D’Anna, V. Micale, and A. Sammartano, *On the associated graded ring of a semigroup ring*, J. Commut. Algebra 3 (2011), 147–168. doi:10.1216/JCA-2011-3-2-147.
- [18] W. Bruns and J. Herzog, *Semigroup rings and simplicial complexes*, J. Pure Appl. Algebra 122 (1997), no. 3, 185–208. doi:10.1016/S0022-4049(97)00051-0.
- [19] D. I. Stamate, *Betti numbers for numerical semigroup rings*, in *Multigraded Algebra and Applications*, Springer Proc. Math. Stat. 238 (2018), 133–157. arXiv:1801.00153v2.
- [20] A. Biere, K. Fazekas, M. Fleury, and M. Heisinger, *CaDiCaL, Kissat, Paracooba, Plingeling and Treengeling entering the SAT Competition 2020*, with current CaDiCaL documentation and sources. <https://github.com/arminbiere/cadical>.

- [21] N. Wetzler, M. J. H. Heule, and W. A. Hunt Jr., *DRAT-trim: efficient checking and trimming using expressive clausal proofs*, SAT 2014, LNCS 8561, 422–429. doi:[10.1007/978-3-319-09284-3_31](https://doi.org/10.1007/978-3-319-09284-3_31).