The Bottom of the Shub–Smale Tau Conjecture: an Exact Census of Integer Roots for Constant-Free Straight-Line Programs of Length at Most Nine

Felipe Santibáñez-Leal
CAOS Research program, `github.com/fsantibanezleal/CAOS_RESEARCH`

5 November 2026

Preprint · v0.06, 5 November 2026 · Licence CC BY 4.0
CAOS Research open-problems programme · problem `tau-conjecture`
Version DOI 10.5281/zenodo.23980033 · concept DOI (always latest) 10.5281/zenodo.21753438
Not peer reviewed. Source code, data and computational records: github.com/fsantibanezleal/CAOS_RESEARCH

Abstract

For a univariate integer polynomial f, let $\tau(f)$ be the minimum number of $+, -, \times$ gates needed to compute f from x and the constants $-1, 0, 1$, and let $z(f)$ be its number of distinct integer roots. The Shub–Smale tau conjecture (Smale's fourth problem) asserts $z(f) \le (1+\tau(f))^{\kappa}$ for a universal κ; it implies $\mathrm{P}_{\mathbb{C}} \neq \mathrm{NP}_{\mathbb{C}}$ in the Blum–Shub–Smale model and $\mathrm{VP}^0 \neq \mathrm{VNP}^0$, and it is open even for $\kappa = 1$. We report the first exact census of the conjecture's growth function at the bottom of its ladder: writing $z_{\max}(\tau)$ for the maximum of $z(f)$ over all nonzero f with $\tau(f) \le \tau$, we prove by exhaustive, exactly verified computation that

$$z_{\max}(1), \ldots, z_{\max}(9) \;=\; 1,\ 2,\ 3,\ 3,\ 4,\ 5,\ 5,\ 6,\ 6.$$

In particular the minimum cost of 4 distinct integer roots is 5 gates, of 5 roots is 6 gates, and of 6 roots is exactly 8 gates, by an explicit verified witness: $q(q-2)(q-6)$ for $q = x^2 - x$, computable in 8 gates by chained subtraction sharing; the growth function has plateaus at $\tau = 4$, $\tau = 7$ and $\tau = 9$, so an extra gate does not always buy an extra root. No program with at most nine gates computes a polynomial with 7 distinct integer roots, and ten gates suffice, so the seven-root threshold is exactly 10. At the ninth gate a multiplicative final gate is decided through unions of known root sets, and an additive one by an exhaustive scan of all 4.19×10^{12} candidates with a 2-adic and 3-adic divisibility filter that is valid for roots anywhere in $\mathbb{Z}$, so no evaluation window is involved. The censuses are decision-complete: depth 6 required the exact construction of all 25,844,905 reachable computation states; depth 7 a complete scan of 2,013,706 new polynomials, using a "last-gate" lemma that decides one depth beyond any exhausted frontier without storing the next frontier; and depth 8 the first complete construction of the depth-7 frontier itself: 1,048,460,912 states, built out-of-core by hash-partitioned external deduplication and scanned exhaustively ($\sim 2 \times 10^{11}$ last-gate applications) by a validated multiprocess engine. The enumerator is anchored on Markström's published exhaustive census of integer targets (all fourteen anchor values reproduced exactly) and cross-checked against an independent computer-algebra root

counter on 284 polynomials. The six-root threshold is decided by a case split: the last gate of an 8-gate 6-rooter must involve the seventh value, and if it is a multiplication its root count is a union of two known root sets, so that case reduces to a co-occurrence query over the depth-6 frontier; the scan found 408 witnesses. We further prove two elementary stall theorems explaining the observed record mechanisms: for any monic $h \in \mathbb{Z}[x]$ of degree at least 2, towers built by iterating h alone have depth-independent integer root counts (for $h = x^2 - 2$, the iterates $h^{\circ k}(x) - x$ keep exactly 2 integer roots against 2^k real roots), and across the quadratic family $h_c = x^2 - c$ with $c \le 200$ the maximum tower yield is 5, attained only at $c = 2$; a second yield series at $c = m^2 + m + 1$, produced by genuine integer 2-cycles, is identified and closed by the classical fact that integer polynomial cycles have length at most 2. All computations are exact integer arithmetic, reproducible from the released code. We also report the exact ladders of the digit-restricted census (the form of the conjecture Rojas proved sufficient): the maximum number of *odd* roots grows as $1, 2, 2, 2, 2, 3, 4$ through $\tau = 7$, with its own extremal family $(x^2 - 1)(x^2 - 9)$, distinct from the unrestricted records. Finally we place the census against its two neighbouring worlds, measured on the same enumerated programs. Over $\mathbb{F}_p$ the analogous maximum is *exactly* $2^{\tau-1}$ for every τ (proved here; the extremal program is k squarings and one subtraction, and the Fermat primes that the small cases suggest are not required, only a prime $p \equiv 1 \bmod 2^k$). Over $\mathbb{R}$ it is $1, 2, 3, 4, 6, 8$ through $\tau = 6$, exhaustively, by exact Sturm counting with no floating point; its $\tau = 4$ and $\tau = 6$ records are linked by a step $g \mapsto g \cdot (g - x^2)$ that costs two gates and doubles the real root count, beating a Chebyshev tower's three gates per doubling, and that step itself stalls after three iterations. The three ladders agree at $\tau \le 2$ and separate at $\tau = 3$ and $\tau = 4$ respectively; over $\mathbb{F}_p$ the degree ceiling $2^{\tau-1}$ is saturated at every τ, over $\mathbb{R}$ it is not, and over $\mathbb{Z}$ the census stays linear. A five-gate program with six distinct *real* roots and only two integer ones exhibits the difference concretely. The work is experimental mathematics in the tradition of Markström's census; it decides nothing about the conjecture asymptotically.

1 Introduction

1.1 The conjecture

A *constant-free straight-line program* (SLP) over $\mathbb{Z}[x]$ is a sequence $(1, x, f_2, \ldots, f_N)$ in which every f_i $(i \ge 2)$ is a sum, difference, or product of two earlier entries; following [1, 8] we write $\tau(f)$ for the least $N - 1$ over all such programs with $f_N = f$. Equivalently [6], $\tau(f)$ is the minimal number of operation gates of an arithmetic circuit over the inputs x and the free constants $-1, 0, 1$. Let $z(f)$ denote the number of distinct roots of f in $\mathbb{Z}$.

Conjecture 1.1 (Shub–Smale [1]; Smale's fourth problem [2])**.** *There is a universal constant $\kappa \ge 1$ such that every nonzero $f \in \mathbb{Z}[x]$ satisfies $z(f) \le (1 + \tau(f))^\kappa$.*

The conjecture fails for $\kappa < 1$: the geometric-progression products $(x-2)(x-4)\cdots(x-2^{2^j})$ acquire one root per $O(1)$ gates [8]. It is open even for $\kappa = 1$. Its significance is structural: Shub and Smale proved that it implies $\mathrm{P}_\mathbb{C} \ne \mathrm{NP}_\mathbb{C}$ in the Blum–Shub–Smale model [1, 3], the proof reducing to the hardness of computing all sequences $(m_n \cdot n!)$; Bürgisser [5], building on Koiran [4], proved that it implies $\mathrm{VP}^0 \ne \mathrm{VNP}^0$, and that the single sequence $(n!)$ being hard to compute already yields that separation. Easy factorials would give nonuniform polynomial-time integer factoring [3, 9]; with integer division allowed, $n!$ *is* easy [11], so the restriction to $+, -, \times$ carries the entire content. The real-zeros analogue of the conjecture is false: iterating the logistic map produces 2^j real roots at cost $O(j)$ [8], which motivated Koiran's real tau conjecture for structured expressions [13, 14, 15, 16, 17]. On the lower-bound side, no nontrivial lower bound on $\tau(n!)$ is known [4]; on the reduction side, Rojas proved that a p-adic digit version

of the conjecture, for any fixed prime, already implies the full statement, via a p-independent subquadratic bound on the valuation spectrum of roots [8]. Recent work applies the conjecture to explicit exponential lower bounds [18] and tightens the bridge from BSS separations to constant-free Valiant separations [19]; the current survey reference is [6, §4.6].

1.2 What this paper does

The literature contains, to our knowledge, exactly one exhaustive computation in this model: Markström's census of *integer targets* (the straight-line complexity of small factorials and primorials), complete to program length 11 on 2013-era hardware [7]. No published computation addresses the *polynomial* side, i.e. the actual growth function of the conjecture. We define

$$z_{\max}(\tau) \;=\; \max\{\, z(f) \;:\; f \in \mathbb{Z}[x] \setminus \{0\},\ \tau(f) \le \tau \,\}$$

and determine it exactly for $\tau \le 9$, together with the full record galleries through $\tau = 7$, explicit optimal witnesses, and the exact cost thresholds for $3, 4, 5, 6$ and 7 distinct integer roots. We also prove two elementary theorems that explain the observed record mechanisms and close entire construction classes (Section 6), and we measure the quadratic tower family exhaustively (Section 7).

Our contributions:

1. **The census** (Section 4): for $\tau = 1, \ldots, 9$, the values of $z_{\max}(\tau)$ are 1, 2, 3, 3, 4, 5, 5, 6, 6, decision-complete, with per-depth state and polynomial counts, record galleries and replayed witness programs; plateaus occur at $\tau = 4$, 7 and 9.

2. **Exact thresholds**: the minimum τ admitting 3 distinct integer roots is 3 ($x^3 - x$); for 4 roots it is 5; for 5 roots it is 6; for 6 roots it is exactly 8 (Section 4.1); for 7 roots it is exactly 10 (Section 8: the complete depth-8 census excludes 8, the two halves of the ninth gate exclude 9, and two 10-gate witnesses exist).

3. **A method lemma** (Lemma 2.5): every polynomial of cost $d + 1$ is one gate over a normalized depth-d state, so $z_{\max}(d+1)$ is computable from an exhausted depth-d frontier without storing the depth-$(d + 1)$ frontier. This bought depth 7 (a $\sim 10^9$-state frontier avoided).

4. **Stall theorems** (Section 6): iterating any fixed monic map of degree ≥ 2 yields depth-independent integer root counts; for the Chebyshev-conjugate map $x^2 - 2$, whose difference-of-squares splittings generate all census records at depths 5–7, the exact stable core is $\{0, \pm 1, \pm 2\}$.

5. **The family measurement** (Section 7): across $h_c = x^2 - c$, $c \le 200$, and all tower shapes to four iterations, the maximum integer-root yield is 5, attained only at $c = 2$; two yield-4 series exist, $c = m(m + 1)$ (fixed and anti-fixed points) and $c = m^2 + m + 1$ (genuine 2-cycles), and the classical bound on integer polynomial cycle lengths shows this inventory is complete.

6. **The seven-root threshold** (Section 8): the upper bound is verified by two structurally different 10-gate programs, executed gate by gate, one folding $(q(q-2)(q-6)\cdot(x-4)$, root set the interval $\{-2, \ldots, 4\}$) and one squaring $(x^2(x^2-1)(x^2-4)(x^2-16)$, root set the tower $\{0, \pm 1, \pm 2, \pm 4\}$); the squaring shape generalizes to a family giving $2m+3$ roots in $3m+4$

gates. On the lower side, no 9-gate program whose final gate is a *multiplication* reaches 7 roots, over all 1,048,460,912 depth-7 states (maximum union size exactly 6). For an *additive* final gate we prove a window-free divisibility criterion: if f has k distinct integer roots, then on any block of p^a consecutive integers the p-adic valuations of the values of f, corrected for a proven power of x, must sum to at least a number that depends only on which values vanish modulo p^e. Applied with $p = 2$ and $p = 3$ to every one of the 4.19×10^{12} additive candidates over the same states, with every survivor built and root-counted exactly, it finds no seven-rooter. Hence $z_{\max}(9) = 6$ and the threshold is exactly 10. An earlier scan on the window $[-32, 32]$, with a sharp confinement lemma pricing what the window misses, is kept as an independent partial check.

7. **The three worlds** (Section 9): on the same enumerated programs, $z^{p}_{\max}(\tau) = 2^{\tau-1}$ exactly for every τ [D], while $z^{\mathbb{R}}_{\max}(1..6) = 1, 2, 3, 4, 6, 8$ [MV] and the integer census stays linear. What limits $\mathbb{Z}$ is not degree and not splitting, both of which cheap programs reach at exponential scale, but *distinctness*: five gates already give a completely split polynomial of degree 16 over $\mathbb{Z}$ with two distinct roots.

8. **Anchoring and cross-checks** (Section 3): the enumerator reproduces all fourteen of Markström's published anchor values; an independent computer-algebra cross-check agrees on 284 polynomials; and two general-purpose solver engines, validated on known witnesses, do not complete the smallest free-structure search within their time limits (Section 4.2).

Every claim in this paper is labeled: **[MV]** machine-verified in exact arithmetic from the released code, **[D]** proved here (elementary, self-contained), or **[C]** conjectural/observational. Nothing here decides the conjecture in either direction, and no asymptotic consequence is claimed: the census maps where the conjecture lives at the bottom of its ladder, in the same genre as [7].

2 Model, normalization, and the census method

Definition 2.1. Programs operate on values in $\mathbb{Z}[x]$. The inputs are -1, 1 and x; each gate applies $+$, $-$ or $\times$ to two (not necessarily distinct) earlier values; the length of a program is its number of gates; $\tau(f)$ is the minimal length of a program computing f at some gate.

Lemma 2.2 (free 0 is redundant)**.** *Adding* 0 *as a fourth free input does not change* τ*.* [D]

Proof. In any program using the input 0: $a + 0$ and $a - 0$ duplicate a; $a \times 0$ computes 0; $0 - a$ equals $(-1) \times a$, one gate either way. Replacing each use and deleting duplicated gates never lengthens the program. □

Lemma 2.3 (normalization)**.** *Every* f *has an optimal program in which no gate recomputes an input or an earlier value, and no gate computes* 0*.* [D]

Proof. A gate duplicating an available value can be deleted, rewiring its uses to the original; a computed 0 is usable only in the redundant patterns of Lemma 2.2. Each deletion shortens the program, so an optimal one contains neither. □

Lemma 2.4 (reached-set sufficiency)**.** *The extensions available to a program depend only on the* set *of values it has computed. Consequently a breadth-first search over reached sets ("states"),*

with set-level deduplication, visits every normalized reachable state of each depth exactly once. [D]

This is the polynomial analogue of the range-isomorphism reduction of [7]. The census algorithm is exactly this BFS, in exact integer arithmetic on dense coefficient tuples, with each new polynomial's first appearance depth recorded (that depth is $\tau(f)$, by Lemmas 2.3 and 2.4). Integer roots are counted exactly: strip the power of x, then test the divisors of the trailing coefficient (both signs).

Lemma 2.5 (last gate)**.** *Let $\mathcal{F}_d$ be the set of normalized reached states of depth d. Every f with $\tau(f) = d+1$ is $a \circ b$ for a single gate $\circ$ and operands a, b available in some state of $\mathcal{F}_d$. Hence $z_{\max}(d+1) = \max\big(z_{\max}(d),\ \max z(a \circ b)\big)$ over one op over every state of $\mathcal{F}_d$, and this maximum is computable without constructing $\mathcal{F}_{d+1}$.* [D]

Proof. Take an optimal program of length $d+1$ for f and normalize it (Lemma 2.3; deletions would shorten it, contradicting optimality, so it is already normalized). Its first d gates form a normalized reached state of depth d, and its last gate is one op over that state's available values. □

3 Anchoring and cross-checks

Markström anchor [MV]. Restricted to the integer model (start value 1, positive normalized values), our enumerator reproduces Figure 1 of [7] exactly at every compared depth: reached-set sizes $2, 4, 9, 26, 102, 562, 4363$ and complete initial intervals $2, 4, 6, 12, 40, 112, 310$ for lengths 1 through 7: fourteen of fourteen anchor values.

Independent cross-check [MV]. An independent root counter (SymPy's exact factorization) agrees with ours on all 284 polynomials checked: every polynomial of $\tau \le 3$ and every record polynomial of the censuses.

Engine re-derivation [MV]. The depth-7 computation uses a reimplemented, memoized engine, which re-derives every prior census value (state counts 9/98/1462/29506/778087, new-polynomial counts through depth 6, and the record sets) before any new value is taken from it.

Root finding for tower shapes. Divisor-based root counting is infeasible against tower constant terms of size c^{2^k}; for the family measurement only, it is replaced by direct evaluation over a window proved correct in advance (Section 7), cross-checked against the divisor method where both run.

4 The census

Theorem 4.1 (the census; MV)**.** *In the model of Definition 2.1,*

τ	1	2	3	4	5	6	7	8	9
$z_{\max}(\tau)$	1	2	3	3	4	5	5	6	6

and the censuses behind each value are decision-complete. The value at $\tau = 9$ is Corollary 8.8 of Section 8; it rests on scans of the ninth gate, not on a constructed depth-8 frontier.

The supporting exhaustive counts are:

τ	1	2	3	4	5	6	7	8
reached states	9	98	1,462	29,506	778,087	25,844,905	1,048,460,912	(scan)
new polynomials	9	34	177	1,249	11,377	134,494	2,013,706	(see text)
$z_{\max}$	1	2	3	3	4	5	5	6

Depth 7 was decided by Lemma 2.5 over the exhausted depth-6 frontier; among its 2,013,706 new polynomials, the histogram of distinct integer roots is 742,635 rootless, 904,421 with one, 299,488 with two, 63,903 with three, 3,196 with four, 63 with five, and none with six or more.

Theorem 4.2 (cost thresholds; MV)**.** *The minimum τ admitting k distinct integer roots is: $k = 1{:}\,1$; $k = 2{:}\,2$; $k = 3{:}\,3$; $k = 4{:}\,5$; $k = 5{:}\,6$; $k = 6{:}\,8$; $k = 7{:}\,10$.*

The $k = 6$ lower bound is Theorem 4.1 ($z_{\max}(7) = 5$); the upper bound is the explicit 8-gate witness of Section 4.1; the $k = 7$ value is Corollary 8.8. Representative optimal witnesses, replayed gate by gate in exact arithmetic from the released artifacts:

τ	record	witness program
3	$x^3 - x$, roots $\{0, \pm 1\}$	$x{\cdot}x$; $x^2{-}1$; $x{\cdot}(x^2{-}1)$
4	$-x(x{+}1)(x{+}2)$, roots $\{0, -1, -2\}$	$-1{-}x$; $1{-}(-1{-}x)$; $x(-1{-}x)$; $(x^2{+}x)\ldots$
5	$x^2 - (x^2{-}2)^2 = -(x^2{-}1)(x^2{-}4)$	-2; x^2; $x^2{-}2$; $(x^2{-}2)^2$; $x^2 - (x^2{-}2)^2$
6	$\mp x(x^2{-}1)(x^2{-}4)$, roots $\{0, \pm 1, \pm 2\}$	the $\tau{=}5$ record, then $\times\, x$

Plateaus. $z_{\max}$ pauses at $\tau = 4$ and $\tau = 7$, and again at $\tau = 9$ (Corollary 8.8). The record galleries explain the asymmetry: the step $5 \to 6$ succeeds because multiplying by the input x costs one gate and adjoins the root 0 to a record avoiding it; the steps $3 \to 4$ and $6 \to 7$ fail because every candidate next root beyond the "free" set $\{0, \pm 1, \pm 2\}$ requires a built constant, and constants cost gates. The $\tau = 5$ records (ten polynomials) and $\tau = 6$ records (four) are all difference-of-squares splittings over the map $x^2 - 2$ (Section 6); at $\tau = 7$ the sixty-three five-rooters appear, but no six-rooter.

4.1 Closing the six-root window

The six-root threshold is exactly 8:

Theorem 4.3 (MV, both directions)**.** *There is an 8-gate constant-free SLP whose output has 6 distinct integer roots, and none with 7 gates. An explicit witness (replayed gate by gate in exact arithmetic) is, with $q = x(x-1)$:*

$$x{-}1; \quad 2; \quad q = x \cdot (x{-}1); \quad 4; \quad q{-}2; \quad (q{-}2){-}4 = q{-}6; \quad q(q{-}2); \quad f = q(q{-}2) \cdot (q{-}6),$$

whose root set is $\{-2, -1, 0, 1, 2, 3\}$. Hence $z_{\max}(8) \geq 6$.

The method is a case split. By Lemma 2.5 at depth 8, an 8-gate 6-rooter is one gate over a depth-7 state $S \cup \{v\}$; since $z_{\max}(7) = 5$, the final gate must involve the seventh value v. If that gate is a multiplication, $f = v \cdot b$ and $z(f) = |R_v \cup R_b|$: a pure co-occurrence condition on known root sets, decided exactly by one scan of the 25,844,905 depth-6 states: 408 hits, three reconstructed as explicit programs and verified independently of the scan machinery. A construction search over program schemas, which builds the constants $\{2, 6\}$ independently, reaches this witness family only at 9 gates; the scan's witness builds $\{2, 4\}$ and reuses the

subtraction chain $q \mapsto q-2 \mapsto q-6$, saving one gate. The five-rooters with $\tau \le 7$ do not share one root set: the 67 five-rooters realize seven root-set patterns, including the non-consecutive $\{-1,0,1,2,4\}$ and $\{-4,-2,-1,0,1\}$, and it is precisely such shifted and punctured sets that leave room for a co-occurring factor to complete six roots.

4.2 The depth-eight resolution

Depth 8 is decided completely:

Theorem 4.4 (MV)**.** $z_{\max}(8) = 6$*, and no* 8*-gate program of* any *final-gate shape computes a polynomial with* 7 *distinct integer roots. Hence the seven-root threshold is* 9 *or* 10*.*

The lower side ($z_{\max}(8) \ge 6$) is Theorem 4.3. The upper side required the whole depth-7 frontier: it was *constructed*, for the first time, as 1,048,460,912 reached-set states (out-of-core: 2.43 billion raw 28-byte successor rows, hash-partitioned into 256 files that deduplicate independently in memory), with an exact internal cross-anchor: the build's count of new depth-7 polynomials, 2,013,706, equals the value computed independently by the depth-7 scan of Section 4. The frontier was then scanned exhaustively by the last-gate lemma ($\sim 2 \times 10^{11}$ gate applications): zero results with 7 or more distinct integer roots. Every pipeline stage was first validated on a known answer: the identical machinery one level down reproduces the depth-6 frontier and the $z_{\max}(7) = 5$ census exactly. For completeness of the methods: two general-purpose solver engines (nonlinear integer arithmetic and pure bit-vector SAT over a gate-evaluation encoding) were validated semantically on pinned witnesses, yet neither completed even the smallest free-structure known-answer search (six gates, five roots) within time limits of 30 minutes and one hour respectively; at this problem's scale, exhaustive construction beat solver search.

4.3 The digit-restricted census

Rojas' reduction makes a digit-restricted bound *sufficient* for the full conjecture: bounding only roots $\equiv 1 \pmod p$ for one fixed prime suffices. We measured that restricted growth function exactly through $\tau = 7$: the maximum number of odd roots is $1,2,2,2,2,3,4$, and for roots $\equiv 1 \pmod 3$: $1,1,1,2,2,3,3$. The digit-restricted world has its *own* extremal family, e.g. $(x^2-1)(x^2-9)$ at exactly 7 gates with all four roots odd: symmetric products over odd squares, distinct from the unrestricted records (which crowd $\{0,\pm1,\pm2\}$ and are digit-poor). The sufficient form of the conjecture is thus measurably harder to grow *and* mechanically independent at the bottom of the ladder.

5 Record mechanisms

The census records decompose into a small move inventory [MV for the instances; observational as a classification]: linear factors $x-a$ for available constants a; block translation at one gate per unit shift; multiplication by the input x (one gate, adjoins 0); and difference-of-squares splittings $B^2 - A^2 = (B-A)(B+A)$ at two gates given A and B. The depth-5 record $x^2-(x^2-2)^2 = -(x-1)(x+1)(x-2)(x+2)$ is the fundamental instance: its inner map $C(x) = x^2-2$ is the Chebyshev-conjugate doubling map ($h(z) = z+1/z$ satisfies $h(z)^2-2 = h(z^2)$), i.e. the record is the integer shadow of exactly the iteration that makes the *real* analogue of the conjecture false. Over $\mathbb{Z}$, that factory is sterile beyond one splitting, which the next section proves.

6 Stall theorems

Lemma 6.1 (escape)**.** *Let $h = x^d + \sum_{i<d} a_i x^i \in \mathbb{Z}[x]$, $d \geq 2$, and $R = 1 + \sum_{i<d} |a_i|$. If $|x| \geq R+1$ then $|h(x)| \geq |x| + 1$.* [D]

Proof. $\sum_{i<d} |a_i|\,|x|^i \leq (R-1)|x|^{d-1}$ for $|x| \geq 1$, so $|h(x)| \geq |x|^{d-1}(|x| - (R-1)) \geq 2|x|^{d-1} \geq 2|x| \geq |x| + 1$ for $|x| \geq R+1$. □

Theorem 6.2 (monic stall; D)**.** *Fix a monic $h \in \mathbb{Z}[x]$ of degree ≥ 2. There is a constant $Z(h)$ such that for every $k \geq 1$, both $h^{\circ k}(x) - x$ and the difference-of-squares towers $G_k = h^{\circ(k-1)}(x)^2 - h^{\circ k}(x)^2$ have at most $Z(h)$ distinct integer roots. One may take $Z(h) = \#([-M, M] \cap \mathbb{Z})$ where $M = \max(R+1, \max|a|)$ over the integer solutions a of $h(y) = \pm y$.*

Proof. Integer roots of $h^{\circ k}(x) - x$ are periodic points of h; by Lemma 6.1 all integer periodic points lie in $[-R-1, R+1]$. For G_k, factor $G_k = (h^{\circ(k-1)} - h^{\circ k})(h^{\circ(k-1)} + h^{\circ k})$: an integer root x has $y = h^{\circ(k-1)}(x)$ solving $h(y) = \pm y$, so x lies in the increasing union $\bigcup_j (h^{\circ j})^{-1}(A) \cap \mathbb{Z}$ for the finite set A of such y. If $|x| > M$ then $|h^{\circ j}(x)| \geq |x| + j > \max_{a \in A} |a|$ for all j by Lemma 6.1 and induction, so no element of the union exceeds M in absolute value; the union is contained in a fixed finite set and stabilizes. □

Proposition 6.3 (the Chebyshev core; D, MV spot-checks)**.** *For $C(x) = x^2 - 2$: the integer periodic points are exactly $\{2, -1\}$; for every $k \geq 1$ the polynomial $C^{\circ k}(x) - x$ has exactly 2 distinct integer roots while having 2^k distinct real roots and $\tau \leq 2k + 2$; and the towers G_k have root set $\{\pm 1, \pm 2\}$ for $k = 1$ and exactly $\{0, \pm 1, \pm 2\}$ for every $k \geq 2$.*

Proof. Orbits: $0 \mapsto -2 \mapsto 2 \mapsto 2$ and $1 \mapsto -1 \mapsto -1$; by Lemma 6.1 (with $R = 3$; direct check for $|x| \geq 3$) all other integer orbits escape, so the only cycles are the fixed points 2 and -1, giving the first two claims (the real-root count is the classical Chebyshev conjugacy). For G_k: $h(y) = y$ has integer solutions $\{2, -1\}$ and $h(y) = -y$ has $\{-2, 1\}$; integer preimages are $C^{-1}(2) = \{\pm 2\}$, $C^{-1}(-1) = \{\pm 1\}$, $C^{-1}(-2) = \{0\}$, $C^{-1}(1) = C^{-1}(0) = \emptyset$, so the preimage closure of $\{\pm 1, \pm 2\}$ is $\{0, \pm 1, \pm 2\}$ and is stable. □

Theorem 6.2 closes a construction class: no family built by iterating one fixed monic map can witness superpolynomial z versus τ; a refutation of the conjecture, if one exists, must build fresh constants or mix maps, paying gates for them. That is precisely the regime the census measures.

7 The quadratic family

We measured all tower shapes ($h^{\circ k}(x) - x$ and $h^{\circ i}(x)^2 - h^{\circ j}(x)^2$, $0 \leq i < j \leq 4$) across $h_c = x^2 - c$ for $1 \leq c \leq 200$, in exact arithmetic. Root finding here cannot use trailing-coefficient divisors (those constants reach $c^{2^k} \sim 10^{18}$); instead, by Lemma 6.1, every integer root of such a shape lies in $[-(c+1), c+1]$, and direct evaluation over that window is exact. The two finders agree on every shape where both are feasible ($c \leq 10$, $j \leq 2$) [MV].

Theorem 7.1 (family census; MV)**.** *For $1 \leq c \leq 200$ and the shapes above: the maximum number of distinct integer roots is 5, attained only at $c = 2$ (the set $\{0, \pm 1, \pm 2\}$ of Proposition 6.3). Nonzero yields occur exactly on two series: $c = m(m+1)$, yield 4 for $m \geq 2$ with root set $\{\pm m, \pm(m+1)\}$ (from the fixed points $\{m+1, -m\}$ and anti-fixed points*

$\{m, -m-1\}$*), and* $c = m^2 + m + 1$*, yield* 4 *with the same root sets, produced by the genuine* 2*-cycles* $m \mapsto -m-1 \mapsto m$ *(for* $c = 1$*: yield* 3*, root set* $\{0, \pm 1\}$*, from the cycle* $0 \leftrightarrow -1$*).*

The fixed-point series is therefore not the only series that yields roots. The second series is, however, the last one available to this construction class:

Proposition 7.2 (cycle ceiling; D, classical)**.** *For any* $f \in \mathbb{Z}[x]$*, every cycle of* f *in* $\mathbb{Z}$ *has length at most* 2*. Consequently fixed points, anti-fixed points and* 2*-cycles are the complete periodic inventory harvestable by tower shapes over any integer polynomial map.*

Proof. For integers $a \neq b$, $(a-b) \mid (f(a) - f(b))$. Around a cycle $a_0 \mapsto a_1 \mapsto \cdots \mapsto a_r = a_0$ of distinct points, the differences $d_i = a_{i+1} - a_i$ satisfy $d_i \mid d_{i+1}$ cyclically, so all $|d_i|$ equal one value $d > 0$, and the cycle is a closed walk on the arithmetic line $a_0 + d\,\mathbb{Z}$ with steps $\pm d$. If two consecutive steps have opposite signs then $a_{i+2} = a_i$, which forces $r = 2$; if all steps have the same sign the walk never closes. Hence $r \leq 2$. The fact is classical in the polynomial-cycles literature (see Narkiewicz [21]); the proof is included to keep the paper self-contained. □

Yield per gate across the family is maximized at $c = 2$ and decreases in c (the yield is bounded by 5 while building c costs $\tau(c) \sim \log c$ gates, measured exactly against our anchored integer census): within this family, the "parameterized loophole" left open by Theorem 6.2 is empty [MV for $c \leq 200$; C beyond].

8 The seven-root threshold

Theorem 4.4 left the threshold for 7 distinct integer roots at 9 or 10. This section closes the upper side constructively, decides the multiplicative half of the ninth gate exhaustively, decides the additive half first on an evaluation window and then without one, and concludes that the threshold is exactly 10.

8.1 Two ten-gate witnesses

We give two 10-gate witnesses, written as explicit programs and executed gate by gate in the same exact arithmetic as the census; they are structurally different, and Section 8.2 compares them.

Proposition 8.1 (upper bound; MV)**.** *Each of the following constant-free programs has length* 10 *and computes a polynomial with* 7 *distinct integer roots.*

(A) With $q = x^2 - x$*:* $p_A = q(q-2)(q-6) \cdot (x-4)$*, root set the interval* $\{-2, \ldots, 4\}$*, height* 56*.*

(B) $p_B = x^2(x^2-1)(x^2-4)(x^2-16)$*, root set the tower* $\{0, \pm 1, \pm 2, \pm 4\}$*, height* 84*.*

Hence the seven-root threshold is at most 10*.*

Both gate counts are exact against the model's conventions because the programs are run as data rather than counted by hand; neither is claimed to be optimal, which is the business of the scan below. Witness (A) costs nothing extra for its third factor, since $q - 6 = (q-2) - 4$ and the constant 4 is already in the state of the exhaustive 8-gate six-rooter $q(q-2)(q-6)$ of Theorem 4.3; appending $(x-4)$ costs one subtraction and one multiplication. Witness (B)

is cheap for a different reason: each of its constants is a single squaring past the previous one ($2 \to 4 \to 16$), so it costs one gate, and each is a perfect square, so it contributes a *pair* of integer roots.

That second mechanism generalizes.

Proposition 8.2 (square-tower family; MV for $m \le 4$)**.** *Let $t_1 = 2$ and $t_{i+1} = t_i^2$, and set*

$$p_m(x) \;=\; x^2\,(x^2-1)\prod_{i=2}^{m+1}\bigl(x^2 - t_i\bigr).$$

Then p_m has $z = 2m+3$ distinct integer roots and is computed by $\tau = 3m+4$ gates for $m \ge 1$ (3 gates when $m = 0$): three gates per two roots.

The family is a useful calibration precisely because it is *not* uniformly optimal. At $m = 0$ it gives 3 roots in 3 gates and the census says the minimum for 3 roots is exactly 3; at $m = 1$ it gives 5 roots in 7 gates where the census achieves 5 roots in 6. It therefore supplies upper bounds only, and it is the exhaustive scan below, not the construction, that decides optimality at seven.

8.2 Two arithmetics of opposite parity

Witnesses (A) and (B) exploit different mechanisms, and the mechanisms differ in parity. Folding by $q = x^2 - x$ identifies x with $1 - x$, so every value with integer preimages contributes exactly two roots: folding natively produces *even* root counts, and an odd count costs an extra linear factor. Squaring by $y = x^2$ has one self-paired value, 0, whose preimage is the single point $x = 0$: squaring natively produces *odd* counts, $2m + 3$.

Seven is odd, so the tower is native there and the interval needs its patch, and the two come out equal at ten gates. Across the exhaustive range an interval is always *among* the minimal-gate record sets, though it is not the only one: at $\tau = 5$ the four-root records include the interval $\{-1, 0, 1, 2\}$ and also the difference-of-squares set $\{\pm 1, \pm 2\}$, computed by $x^2 - (x^2 - 2)^2$, which is not an interval [MV]. The census therefore ends exactly where the two mechanisms tie, and never separates them.

Height does separate them. Witness (A) has height 56 against (B)'s 84, and at six roots the interval record has height 15 against 84 for the corresponding tower. This matches a direct measurement over the census witnesses: the minimal height of a polynomial in our catalog with z distinct integer roots is $1, 1, 2, 4, 15$ for $z = 2, \ldots, 6$ [MV], attained by intervals. Additively structured root sets are the height-cheap ones; multiplicatively structured ones buy their constants at doubly exponential magnitude. We record this because it constrains proof strategies: a lower-bound argument tuned to spread, multiplicatively built root sets will be blind to interval sets, and conversely.

8.3 The ninth gate, multiplicative half

By the last-gate lemma a 9-gate program is one gate over a depth-8 state, and since $z_{\max}(8) = 6$ the final gate of any 9-gate seven-rooter must involve the eighth value. When that gate is a multiplication, $f = v \cdot b$ and $z(f) = |R_v \cup R_b|$, so the question reduces to unions of root sets already known from the frontier, with no polynomial arithmetic in the inner loop.

Theorem 8.3 (MV)**.** *No 9-gate constant-free program whose final gate is a multiplication computes a polynomial with 7 distinct integer roots. Over all* 1,048,460,912 *depth-7 states, every one-gate extension and every operand, the maximum union size is exactly* 6*.*

The scan was validated before the full run: the same machinery run at threshold 6 returns 793 unions on a single partition, so it detects unions when they exist. This half is unconditional; root-set unions require no evaluation window.

8.4 The additive half, and what "windowed" costs

An additive final gate admits no such reduction, since root sets do not compose across addition. The instrument works with values instead: every operand and extension is evaluated on the window $[-32, 32]$ modulo a 31-bit prime, and $f = v \pm b$ vanishes at a point only if the residues agree, so counting modular agreements never misses a witness; candidates reaching seven agreements are promoted to exact construction and exact root counting. Two numerical points matter. First, 61-bit primes overflow `int64` under modular multiplication, so the modulus is 31-bit. Second, identically-zero polynomials agree at every window point and would otherwise flood the promotion stage. They are excluded, but only under a degree side-condition: agreement at all 65 window points is discarded when the candidate's degree bound is below 65, since a nonzero polynomial of degree below 65 cannot vanish at 65 points, and is *retained* when the degree bound is 65 or more. The side-condition matters, because a 9-gate polynomial can have degree up to 2^8, and discarding full agreement unconditionally would throw away exactly the high-degree witnesses the instrument is designed to find.

An empty result from this instrument is *windowed*: on its own it excludes only witnesses all of whose roots lie in $[-32, 32]$, and a 9-gate program can build constants as large as 2^{256} by repeated squaring. The following turns that caveat from an unknown into a decidable criterion.

Lemma 8.4 (confinement; D, with the bound MV)**.** *Let $f \in \mathbb{Z}[x]$ be nonzero with distinct integer roots $r_1, \ldots, r_k$. Then $\prod_i (n - r_i)$ divides $f(n)$ for every $n \in \mathbb{Z}$. Consequently, if f has 7 distinct integer roots and c denotes its trailing (first nonzero) coefficient, then every integer root of f lies in $[-32, 32]$ whenever $|c| \le 395$, and, if 0 is not a root of f, whenever $|c| \le 1187$.*

Proof. The r_i are distinct integers, so $\prod_i (x - r_i)$ is monic in $\mathbb{Z}[x]$ and divides f in $\mathbb{Q}[x]$; by Gauss the cofactor lies in $\mathbb{Z}[x]$, and evaluating at n gives the divisibility. For the bound, suppose some root satisfies $|r_j| \ge 33$. If 0 is not a root then $c = f(0)$ and $|c| \ge \prod_i |r_i|$; the six remaining roots are distinct and nonzero, so their product has absolute value at least $|(-1)(1)(-2)(2)(-3)(3)| = 36$, giving $|c| \ge 33 \cdot 36 = 1188$. If 0 is a root, write $f = x^m g$ with $c = g(0)$; the six other roots are distinct and nonzero, and the five non-escaping ones give a product of absolute value at least 12, so $|c| \ge 33 \cdot 12 = 396$. An escaping root therefore forces $|c| \ge 1188$ when 0 is not a root and $|c| \ge 396$ in every case, contradicting either hypothesis. □

The constant 395 is sharp: searching seven-root sets containing an escaping root returns minimum trailing coefficient exactly 396, at $\{-3, -2, -1, 0, 1, 2, 33\}$ [MV]. We note that the two witnesses of Proposition 8.1 have trailing coefficients 48 and -64, an order of magnitude inside the criterion: the shapes that actually reach seven roots cheaply are exactly the ones the window sees.

Lemma 8.4 does not make the windowed instrument unconditional. It prices the gap, and we measured the price. Constructing every candidate $f = t \pm b$ exactly on a sample of 6,400,000

candidates drawn from four hash-partitioned frontier files, and reading each one's true trailing coefficient [MV]:

identically zero (excluded)	93,898	1.467%
confined by Lemma 8.4	6,301,361	99.9248%
excluded by $\deg f < 7$	4,619	0.0732%
window-limited, $\deg f \geq 7$	122	0.0019%

the last three percentages being of the 6,306,102 nonzero candidates. The degree row is arithmetic rather than a lemma: seven distinct roots require degree at least seven, so those candidates are excluded with no reference to the window at all. Together, 99.998% of nonzero candidates are decided independently of the evaluation window, and the residual is a structurally narrow class (degrees 8, 9 and 12; 64 distinct trailing coefficients between 512 and 38,220).

The fractions above are sample estimates from four of 256 partitions, not exhaustive counts; the scan of the next theorem is exhaustive.

Theorem 8.5 (MV, windowed)**.** *No* 9*-gate constant-free program whose final gate is an addition or a subtraction computes a polynomial with* 7 *distinct integer roots that all lie in* $[-32, 32]$*. Over all* 1,048,460,912 *depth-*7 *states, every one-gate extension, every operand and both signs,* 1,272,725 *candidates reach seven agreements on the window, and none of them has* 7 *distinct integer roots.*

The scan was gated before production: at threshold 5 on 20,000 states of one partition it returns 41 witnesses, each verified exactly, so it finds witnesses when they exist. Its 256 partition records each match the state count of the corresponding frontier file in the frontier's SHA-256 manifest, so no partition was scanned partially.

Together with Theorem 8.3, $z_{\max}(8) = 6$ and Lemma 8.4, a 9-gate program computing a polynomial with 7 distinct integer roots, if one exists, has an additive final gate and a root outside $[-32, 32]$, hence a trailing coefficient with $|c| \geq 396$, and $|c| \geq 1188$ when 0 is not a root. The seven-root threshold is therefore 10 unless this residual class contains a seven-rooter; strictly, it remains in $\{9, 10\}$ on the strength of this scan alone. The next subsection decides it.

8.5 The additive half without a window

The confinement lemma bounds where a missed witness could be; the following criterion removes the window altogether. It uses the same divisibility, read p-adically on a block of consecutive integers rather than archimedeanly at 0.

Lemma 8.6 (p-adic block criterion; D)**.** *Let* $f \in \mathbb{Z}[x]$ *be nonzero with distinct integer roots* $r_1, \ldots, r_k$*, let* x^m *be the exact power of* x *dividing* f*, and let* $0 \leq m' \leq m$*. For every prime* p *and every integer* n *with* $f(n) \neq 0$*,*

$$v_p\big(f(n)\big) - \max(m' - 1, 0)\, v_p(n) \;\geq\; \sum_{i=1}^{k} v_p(n - r_i).$$

Let B *be a block of* p^a *consecutive integers and* $e \geq 1$*, and let* Z *be the set of* $n \in B$ *with* $f(n) \equiv 0 \bmod p^e$*. Put* $w(t) = v_p(t)$ *if* $p^a \nmid t$ *and* $w(t) = a$ *otherwise, and*

$$C(z) = \sum_{n \in B \setminus Z} v_p(n - z) \;\; (z \in Z), \qquad M = \min_{\rho \bmod p^a} \sum_{n \in B \setminus Z} w(n - \rho).$$

If $L_k(Z)$ is the sum of the k smallest numbers in the multiset $\{C(z) : z \in Z\} \cup \{M, \dots, M\}$ (k copies of M), then

$$S(f) := \sum_{n \in B \setminus Z} \Big(v_p\big(f(n)\big) - \max(m' - 1, 0)\, v_p(n) \Big) \;\geq\; L_k(Z). \tag{1}$$

Proof. As in Lemma 8.4, $f = g \prod_i (x - r_i)$ with $g \in \mathbb{Z}[x]$. If $m \geq 1$ then 0 is one of the r_i and the remaining factor x^{m-1} divides g, which gives the first inequality; for $m' = 0$ nothing is subtracted. On $B \setminus Z$ the value $f(n)$ is nonzero, so the inequality holds there; sum it. A root cannot lie in $B \setminus Z$. A root $z \in Z$ contributes exactly $C(z)$. A root $r \notin B$ contributes at least $\sum_n w(n-\rho)$ with $\rho = r \bmod p^a$, because $v_p(n-r) = w(n-\rho)$ when $p^a \nmid n - r$ and $v_p(n-r) \geq a$ otherwise; distinct roots outside B may share a residue, so each contributes at least M. The least total over all placements of k distinct roots is $L_k(Z)$. □

The criterion is cheap because residues are exact: $f(n) \bmod p^e$ is computed from the residues of the operands, and $L_k(Z)$ depends only on the zero mask of the block, so it is tabulated once (2^{16} masks for $p = 2$, $B = \{-8, \dots, 7\}$, $e = 64$, where unsigned 64-bit arithmetic is exactly arithmetic mod 2^{64}; 2^9 masks for $p = 3$, $B = \{-4, \dots, 4\}$, $e = 20$). The power of x is bounded below by the operands' trailing indices (their sum for a product, their minimum otherwise). Without that correction, candidates of the form $x^j h$ or $2^j h$ pass the 2-adic block easily; the 3-adic block is blind to powers of 2.

Theorem 8.7 (MV, no window)**.** *No 9-gate constant-free program whose final gate is an addition or a subtraction computes a polynomial with 7 distinct integer roots.*

The scan covers the same space as Theorem 8.5: all 1,048,460,912 depth-7 states, all 200 eighth gates on their 10 operands, all 10 operands b and both signs. Every candidate $f = t \pm b$ is dropped only by a proof or decided exactly:

candidates	4,193,843,648,000
dropped: $\deg f \leq \max(\deg t, \deg b) < 7$	4,026,692,708,694
pass the 2-adic block	3,840,174,256
pass the 3-adic block	2,937,472,943
identically zero ($t = \mp b$, built exactly)	92,943,826
built exactly and root-counted	2,345,860,018
with 7 distinct integer roots	0

Before production, the engine at thresholds $k = 4, 5, 6$ was compared with building *every* candidate exactly, with no filter, on samples of 150 to 3,000 states from five partitions: the counts agree in every case (1535, 864, 6, 15 and 0). Planted polynomials with 4 to 7 distinct roots of absolute value up to 5000, with powers of x and of 2 and 3 in their content, pass both blocks for every admissible m'. Each partition's frontier file was hashed against the frontier's SHA-256 manifest before its scan, and the 256 partition records total the manifest's state count.

Corollary 8.8 (MV)**.** $z_{\max}(9) = 6$*, and the minimum number of gates of a constant-free program computing a polynomial with* 7 *distinct integer roots is exactly* 10*.*

Proof. By the last-gate lemma a 9-gate program is one gate over a depth-8 state, and since $z_{\max}(8) = 6$ a seven-rooter's final gate uses the eighth value; it is multiplicative or additive, excluded by Theorems 8.3 and 8.7. Hence $z_{\max}(9) \leq 6$, and $z_{\max}(9) \geq z_{\max}(8) = 6$. Proposition 8.1 gives the upper side of the threshold. □

9 The three worlds, measured

It is folklore that the conjecture's analogue fails over finite fields, and that $x^{p-1}-1$ is the reason; this is why the statement is always made in characteristic zero. We record here that the finite-field question has an exact answer, obtained on the *same* enumerated programs as the integer census, so that the two ladders are comparable term by term.

For a prime p write $z_p(f)$ for the number of distinct roots of f in $\mathbb{F}_p$, defined only when $f \not\equiv 0 \pmod p$, and set

$$z^p_{\max}(\tau) \;=\; \max\big\{\, z_p(f) \;:\; \tau(f) \le \tau,\ p \text{ prime},\ f \not\equiv 0 \pmod p \,\big\}.$$

Proposition 9.1 (D, with both halves MV)**.** $z^p_{\max}(\tau) = 2^{\tau-1}$ *for every* $\tau \ge 1$.

Proof. Upper bound. For a value v of a constant-free program let $\mu(v)$ count the *distinct* multiplicative gates in its sub-DAG. We claim $\deg v \le 2^{\mu(v)}$, by induction. The input has degree 1 with $\mu = 0$, and constants have degree 0. For an additive gate $v = a \pm b$, the sub-DAG of v contains those of a and b, so $\mu(v) \ge \max(\mu(a), \mu(b))$ and $\deg v \le \max(\deg a, \deg b) \le 2^{\max(\mu(a),\mu(b))} \le 2^{\mu(v)}$. For a multiplicative gate $v = ab$ we have $\mu(v) = |M(a) \cup M(b)| + 1 \ge \max(\mu(a), \mu(b)) + 1$, where $M(\cdot)$ denotes the set of multiplicative gates in a sub-DAG, and hence

$$\deg v = \deg a + \deg b \le 2^{\mu(a)} + 2^{\mu(b)} \le 2^{\max(\mu(a),\mu(b))+1} \le 2^{\mu(v)}.$$

Counting *distinct* gates, so that shared subexpressions are not charged twice, is what makes the multiplicative step close: bounding the exponent by $\mu(a) + \mu(b) + 1$ instead would exceed $\mu(v)$ whenever a and b share gates.

Now a sub-DAG with no additive gate computes only $\pm x^k$ or 0, since the free constants are $-1, 0, 1$ and products of those remain in $\{-1, 0, 1\}$; such a polynomial has at most one distinct root in any field. So if f is not of that form its sub-DAG contains an additive gate, whence $\mu(f) \le \tau(f) - 1$ and $\deg f \le 2^{\tau-1}$. Finally a polynomial that is nonzero over a field has at most its degree many roots, and reduction mod p can only lower the degree.

Lower bound. By Dirichlet there is a prime $p \equiv 1 \pmod{2^k}$ for every k. Modulo such p the polynomial $x^{2^k} - 1$ has exactly $\gcd(2^k, p-1) = 2^k$ roots, and it is computed by k squarings and one subtraction, so $\tau \le k + 1$. Taking $k = \tau - 1$ gives $2^{\tau-1}$. □

Both halves were checked mechanically: the degree bound holds on every non-monomial in the complete census to depth 6, where it is moreover attained exactly ($\max \deg = 2^{\tau-1}$ at each τ, against 2^τ for monomials), and the construction was verified for $k = 1, \ldots, 8$ at $p = 3, 5, 17, 17, 97, 193, 257, 257$ with root counts $2, 4, 8, 16, 32, 64, 128, 256$ [MV]. We note that the Fermat primes, which the small cases suggest, are not needed: only $p \equiv 1 \pmod{2^k}$ is.

9.1 The real ladder

The real analogue of the conjecture is known to fail. We measured how far, on the same enumerated programs, counting distinct real roots exactly: take the square-free part $f/\gcd(f, f')$ and count real roots by Sturm's theorem in exact rational arithmetic, with no floating point at any stage. The method was first validated on seven known answers, among them $x^2 + 1$ (none), x^2 (a double root counted once), T_4 (four), and $(x^2-1)(x^2-4)(x^2-16)$ (six). Writing $z^{\mathbb{R}}_{\max}(\tau)$

for the maximum over τ-gate polynomials:

τ	1	2	3	4	5	6	7	8
$z_{\max}(\tau)$	1	2	3	3	4	5	5	6
$z^{\mathbb{R}}_{\max}(\tau)$	1	2	3	4	6	8	–	–
$z^{p}_{\max}(\tau)$	1	2	4	8	16	32	64	128

(The real row is exhaustive through $\tau = 6$. That row was obtained without a depth-6 census: the last-gate lemma applied to the 778,087-state depth-5 frontier enumerates all 134,494 new $\tau = 6$ polynomials, and the same run is validated on that set by requiring the *integer* maximum to come out at 5. Reaching $\tau = 7$ needs the 25.8M-state depth-6 frontier resident in memory, which is not attempted here.) The three worlds separate at three different points: $\mathbb{F}_p$ leaves $\mathbb{Z}$ at $\tau = 3$ and saturates the degree ceiling $2^{\tau-1}$ at every τ; $\mathbb{R}$ leaves $\mathbb{Z}$ at $\tau = 4$ and does *not* saturate it (4 against 8, then 6 against 16); $\mathbb{Z}$ trails both.

The real record at $\tau = 5$ is worth writing out, because it is produced by the same difference-of-squares mechanism that Section 5 identifies as the integer record-maker:

$$a = x \cdot x, \quad b = a - 1, \quad c = b \cdot b, \quad d = c - a, \quad e = b \cdot d,$$

$$e \;=\; (x-1)(x+1)(x^2 - x - 1)(x^2 + x - 1).$$

Five gates, *six* distinct real roots $\pm 1, \pm\varphi, \pm\varphi^{-1}$ with φ the golden ratio, and exactly *two* distinct integer roots [MV]. The program is cheap and the roots are there; four of them are simply irrational. This is the conjecture's content in one line: what limits $\mathbb{Z}$ is arithmetic, not the cost of manufacturing roots. It also shows the real ladder is not merely the Chebyshev ladder: that tower reaches $T_2 = 2x^2 - 1$ with two real roots at three gates and needs six for T_4's four, so at five gates it is still at two, against six for the $\tau = 5$ record [MV].

The $\tau = 6$ record exposes the mechanism. It is

$$x^8 - 7x^6 + 14x^4 - 7x^2 + 1 \;=\; (x^4 - 3x^2 + 1)(x^4 - 4x^2 + 1),$$

whose left factor is exactly the $\tau = 4$ record $g_1 = (x^2 - 1)^2 - x^2$. So the step from four gates to six is $g \mapsto g \cdot (g - x^2)$, costing *two* gates because x^2 is already in the state, and *doubling* the real root count. It beats the obvious construction: a Chebyshev tower costs three gates per doubling ($T_2 = 2x^2 - 1$ is three gates and composition repeats it), so it reaches only about $2^{\tau/3}$. And it is not merely a good construction here, it matches the exhaustive census at both $\tau = 4$ and $\tau = 6$.

It also stalls, in the same manner as the integer-side towers of Section 6. Iterating the step gives distinct real roots against degree

gates	4	6	8	10	12
deg	4	8	16	32	64
real	4	8	16	28	48

[MV]: three clean doublings, reaching 16 distinct real roots at 8 gates, after which the iterate ceases to be totally real and the successive ratios fall to 1.75 and 1.71.

We claim no rate for $z^{\mathbb{R}}_{\max}$: six points cannot distinguish $\Theta(2^{\tau/2})$ from anything nearby, the family's own rate decays past $\tau = 8$, and the paper needs only the known failure of the real analogue.

One structural point is worth recording. The $\tau = 6$ real record has *zero* integer roots, while the $\tau = 6$ integer record has five: at equal cost the two worlds are optimized by entirely disjoint

polynomials. The real ladder is not the integer ladder with further roots attached, but a different extremal problem that happens to share the difference-of-squares mechanism at the bottom.

Of the three ladders, the $\mathbb{F}_p$ row is known exactly for every τ (Proposition 9.1). The ladders agree at $\tau \leq 2$ and separate from $\tau = 3$ onward. Over $\mathbb{F}_p$ the count doubles with every gate, which is the fastest the degree bound permits, and the extremal programs are the cheapest available: repeated squaring, then subtract one. The same polynomial $x^{2^k} - 1$ has exactly two integer roots, ± 1, for every k.

9.2 A fourth column: the rational ladder

The conjecture counts *integer* roots. Its natural algebraic relaxation counts rational ones, and it is not obvious which of the two constraints is doing the work. Writing $z^{\mathbb{Q}}_{\max}(\tau)$ for the maximum number of distinct rational roots, computed exactly by the rational root theorem over $\mathbb{Z}$:

τ	1	2	3	4	5	6
$z_{\max}(\tau)$	1	2	3	3	4	5
$z^{\mathbb{Q}}_{\max}(\tau)$	1	2	3	3	4	5

exhaustively through $\tau = 6$ [MV], the $\tau = 6$ row again obtained by the last-gate lemma over the depth-5 frontier and validated by requiring the integer maximum on the same set to reproduce $z_{\max}(6) = 5$.

The two ladders coincide: allowing denominators buys nothing at all. This is not because the rational world is a copy of the integer one, and the extremal objects are in fact different. The $\tau = 6$ rational record is

$$8x^6 - 10x^4 + 2x^2 \;=\; 2x^2(x-1)(x+1)(2x-1)(2x+1),$$

with five distinct rational roots $\{0, \pm\frac{1}{2}, \pm 1\}$ but only *three* integer ones, while the $\tau = 6$ integer record attains five integer roots outright. Half-integer roots are cheap here, since the constants 2 and 4 are cheap, and they still do not raise the ceiling.

So integrality is not where the difficulty lives, at least at the bottom of the ladder: the same growth function appears over $\mathbb{Q}$. Whatever separates $\mathbb{Z}$ from $\mathbb{R}$ and $\mathbb{F}_p$ in the table above already separates $\mathbb{Q}$ from them too. We state this as a measurement over the census range and claim nothing asymptotic from it.

The mechanism is the one the three-worlds reading predicts. In $\mathbb{F}_p^\times$, cyclic of order $p-1$, the equation $x^d = 1$ has $\gcd(d, p-1)$ solutions, up to d of them; in $\mathbb{Z}^\times = \{\pm 1\}$ it has at most two, for every d. The exponential ladder is precisely the 2^k-torsion that the cyclic group carries and the integers do not. In this light the conjecture asserts that no program shape recovers over $\mathbb{Z}$ what the cyclotomic one achieves over $\mathbb{F}_p$, and the census is consistent with that: through $\tau = 8$ nothing over $\mathbb{Z}$ does better than linear growth. What Proposition 9.1 contributes is that the comparison is exact on one side, so the gap being conjectured is a gap between a proved exponential and a measured linear rate rather than between two informal impressions.

10 Limits, and what would count as progress

Nothing above bears on the conjecture asymptotically: censuses to depth 8 cannot distinguish polynomial from superpolynomial growth, and our theorems close construction classes rather than open the bounding problem. In the measured landscape, every mechanism observed or

excluded so far pays $\Theta(1)$ gates per root, the growth function's first structure (plateaus at 4, 7 and 9) is the visible cost of constant-building, and the geometric-progression family remains the best known factory, linear-rate, exactly as when Rojas noted that the conjecture fails for $\kappa < 1$ [8].

The census makes the following open items precise: (i) $z_{\max}(10)$ and the eight-root threshold, which lies in $\{10, \ldots, 13\}$ by the square-tower family ($m = 3$ gives 9 roots in 13 gates) and the census; the block criterion of Lemma 8.6 applies unchanged at the tenth gate, but the depth-8 frontier it would need has not been built; (ii) extend the integer frontier past Markström's length 11 [7], including his monotonicity question for $\tau(n!)$; (iii) measure the growth of the valuation-spectrum bound $s \le N_p(s) \le s(s+1)/2$ of [8], where the true rate is stated as open; (iv) the census's 2-adic instrumentation records that all depth-5–7 record polynomials concentrate their roots on the valuations $\{0, 1\}$ [MV], consistent with the pressure sitting on the digit-conjecture side of Rojas' reduction [C].

Data and code availability

All code, run artifacts, experiment records and the problem wiki are available at `https://github.com/fsantibanezleal/CAOS_RESEARCH` under `problems/computation-complexity/tau-conjecture/`, together with the package `tclib` and its test suite. The computations are archived there as the experiment records `EXP-001`–`EXP-015` (folder `experiments/`); each record holds the code, the artifacts and the result.

Every root count reported here is exact. The census enumerator and all integer root counting are exact integer arithmetic in the Python 3.13 standard library. Two later instruments add libraries without giving up exactness. The depth-8 and nine-gate scans use NumPy for a modular pre-filter over a fixed evaluation window, whose surviving candidates are reconstructed and root-counted exactly by the standard-library engine, so the filter can over-admit but never discard a witness; the window-free scan uses NumPy for the 2-adic and 3-adic residues of Lemma 8.6 in the same way. The real and rational ladders use SymPy for square-free decomposition with Sturm sequences and for the rational root theorem, both in exact rational arithmetic with no floating point at any stage.

Compute is dominated by the two largest scans, on one desktop machine with 32 cores. The depth-8 pipeline took 2h50m to build the frontier and 6h48m to scan it with 20 processes; the multiplicative nine-gate scan took 13.1h, and the additive nine-gate scan about 74h of wall-clock time with 8 to 20 processes, interruptions included, and the window-free additive scan 21.4h of wall-clock time with 20 processes (381 process-hours); the earlier experiments together add roughly 17h, of which 6.7h went to the inconclusive solver runs. The co-occurrence scan that decides the six-root threshold took 2h23m on one core, and the depth-6 real ladder 180 s. Peak RAM was ∼12 GB.

The constructed depth-7 frontier (1,048,460,912 states, 29,356,905,536 bytes, that is 27.3 GiB) and the polynomial catalog (2,161,049 entries) are retained as reusable data assets with SHA-256 manifests recorded in the repository. The frontier's manifest carries a digest for each of its 256 files together with an aggregate fingerprint (`7a3b4484...6baaf174`, the SHA-256 of the concatenated per-file digests in name order), and the state count in that manifest is derived from the file sizes rather than restated, so the asset and the census figure check against each other.

References

[1] M. Shub, S. Smale, *On the intractability of Hilbert's Nullstellensatz and an algebraic version of "NP ≠ P?"*, Duke Math. J. 81(1):47–54, 1995. doi:10.1215/S0012-7094-95-08105-8.

[2] S. Smale, *Mathematical problems for the next century*, Math. Intelligencer 20(2):7–15, 1998. doi:10.1007/BF03025291.

[3] L. Blum, F. Cucker, M. Shub, S. Smale, *Complexity and Real Computation*, Springer, 1998.

[4] P. Koiran, *Valiant's model and the cost of computing integers*, Comput. Complexity 13(3–4):131–146, 2004. doi:10.1007/s00037-004-0186-2.

[5] P. Bürgisser, *On defining integers and proving arithmetic circuit lower bounds*, Comput. Complexity 18(1):81–103, 2009. doi:10.1007/s00037-009-0260-x.

[6] P. Bürgisser, *Completeness classes in algebraic complexity theory*, arXiv:2406.06217, 2024.

[7] K. Markström, *The straight line complexity of small factorials and primorials*, INTEGERS 14, #A67, 2014. arXiv:1306.3091.

[8] J. M. Rojas, *A direct ultrametric approach to additive complexity and the Shub–Smale tau conjecture*, arXiv:math/0304100, 2003.

[9] Q. Cheng, *Straight-line programs and torsion points on elliptic curves*, Comput. Complexity 12(3–4):150–161, 2003. doi:10.1007/s00037-003-0180-0.

[10] Q. Cheng, *On the ultimate complexity of factorials*, Theoret. Comput. Sci. 326(1–3):419–429, 2004. doi:10.1016/j.tcs.2004.07.032.

[11] A. Shamir, *Factoring numbers in $O(\log n)$ arithmetic steps*, Inform. Process. Lett. 8(1):28–31, 1979. doi:10.1016/0020-0190(79)90087-5.

[12] R. J. Lipton, *Straight-line complexity and integer factorization*, ANTS-I, LNCS 877, 1994. doi:10.1007/3-540-58691-1_50.

[13] P. Koiran, *Shallow circuits with high-powered inputs*, Innovations in Computer Science, 2011. arXiv:1004.4960.

[14] P. Koiran, N. Portier, S. Tavenas, S. Thomassé, *A tau-conjecture for Newton polygons*, Found. Comput. Math. 15:185–197, 2015. arXiv:1308.2286.

[15] P. Hrubeš, *On the real τ-conjecture and the distribution of complex roots*, Theory of Computing 9(10):403–411, 2013. doi:10.4086/toc.2013.v009a010.

[16] I. Briquel, P. Bürgisser, *The real tau-conjecture is true on average*, Random Struct. Algorithms 57(2):279–303, 2020. doi:10.1002/rsa.20926.

[17] P. Dutta, *Real τ-conjecture for sum-of-squares: a unified approach to lower bound and derandomization*, CSR 2021. doi:10.1007/978-3-030-79416-3_5.

[18] S. Bhattacharjee, M. Bläser, P. Dutta, S. Mukherjee, *Exponential lower bounds via exponential sums*, ICALP 2024; extended version arXiv:2601.00387, 2026.

[19] P. Bürgisser, *Intractability of Hilbert's Nullstellensatz implies algebraic hardness of the permanent*, arXiv:2606.25121, 2026.

[20] C. Fuhs, P. Schneider-Kamp, *Synthesizing shortest linear straight-line programs over GF(2) using SAT*, SAT 2010, LNCS 6175. doi:10.1007/978-3-642-14186-7_8.

[21] W. Narkiewicz, *Polynomial Mappings*, Lecture Notes in Mathematics 1600, Springer, 1995.