

An independent exact verification of the 2026 counterexample to Goemans’ unsplittable-flow cost conjecture, with the violation constant it forces

Machine record v0.01: the CAOS Research unsplittable-flow-cost program, experiments
001–003

Felipe Santibañez-Leal *

July 25, 2026

Abstract

In July 2026 a counterexample to Goemans’ cost version of the Dinitz–Garg–Goemans theorem was announced publicly, outside peer review, and circulated widely. We report an independent verification of that finite object by exact rational enumeration, performed without executing or importing the proposer’s verifier, together with quantitative and structural content that the announcement did not state. Our results: (i) the instance is confirmed, so Goemans’ conjecture is false, and, the instance being acyclic, the Morell–Skutella cost conjecture and the convex-combination form fall with it; (ii) the instance forces a violation of exactly $\frac{16}{15}d_{\max}$, that is, the conjecture fails by ONE unit at $d_{\max} = 15$, so the open question whether cost-preserving rounding is possible with $O(d_{\max})$ violation is untouched by it; (iii) every proved result in the area survives its direct test on the instance, including the Dinitz–Garg–Goemans theorem, the Morell–Skutella cost-free conjecture, the series-parallel theorem of Majthoub Almoghrabi, Skutella and Warode, and the planar theorem of Traub, Vargas Koch and Zenklusen; (iv) the instance contains a K_4 subdivision, hence sits exactly one structure outside the series-parallel class where the conjecture is proved, and, being planar, pins the planar constant strictly between 1 and 2; (v) an exact separation linear program shows that the published cost vector is an optimal separator for that graph and fractional flow, so its cost gap cannot be improved by any reweighting; and (vi) a first minimality result, that no single-terminal instance is a counterexample for any nonnegative cost vector, together with a necessary condition at two terminals. The verification is a replication; items (ii) and (iv)–(vi) are, to our knowledge, recorded here for the first time. A prediction of ours about two terminals was refuted by our own machine and is retracted in Section 7.

1 The statement

Let $G = (V, A)$ be a digraph with a source s and terminals T carrying demands $d \in \mathbb{Q}_{\geq 0}^T$, and let $x \in \mathbb{Q}_{\geq 0}^A$ be a *fractional flow* routing d_t to each $t \in T$. An *unsplittable flow* $\mathcal{P} = \{P^t\}_{t \in T}$

*CAOS open-research programme, Santiago, Chile. With machine-executed computations; all code, declared hypotheses, artifacts and verdicts are versioned at https://github.com/fsantibanezleal/CAOS_RESEARCH under `problems/optimization-geometry/unsplittable-flow-cost/`.

assigns one s - t path per terminal, with arc load

$$f^{\mathcal{P}}(a) = \sum_{t \in T: a \in P^t} d_t, \quad d_{\max} = \max_{t \in T} d_t.$$

Throughout this literature the capacity of an arc is taken to be $x(a)$ itself, so violation is measured against the flow being rounded [5].

Theorem 1 (Dinitz, Garg, Goemans [1]). *Every such instance admits, in polynomial time, an unsplittable flow with $f^{\mathcal{P}}(a) \leq x(a) + d_{\max}$ for every arc a .*

Call such a routing *congestion-good*, and call it *cost-good* for a cost vector $c \in \mathbb{Q}_{\geq 0}^A$ if $c^{\top} f^{\mathcal{P}} \leq c^{\top} x$.

Conjecture (Goemans). *For every instance and every $c \geq 0$ there is an unsplittable flow that is simultaneously congestion-good and cost-good.*

Each condition alone is easy or known: congestion-goodness alone is the theorem above, and cost-goodness alone is achieved by routing every terminal on a cheapest path. The content of the conjecture is the simultaneity. Its status before 2026: proved when all demands are multiples of one another [2]; proved for series-parallel digraphs, in the stronger convex-combination form and with strict deviation below $d_{\max}$ [6]; and known for planar graphs only at the doubled violation $2d_{\max}$ [5]. As of October 2025 the primary literature recorded that whether the conjecture holds even with $O(d_{\max})$ violation “remains wide open” and that proving such a weaker form “would already be considered a breakthrough” [7].

2 The instance

Vertices $\{s, u, v, w, t_1, t_2, t_3\}$, demands $d = (15, 10, 15)$ so $d_{\max} = 15$, and nine arcs:

arc a	x_a	c_a	$x_a + d_{\max}$
$s \rightarrow t_1$	10	2	25
$s \rightarrow t_2$	6	3	21
$s \rightarrow u$	24	0	39
$u \rightarrow t_3$	10	2	25
$u \rightarrow v$	14	0	29
$v \rightarrow t_1$	5	0	20
$v \rightarrow w$	9	0	24
$w \rightarrow t_2$	4	0	19
$w \rightarrow t_3$	5	0	20

The flow is feasible (conservation verified at all seven vertices) with $c^{\top} x = 2 \cdot 10 + 3 \cdot 6 + 2 \cdot 10 = 58$.

3 Verification

Method and independence. We wrote an exact checker from the conjecture statement as printed in [7]. It models arcs by index, so parallel arcs are representable; it enumerates simple

s – t paths by its own depth-first search, never from a supplied list; it decides both conditions with inclusive inequalities in `fractions.Fraction` arithmetic, with floats refused by the instance constructor. It was calibrated first on instances whose answers are fixed by inspection, including a parallel-arc instance, an instance with a directed cycle, and an instance engineered to load an arc to exactly $x_a + d_{\max}$. The proposer’s verifier was archived and hashed but never imported or executed, and the instance was re-entered by hand rather than parsed from their file, so that agreement is evidence rather than a shared-code artifact. The historical failure mode of candidate counterexamples in this area is incomplete path enumeration, so the path count is asserted, and the Dinitz–Garg–Goemans theorem is used as a per-instance oracle: an enumeration reporting no congestion-good routing would indicate an error in our data or code, never in the theorem.

Result. Each terminal has exactly two simple paths, an expensive direct choice E_i and a free detour Z_i :

$$E_1 = st_1, \quad Z_1 = suvt_1; \quad E_2 = st_2, \quad Z_2 = suvw t_2; \quad E_3 = sut_3, \quad Z_3 = suvw t_3,$$

each E_i costing exactly 30 when carrying its demand and each Z_i costing zero. All eight routings:

t_1	t_2	t_3	cost	α	status	violated arcs
E_1	E_2	E_3	90	1/3	congestion-good	–
E_1	E_2	Z_3	60	2/3	congestion-good	–
E_1	Z_2	E_3	60	2/5	congestion-good	–
E_1	Z_2	Z_3	30	16/15	cost-good only	$v \rightarrow w$ by 1
Z_1	E_2	E_3	60	2/3	congestion-good	–
Z_1	E_2	Z_3	30	16/15	cost-good only	$u \rightarrow v$ by 1
Z_1	Z_2	E_3	30	16/15	cost-good only	$s \rightarrow u$ by 1
Z_1	Z_2	Z_3	0	26/15	cost-good only	$s \rightarrow u$ by 1, $u \rightarrow v$ by 11, $v \rightarrow w$ by 1

Here α denotes the least budget, in units of $d_{\max}$, at which the routing becomes congestion-good. The congestion-good and cost-good sets are disjoint.

Theorem 2 (machine-verified). *For the instance above, no unsplittable flow satisfies both $f^P(a) \leq x(a) + d_{\max}$ for all a and $c^\top f^P \leq c^\top x$. Goemans’ conjecture is false.*

An independent second route. The same conclusion follows without costing the eight routings. Call two free choices *in conflict* when every completion of the remaining terminal violates some arc bound. Then Z_1Z_3 conflict ($u \rightarrow v$ carries $30 > 29$), Z_2Z_3 conflict ($v \rightarrow w$ carries $25 > 24$), and Z_1Z_2 conflict ($s \rightarrow u$ carries $40 > 39$, because t_3 enters through $s \rightarrow u$ on either of its paths). The conflict graph is a triangle, its independence number is 1, so every congestion-good routing pays for at least two expensive choices and costs at least $30 + 30 = 60 > 58$. Our implementation computes the conflict graph and this bound in code independent of the enumeration, and the two agree.

The mechanism is a stable-set gap. Writing ρ_i for the fraction of terminal i ’s demand that the fractional flow sends on its free choice,

$$\rho = \left(\frac{5}{15}, \frac{4}{10}, \frac{5}{15}\right) = \left(\frac{1}{3}, \frac{2}{5}, \frac{1}{3}\right), \quad \sum_i \rho_i = \frac{16}{15} > 1.$$

A congestion-good routing selects an independent set of the conflict graph and so buys at most one unit of free routing, while the fractional flow buys 16/15. The instance is the linear-programming integrality gap of the stable-set polytope on a triangle, transported into flow language, with the arc costs acting as a nonnegative separator of the violated triangle inequality.

4 How much is actually forced

Define, for an instance, α_{inst} as the least α such that some cost-good routing satisfies $f^{\mathcal{P}}(a) \leq x(a) + \alpha d_{\max}$ everywhere.

Proposition 1 (machine-verified). *For the instance above, $\alpha_{\text{inst}} = 16/15$.*

In absolute terms, a cost-preserving rounding of this instance needs 16 units of slack on one arc where the conjecture allows 15: the conjecture is refuted by one unit. Consequently the instance says nothing about the question the literature identifies as the real target, namely whether a cost-good routing always exists within $x + O(d_{\max})$; the only known upper bound remains 2, for planar graphs [5], and no finite bound is known in general. We stress this because the distinction is easily lost: refuting the constant 1 and refuting $O(d_{\max})$ are very different statements, and only the former has happened.

5 Consistency with everything proved, and the class boundary

A counterexample must contradict no theorem. We tested each proved result directly on the instance; all pass.

Result	Test	Outcome
Dinitz–Garg–Goemans [1]	a congestion-good routing exists	4 exist
Morell–Skutella cost-free [4]	some routing meets $x \pm d_{\max}$	4 witnesses
Conjecture 1.5 and planar [5]	cost-good within $2d_{\max}$	witnesses exist
Skutella [2]	demands not all multiples	$\{10, 15\}$
Majthoub Almoghrabi et al. [6]	not series-parallel	K_4 subdivision
Two-layer case [5]	not a two-layer network	longest path 4 arcs

Two consequences deserve statement. First, since the digraph is acyclic (verified), any routing satisfying the Morell–Skutella cost conjecture (Conjecture 1.4 of [7]) would satisfy Goemans’ conclusion, so that conjecture is false as well; and if x were a convex combination of congestion-good unsplittable loads y_i , then $\min_i c^T y_i \leq c^T x$ would give a congestion-good routing of cost at most 58, so the convex-combination form [3, 6] is false too. The cost-free conjecture (Conjecture 1.3) is untouched, and with it Theorem 1.6 of [7], which makes the cost-free statement the live route to a $2d_{\max}$ cost result.

Second, the branch vertices $\{s, u, v, w\}$ carry six internally disjoint connecting paths, so the underlying graph contains a K_4 subdivision. Series-parallel digraphs are K_4 -minor-free, and the conjecture is proved for them [6]. The counterexample therefore sits exactly one structure past the class where the statement holds. Moreover the graph has no vertex of degree 4 or more and only four vertices of degree 3, so by Kuratowski’s theorem it is planar with no embedding computation; since [5] proves the cost statement for planar graphs at $2d_{\max}$, the planar constant lies strictly between 1 and 2.

6 The separation LP, and the optimality of the published prices

Let $\mathcal{U}(x)$ be the set of loads of congestion-good routings: finite, nonempty by [1], and independent of c . An instance admits some nonnegative cost vector making it a counterexample precisely when

$$(\text{SEP}) \quad \max \delta \quad \text{s.t.} \quad c^\top(y - x) \geq \delta \quad \forall y \in \mathcal{U}(x), \quad \sum_a c_a = 1, \quad c \geq 0$$

has optimum $\delta^* > 0$. (If $\delta^* > 0$ with witness c , every congestion-good routing costs more than x ; conversely a witnessing $c \neq 0$ has $\min_y c^\top(y - x) > 0$ over a finite nonempty set, and rescaling by $1/\sum_a c_a$ gives a feasible point with positive objective.) We solve (SEP) in exact rational arithmetic.

Proposition 2 (machine-verified). *For the instance above, $\delta^* = 2/7$, attained by $c = \frac{1}{7}(2, 3, 0, 2, 0, 0, 0, 0, 0)$, which is the published cost vector normalised.*

Hence the published prices are an optimal separator for that graph and fractional flow: no reweighting of the arcs produces a larger normalised gap, and the 58-versus-60 separation is the best that instance can do. Practically, (SEP) removes the cost vector from any future search: an exhaustive hunt ranges over graphs, demands and fractional flows only, with one exact linear program per instance.

7 Minimality, and a retraction

Theorem 3. *No single-terminal instance is a counterexample, for any nonnegative cost vector.*

Proof. With $T = \{t\}$ and demand d we have $d_{\max} = d$. Any routing loads its chosen path with d and the rest with 0, and $d \leq x_a + d$ holds on every arc because $x_a \geq 0$, so every routing is congestion-good. The flow x has value d from s to t , hence decomposes into s – t paths of total weight d plus circulations of nonnegative cost; each path costs at least the cheapest path cost γ , so $c^\top x \geq d\gamma$, which is the cost of routing on a cheapest path. That routing is therefore congestion-good and cost-good. $\square$

Remark 1 (retraction). In an earlier internal record we asserted, as a derivation, that no counterexample exists with at most two terminals, on the grounds that a conflict graph on at most two nodes has no odd cycle. That argument is invalid: a terminal may have many path choices, so the conflict graph need not have two nodes, and the step from an integral stable-set polytope to the absence of a separating nonnegative cost vector was asserted rather than proved. We flagged the claim as suspect in the declared hypothesis of the experiment that tested it, and we withdraw it here. The two-terminal case is open.

What does hold at two terminals, and in fact for any number of terminals in its first half:

Proposition 3. *Routing every terminal on a cheapest path is always cost-good. With two terminals, that routing is congestion-good if and only if every arc lying on both cheapest paths carries $x_a \geq \min(d_1, d_2)$. Consequently any two-terminal counterexample contains an arc on both cheapest paths with $x_a < \min(d_1, d_2)$.*

Proof. Cost-goodness follows by applying the decomposition argument of the theorem to each per-terminal flow in a decomposition $x = \sum_i x^i$. For the second part, an arc used by exactly one of the two paths carries at most $d_{\max} \leq x_a + d_{\max}$; a shared arc carries $d_1 + d_2$, and $d_1 + d_2 \leq x_a + \max(d_1, d_2)$ is exactly $x_a \geq \min(d_1, d_2)$. $\square$

We swept 184 two-terminal instances with a shared spine and two choices per terminal, over four demand pairs, every integer split of each demand, and four cost pairs, solving (SEP) exactly on each: none admits any nonnegative cost vector making it a counterexample, and the characterisation above held on every one. This is evidence over a bounded box, not a proof, and we state it as such.

8 What remains open

The refutation settles the constant 1 and leaves the quantitative question essentially untouched. Writing α^* for the least α such that every instance admits a cost-good routing within $x + \alpha d_{\max}$, the record after this work is $\alpha^* \geq 16/15$ and, for planar instances, $\alpha_{\text{planar}}^* \leq 2$; in general no finite upper bound is known. Three concrete questions follow. Can the conflict-triangle mechanism be extended to longer odd cycles or to conflict cliques, whose stable-set gaps are larger, and does flow structure permit the corresponding instances? What is the minimum size of a counterexample in terminals, vertices and arcs? And does the equality $\sum_i \rho_i = \alpha_{\text{inst}} = 16/15$ observed here reflect a general relation between the stable-set violation and the forced violation budget, or is it a coincidence of this instance’s calibration? We record it as undecided.

Provenance and scope

The counterexample instance is not ours. It was announced publicly on 22–23 July 2026, attributed to D. Rybin working with a large language model, and was supplied to this programme as a file bundle which we archived with SHA-256 hashes. What is ours is the independent exact verification, the quantitative constant 16/15, the consistency battery against the proved results, the K_4 and planarity placements, the separation-LP optimality statement, the single-terminal theorem, the two-terminal characterisation, and the retraction recorded above. We make no claim about priority or attribution, which are not mathematical facts, and we take no position on how the community will receive the announcement. Every computation is exact and reproducible from the committed code; hypotheses were declared and committed before each run, and the verdicts, including the refuted prediction, are part of the public record.

References

- [1] Y. Dinitz, N. Garg, M. X. Goemans, *On the single-source unsplittable flow problem*, *Combinatorica* **19**(1) (1999) 17–41. doi:10.1007/s004930050043
- [2] M. Skutella, *Approximating the single source unsplittable min-cost flow problem*, *Mathematical Programming* **91** (2002).
- [3] M. Martens, F. Salazar, M. Skutella, *Convex combinations of single source unsplittable flows*, *ESA 2007*. doi:10.1007/978-3-540-75520-3_36
- [4] A. Morell, M. Skutella, *Single source unsplittable flows with arc-wise lower and upper bounds*, *Mathematical Programming* (2022).
- [5] V. Traub, L. Vargas Koch, R. Zenklusen, *Single-source unsplittable flows in planar and bounded-genus graphs*, *Mathematical Programming* (2026); preprint arXiv:2308.02651. doi:10.1007/s10107-026-02365-x

- [6] M. Majthoub Almoghrabi, M. Skutella, P. Warode, *Integer and unsplittable multiflows in series-parallel digraphs*, IPCO 2025; Mathematical Programming (2026); preprint arXiv:2412.05182. doi:10.1007/s10107-026-02392-8
- [7] C. Swamy, V. Traub, L. Vargas Koch, R. Zenklusen, *Unsplittable cost flows from unweighted error-bounded variants*, preprint arXiv:2510.21287 (2025). doi:10.48550/arXiv.2510.21287